Convergent evolution of seasonal camouflage in response to reduced snow cover across the

snowshoe hare range

Matthew R. Jones[1,2], L. Scott Mills[3,4], Jeffrey D. Jensen[2], Jeffrey M. Good[1,3]

[1]Division of Biological Sciences, University of Montana, Missoula, MT 59812, USA.

[2]School of Life Sciences, Arizona State University, Tempe, AZ 85281, USA

[3]Wildlife Biology Program, University of Montana, Missoula, MT 59812, USA.

[4]Office of Research and Creative Scholarship, University of Montana, Missoula, MT 59812,

USA.

Corresponding authors: jeffrey.good@umontana.edu, matthew.r.jones.1@asu.edu

Abstract

Determining how different populations adapt to similar environments is fundamental to understanding the limits of adaptation under rapidly changing environments. Snowshoe hares (*Lepus americanus*) molt into white winter coats to remain camouflaged in snowy environments. In warmer climates across a small portion of their range hares have evolved brown winter camouflage – an adaptation that may spread under climate change. We used extensive genotype, exome, and whole genome data to 1) resolve range-wide patterns of population structure and gene flow in snowshoe hares and 2) investigate the factors shaping the origins and distribution of winter-brown camouflage variation across the range. In coastal Pacific Northwest (PNW) populations, winter-brown camouflage is known to be determined by a single recessive haplotype at the *Agouti* pigmentation gene. Our phylogeographic analyses revealed deep historical structure and limited gene flow between PNW and more northern Boreal populations, where winter-brown camouflage is rare but widespread along the range edge. We sequenced the genome of a winter-brown Alaskan hare and show that it does not possess the winter-brown PNW *Agouti* haplotype, and thus represents a convergent phenotype that arose through independent mutations. However, the winter-brown PNW haplotype is present at low frequency in a distant winter-white population from Montana, consistent with a model in which strongly deleterious recessive variants spread easily across space because they are masked from selection at low frequency. Simulations show that if annual snow cover were to dramatically reduce, positive selection would likely drive the *Agouti* allele to eventual fixation in Montana, although the initial increase in allele frequency would be extremely slow due to the same masking effect. Our findings underscore how allelic dominance can shape the geographic extent and rate of convergent adaptation in response to rapidly changing environments.

Introduction

In response to a shared selection pressure, populations may adapt through the migration of beneficial alleles or through independent mutations that result in the evolution of convergent phenotypes (Haldane, 1932; Ralph & Coop, 2015; Wright, 1931). Distinguishing between these scenarios is crucial to understand the capacity of populations to adapt rapidly to environmental change (Bridle & Vines, 2007; Hoffmann & Sgrò, 2011; Ralph & Coop, 2015). If gene flow between populations is sufficiently high, then beneficial variation may spread quickly (Fisher, 1937) and potentially allow for rapid adaptive responses to changing environments (Bay et al., 2017). However, the spread of adaptive variation may be limited at broad geographic scales, and populations may have to rely on independent standing genetic variation or new mutations to evolve convergent traits. While there is considerable evidence for both adaptation through gene flow and independent mutations in nature (Dobler, Dalla, Wagschal, & Agrawal, 2012; Hoekstra, Krenz, & Nachman, 2005; Hoekstra & Nachman, 2003; Kreiner et al., 2019; Marques et al., 2017; Rosenblum, Römpler, Schöneberg, & Hoekstra, 2010; Steiner, Rompler, Boettger, Schoneberg, & Hoekstra, 2008), few empirical studies have examined the specific factors that influence these outcomes in natural populations (Ralph & Coop, 2015).

When a species range encompasses a mosaic of habitats, the relative probability of adaptation through gene flow versus independent mutation is predicted to be primarily a function of the distance between habitat patches (in units of dispersal distance) and the strength of selection against locally adaptive alleles in intervening habitats (Ralph & Coop, 2010, 2015; Slatkin, 1973). In general, as distance between patches and the strength of purifying selection in intervening habitats increases, so does the relative probability of adaptation through independent mutations. Adaptation via independent mutations is therefore predicted to be more common in

widespread populations where dispersal distance is short relative to the total range size (Ralph &

Coop, 2010). For example, rock pocket mice (*Chaetodipus intermedius*) have repeatedly evolved

melanistic coats across patchy lava flows in the southwestern United States. Although substantial

gene flow between adjacent lava flows has likely resulted in the migration of melanic alleles

(Hoekstra et al., 2005), melanism is attributed to different mutations across disparate lava flows

(Harris et al., 2019; Hoekstra & Nachman, 2003; Nachman, Hoekstra, & D'Agostino, 2003a).

Theoretical models further predict that the relative probability of adaptation via independent

mutations increases rapidly with distance between lava flows (over the scale of tens to hundreds

of kilometers; Ralph and Coop 2015), due in large part to strong selection against coat color

mismatch (Barrett et al., 2019; Hoekstra, Drumm, & Nachman, 2004; Pfeifer et al., 2018). Thus,

there is a clear tradeoff between dispersal distance and the strength of purifying selection

strength that strongly dictates the probability of adaptation through convergent evolution or gene

flow.

The effect of genetic dominance on the probability of convergent evolution has not yet

been thoroughly explored (Ralph & Coop, 2015). 'Haldane's sieve' (Turner, 1981) predicts that

de novo dominant mutations enjoy a much greater probability of fixation compared to *de novo*

recessive mutations (probability of fixation $\approx 2hs$), because rare dominant mutations are visible to

selection (Haldane, 1924). As *de novo* beneficial recessives are masked to selection when rare,

those that do ultimately reach fixation may spend a longer period of time drifting at low

frequency (Teshima & Przeworski, 2006). Likewise, rare recessive migrant alleles are expected

to exhibit the same behavior, which may allow more time for mutation to generate 'competing'

convergent phenotypes upon which selection can act. However, Orr and Betancourt (2001)

demonstrated that genetic dominance has no effect on the probability of fixation for alleles in

mutation-selection balance because recessive alleles have a higher mutation-selection balance

frequency. Thus, for relatively high amounts of gene flow, genetic dominance may have little to

no effect on the probability of adaptation via migration versus *de novo* mutation. In fact, the

masking of low frequency recessive alleles may result in weaker purifying selection in

intervening habitats therefore facilitating the spread of adaptive variation through gene flow

(Ralph & Coop, 2015), although this hypothesis remains to be tested.

Snowshoe hares (*Lepus americanus*) are one of at least 21 species that have evolved

seasonal molts to white winter coats to maintain camouflage in snowy winter environments.

Because color molts are cued by photoperiod, and hence may become mismatched under rapidly

changing environments (Mills et al., 2013; Zimova et al., 2018), seasonal camouflage provides a

useful trait to understand adaptation to climate change (Jones et al., 2018; Mills et al., 2018,

2013; Zimova et al., 2018; Zimova, Mills, & Nowak, 2016). Reduction in the extent and duration

of snow cover is one of the strongest signatures of climate change in the Northern hemisphere

(Brown & Mote, 2009; Knowles et al., 2006), suggesting that selection should increasingly favor

delayed winter-white molts in snowshoe hares to reduce the total duration of coat color

mismatch. However, seasonal color molts develop relatively slowly in most seasonally changing

species (e.g., 40 days in hares, Mills et al. 2013) and appear to require some minimum duration

of annual snow cover to be beneficial. Below a species-specific threshold, populations are

predicted to maintain brown coloration during the winter (Mills et al., 2018). Consistent with

this, snowshoe hares maintain brown winter camouflage in some temperate environments with

reduced snow cover (Nagorsen, 1983), a strategy that should be increasingly favored under

climate change (Jones et al., 2018; Mills et al., 2018).

Winter-brown snowshoe hares are common in coastal regions of the Pacific Northwest

(PNW), where this Mendelian trait is determined by a recessive variant of the *Agouti*

pigmentation gene (Jones et al. 2018). The locally adaptive *Agouti* variant was introduced into

snowshoe hares by introgression with black-tailed jackrabbits ~9-18 thousand years ago (kya)

and subsequently experienced a local selective sweep within the last few thousand years (Jones

et al. 2019). Occasional records of winter-brown camouflage also occur north of the PNW along

the Pacific coast of Canada and southern Alaska, and in some eastern North American

populations (Gigliotti et al. 2017; Mills et al. 2018). Although hares are expected to be more or

less continuously distributed along suitable coastal environments of western North America,

there is a deep genetic split between northern populations of 'Boreal' hares and populations from

the PNW and southern Rocky Mountains with little evidence of recent gene flow (Cheng,

Hodges, Melo-Ferreira, Alves, & Mills, 2014; Melo-Ferreira, Seixas, Cheng, Mills, & Alves,

2014). Given this historic population structure, it remains unclear whether the distribution of

winter-brown camouflage across populations from disparate parts of the range reflects

independent genetic origins (i.e. trait convergence) or spread of the introgressed PNW *Agouti*

allele.

Here, we use new and previously published genetic data to investigate the roles of gene

flow and mutation in shaping the evolution of winter-brown camouflage across populations of

snowshoe hares. We first combined previously published microsatellite (*n*=853 individuals, 8

loci) with new and published whole exome data (*n*=95 individuals) to resolve range-wide

patterns of population history and gene flow in snowshoe hares, which provides crucial context

for understanding the historical spread and adaptive potential for winter-brown camouflage to

climate change. We then generated whole genome sequence (WGS) data of a winter-brown

Alaska (AK) hare to test whether winter-brown camouflage in Boreal snowshoe hares arose

independently from PNW populations, located ~3000 km away. Next, to understand the

geographic limits of the recessive PNW *Agouti* allele, we used pooled WGS data to estimate its

frequency in a winter-white population from Montana (MT), approximately 600 km in a straight-

line from the closest winter-brown PNW population. Finally, we used both theoretical

predictions and simulations to understand the factors influencing the geographic scope of the

PNW *Agouti* allele and its potential to contribute to rapid adaptation in response to warming

climates.

Methods

<u>Samples and genomic data generation</u>

To resolve patterns of range-wide population structure, we performed targeted whole exome

enrichment of 12 snowshoe hares previously sampled from 12 localities across Canada, Alaska,

and the eastern United States and three hares from the southern Rocky Mountains (representing

the Boreal and Rockies lineages as defined by Cheng et al. (2014)). The targeted whole exome

capture was designed by Jones et al. (2018) to enrich for ~99% of genic regions annotated in the

European rabbit (*Oryctolagus cuniculus*) genome (61.7-Mb spanning 213,164 intervals; ~25-Mb

protein-coding exons, ~28-Mb untranslated region, ~9-Mb intronic or intergenic). We followed

the library preparation protocols outlined in Jones et al. (2018) and sequenced libraries on one

lane of Illumina HiSeq2500 with paired-end 100 bp reads (HudsonAlpha Institute for

Biotechnology; Huntsville, AL). Exome sequences for Boreal and Rockies snowshoe hares were

combined with published whole exome data (Jones et al. 2018) from 80 PNW snowshoe hares

(*n*=95 total, Table S1), including a monomorphic winter-brown population from southeast

British Columbia (BC1, *n*=14), a monomorphic winter-white population from Seeley Lake area

in western MT (*n*=14), and two polymorphic coat color populations from Oregon (OR, *n*=26;

two localities) and Washington (WA, *n*=26; Jones et al. 2018).

To resolve the historical spread of the winter-brown PNW *Agouti* allele, we performed

whole genome sequencing of a winter-brown snowshoe hare museum specimen (AK) from

southwest Alaska first noted by Link Olson and later obtained through loan from University of

Alaska Museum of the North (UAM 116170, collected on 4 January 2013). We also performed

whole genome sequencing on a DNA pool of 81 snowshoe hares from two localities in Glacier

National Park, MT (Table S1). We extracted genomic DNA from muscle tissue of the AK hare

sample following the Qiagen DNeasy Blood and Tissue kit protocol (Qiagen, Valencia, CA). For

the pooled MT whole genome sequencing, we pooled previously extracted DNA samples (Cheng

et al., 2014) in approximately equimolar quantities based on Qubit concentrations (Invitrogen

Qubit Quantitation system LTI). We then prepared genomic libraries for all samples following

the KAPA Hyper prep kit manufacturer's protocol. We sheared genomic DNA to ~300 bp using

a Covaris E220evolution ultrasonicator and performed a stringent size selection using a custom-

prepared carboxyl-coated magnetic bead mix (Rohland & Reich, 2012). We determined indexing

PCR cycle number for each library with quantitative PCR (qPCR) on a Stratagene Mx3000P

thermocycler (Applied Biosystems) using a DyNAmo Flash SYBR Green qPCR kit (Thermo

Fisher Scientific). Final libraries were size-selected again with carboxyl-coated magnetic beads,

quantified with a Qubit (Thermo Fisher Scientific), and pooled for sequencing by Novogene

(Novogene Corporation Ltd.; Davis, CA) on two lanes of Illumina HiSeq4000 using paired-end

150bp reads. The whole genome sequence from the AK hare was combined with previously

published whole genome sequencing of five black-tailed jackrabbits, three winter-brown

snowshoe hares from OR, WA, and BC1, and three winter white snowshoe hares from MT, a

hare from Utah (UT), and a hare from Pennsylvania (PA; Jones et al. 2018, Table S1).

<u>Read processing and variant calling</u>

For all raw Illumina sequence data, we trimmed adapters and low-quality bases (mean phred-

scaled quality score <15 across 4 bp window) using Trimmomatic v0.35 (Bolger, Lohse, &

Usadel, 2014) and merged paired-end reads overlapping more than 10 bp and with lower than

10% mismatched bases using FLASH2 (Magoč & Salzberg, 2011).

179   Whole genome sequence data were mapped to either a snowshoe hare or black-tailed

jackrabbit pseudoreference (see Jones et al. 2018 for details) using default settings in BWA-

MEM v0.7.12 (Li, 2013). We used *PicardTools* to remove duplicate reads with the

MarkDuplicates function and assigned read group information with the

AddOrReplaceReadGroups function. Using GATK v3.4.046 (McKenna et al., 2010), we then

identified poorly aligned genomic regions with RealignerTargetCreator and locally realigned

sequence data in these regions with IndelRealigner. We performed population-level multi-sample

variant calling using default settings with the GATK UnifiedGenotyper. Here, we called variants

separately for each previously defined snowshoe hare population genetic cluster (i.e., Boreal,

Rockies, BC1, MT, OR, and WA) and for black-tailed jackrabbits. We performed variant

filtration in VCFtools v0.1.14 (Danecek et al., 2011). For whole exome and whole genome data,

we filtered genotypes with individual coverage <5× or >70× or with a phred-scaled quality score

<30. Additionally, we removed all indel variants and filtered SNPs with a phred-scaled quality

score <30, Hardy-Weinberg $P<0.001$. We required that sites have no missing data across

individuals.

Range-wide population genetic structure and gene flow

We inferred a maximum likelihood tree with a general time reversible model in RAxML v8.2.8

(Stamatakis, 2014) using a subset of the concatenated snowshoe hare exome data (n=12 Boreal

hares, n=3 Rockies hares, n=12 PNW hares; 21,167,932 total sites) with European rabbit

(*Oryctolagus cuniculus*) as the outgroup. Using this maximum likelihood tree as a starting tree,

we estimated a maximum clade credibility tree and node ages with a constant population size

coalescent model in BEAST 2 (Bouckaert et al., 2014). We assumed a strict molecular clock and

an HKY substitution model using empirical base frequencies. We specified default priors for the

kappa parameter, gamma shape parameter, and population size parameter and used a gamma

distribution (alpha=0.0344, beta=1) as a prior for the clock rate parameter. We ran the MCMC

for 5 million steps and calibrated divergence times using a log-normal distribution for the

Oryctolagus-Lepus node age with a median of 11.8 million years (95% prior density: 9.8–14.3;

Matthee et al. 2004). We also performed a species-tree analysis using gene trees in ASTRAL

v5.6.3 (Zhang, Rabiee, Sayyari, & Mirarab, 2018). Gene trees were generated across 200 kb

windows (excluding windows wither fewer than 500 sequenced bases) using RAxML with a

GTR+gamma model and rapid bootstrap analysis (-f a -# 10). We collapse nodes on gene trees

with low bootstrap support values (<=10) and performed ASTRAL analyses on the collapsed

gene trees using default settings.

To test for signatures of population structure and gene flow among lineages, we

performed a population cluster analysis in ADMIXTURE v1.3.0 (Alexander, Novembre, & Lange,

2009). We tested K values from 1-10 (representing the number of population clusters) and

selected the K value with the lowest cross-validation error. We also estimated range-wide

effective migration and diversity surfaces with EEMS (Petkova, Novembre, & Stephens, 2016)

using extensive microsatellite data (*n*=853 individuals, 8 loci) generated by Cheng et al. (2014).

Varying the number of demes (50, 100, and 200) had little effect on estimates of effective rates

of migration and diversity, therefore we only report results for 200 demes. We used default

hyper-parameter values and tuned the proposal variances such that proposals were accepted

approximately 30% of the time. We ran EEMS for 2 million iterations with a burn-in of 1 million

iterations and thinning iteration of 9999. Runs produced strong correlations between observed

and expected genetic dissimilarity both within and between demes, indicating good model fit.

Geographic distribution of the PNW *Agouti* allele

The winter-brown AK snowshoe hare was collected approximately 3000 km (via the Pacific

coast) from Vancouver, BC, the northern limit of winter-brown PNW hare populations (Jones et

al., 2018). To test whether the introgressed PNW *Agouti* allele has seeded winter-brown

camouflage in southwest AK, we generated a tree of the PNW *Agouti* haplotype region from our

whole genome sequence data of winter-brown and winter-white snowshoe hares (*n*=7) and black-

tailed jackrabbits (*n*=5). We defined the PNW *Agouti* haplotype region (chr4:5,340,275 –

5,633,020, oryCun2 coordinates) based on the location of the introgressed black-tailed jackrabbit

tract identified by Jones et al. (2018) using a hidden Markov model. We then inferred a

maximum likelihood phylogeny for this interval using RAxML (Stamatakis, 2014) as above.

Node support values were generated from 1000 replicate bootstrap runs.

We also estimated the frequency of the PNW *Agouti* haplotype among pooled individuals

from two localities in MT (*n*=81) that are 575 km and 627 km from the polymorphic sampling

locality in WA, where winter-brown camouflage is relatively common. We used PoPoolation2

(Kofler et al., 2011) to calculate the frequency of the winter brown-associated alleles at 555 sites

across the introgressed *Agouti* haplotype (chr4: 5,340,275 – 5,633,020). These positions were

previously shown to be strongly associated with coat color based on a likelihood ratio test of

allele frequency differences between winter-brown and winter-white individuals from low

coverage whole genome sequence data (Jones et al. 2018). We summed the counts of winter-

white and winter-brown alleles across all positions to estimate both the mean and standard

deviation of winter-brown allele frequency. We excluded seven positions with unusually high

frequencies of the winter-brown allele (~45-100%) supported by reads that did not otherwise

carry brown-associated alleles (i.e., incongruent haplotype information), as these likely reflect

sequencing errors.

<u>Probability of convergent adaptation through independent mutation</u>

To compare the observed spread of the PNW *Agouti* allele to theoretical expectations, we

estimated the relative probability of adaptation via independent mutations with distance from a

focal habitat patch using the model developed by Ralph and Coop (2015, equation 12) under a

range of conditions. Since the relative probability of adaptation through migration or *de novo*

mutation depends strongly on the rate at which mutations generate convergent phenotypes

(Ralph & Coop, 2015), we tested a range of plausible mutation rates. A previous study using

over 7 million house mice estimated a mean rate of spontaneous visible coat color mutations of

$11e^{-6}$ per locus/gamete (Schlager & Dickie, 1971), which is ~1930× higher than the average

genome-wide germline per site mutation rate ($5.7e^{-9}$ per site/gamete; Milholland et al. 2017).

Given a genome-wide mutation rate of $2.35e^{-9}$ per site/generation for rabbits (Carneiro et al.,

2012), we tested a high mutation rate of $\mu=4.54e^{-6}$ (i.e., the overall expected rate of visible coat

color mutations) and a low mutation rate of μ=4.54e^{-8} (i.e., assuming only 1% of coat color

mutations would lead to brown winter camouflage). We also tested how strong (s_m=0.01),

moderate (s_m=0.001), or weak (s_m=0.0001) purifying selection against the winter-brown allele in

intervening (i.e., snow-covered) habitats influences the probability of adaptation through

independent mutations. We assumed a relatively high mean dispersal distance of 2 km (Gillis &

Krebs, 1999), which is likely to produce a conservative estimate of the probability of adaptation

via independent mutation. We also assumed that the width of the secondary habitat patch was

relatively small (w=1 or ~6-45 km, depending on s_m), which is consistent with observations of

winter-brown camouflage at low frequency along portions of the range edge. Finally, we

assumed d=1 (dimension of the habitat), C=1, and γ=0.5.

<u>Simulations of selection on migrant alleles with genetic dominance</u>

Theory predicts that allelic dominance should not influence a mutation's probability of fixation

under positive selection if it starts in mutation/migration-selection balance (Orr & Betancourt,

2001). Nonetheless, rare recessive mutations may increase in frequency more slowly relative to

rare dominant mutations because they are initially invisible to selection, allowing more

opportunity for competition from convergent phenotypes arising through independent mutations.

We used simulations to test how the genetic dominance of alleles at equivalent mutation-

selection equilibrium frequencies influences the probability and rate of adaptation to new

habitats when environmental conditions change. In SLiM 3.0 (Haller & Messer, 2019) we

performed 100 simulations of the MT population (estimated N_e=245430; Jones et al. 2019) with

a positively selected recessive mutation (s=0.01, h=0) starting at the inferred frequency of the

PNW *Agouti* allele in MT (p). We assumed that p reflects the equilibrium frequency (x) between

the migration rate of the allele into the environment (m) and the selection coefficient against the

allele (s), which is given as $\sqrt{\frac{m}{s}}$ for a recessive mutation. Given that x for a dominant mutation is

simply $\frac{m}{s}$, we calculated x of an equivalent dominant mutation (same values of m and s) as simple

p^2. We then repeated these simulations for a positively selected ($s=0.01$), completely dominant

allele ($h=1$) starting at its expected mutation-selection balance frequency. For each simulation we

tracked the frequency of the selected allele until it was either fixed or lost. To validate these

simulations, we compared the probabilities of allelic fixation from simulations to fixation

probabilities analytically derived by Kimura (1957). To test for significant differences in fixation

probabilities and rates of adaptation between recessive and dominant mutations we used

Student's t-tests in R (R Core Team, 2018).

Results

<u>Snowshoe hare phylogeography</u>

We combined previously generated whole exome data for 80 PNW snowshoe hares (mean

coverage $21\times \pm 7.6$ per individual) with new whole exomes of 15 hares across the range

sequenced to a mean coverage of $20.2\times \pm 8.2$ (Fig. 1A). Exome-wide phylogenetic analyses

show three broad phylogeographic groups of snowshoe hares (Fig. 1B) with unambiguous

ASTRAL support scores using 6582 gene trees, consistent with previous studies (Cheng et al.

2014; Melo-Ferreira et al. 2014). However, ~12-25% of SNPs within phylogeographic groups

appear to be shared across some population boundaries (Table 1), indicating some amount of

shared ancestral variation or gene flow. We estimated that snowshoe hares from Canada, Alaska,

and the eastern United States—representing the Boreal lineage—diverged approximately 97.2

kya (95% posterior density: 77.4-120.8 kya) from PNW and Rockies snowshoe hares (Fig. 1B).

PNW and Rockies hare populations are estimated to have split approximately 47.1 kya (95%

posterior density: 37.2-58.2 kya).

Population structure analyses in ADMIXTURE also found strongest support for three

population clusters, although MT snowshoe hares (in the PNW lineage) showed 11-16% of their

ancestry assigned to the Rockies lineage, indicating ongoing gene flow or continuous genetic

structure (Fig. 1C). The ADMIXTURE analysis also indicated little apparent gene flow between the

Boreal lineage and PNW or Rockies lineages. Likewise, microsatellite-based estimates of

effective migration surfaces revealed that effective migration rates are approximately 100-fold

lower than the range-wide average near the apparent zone of contact between PNW and Boreal

lineages in western North America (Fig. 2A). We also estimated relatively low effective

migration rates across the southwestern edge of snowshoe hare range (western US; $\log(m) \approx -1$,

or 10-fold lower than the mean). In contrast, effective migration rates in the northwest part of the

range (Alaska and western Canada) appeared relatively high (~10-fold higher than the mean).

Effective diversity surfaces show that the highest relative genetic diversity occurs in the eastern

extent (Boreal lineage) and southwestern extent (PNW lineage) of the snowshoe hare range (Fig.

2B), consistent with previous studies suggesting that these regions were likely glacial refugia

(Cheng et al., 2014). Relative genetic diversity gradually decreased moving from east to west

across the Boreal range and the lowest genetic diversity was observed in the Rocky Mountain

lineage, distributed across the western US.

<u>Convergent evolution and the distribution of winter-brown camouflage variation</u>

Under a single origin of winter brown camouflage, we would expect the winter-brown AK

individual to nest within the black-tailed jackrabbit clade with the other winter-brown snowshoe

hares from the PNW. However, our phylogenetic analysis indicates the winter-brown AK

individual unambiguously groups with a winter-white Boreal individual from PA (100%

bootstrap support), and more broadly with other winter-white hares across the range (Fig. 3).

These results demonstrate that the winter-brown phenotype of this AK hare is caused by

mutations that are independent from the introgressed *Agouti* haplotype that encodes winter-

brown camouflage in the PNW, approximately 3000 km away. Determining the genetic basis of

this independent origin of winter brown camouflage will likely require an independent genetic

association study in these populations.

Pooled whole genome sequencing detected that the recessive winter-brown *Agouti* allele

occurs at an estimated frequency of 1.24% ($\pm$ 0.01%) across predominately winter-white MT

localities ~600 km from the polymorphic zone. Interestingly, long-term live-trapping and

radiotelemetry-based field work from this region has resulted in a single observation of a winter-

brown hare in approximately 300 winter hare observations (0.33%; Scott Mills pers. obs.),

although this observed frequency is slightly higher than the Hardy-Weinberg expectation under

the estimated frequency of the winter-brown *Agouti* allele (p^2=0.015%, binomial test *p*-

value=0.044).

To understand the factors that have shaped the geographic distribution of the PNW

Agouti allele, we modelled the relative probability of adaptation through independent mutation

with distance from a focal habitat using the theoretical framework of Ralph and Coop (2015).

The change in relative probability of adaptation through independent mutation depended strongly

on both the mutation rate and selection coefficient against the allele in intervening habitats (Fig.

4). Stronger selection coefficients produced a more rapid shift in the convergence probability

with distance relative to weaker selection (Fig. 4). For instance, under a high mutation rate to the

winter-brown phenotype (μ=4.54e^{-6}) and strong purifying selection (s_m=0.01) the probability of

independent adaptation transitioned sharply from P=0.1 at 78 km to P=0.9 at 140 km from a

focal habitat. Under a low mutation rate to the winter-brown phenotype (μ=4.54e^{-8}) and weak

selection (s_m=0.0001), independent adaptation was much less likely at close distances, as

expected, and the probability increased more gradually (P=0.1 at 778 km to P=0.9 at 1400 km).

<u>Migration-selection balance</u>

Assuming an environmental shift towards less snow results in strong positive selection (s=0.01)

on standing variation of the recessive (winter-brown) *Agouti* allele in MT, we used simulations

to estimate that the corresponding probability of fixation would be ~81% (95% confidence

intervals: 72.2-87.5%; Fig. 5A), which is consistent with an analytically-derived probability of

78.1%. For an equivalent dominant mutation starting in mutation-selection balance frequency

(0.015%) the simulated probability of fixation was approximately 77% using simulations (95%

confidence intervals: 67.8-84.2%) or 77.9% using the Kimura (1957) equation. Neither estimate

was significantly different from the fixation probability of a recessive mutation (p=0.48, two-

tailed test of equality of proportions). However, we observed striking differences in the rates of

increase, conditional on fixation, of dominant and recessive mutations following a sudden

environmental change (Fig. 5B). The mean time to fixation was significantly faster (p<2.2e-16,

Student's T-test) for a recessive mutation (9645 generations, SD=3609 generations) compared to

a dominant mutation (19445 generations, SD=7273 generations, p<2.2e^{-16}). However, as

expected, we see that initial rates of allele frequency change and phenotypic adaptation are

considerably faster for a positively selected dominant mutation (Fig. 5B). For instance, the mean

time for the selected phenotype determined by the recessive mutation to reach a frequency of

75% (*p*=0.866) was 8837 generations (SD=3630 generations), compared to 1007 generations

(SD=116 generations) if determined by the dominant mutation. The striking difference in the rate

at which beneficial dominant versus recessive alleles contribute to adaptation is maintained even

up to the point at which the selected phenotype reaches a frequency of 99%, which takes

approximately 9262 generations (SD=3624 generations) for a recessive mutation and only 2030

generations for dominant mutation (SD=141 generations).

Discussion

A growing number of studies have found evidence for convergent adaptation within and between

species (Dobler et al., 2012; Giska et al., 2019; Harris et al., 2019; Hoekstra & Nachman, 2003;

Marques et al., 2017; Nelson, Jones, Velotta, Dhawanjewar, & Schweizer, 2019; Rosenblum et

al., 2010; Steiner et al., 2008), although we often lack an understanding of the forces that

determine whether local adaptation occurs through independent *de novo* mutations or migration

of pre-existing alleles from other populations (Ralph & Coop, 2015). Our study provides rare

empirical insights into how gene flow, mutation, allelic dominance, and selection interact to

shape the spatial scale and pace of local adaptation to new or changing environments.

Understanding the potential for adaptive variation to spread between populations across

large spatial scales requires basic insights into range-wide patterns of population structure and

gene flow. Winter-brown camouflage in snowshoe hares occurs across only ~5% of the total

range but is broadly-distributed from the western edge of the range along the Pacific coast to the

eastern extent of the range in Pennsylvania (Gigliotti et al., 2017; Mills et al., 2018; Nagorsen,

1983). Previous phylogeographic studies based on limited genetic data suggest that this

distribution spans a deep phylogenetic boundary between Boreal and PNW lineages (~2 Myr

divergence; Cheng et al. 2014; Melo-Ferreira et al. 2014) with little evidence for gene flow. With

whole exomes (n=95), our phylogenetic analysis also supports a deep split but suggests a

substantially more recent divergence time between Boreal and Rockies/PNW lineages than

previously estimated (~97 kya; Fig. 1B). Our more recent estimates may be attributable to the

increased power of whole exomes or heavy reliance on mitochondrial DNA in previous studies

(Melo-Ferreira et al., 2014), which can lead to overestimation of divergence times (Zheng, Peng,

Kuro-o, & Zeng, 2011). Consistent with previous studies, our ADMIXTURE analyses and range-

wide effective migration surfaces suggest low gene flow between the PNW and Boreal lineages

(Fig. 1C, Fig. 2A). The absence of observable gene flow is perhaps striking given their close

proximity to each other in western North America. Melo-Ferreira et al. (2014) hypothesized that

these lineages may exhibit incipient reproductive isolation. The evolution of reproductive

isolation between these populations is possible, but given our more recent divergence time

estimates we suggest other factors are likely to contribute to the estimated lack of gene flow. For

example, the low effective migration rates (Nm; Fig. 2A) could be partially attributable to

reductions in either population densities (N) or dispersal capability near the geographic boundary

of Boreal and PNW lineages (m; Petkova et al. 2016), although this region appears to be in core

snowshoe hare habitat. More likely, reduced gene flow may reflect quite recent secondary

contact between these populations in western North America. Genetic and fossil evidence

suggests the common ancestors of PNW and Boreal populations occupied separate refugia in

southern and eastern North America, respectively, during the last glacial maximum (Cheng et al.,

2014). The east-to-west phylogenetic clustering (Fig. 1B) and genetic diversity gradient (Fig.

2B) of the Boreal population is consistent with a recent range expansion from this eastern refugia

(Cheng et al. 2014), which implies the PNW and Boreal hares have only recently experienced

secondary contact. However, more detailed population history modeling and sampling of the

putative contact zone is required to test these hypotheses.

Consistent with low historical gene flow, the AK winter-brown hare lacks the PNW

winter-brown *Agouti* haplotype (Fig. 3), indicating a role for independent mutation leading to the

convergent evolution of brown winter coats. This likely reflects constraints on the migration of

adaptive variation between populations. As previously mentioned, it is unclear if population

structure between Boreal and PNW hares reflects nascent barriers to gene flow (Melo-Ferreira et

al., 2014). Even in the absence of intrinsic or extrinsic reproductive isolation, the rate at which a

positively selected variant spreads across a uniform environment is constrained by dispersal

distance and the strength of positive selection (Fisher 1937). Across spatially or temporally

heterogenous landscapes, the spread of an adaptive allele between patches is further constrained

by the strength or frequency of purifying selection in maladaptive habitats or climatic periods

(Ralph & Coop, 2015). Given that previous models suggest a low probability of winter-brown

camouflage along the majority of the Pacific coast (Mills et al., 2018), we suspect that the

winter-brown PNW variant would have to traverse snowy habitats where it is strongly disfavored

in order to reach Alaska, ~3000 away via the coast. At this distance, the probability of adaptation

through gene flow is virtually zero under a range of model assumptions (Fig. 4). In fact, dispersal

limitations and patchy range-edge habitats should generally favor independent evolution of

winter-brown camouflage at scales beyond 100-1000 km (depending on the assumed values of μ

and s_m; Fig. 4). Further research is needed to dissect the genetic basis of winter-brown

camouflage in the northwestern and eastern edges of the range (e.g., Alaska and Pennsylvania).

Subtle phenotypic similarities between observed winter-brown hares in Alaska and Pennsylvania

could imply a shared genetic basis (e.g., white feet versus brown feet in the PNW, unpublished

data). However, our theoretical modelling would suggest that they likely reflect independent

mutations as well (i.e., ~5900 km between AK and PA sampling sites).

449   Convergent evolution is thought to be more common for 'loss-of-function' traits because

they may have a larger mutational target size relative to 'gain-of-function' traits (Manceau,

Domingues, Mallarino, & Hoekstra, 2011; Rosenblum et al., 2010). Convergent color adaptation

involving loss-of-function mutations has been shown between different species of lizard (Laurent

et al., 2016; Rosenblum et al., 2010) and cavefish (Protas et al., 2006) and the repeated evolution

of melanism across populations of deer mice involves convergent loss-of-function *Agouti*

mutations (Kingsley, Manceau, Wiley, & Hoekstra, 2009). In PNW hares, adaptive introgression

of *Agouti* variation from black-tailed jackrabbits appears to have caused a reversion to the

ancestral winter-brown condition in *Lepus* (i.e., the likely ancestral state before the common

ancestor of winter-white *Lepus* species). As the ancestral winter-brown variant is recessive, this

implies that derived winter-white camouflage in snowshoe hares evolved through dominant gain-

of-function mutations, consistent with the seasonal upregulation of *Agouti* during the

development of white coats (Jones et al., 2018). Independent origins of winter-brown

camouflage across the snowshoe hare range could similarly involve relatively simple loss-of-

function mutations that break the molecular pathways contributing to the development of white

winter coats. Indeed, the evolution of darkened winter coats in some populations of mountain

hares (*Lepus timidus*) appears to have also involved introgression of a recessive, loss-of-function

Agouti variant (Giska et al., 2019). Collectively, our results suggest that geographic distance and

mutational target size, in addition to population structure and history, should play a crucial role

in generating hypotheses about the relative roles of independent mutation and gene flow in

adaptation.

Despite the evidence for independent winter-brown mutations at broad spatial scales, we

also find that the PNW *Agouti* allele has traversed a substantial distance (~600 km) to western

Montana – a predominately winter-white locality – where it occurs at a frequency of 1.24%. At

this starting frequency, both theory and simulations show that a shift in environmental conditions

towards favoring the brown allele results in a ~80% probability of fixation (Fig. 5). The high

probability of adaptation through gene flow ~600 km from a focal patch is consistent with a

scenario of very weak purifying selection (s_m=0.0001) against the winter-brown allele in snowy

environments under the Ralph and Coop (2015) model (P(independent mutation)=~3%

(μ=4.54e-8) or ~76% (μ=4.54e-6); Fig. 4). Although winter-brown camouflage is likely more

strongly deleterious in winter-white environments given the known fitness consequences of

mismatch (Zimova et al., 2016), purifying selection against the winter-brown *Agouti* allele in

winter-white environments may be weak or absent because it is recessive and thus hidden to

selection at low frequency. In agreement with Ralph and Coop (2015), we suggest that

adaptation between distant populations via gene flow may be relatively more common through

recessive variation compared to dominant variation when purifying selection is strong in

intervening habitats.

Climate change is expected to reduce winter snow cover across the snowshoe hare range

(Mills et al., 2013), which could potentially result in winter-brown camouflage being favored in

predominately winter-white populations (Mills et al., 2018). Under this scenario, positive

selection could operate on convergent *de novo* winter-brown mutations or on standing genetic

variation for winter-brown camouflage to facilitate rapid adaptation. Using simulations of

positive selection, we show that although dominant and recessive mutations in

mutation/migration-selection balance share a similar fixation probability (Fig. 5A), they are

likely to experience vastly different frequency change dynamics (Teshima and Przeworski 2006;

Fig. 5B) and thus lead to very different rates of adaptation following environmental change. For

instance, we show that in MT the initial rate of adaptation from segregating winter-brown *Agouti*

variation is likely to be slow relative to an equivalently beneficial dominant mutation. While

recessive mutations tended to fix more quickly than dominant mutations, they also took

substantially longer to reach high frequencies in populations (Fig. 5B; e.g., 8837 vs. 1007

generations for a phenotype determined by recessive vs. dominant mutation to reach a population

frequency of 75%, respectively). This pattern can be explained by the different allele frequency

phases most strongly affected by genetic drift. For instance, positive selection can readily act on

beneficial dominant mutations at low frequency, but at high frequency the recessive wildtype

allele is hidden in heterozygotes and genetic drift governs (and generally slows) fixation

dynamics. The opposite is true for beneficial recessive mutations, which are governed by genetic

drift at low frequency but driven to fixation by selection. Overall, selection on low frequency

recessive variation may be a relatively ineffective way to adapt rapidly to changing

environments. In fact, the temporal lag for the spread of beneficial recessive variation may be

sufficient enough to allow time for dominant independent mutations (e.g., gain-of-function

MC1R mutations that result in melanism; Nachman et al. 2003*b*) to appear and spread before the

recessive allele increases appreciably. These findings highlight the important roles of genetic

dominance in shaping both the geographic scope and rate of convergent adaptation and

underscores the need for further study.

Collectively, our study provides important insights into long-standing evolutionary

questions, as well as the potential for adaptation to climate change in snowshoe hares. A key to

understanding the potential of adaptive variation to spread under climate change is revealing how

it has spread in the past. We show that adaptive responses to reduced snow cover in snowshoe

hares have involved both the local spread of winter-brown camouflage in the PNW as well as

convergent evolution elsewhere in the range. In snowshoe hares and other seasonally changing

species, regions of winter-camouflage polymorphism may be crucial areas to maintain

population connectivity as a conduits for the spread of winter-brown camouflage across broader

portions of the range (Mills et al., 2018). However, facilitating gene flow alone may not be

sufficient to facilitate rapid adaptive responses, given the broad range of snowshoe hares and the

genetic architecture of winter-brown camouflage variation. Rather, we suspect that rapid climate

change responses in winter-white populations distant from winter-brown habitat may also likely

involve independent origins of winter-brown camouflage and selection on variation in coat color

phenology (i.e., shifts in the timing of winter-white molts). Although the genetic basis of

variation in camouflage phenology remains unresolved in this system, it is likely quantitative and

perhaps more responsive to selection. Regardless, this study represents an important step towards

making predictions about evolutionary responses under climate change in this species.

Data availability

Original sequence data are available in the Sequence Read Archive (www.ncbi.nlm.nih.gov/sra).

Previously generated whole exome and genome sequence data of snowshoe hare (BioProject

PRJNA420081, SAMN02782769, SAMN07526959) are also available in the Sequence Read

Archive.

Acknowledgements

We thank E. Cheng and K. Garrison for assistance with sample collection. We thank L. Olson for

informing us on the recent collection of a winter brown snowshoe hare, and L. Olson and the

University of Alaska Museum of the North facilitating a tissue loan of this specimen (UAM

116170). Our research would not be possible without the incredible and irreplaceable support of

natural history museums. We thank J. Melo-Ferreira, P. C. Alves, M. S. Ferreira, N. Herrera, E.

Kopania, A. Kumar, M. Zimova, K. Garrison, and the UNVEIL network for helpful discussions.

Funding and support for this research was provided a National Science Foundation (NSF)

Graduate Research Fellowship (DGE-1313190), NSF Doctoral Dissertation Improvement Grant

(DGE-1702043), NSF Graduate Research Opportunities Worldwide, NSF EPSCoR (OIA-

1736249), and NSF (DEB-0841884; DEB-1907022), the Drollinger-Dial Foundation, American

Society of Mammalogists Grant-in-aid of Research, and a Swiss Government Excellence

Scholarship.

Figure 1. (**A**) Range-wide phylogeography of snowshoe hares based on whole exome sequences. The snowshoe hare range is colored brown to gray according to the modelled probability of winter-brown camouflage from Mills et al. (2018). Sizes of sampling locality circles are scaled to sample size and are colored according to their population assignment (see C). (**B**) A maximum credibility phylogenetic tree estimated with BEAST 2 (Bouckaert et al., 2014). All nodes have posterior probabilities >99%. Each major node shows the median node age in thousands of years (95% posterior density in parentheses and gray rectangles) and the ASTRAL support score. (**C**) The ADMIXTURE plot shows the proportion of ancestry across samples based on a *K*=3 clustering, which had the lowest cross validation error.

Figure 2. (**A**) Effective migration rates for snowshoe hares inferred from range-wide

microsatellite data set (853 individuals, 8 loci) from Cheng et al. (2014). The sizes of circles are

scaled to the number of samples collected at that location. Effective migration rate is measured

as the rate of decay in genetic similarity of individuals across space. Regions that are colored

white are characterized by isolation-by-distance while regions that are colored blue or red have

higher or lower effective migration, respectively. (B) Effective diversity rates based on the same

microsatellite data set. Here effective diversity rates measure the genetic dissimilarity between

individuals in the same deme, where blue regions have higher than average diversity and red

regions have lower than average diversity.

Figure 3. A maximum clade credibility tree of the introgressed *Agouti* locus (~293 kb) based on

whole genome sequencing of black-tailed jackrabbits (BTJR; black circles) and snowshoe hares.

Values indicate node support based on bootstrapping. Colored stars indicate winter-brown

snowshoe hares from Alaska (yellow, UAM 116170 pictured top) or the Pacific Northwest

(purple, pictured bottom).

Figure 4. The probability of adaptation through independent mutations in snowshoe hares as a

function of distance in kilometers from a focal habitat patch harboring a locally adaptive variant.

The probability of independent mutation is calculated according to equation 12 in Ralph and

Coop (2015). Here we varied the mutation rate to the winter-brown phenotype (μ; black=4.54e^{-8},

gray=4.54e^{-6}) and the negative selection coefficient in intervening habitats (s_m; solid line=0.0001,

dashed line=0.001, dotted line=0.01).

Figure 5. (**A**) The simulated probability of fixation of a completely dominant (red, mean=77%,

N=100) or recessive (blue, mean=81%, *N*=100) mutation experiencing positive selection and

starting in migration-selection balance frequency (0.015% for dominant, 1.24% for recessive).

(**B**) The simulated allele frequency trajectories of the same dominant (blue) and recessive (red)

mutations starting in migration-selection balance.

Table 1. Counts of the number of private (e.g., Boreal/Boreal) versus shared SNPs (e.g.,

Rockies/Boreal) for each major snowshoe hare clade. Individuals from WA represent the PNW.

	Boreal (N=120978 SNPs)	Rockies (N=158207 SNPs)	PNW (N=275614 SNPs)
Boreal	106051	3760	5820
Rockies & PNW	5347	-	-
Rockies	-	117863	31237
PNW	-	-	233210

References

Alexander, D. H., Novembre, J., & Lange, K. (2009). Fast model-based estimation of ancestry in unrelated individuals. *Genome Research*, *19*(9), 1655–1664. doi: 10.1101/gr.094052.109

Barrett, R. D. H., Laurent, S., Mallarino, R., Pfeifer, S. P., Xu, C. C. Y., Foll, M., … Hoekstra, H. E. (2019). Linking a mutation to survival in wild mice. *Science*, *363*(6426), 499–504. doi: 10.1126/science.aav3824

Bay, R. A., Rose, N., Barrett, R., Bernatchez, L., Ghalambor, C. K., Lasky, J. R., … Ralph, P. (2017). Predicting responses to contemporary environmental change using evolutionary response architectures. *The American Naturalist*, *189*, 463–473. doi: 10.1086/691233

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*(15), 2114–2120. doi: 10.1093/bioinformatics/btu170

Bouckaert, R., Heled, J., Kühnert, D., Vaughan, T., Wu, C.-H., Xie, D., … Drummond, A. J. (2014). BEAST 2: a software platform for bayesian evolutionary analysis. *PLoS Computational Biology*, *10*(4), e1003537. doi: 10.1371/journal.pcbi.1003537

Bridle, J. R., & Vines, T. H. (2007). Limits to evolution at range margins: when and why does adaptation fail? *Trends in Ecology & Evolution*, *22*(3), 140–147. doi: 10.1016/j.tree.2006.11.002

Brown, R. D., & Mote, P. W. (2009). The response of northern hemisphere snow cover to a changing climate. *Journal of Climate*, *22*(8), 2124–2145. doi: 10.1175/2008JCLI2665.1

Carneiro, M., Albert, F. W., Melo-Ferreira, J., Galtier, N., Gayral, P., Blanco-Aguiar, J. A., … Ferrand, N. (2012). Evidence for widespread positive and purifying selection across the European rabbit (*Oryctolagus cuniculus*) genome. *Molecular Biology and Evolution*, *29*(7), 1837–1849. doi: 10.1093/molbev/mss025

Cheng, E., Hodges, K. E., Melo-Ferreira, J., Alves, P. C., & Mills, L. S. (2014). Conservation implications of the evolutionary history and genetic diversity hotspots of the snowshoe hare. *Molecular Ecology*, *23*(12), 2929–2942. doi: 10.1111/mec.12790

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., … Durbin, R. (2011). The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–2158. doi: 10.1093/bioinformatics/btr330

Dobler, S., Dalla, S., Wagschal, V., & Agrawal, A. A. (2012). Community-wide convergent evolution in insect adaptation to toxic cardenolides by substitutions in the Na,K-ATPase. *Proceedings of the National Academy of Sciences of the United States of America*, *109*(32), 13040–13045. doi: 10.1073/pnas.1202111109

Fisher, R. A. (1937). The wave of advance of advantageous genes. *Annals of Eugenics*, *7*(4), 355–369. doi: 10.1111/j.1469-1809.1937.tb02153.x

Gigliotti, L. C., Diefenbach, D. R., & Sheriff, M. J. (2017). Geographic variation in winter adaptations of snowshoe hares (*Lepus americanus*). *Canadian Journal of Zoology*, *95*(8), 539–545. doi: 10.1139/cjz-2016-0165

Gillis, E. A., & Krebs, C. J. (1999). Natal dispersal of snowshoe hares during a cyclic population increase. *Journal of Mammalogy*, *80*, 933–939. doi: 10.2307/1383263

Giska, I., Farelo, L., Pimenta, J., Seixas, F. A., Ferreira, M. S., Marques, J. P., … Melo-Ferreira, J. (2019). Introgression drives repeated evolution of winter coat color polymorphism in hares. *Proceedings of the National Academy of Sciences of the United States of America*. doi: 10.1073/pnas.1910471116

Haldane, J. B. S. (1924). A mathematical theory of natural and artificial selection (Part I). *Transactions of the Cambridge Philosophical Society*, *23*, 19–41.

Haldane, J. B. S. (1932). *The causes of evolution*. London: Longmans, Green and Co.
Haller, B. C., & Messer, P. W. (2019). SLiM 3: Forward genetic simulations beyond the Wright-
Fisher model. *Molecular Biology and Evolution*, *36*(3), 632–637. doi: 10.1101/418657
Harris, R. B., Irwin, K., Jones, M. R., Laurent, S., Barrett, R. D. H., Nachman, R. D. H., …
Pfeifer, S. P. (2019). The population genetics of crypsis in vertebrates: recent insights from
mice, hares, and lizards. *Heredity*, doi:10.1038/s41437-019-0257-4.
Hoekstra, H. E., Drumm, K. E., & Nachman, M. W. (2004). Ecological genetics of adaptive
color polymorphism in pocket mice: Geographic variation in selected and neutral genes.
Evolution, *58*(6), 1329–1341. doi: 10.1111/j.0014-3820.2004.tb01711.x
Hoekstra, H. E., Krenz, J. G., & Nachman, M. W. (2005). Local adaptation in the rock pocket
mouse (Chaetodipus intermedius): Natural selection and phylogenetic history of
populations. *Heredity*, *94*(2), 217–228. doi: 10.1038/sj.hdy.6800600
Hoekstra, H. E., & Nachman, M. W. (2003). Different genes underlie adaptive melanism in
different populations of rock pocket mice. *Molecular Ecology*, *12*(5), 1185–1194. doi:
10.1046/j.1365-294X.2003.01788.x
Hoffmann, A. A., & Sgrò, C. M. (2011). Climate change and evolutionary adaptation. *Nature*,
470(7335), 479–485. doi: 10.1038/nature09670
Jones, M. R., Mills, L. S., Alves, P. C., Callahan, C. M., Alves, J. M., Lafferty, D. J. R., …
Good, J. M. (2018). Adaptive introgression underlies polymorphic seasonal camouflage in
snowshoe hares. *Science*, *360*(6395), 1355–1358. doi: 10.1126/science.aar5273
Kimura, M. (1957). Some Problems of Stochastic Processes in Genetics. *The Annals of
Mathematical Statistics*, *28*(4), 882–901. doi: 10.1214/aoms/1177706791
Kingsley, E. P., Manceau, M., Wiley, C. D., & Hoekstra, H. E. (2009). Melanism in *Peromyscus*
is caused by independent mutations in *Agouti*. *PLoS ONE*, *4*(7), e6435. doi:
10.1371/journal.pone.0006435
Knowles, N., Dettinger, M. D., Cayan, D. R., Knowles, N., Dettinger, M. D., & Cayan, D. R.
(2006). Trends in snowfall versus rainfall in the western United States. *Journal of Climate*,
19(18), 4545–4559. doi: 10.1175/JCLI3850.1
Kofler, R., Orozco-terWengel, P., De Maio, N., Pandey, R. V., Nolte, V., Futschik, A., …
Schlötterer, C. (2011). PoPoolation: a toolbox for population genetic analysis of next
generation sequencing data from pooled individuals. *PLoS ONE*, *6*(1), e15925. doi:
10.1371/journal.pone.0015925
Kreiner, J. M., Giacomini, D. A., Bemm, F., Waithaka, B., Regalado, J., Lanz, C., … Wright, S.
I. (2019). Multiple modes of convergent adaptation in the spread of glyphosate-resistant
Amaranthus tuberculatus . *Proceedings of the National Academy of Sciences*, 201900870.
doi: 10.1073/pnas.1900870116
Laurent, S., Pfeifer, S. P., Settles, M. L., Hunter, S. S., Hardwick, K. M., Ormond, L., …
Rosenblum, E. B. (2016). The population genomics of rapid adaptation: disentangling
signatures of selection and demography in white sands lizards. *Molecular Ecology*, *25*(1),
306–323. doi: 10.1111/mec.13385
Li, H. (2013). Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
ArXiv, 1303.3997. Retrieved from http://arxiv.org/abs/1303.3997
Magoč, T., & Salzberg, S. L. (2011). FLASH: fast length adjustment of short reads to improve
genome assemblies. *Bioinformatics*, *27*(21), 2957–2963. doi: 10.1093/bioinformatics/btr507
Manceau, M., Domingues, V. S., Mallarino, R., & Hoekstra, H. E. (2011). The developmental
role of *Agouti* in color pattern evolution. *Science*, *331*(6020), 1062–1065. doi:

10.1126/science.1200684

Marques, D. A., Taylor, J. S., Jones, F. C., Di Palma, F., Kingsley, D. M., & Reimchen, T. E. (2017). Convergent evolution of SWS2 opsin facilitates adaptive radiation of threespine stickleback into different light environments. *PLOS Biology*, *15*(4), e2001627. doi: 10.1371/journal.pbio.2001627

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., … DePristo, M. A. (2010). The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data. *Genome Research*, *20*(9), 1297–1303. doi: 10.1101/gr.107524.110

Melo-Ferreira, J., Seixas, F. A., Cheng, E., Mills, L. S., & Alves, P. C. (2014). The hidden history of the snowshoe hare, *Lepus americanus*: extensive mitochondrial DNA introgression inferred from multilocus genetic variation. *Molecular Ecology*, *23*(18), 4617–4630. doi: 10.1111/mec.12886

Milholland, B., Dong, X., Zhang, L., Hao, X., Suh, Y., & Vijg, J. (2017). Differences between germline and somatic mutation rates in humans and mice. *Nature Communications*, *8*, 15183. doi: 10.1038/ncomms15183

Mills, L. S., Bragina, E. V, Kumar, A. V, Zimova, M., Lafferty, D. J. R., Feltner, J., … Fay, K. (2018). Winter color polymorphisms identify global hot spots for evolutionary rescue from climate change. *Science*, *359*(6379), 1033–1036. doi: 10.1126/science.aan8097

Mills, L. S., Zimova, M., Oyler, J., Running, S., Abatzoglou, J. T., & Lukacs, P. M. (2013). Camouflage mismatch in seasonal coat color due to decreased snow duration. *Proceedings of the National Academy of Sciences of the United States of America*, *110*(18), 7360–7365. doi: 10.1073/pnas.1222724110

Nachman, M. W., Hoekstra, H. E., & D'Agostino, S. L. (2003a). The genetic basis of adaptive melanism in pocket mice. *Proceedings of the National Academy of Sciences of the United States of America*, *100*(9), 5268–5273. doi: 10.1073/pnas.0431157100

Nachman, M. W., Hoekstra, H. E., & D'Agostino, S. L. (2003b). The genetic basis of adaptive melanism in pocket mice. *Proceedings of the National Academy of Sciences of the United States of America*, *100*(9), 5268–5273. doi: 10.1073/pnas.0431157100

Nagorsen, D. W. (1983). Winter pelage colour in snowshoe hares (*Lepus americanus*) from the Pacific Northwest. *Canadian Journal of Zoology*, *61*(10), 2313–2318. doi: 10.1139/z83-305

Nelson, T. C., Jones, M. R., Velotta, J. P., Dhawanjewar, A. S., & Schweizer, R. M. (2019). UNVEILing connections between genotype, phenotype, and fitness in natural populations. *Molecular Ecology*, *28*(8), 1866–1876. doi: 10.1111/mec.15067

Orr, H. A., & Betancourt, A. J. (2001). Haldane's sieve and adaptation from the standing genetic variation. *Genetics*, *157*(2), 875–884.

Petkova, D., Novembre, J., & Stephens, M. (2016). Visualizing spatial population structure with estimated effective migration surfaces. *Nature Genetics*, *48*(1), 94–100. doi: 10.1038/ng.3464

Pfeifer, S. P., Laurent, S., Sousa, V. C., Linnen, C. R., Foll, M., Excoffier, L., … Jensen, J. D. (2018). The evolutionary history of Nebraska deer mice: local adaptation in the face of strong gene flow. *Molecular Biology and Evolution*, *35*, 792–806. doi: 10.1093/molbev/msy004

Protas, M. E., Hersey, C., Kochanek, D., Zhou, Y., Wilkens, H., Jeffery, W. R., … Tabin, C. J. (2006). Genetic analysis of cavefish reveals molecular convergence in the evolution of albinism. *Nature Genetics*, *38*(1), 107–111. doi: 10.1038/ng1700

R Core Team. (2018). *R: A language and environment for statistical computing.* Retrieved from
https://www.r-project.org/
Ralph, P. L., & Coop, G. (2010). Parallel adaptation: One or many waves of advance of an
advantageous allele? *Genetics*, *186*(2), 647–668. doi: 10.1534/genetics.110.119594
Ralph, P. L., & Coop, G. (2015). Convergent evolution during local adaptation to patchy
landscapes. *PLOS Genetics*, *11*(11), e1005630. doi: 10.1371/journal.pgen.1005630
Rohland, N., & Reich, D. (2012). Cost-effective, high-throughput DNA sequencing libraries for
multiplexed target capture. *Genome Research*, *22*(5), 939–946. doi: 10.1101/gr.128124.111
Rosenblum, E. B., Römpler, H., Schöneberg, T., & Hoekstra, H. E. (2010). Molecular and
functional basis of phenotypic convergence in white lizards at White Sands. *Proceedings of*
the National Academy of Sciences of the United States of America, *107*(5), 2113–2117. doi:
10.1073/pnas.0911042107
Schlager, G., & Dickie, M. M. (1971). Natural mutation rates in the house mouse. Estimates for
five specific loci and dominant mutations. *Mutation Research*, *11*(1), 89–96. doi:
10.1016/0027-5107(71)90034-0
Slatkin, M. (1973). Gene flow and selection in a cline. *Genetics*, *75*(4), 733–756. Retrieved from
http://www.ncbi.nlm.nih.gov/pubmed/4778791
Stamatakis, A. (2014). RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics*, *30*(9), 1312–1313. doi: 10.1093/bioinformatics/btu033
Steiner, C. C., Rompler, H., Boettger, L. M., Schoneberg, T., & Hoekstra, H. E. (2008). The
genetic basis of phenotypic convergence in beach mice: similar pigment patterns but
different genes. *Molecular Biology and Evolution*, *26*(1), 35–45. doi:
10.1093/molbev/msn218
Teshima, K. M., & Przeworski, M. (2006). Directional positive selection on an allele of arbitrary
dominance. *Genetics*, *172*(1), 713–718. doi: 10.1534/genetics.105.044065
Turner, J. R. G. (1981). Adaptation and evolution in Heliconius: a defense of NeoDarwinism.
Annual Review of Ecology and Systematics, *12*, 99–121. doi:
10.1146/annurev.es.12.110181.000531
Wright, S. (1931). Evolution in Mendelian populations. *Genetics*, *16*, 97–159.
Zhang, C., Rabiee, M., Sayyari, E., & Mirarab, S. (2018). ASTRAL-III: Polynomial time species
tree reconstruction from partially resolved gene trees. *BMC Bioinformatics*, *19*. doi:
10.1186/s12859-018-2129-y
Zheng, Y., Peng, R., Kuro-o, M., & Zeng, X. (2011). Exploring patterns and extent of bias in
estimating divergence time from mitochondrial DNA sequence data in a particular lineage:
a case study of salamanders (Order Caudata). *Molecular Biology and Evolution*, *28*(9),
2521–2535. doi: 10.1093/molbev/msr072
Zimova, M., Hackländer, K., Good, J. M., Melo-Ferreira, J., Alves, P. C., & Mills, L. S. (2018).
Function and underlying mechanisms of seasonal colour moulting in mammals and birds:
what keeps them changing in a warming world? *Biological Reviews*, *93*(3), 1478–1498. doi:
10.1111/brv.12405
Zimova, M., Mills, L. S., & Nowak, J. J. (2016). High fitness costs of climate change-induced
camouflage mismatch. *Ecology Letters*, *19*(3), 299–307. doi: 10.1111/ele.12568
772