1 **Title:**

2 **Evolution of gastrointestinal tract morphology and plasticity in cave-adapted**

3 **Mexican tetra, *Astyanax mexicanus***

5 **Authors:**

Misty R. Riddle[1], Fleur Damen[1], Ariel Aspiras[1,*], Julius A. Tabin[1], Suzanne McGaugh[2], Clifford J. Tabin[1,#]

6 **Affiliations:**

1. Department of Genetics, Blavatnik Institute, Harvard Medical School, Boston, MA 02115
7 2. Ecology, Evolution, and Behavior, University of Minnesota, Saint Paul, MN
8
* Current affiliation: Department of Stem Cell and Regenerative Biology, Harvard
University, Cambridge, MA 02138
For correspondence: tabin@genetics.med.harvard.edu

Abstract

The gastrointestinal tract has evolved in numerous ways to allow animals to

optimally assimilate energy from different foods. The morphology and physiology of the

gut is plastic and can be greatly altered by diet in some animals. In this study, we

investigated the evolution and plasticity of gastrointestinal tract morphology by comparing

laboratory-raised cave- and river-adapted forms of the Mexican tetra, *Astyanax mexicanus,*

reared under different dietary conditions. In the wild, river-dwelling populations (surface

fish) consume plants and insects throughout the year, while cave-dwelling populations

(cavefish) live in a perpetually dark environment and depend on nutrient-poor food

brought in by bats or seasonal floods. We found that multiple cave populations converged

on a reduced number of digestive appendages called pyloric caeca and that some cave

populations have a lengthened gut while others have a shortened gut. Moreover, we

identified differences in how gut morphology and proliferation respond to diet between

surface fish and cavefish. Using a combination of quantitative genetic mapping, population

genetics, and RNA sequencing, we found that changes to the molecular and genetic

pathways that influence cell proliferation, differentiation, and immune system function

may underlie evolution of the cavefish gut.

Introduction

The gastrointestinal tract consists of functional regions distinguished by the organization of muscle and mucosal layers and cellular composition of epithelium lining the lumen. The secretory and absorptive cell types of the epithelium are constantly being replaced by self-renewing stem cells. Gut homeostasis requires integration of intrinsic signaling pathways set up during development, with external cues. For example, in the mammalian gut, Notch and WNT signaling is essential for establishing the stem cell niche and balancing stem cell renewal and differentiation of cell types (reviewed in 1). High-fat diet alters these pathways to favor self-renewal and secretory cell formation (2, 3). Cytokine signals from tissue-resident immune cells can also promote stem cell renewal in response to infection (4, 5). While studies utilizing model organisms and organoid culture have begun to reveal how homeostasis of the epithelium is maintained, there is a limited understanding of how these pathways evolve as animals adapt to different diets.

Evolutionary changes to gut morphology, such as expansion of functional domains, have been correlated to altered patterns of gene expression during development (6). The organization of the gut epithelium into crypts that support villi is important for stem cell maintenance in mammals, but in other species the epithelium is characterized by irregular folds, zig-zags, honeycomb patterns, or a spiral fold (7, 8). The presence of spatially restricted gut stem cells is common across distant phyla (i.e. arthropods and chordates) (9–11). Although it has been demonstrated that diet influences gut morphology in a number of species (reviewed in 12), evolution of the signaling pathways that integrate internal and external cues is not well understood. For animals that survive months without food, like snakes and migrating birds (13–15), altering the pathways that control homeostasis of the epithelium was likely critical to balance nutrient assimilation and energy conservation.

Astyanax mexicanus is a species of fish that exists as river- and cave-adapted populations that evolved on very different diets (16, 17). The river (surface) fish have access to insects and plants in abundance, while the cavefish are dependent on predominantly bat droppings and material brought in by seasonal floods (16, 17). There are a number of cave populations, named for the caves they inhabit (i.e. Tinaja, Pachón, Molino). Based on whole genome sequencing analysis, these populations reflect two independent cave derivations from surface fish within the last 200,000 years; Tinaja and

Pachón cavefish populations form a separate clade from Molino cavefish (18). Cavefish

have converged on similar morphological changes, such as eye loss and increased sensory

structures (19–21), behavioral changes such as reduced sleep (22–25), and metabolic

changes such as increased fat accumulation and starvation resistance (26–28). We

previously showed that Pachón cavefish have altered gastrointestinal motility during post-

larval growth to slow food transit, possibly to achieve increased nutrient absorption (29).

In this study, we explore whether cave-adaptation has led to changes in the adult

gut. First, we describe the anatomy and histology of the *A. mexicanus* gastrointestinal tract,

noting distinct functional regions. Second, we compare *A. mexicanus* surface and cave

populations fed the same diet and discover differences in pyloric caeca number and gut

length. We show that the homeostasis and morphology of the gut responds differently to

changes in diet comparing Tinaja cavefish to surface fish. To investigate the genetic

architecture controlling gut length, we carry out a Quantitative Trait Locus (QTL) analysis,

and identify a significant QTL associated with hindgut length. We examine genes within the

QTL using population genetics and RNA sequencing and reveal genetic pathways that have

been altered during the evolution of gut morphology in *A. mexicanus*.

Results

Anatomy and histology of the adult Astyanax mexicanus gastrointestinal (GI) tract

The GI tract of teleost fish is commonly divided into four sections from anterior to

posterior: head gut, foregut, midgut, and hindgut. These sections are easily defined in *A.*

mexicanus and can be further subdivided based on distinct morphology and histology. The

foregut consists of the esophagus, stomach, and pylorus (Figure 1A, B). The esophagus has

two perpendicular layers of striated muscle and a stratified epithelium of mucus secreting

cells, identifiable by a lack of staining with hematoxylin and eosin (Figure 1A). It connects

to the J-shaped stomach that has three muscle layers: outermost longitudinal, the middle

and thickest circumferential, and innermost oblique (Figure 1B). Gastric glands are evident

in the stomach mucosa; at the base of the glands, pepsin-secreting chief cells are

distinguishable by eosinophilic granules. The lumen-facing epithelium of the stomach

consists of columnar mucus-secreting cells. The stomach ends at the pylorus that has thick

muscle layers and connects the foregut to the midgut.

The midgut is the longest portion of the GI tract. The proximal end of the midgut

contains pyloric caeca: pouch structures that have a thin layer of outer connective tissue

and lamina propria, and an inner epithelium organized into ridges that run from the base,

to the tip of the pouch (Figure 2A). The epithelium is made up of mostly columnar

enterocytes and a few mucus secreting goblet cells (Figure 1C). There are four pyloric caeca

on the right-hand side of the midgut (Figure 2, "1-4"), two on the left-hand side ("5-6"), and

as many as three that are shorter and more proximally restricted ("7-9"). Caeca one

through five are present in fish of all four populations (surface, Tinaja, Pachón, and Molino)

while six through nine are variable. Surface fish are more likely to have 1-9, Tinaja and

Molino 1-6, and Pachón 1-7 (Figure 2C).

The remaining tubular portion of the midgut has inner circumferential and outer

longitudinal muscle layers, and the epithelium is folded circumferentially at irregular

angles. The epithelium of the proximal midgut is mostly mucus secreting goblet cells that

completely lack H&E staining (Figure 1D, D', black arrow), compared to the epithelium of

the distal midgut that has goblet cells with basophilic granules (Figure 1E'). The lumen of

the distal midgut also contains basophilic enteroendocrine cells that are distinguishable by

a smaller size and more apical location (Figure 1E' yellow arrow). A greater number of

eosinophilic immune cells are also evident in the lamina propria of the distal region of the

midgut (Figure 1E' black arrowhead).

The proximal hindgut is wider than the midgut and has thinner muscle and lamina

propria layers (Figure 1F-F'). The enterocytes in the epithelium are highly vacuolated likely

representing absorbed hydrophobic molecules (Figure 1F', grey arrow). Apically located

enteroendocrine cells are also evident in the epithelium (Figure 1F', yellow arrow). The

distal hindgut (rectum) has a thicker muscle wall and connects the GI tract to the anus.

Overall, the organization of the *A. mexicanus* gut is similar to other Characiformes. Having

established the general anatomy and histology of the GI tract using the surface form, we

were next able to explore differences between the surface and cave populations.

Ecotypic differences in gut length

The length of each region of the GI tract is an important variable determining

digestive efficiency and can be altered by diet. To determine the ecotypic differences in gut

length between populations, we compared relative gut length (gut length divided by fish

length) between individually housed fish fed the same diet (38% protein, 7% fat, 5% fiber)

and amount per day (6mg). We found that Tinaja (n=6) and Molino (n=5) cavefish tend to

have a longer midgut, and Pachón (n=6) cavefish tend to have a shorter midgut compared

to surface fish (n=6) (Figure 3A, B). Tinaja and Molino also tend to have a longer hindgut

and Pachón have a significantly shorter hindgut compared to surface fish (Figure 2C,

$p=0.05$, one-way ANOVA with Tukey's post hoc test). Interestingly, while all fish were

dissected 24 hours post-feeding, we noted a difference in the amount of fecal matter in the

gut (Figure 3A). All of the Tinaja cavefish guts were entirely full (n=6), compared to one out

of six Pachón, and four out of five Molino. All of the surface fish guts were mostly empty as

seen in Figure 3A (n=6). In summary, there are moderate differences in adult gut length

between populations that are independent of diet, and some adult cavefish populations

may have slower gastrointestinal transit.

Diet influences gut morphology differently in surface fish and Tinaja cavefish

Surface-adapted *A. mexicanus* consume plants and insects throughout the year. In

contrast, cavefish populations are subject to seasonal fluctuations, including periods of

deprivation. We hypothesized that cavefish may have evolved increased plasticity in gut

morphology to maximize nutrient absorption during times of abundance and conserve

energy when food is scarce. To test this hypothesis, we switched the diet of adult surface

fish and Tinaja cavefish from moderate nutrient content (7% fat, 38% protein, 5% fiber) to

either low-nutrient (4% fat, 32% protein, 3% fiber) or high-nutrient (18% fat, 55% protein,

2% fiber). After eight months, we measured the length of the gut segments, as well as the

circumference and average number of folds in cross sections, and compared the values to

fish that remained on the moderate nutrient diet (Figure 4, n=3 fish per population and

diet, and 3 histological sections per segment). On the moderate nutrient diet fed *ad libitum*,

Tinaja tend to have a shorter midgut that is wider and has more folds compared to surface

fish (relative midgut length: surface = 0.45, Tinaja = .36, circumference: surface = 0.16cm,

Tinaja = 0.20, circumference normalized to fish length: surface = 0.03, Tinaja = 0.04, fold

number: surface = 6, Tinaja = 9, Figure 4 A-C). Tinaja also tend to have a longer hindgut on

this diet as was observed on the controlled diet (surface = 0.17, Tinaja = 0.20).

In fish that were switched to a low-nutrient diet, we observed a 24% increase in the

length of the midgut in both populations (surface = 0.45 to 0.56, p=0.36; Tinaja = 0.36 to

0.44, p = 0.60; one-way ANOVA with HSD post-hoc test, Figure 4A, G). The circumference of

the midgut also increased significantly by 89% in surface fish (0.16 to 0.30 cm, normalized

to fish length 0.03 to 0.06, p<0.005) and 78% in Tinaja cavefish (0.20 to 0.36 cm,

normalized to fish length 0.04 to 0.07, p<0.005, Figure 4B, G). The number of folds in the

midgut epithelium increased 2-fold in Tinaja (9 to 18) and nearly 2-fold in surface fish (6 to

11). The hindgut length increased by 70% in surface fish (0.17 to 0.28) and only 7% in

Tinaja cavefish (0.20 to 0.21, Figure 4D, H). The circumference of the surface fish hindgut

did not change significantly (0.34 to 0.33cm, normalized to fish length = 0.06 to 0.07) and

the circumference of the Tinaja hindgut increased by 45% in Tinaja (0.24 to 0.35cm,

normalized to fish length =0.04 to 0.07, Figure 4D, H). The number of folds in the hindgut

epithelium changed only slightly in both populations (Tinaja = 13 to 12, surface = 11 to 13).

In summary, midgut length, width, and fold number increased similarly in both populations

in response to a low-nutrient diet (Figure 4G). In contrast, the hindgut responds by

growing in length in surface fish and, growing in width in cavefish.

In fish switched from a moderate to a high-nutrient diet, the length of the midgut did

not change in surface fish (0.45 to 0.46, Figure 4A) and decreased in Tinaja cavefish by 20%

(0.36 to 0.29). The circumference of the midgut increased in surface fish (0.16 to 0.23cm,

normalized to fish length = 0.03 to 0.05) and did not change in cavefish (0.20 to 0.20cm,

normalized to fish length = 0.04 to 0.04; Figure 4C). The average number of epithelial folds

increased slightly in surface fish (6 to 7) and decreased slightly in the cavefish (9 to 8,

Figure 4C). Normalized hindgut length decreased in both populations in response to a high-

nutrient diet, although the decrease was greater in Tinaja: 39% (0.20 to 0.12) versus 31%

in Surface fish (0.17 to 0.12, Figure 4B). The circumference of the hindgut also decreased,

by 26% in surface fish (0.34 to 0.25cm, normalized to fish length = 0.06 to 0.05) and by

21% in cavefish (0.24 to 0.19cm) although the value normalized to fish length did not

change in cavefish (0.04 to 0.04, Figure 4D). The number of folds in the hindgut decreased

by one in surface fish (11 to 10) but more strikingly in Tinaja cavefish reduced to almost

half (13 to 7, Figure 4C). In summary, the Tinaja cavefish gut is more responsive than

surface fish to a high-nutrient diet since Tinaja cavefish exhibit a greater reduction in

midgut and hindgut length and hindgut fold number. Overall, the results suggest that

plasticity of the gut is dependent on the type of diet and region of the gut and reveal

fundamental differences between the populations.

Differences in the regulation of gut proliferation between surface fish and Tinaja cavefish

The gastrointestinal epithelium undergoes constant turnover, being replenished by

populations of dividing cells that are restricted to the base of villi or epithelial folds (1). We

next sought to understand how differences in morphology are achieved by investigating

homeostasis of the gut epithelium. We injected fish on the low-nutrient diet with a

thymidine analog (5-ethynyl-2'-deoxyuridine, EdU) and, after 24 hours, determined the

number of EdU positive cells as a measure of the amount of cell division (Figure 5A, B, F, J).

We found that Tinaja cavefish had on average over four times as many EdU positive cells in

the midgut (Figure 5F, surface = 103, Tinaja = 449, $p = 0.063$ two-tailed t-test). Higher

proliferation corresponds to a significantly greater number of folds in the midgut

epithelium in Tinaja cavefish (Figure 4C, average surface = 11, Tinaja = 18; $p = 0.04$, two-

tailed t-test). However, the number of EdU positive cells per fold in the midgut is also

greater in Tinaja (average surface = 11, Tinaja = 26; $p = 0.09$); EdU positive cells in the

epithelium appear more restricted to the base of the folds in surface fish, whereas they

extend further up the fold in Tinaja cavefish (Figure 5A, B). In the hindgut, the average

number of EdU positive cells is also greater in Tinaja (Figure 5J, surface = 154, Tinaja =

354; $p = 0.17$), although there is only a slight difference in the number of folds (surface =

13, Tinaja = 12). These results suggest that altered homeostasis of the gut epithelium could

drive morphological differences between the populations.

Cavefish evolved in a nutrient-limited environment and unlike surface fish are

subject to periods of starvation. Because cell-turnover is energetically expensive, we

hypothesized that cavefish may have evolved mechanisms to limit the proliferation in the

gut when food is not available. To test this, we starved fish (for a period of two-weeks) that

were previously fed the low-nutrient diet, and compared EdU incorporation over a 24-hour

period (Figure 5F, J, K, L). The number of EdU positive cells in the Tinaja midgut decreased

significantly by 81% (average 449 fed versus 85 starved, p = 0.005, one-way ANOVA

comparing diet and population). In contrast, the number of EdU positive cells nearly

doubled in surface fish (average 103 fed versus 193 fasted, p = 0.63). In both populations,

midgut length, circumference, and fold number decreased in response to starvation (Figure

5C-D). In surface fish, the midgut length shortened significantly by 40% (0.56 to 0.34,

p=0.01, one-way ANOVA) compared to only 19% in Tinaja cavefish (0.44 to 0.36, p=0.42).

The midgut circumference decreased by 25% in surface fish (0.30 to 0.23cm, normalized to

fish length = 0.06 to 0.05) and significantly in cavefish by 34% (0.36 to 0.24cm, p=0.004,

normalized to fish length = 0.07 to 0.05; p=0.03). The number of folds in the midgut

decreased from 11 to 7 in surface fish (p=0.53) and more dramatically, from 18 to 6 in

Tinaja cavefish (p=0.01). In summary, Tinaja cavefish reduce proliferation in response to

starvation and this is associated with a substantial decrease in midgut circumference and

fold number, but a more moderate decrease in length, compared to surface fish.

In the hindgut, the average number of EdU positive cells also reduced significantly in

Tinaja cavefish in response to starvation (Figure 5J, average 354 to 130, p=0.005, one-way

ANOVA), and to a lesser extent in surface fish (38%, average 154 to 95, p= 0.63 one-way

ANOVA). Relative hindgut length reduced by 50% in surface fish (0.28 to 0.14) and only

20% in Tinaja (0.21 to 0.15). We observed a similar reduction in hindgut circumference, by

35% in surface fish (0.33 to 0.21cm, normalized to fish length = 0.07 to 0.04) and 34% in

Tinaja cavefish (0.35 to 0.23cm, normalized to fish length = 0.07 to 0.05). The number of

hindgut folds reduced from 13 to 8 in surface fish and 12 to 6 in Tinaja cavefish. Similar to

what occurs in the midgut, in the hindgut, Tinaja decrease proliferation more substantially

in response to starvation but have a less drastic decrease in length compared to surface

fish. Combined, our findings support the hypothesis that cavefish have mechanisms for

tuning gut homeostasis in response to food intake that differ from surface fish.

Genetic mapping reveals quantitative trait loci associated with hindgut length

We next sought to understand the genetic basis of differences in gut morphology

between surface fish and cavefish. We carried out a quantitative trait loci (QTL) analysis

using F2 surface/Tinaja hybrids (see methods). To eliminate the effect of diet or appetite,

we individually housed the hybrids as adults and ensured they consumed the same diet

(38% protein, 7% fat, 5% fiber) and amount per day (6mg). After four months, we

determined the length and weight of the fish, number of pyloric caeca, and length of the

midgut and hindgut. We identified a QTL for female body condition (fish weight/fish

length) on linkage group 13 with a LOD score of 5.53 accounting for 21% of the variance in

this trait (Figure 6B). We did not identify a QTL for male body condition (fish weight/fish

length Figure 6A). We also did not identify a significant QTL for pyloric caeca number but

observed a peak on linkage group 24 (Figure 6C).

There is a strong positive correlation between fish length and gut length as

anticipated (midgut p-value=2.2×10^{-16}, hindgut p-value=5.2×10^{-14}, Pearson's product-

moment correlation). We therefore performed the QTL analysis using relative gut length.

We did not identify a significant QTL for relative midgut length but observed a peak on

linkage group 14 (Figure 6D). We found a significant QTL for relative hindgut length on

linkage group 10 with a LOD score of 4.82 at the peak marker r172341 accounting for 11%

of the variance in this trait. Interestingly, it is the heterozygous genotype at this position

that is associated with longest relative hindgut length (Figure 7C). Similarly, F1 hybrids

have a longer hindgut than surface fish or Tinaja cavefish (Figure 7A,B). The midgut of F1

hybrids is also longer; however, when we examined the effect plot for the marker with the

highest LOD score for relative midgut length, we found that individuals with cave genotype

have the longest midgut (Figure 7D, E). Combined, the results indicate overdominance or

positive epistasis in the genetic network controlling gut length.

Candidate genes controlling hindgut length

To search for genetic changes that contribute to variation in hindgut length, we used

a combination of population genetics and differential gene expression analysis. We first

used the Pachón cavefish genome assembly to identify the position of markers that define

the 1.5-LOD support interval for the hindgut QTL. We found that two markers are on

scaffold KB882083 and one is on KB871783 (Table 1). We determined several population

genetic metrics for all of the genes on each scaffold using whole genome sequencing data

(Dataset S1, 18). This data set includes Tinaja, Pachón and Molino cave populations and

two surface populations: Río Choy, which is representative of our lab strains and is more

closely related to the stock of fish that invaded the Molino cave, and Rascón, which is more

representative of a separate stock of surface fish that invaded the Tinaja and Pachón caves.

We found that 45 genes within the 1.5-LOD support interval for the hindgut QTL have fixed

differences in the coding regions between Río Choy surface fish (n=9) and Tinaja cavefish

(n=10) populations (Dataset S1, all on scaffold KB882083). Ten of these genes may be

under selection since the haplotype including the genes exhibits evidence of non-neutral

evolution comparing Río Choy to Tinaja (Max pairwise divergence = 1 and significant

HapFLK p-values (< 0.05), Table 2, Figure 8A). Two of the ten genes also fit within the same

statistical parameters comparing Rascón surface fish to Tinaja cavefish (Table 2). These

two genes are predicted to encode complement factor B-like proteins that are involved in

innate immunity (herein referred to as Cfb1 and Cfb2). We found that *cfb1/2* also have

significant HapFLK p-values comparing Pachón to surface populations, but not to Molino

(Table 2). The ten genes with significant HapFLK values in Tinaja have varying ranks of

divergence in the other cave populations (Figure 8A,B). Genes that show high levels of

divergence in all populations may be generally important for cave adaptation (e.g. *cfb1/2*,

Figure 8A, B). Genes that show divergence in only the Tinaja population (e.g. *klf5l*) may be

important for adaptation only in the Tinaja cave.

Next, we explored which genes in the QTL may have regulatory mutations by

comparing levels of expression. For this analysis, we utilized the surface fish genome

assembly that is organized into chromosomes. We found that the markers that define the

estimated confidence interval for the hindgut QTL are on chromosome 14 and span a

region of approximately 15KB (Table 1). This region has 342 genes. We determined if any

of these genes are differentially expressed in the hindgut using RNA sequencing data from

adult surface, Tinaja, and surface/Tinaja F1 hindguts (30) (n=5 hindguts per population).

We found that 31 genes are differentially expressed using a likelihood ratio test to compare

all three sample types (Table S1). Pairwise comparison between Tinaja cavefish and

surface fish revealed 9 additional differentially expressed genes (Table S1). Included in this

list is *cfb1*. Tinaja cavefish exhibit the highest expression of *cfb1* and expression in the F1

hybrid is intermediate (Figure 8C). We reasoned that genes showing either the greatest or

lowest expression in F1 hybrids may be strong candidates for controlling differences in

length since the heterozygous genotype at the QTL is associated with the longest hindgut.

Of the genes that are in the QTL and differentially expressed, five show the highest

expression in the hybrid and eight show the lowest expression in the hybrid. Among the

genes that are lowest in the hybrid is *Notch1A* (Figure 8C, Table S1). Notch signaling

controls stem cell renewal and formation of secretory versus absorptive cell fate in the

intestine (31).

Expression patterns associated with differences in gut length

We next broadened our analysis of the RNA sequencing data to investigate other

pathways that may have been altered during the evolution of the cavefish gut; 3550 genes

are differentially expressed between surface, F1 hybrids, and Tinaja hindguts (30). We

searched for patterns across the differentially expressed genes between sample groups

(degPatterns function in DEseq 2, Figure S1) and identified clusters of genes that show

either greatest (Table S2, group 3, n=55) or lowest (Table S3, group4, n=53) expression in

the F1 hybrid, reasoning that these may be more likely to be associated with differences in

hindgut length. The genes that have the greatest expression in the hybrid include an

additional component of the complement system (*C8G*), a negative regulator of Notch

signaling (*nrarpa*), retinoic acid signaling components (*Cyp26A1, rdh1*)*,* and a circadian

rhythm gene (*timeless*) (Figure 8C). The genes that show a pattern of lowest expression in

the hybrid include tumor suppressor and apoptosis related genes (*casp6, brca2, pak1*), a

suppressor of cytokine activity (not annotated), regulator of retinoic acid production

(*bco1*), and a circadian rhythm gene that typically correlates with timeless expression

(*per1b*) (Figure 8C). In summary, we found that surface fish and cavefish have differences

in the genetic architecture controlling hindgut length, and that cavefish have altered

expression of genes controlling cell proliferation, cell signaling, and immune system

function.

Discussion

A major challenge for *Astyanax mexicanus* when they invaded caves was adapting to the drastically limited availability of nutrients compared to what they experienced in the ancestral river environment. In response, they evolved hyperphagia, starvation resistance, and increased fat accumulation (27, 28, 32). Here, we investigated how the morphology and homeostasis of their gastrointestinal tract adapted as a consequence of nutrient restriction. Our data suggest that specializations for individual caves may have occurred. By comparing fish on the same diet in the laboratory, we found that Tinaja and Molino cave populations converged on a lengthened gut while Pachón have a shortened gut. A plausible explanation for this is differences in cave ecology; the Pachón cave is less impacted by seasonal flooding and has cave-adapted micro-crustaceans that serve as a food source for post-larval fish (16). In all cave populations, we observed a reduction in the number of pyloric caeca, specifically absence of caeca 7-9. The pyloric caeca may have roles in digestion, absorption, and immune system function (33, 34). It is possible that caeca 7-9 are important for digestion of foods, or response to pathogens, that are only present in the surface fish environment, and these structures became vestigial and eventually disappeared in the cave. Other groups of fish like salmonids can have hundreds of pyloric caeca (35). Our study establishes a genetically accessible model to understand pyloric caeca function, development, and evolution.

We hypothesized that cavefish evolved increased plasticity of the gut as an adaptation to save energy during seasonal fluctuation of food. The most striking difference that we observed between populations is that cavefish have much greater proliferation in the gut epithelium compared to surface fish under fed conditions and reduce proliferation in response to food deprivation. Cavefish may have increased sensitivity in the pathways that sense or respond to nutrient shortage and it is plausible that this provides an advantage in the cave. On a high-nutrient diet, cavefish exhibit a more dramatic reduction in gut length compared to surface fish. Reducing length when nutrients are abundant could achieve a more optimal balance between energy extraction and storage; the gut is packaged within the body cavity and its size could limit the amount of visceral fat that can accumulate. How the cellular dynamics of the epithelium translate to changes in gut

morphology are not entirely clear. For example, proliferation increases in surface fish

during starvation, yet gut length decreases. The balance between cell renewal and death is

likely different between the populations and in line with this, we observed that Tinaja have

lower expression of tumor-suppressor and pro-apoptosis genes (i.e. *brca2*, *p21-activated*

kinase, *caspase 6*). Overall, while we observe plasticity in gut morphology in both surface

fish and cavefish, our data support the hypothesis that cavefish have a more dramatic

response to nutrient fluctuations.

At post-larval stages, Pachón cavefish have altered gastrointestinal motility,

resulting in delayed transit of ingested food (29). Here we found evidence of slower transit

of digestive material in adult cavefish populations as well. Tinaja, Pachón, and Molino guts

were full 24-hours after feeding, compared to surface fish guts that were mostly empty.

Slower transit time may have evolved to optimize digestion of decaying material mixed

with mud, a likely way food is obtained in the caves, and/or could result in increased

nutrient absorption. Our results highlight that altered gut function and morphology likely

accompany adaptation to the food sources in the cave.

We found evidence that the genetic architecture controlling gut morphology

diverged between surface fish and Tinaja cavefish. We identified a QTL associated with

hindgut length and found that F1 and heterozygous F2 hybrids have a longer hindgut

compared to surface fish and cavefish. This phenomenon (heterosis) has been observed for

another *A. mexicanus* trait, thickness of the inner retina (36), and is often seen in plant

breeding, yet the genetic and molecular basis is not easy to define (37). One hypothesis is

that each parent has genetic changes that result in an inferior phenotype, and that in the

hybrid these changes are complemented, resulting in a superior phenotype (Dominance).

Another possibility is that there are multiple alleles that interact in a way that produces a

phenotype greater than what could be achieved in a homozygous condition (Over-

dominance). It is possible that there are multiple alleles that have antagonistic effects on

hindgut length and when they are combined the result is a longer gut.

To investigate which genes in the QTL may be important for controlling gut length

we used a combination of population genetics and RNA sequencing. Our results revealed

two genes that are known to regulate intestinal homeostasis in other species, *klf5* (krüppel-

like factor 5) and *Notch1A*. KLF5 is a transcription factor expressed in intestinal crypts that

can promote or inhibit proliferation and is commonly dysregulated in colon cancer (38–

40). Notch signaling balances stem cell renewal and differentiation; activation of Notch

signaling promotes stem cell proliferation and suppresses secretory cell formation (31).

We found evidence that *klf5l* may be under selection in the Tinaja cave as we observed

significant HapFLK p-values comparing Tinaja to Rio Choy surface fish. While we did not

find the same evidence for the *notch1a* coding region, we found that expression of *notch1a*

is significantly different between surface fish, Tinaja cavefish, and F1 hybrids, suggesting

there may be regulatory mutations within the QTL that influence *notch1a* expression.

Tinaja cavefish tend to have higher n*otch1a* expression, in line with greater proliferation in

the epithelium. However, we found that notch-regulated ankyrin repeat protein (*nrarpa*)

that suppresses Notch signaling in other contexts (41, 42) is more highly expressed in both

Tinaja cavefish and F1 hybrids. Since Notch signaling tends to be context dependent,

comparing the components of the Notch signaling pathway with cellular resolution may

give insight into how differences in proliferation and morphology arise.

Cave-adapted *A. mexicanus* not only experience a different diet in the cave, but also a

different microbial landscape (43). We found that complement factor B (*cfb1*) is within the

hindgut QTL, shows significant divergence in all populations, and is differentially expressed

in the hindgut. Complement factor B is part of the innate immune system. It is expressed in

multiple tissues, including the colonic mucosa, and functions in the pathway that

recognizes and eliminates bacterial pathogens by controlling immune cell differentiation

(44, 45). Mounting evidences suggests that immune cells alter self-renewal of stem cells in

the epithelium through cytokine signaling (5). Interestingly, we found that cavefish and F1

hybrids have lower expression of a suppressor of cytokine signaling in the hindgut. Recent

research suggests that Pachón cavefish have fewer pro-inflammatory immune cells in the

visceral adipose tissue (46), but the amount and differences in the immune cells in the gut

have not been explored. Our results suggest that evolution of the immune system in *A.*

mexicanus likely plays a role in how the gut responds to external cues. In the lab, the

external microbial landscape is the same for the populations, yet it is possible that genetic

differences select for specific microbes and this in turn influences intestinal homeostasis.

Cavefish accumulate more carotenoids (retinoic acid (RA) precursors) in the

visceral adipose tissue compared to surface fish and this correlates with decreased

expression of genes that convert carotenoids into retinoids in the gut epithelium (47). We

found that cavefish and F1 hybrids have lower expression of beta-carotene oxygenase

(*bco1*) and higher expression of retinol dehydrogenase (*rhd1*) in the hindgut. The outcome

may be changes in RA production and signaling. RA signaling is important for stem cell

maintenance in a number of contexts (48), but how it regulates intestinal stem cell

homeostasis *in vivo* is not well understood. However, it is clear that RA signaling regulates

innate immune cells in the gut (49). It is possible that altered RA signaling in the cavefish

gut influences the crosstalk between immune cells and stem cells resulting in differences in

morphology.

We found that expression of the circadian clock genes *per1* and *timeless*, which are

typically synchronized, show the opposite expression pattern in the hindguts of cavefish

and F1 hybrids. Proliferation and gene expression in the intestine is under circadian

control and disruption of clock genes alters renewal of the epithelium (50–52). Having

evolved in complete darkness, cavefish have altered circadian rhythm as evidenced by

developmentally delayed and reduced amplitude of clock gene expression, lack of circadian

cycles of metabolism, and reduced sleep (26, 53–55). It is plausible that alterations to the

gene regulatory network controlling circadian rhythm influenced homeostasis of the

intestinal epithelium during cavefish evolution. Overall, our study reveals multiple levels at

which the *A. mexicanus* gut has evolved in response to ecological differences in food

availability including morphology, homeostasis, and plasticity in response to dietary

fluctuations. Furthermore, we implicate genetic changes and molecular pathways that may

have accompanied evolution of these traits.

Methods

Fish husbandry and diet

Fish husbandry was performed according to (56). For fixed diet experiments fish were

housed individually in 1.5L tanks and fed three pellets (approximately 6mg) of New Life

Spectrum TheraA+ small fish formula once per day for at least 4 months. Commercially-

available low nutrient (Hikari Goldfish Staple), moderate nutrient (New Life Spectrum

TheraA+), and high-nutrient (Ken's Premium Growth Meal #3) diets were fed *ad libitum.*

Phenotype quantification and data visualization

Fish were euthanized in 1400ppm Tricane and dissected 24-hours post feeding. Images

were taken using a Cannon Powershot D12 digital camera. ImageJ was used to measure fish

length and gut length. Data visualization was performed using R (57).

Histology and imaging

Figure 1: The fish was euthanized in 1400 ppm tricane, an incision was made along the

ventral body line, and it was placed into 10% buffered formalin and shipped to FishVet

Group (Portland, ME) for further processing. The gut was dehydrated using a graded

ethanol series, followed by several changes in xylene, and then embedded in paraffin.

Sections were taken at 5μm thickness, stained with hematoxylin and eosin (H&E) and

imaged using Aperio ScanScope CS. Figure 4,5: To quantify proliferation, fish were

anesthetized using 400ppm Tricane, then 40μL of EdU (500μM in .1% DMSO) was injected

into the intraperitoneal cavity using an insulin syringe. After 24 hours, fish were

euthanized in 1400 ppm tricane and the gut was immediately removed, incubated

overnight in 10% formalin, then washed in PBS and dehydrated to 100% ethanol through a

graded ethanol series. The gut was incubated briefly in Histo-clear (National Diagnostics).

The midgut and hindgut were separated and embedded in paraffin side by side in a single

block in anterior to posterior orientation to obtain a transverse section of each structure.

Blocks were sectioned at a thickness of 5μm using a Leica RM 2155 rotary microtome.

Slides were dried overnight, deparaffinized using Histo-clear, and rehydrated to water

through a graded ethanol series. The slides were placed in 10% buffered formalin for 10

minutes and washed twice in 3% Bovine Serum Albumin (BSA). EdU was detected using

the Click-IT EdU Cell Proliferation Kit (ThermoFisher) according to the manufacturers

protocol. Slides were mounted in ProLong Gold (ThermoFisher) and images were taken

using Keyence BZ2 microscope and software. We measured tissue morphology and EdU

cell number manually using ImageJ (58).

Quantitative trait loci analysis

F2 hybrid population. We bred a surface fish female with a Tinaja cavefish male to produce

a clutch of F1 hybrids. We generated a population of surface/Tinaja F2 hybrids by

interbreeding individuals from this clutch. The F2 mapping population (n=219) consisted

of three clutches produced from breeding paired F1 surface/Tinaja hybrid siblings.

Genotype by sequencing. We extracted DNA from caudal tail fins using DNeasy Blood and

Tissue DNA extraction kit (Qiagen). DNA was shipped to Novogene (Chula Vista, CA) for

quality control analysis and sequencing. Samples that contained greater than 1.5 ug DNA,

minimal degradation (as determined by gel electrophoresis), and OD_{260}/OD_{280} ratio of 1.8

to 2.0 were used for library construction. Each genomic DNA sample (0.3~0.6 µg) was

digested with Mse1, HaeIII, and EcoR1. Digested fragments were ligated with two barcoded

adapters: a compatible sticky end with the primary digestion enzyme and the Illumina P5

or P7 universal sequence. All samples were pooled and size-selected after several rounds of

PCR amplification to obtain the required fragments needed to generate DNA libraries.

Concentration and insert size of each library was determined using Qubit® 2.0 fluorometer

and Agilent® 2100 bioanalyzer respectively. Finally, quantitative real-time PCR (qPCR) was

used to detect the effective concentration of each library. Qualified DNA libraries had an

effective concentration of greater than 2 nM and were pooled by effective concentration

and expected data production. Pair-end sequencing was then performed on Illumina®

HiSeq platform, with the read length of 144 bp at each end. Raw Illumina genotype-by-

sequencing reads were cleaned and processed through the *process_shortreads* command in

the Stacks software package (59). The cleaned reads were aligned to the *Astyanax*

mexicanus reference genome (AstMex102, INSDC Assembly GCA_000372685.1, Apr 2013)

using the Bowtie2 software (60). The aligned reads of 4 surface fish, 4 Tinaja cavefish and 4

F_1 surface/Tinaja hybrids were manually stacked using the *pstacks* command. We then

assigned the morphotypic origin of each allele and confirmed heterozygosity in the F_1

samples using the *cstacks* command. Finally, we used this catalog to determine the

genotypes at each locus in the F2 samples with the *sstacks* and *genotypes* command. This

genotype database was formatted for use in R/qtl (61)

Linkage map. Using R/qtl, we selected for markers that were homozygous and had opposite

genotypes in cavefish versus surface fish (based on three surface and three cave

individuals). A linkage map was constructed from these loci using only the F2 population.

All markers that were genotyped in less than 180 individuals were omitted, as well as all

individuals that had poor marker genotyping (<1500 markers). Markers that did not

conform to the expected allele segregation ratio of 1:2:1 were also omitted ($p < 1e^{-10}$).

These methods produced a map with 1839 markers, 219 individuals, and 29 linkage

groups. Unlinked markers and small linkage groups (<10 markers) were omitted until our

map consisted of the optimal 25 linkage groups (*Astyanax mexicanus* has 2n = 50

chromosomes (62)). Each linkage group had 20-135 markers and markers were reordered

by default within the linkage groups, producing a total map length of > 5000 cM. Large gaps

in the map were eliminated by manually switching marker order to the best possible order

within each linkage group (error.prob = 0.005). The final linkage map consisted of 1800

markers, 219 individuals and 25 linkage groups, spanning 2871 cM. Maximum spacing was

32.1 cM and average spacing was 1.6 cM.

QTL scan using R/qtl. Genome wide logarithm of the odds (LOD) scores were calculated

using a single-QTL normal model and Haley-Knott regression. We assessed statistical

significance of LOD scores by calculating the 95th percentile of genome-wide maximum

penalized LOD score using 1000 random permutations (scanone). We estimated the

confidence intervals for identified QTL as the 1.5-LOD support interval expanded to the

nearest genotyped marker (lodint).

Population genomic metrics and analysis

We performed the following measures with GATK-processed data, including a core

set of samples analyzed in previous studies (18, 28) which contained: Pachón, N = 10 (9

newly re-sequenced plus the reference reads mapped back to the reference genome);

Tinaja N =10; Molino N = 9; Rascón N = 8; and Río Choy N = 9 and required six or more

individuals have data for a particular site. Details of sequencing and sample processing are

in (18, 28). We used VCFtools v0.1.13 (63) to calculate π, F_{ST} and d_{XY} and custom python

scripts to calculate these metrics on a per gene basis. We identified the allele counts per

population with VCFtools and used these for subsequent d_{XY} and fixed differences (D$_F$)

calculations. We used hapFLK v1.3 https://forge-dga.jouy.inra.fr/projects/hapflk (64) for

genome-wide estimation of the hapFLK statistic of across all 44 *Astyanax mexicanus*

samples and two *Astyanax aeneus* samples. For all of these metrics, we only used sites that

contained six or more individuals per population and calculated the metric or included the

p-values (in the case of hapFLK) for only the coding region of the gene. Candidate genes

were included if (a) the comparison between Río Choy and Tinaja resulted in a maximum

d_{XY} = 1, suggesting that there was at least one fixed difference between Río Choy and Tinaja

and (b) HapFLK resulted in at least one p value less than 0.05, suggesting that the

haplotype surrounding the gene exhibited evidence for non-neutral evolution.

RNA sequencing

RNA extraction and cDNA synthesis. Adult *A. mexicanus* were euthanized in 1400ppm

Tricane and the hindgut was immediately removed and homogenized in 0.3mL Trizol using

a motorized pellet pestle, and then stored at -80°C. Total RNA was extracted using Zymo

Research Direct-zol RNA MicroPrep with DNAse treatment according to the manufacturers

protocol. LunaScript RT supermix kit with 1μg of RNA was used to synthesize cDNA. All

samples were processed on the same day. Diluted cDNA samples (50ng/μL) were used for

sequencing.

RNA sequencing and differential gene expression analysis. HiSeq Illumina sequencing was

performed by Novogene (Chula Vista, CA). All samples were indexed and run as pools,

providing an estimated 20-30 million single-end reads per sample. Samples were

processed using an RNA-seq pipeline implemented in the bcbio-nextgen project

(https://bcbio-nextgen.readthedocs.org/en/latest/). Raw reads were examined for quality

issues using FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) to

ensure library generation and sequencing are suitable for further analysis.

Reads were aligned to Ensembl build 2 of the *Astyanax mexicanus* genome, augmented with

transcript information from Ensembl release 2.0.97 using STAR (65) with soft trimming

enabled to remove adapter sequences, other contaminant sequences such as polyA tails

and low quality sequences. Alignments were checked for evenness of coverage, rRNA

content, genomic context of alignments and complexity using a combination of FastQC,

Qualimap (66), MultiQC (https://github.com/ewels/MultiQC), and custom tools. Counts of

reads aligning to known genes were generated by featureCounts (67) and used for further

QC with the R package bcbioRNASeq [Steinbaugh MJ, Pantano L, Kirchner RD et al.

bcbioRNASeq: R package for bcbio RNA-seq analysis [version 2; peer review: 1 approved, 1

approved with reservations]. F1000Research 2018, 6:1976

(https://doi.org/10.12688/f1000research.12093.2)]. In parallel, Transcripts Per Million

(TPM) measurements per isoform were generated by quasialignment using Salmon (68)

for downstream differential expression analysis as quantitating at the isoform level has

been shown to produce more accurate results at the gene level (69). Salmon output was

imported into R using tximport (70) and differential gene expression analysis and data

visualization was performed using R with the DEseq 2 package (57, 71).

Acknowledgments

Brian Martineu and Megan Peavey for fish husbandry. This work was supported by grants

from the National Institutes of Health [HD089934, DK108495].

References

1.   H. Gehart, H. Clevers, Tales from the crypt: new insights into intestinal stem cells.

Nat. Rev. Gastroenterol. Hepatol. **16**, 19–34 (2019).

2.   S. Beyaz, *et al.*, High-fat diet enhances stemness and tumorigenicity of intestinal

progenitors. *Nature* **531**, 53–58 (2016).

3.   C.-W. Cheng, *et al.*, Ketone Body Signaling Mediates Intestinal Stem Cell Homeostasis

and Adaptation to Diet. *Cell* **178**, 1115-1131.e15 (2019).

4.   C. A. Lindemans, *et al.*, Interleukin-22 promotes intestinal-stem-cell-mediated

epithelial regeneration. *Nature* **528**, 560–564 (2015).

5.   M. Biton, *et al.*, T Helper Cell Cytokines Modulate Intestinal Stem Cell Renewal and

Differentiation. *Cell* **175**, 1307-1320.e22 (2018).

6.   D. M. Smith, R. C. Grasty, N. A. Theodosiou, C. J. Tabin, N. M. Nascone-Yoder,

Evolutionary relationships between the amphibian, avian, and mammalian stomachs.

Evol. Dev. **2**, 348–359 (2000).

7.   K. D. Walton, D. Mishkind, M. R. Riddle, C. J. Tabin, D. L. Gumucio, Blueprint for an

intestinal villus: Species-specific assembly required. *Wiley Interdiscip. Rev. Dev. Biol.*

7, e317 (2018).

8.   N. A. Theodosiou, E. Oppong, 3D morphological analysis of spiral intestine

morphogenesis in the little skate, <scp> *Leucoraja erinacea* </scp>. *Dev. Dyn.* **248**,

688–701 (2019).

9.   S. Takashima, D. Gold, V. Hartenstein, Stem cells and lineages of the intestine: a

developmental and evolutionary perspective. *Dev. Genes Evol.* **223**, 85–102 (2013).

10.   I. Miguel-Aliaga, H. Jasper, B. Lemaitre, Anatomy and Physiology of the Digestive

Tract of Drosophila melanogaster. *Genetics* **210**, 357–396 (2018).

11.   N. Aghaallaei, *et al.*, Identification, visualization and clonal analysis of intestinal stem

cells in fish. *Development* **143**, 3470–3480 (2016).

12.   S. C. Leigh, B.-Q. Nguyen-Phuc, D. P. German, The effects of protein and fiber content

on gut structure and function in zebrafish (Danio rerio). *J. Comp. Physiol. B* **188**, 237–

253 (2018).

13.   S. M. Secor, Digestive physiology of the Burmese python: broad regulation of

integrated performance. *J. Exp. Biol.* **211**, 3767–3774 (2008).

14.   S. R. McWilliams, W. H. Karasov, Phenotypic flexibility in digestive system structure

and function in migratory birds and its ecological significance. *Comp. Biochem.*

Physiol. Part A Mol. Integr. Physiol. **128**, 577–591 (2001).

15.   A. L. Andrew, *et al.*, Growth and stress response mechanisms underlying post-feeding

regenerative organ growth in the Burmese python. *BMC Genomics* **18**, 338 (2017).

16.   L. Espinasa, *et al.*, Contrasting feeding habits of post-larval and adult Astyanax

cavefish. *Subterr. Biol.* **21**, 1–17 (2017).

17.   G. Spicher, [Cytochrome absorption spectra of bacteria as aid for solving taxonomic

problems. 1. Elaboration and description of a method for measuring the redox and

carbon monoxide difference spectra of suspensions of intact live bacterial cells

(author's transl)]. *Zentralbl. Bakteriol. Orig. A.* **226**, 524–40 (1974).

18.   A. Herman, *et al.*, The role of gene flow in rapid and repeated evolution of cave-

related traits in Mexican tetra, Astyanax mexicanus. *Mol. Ecol.* **27**, 4397–4416 (2018).

19. M. E. Protas, *et al.*, Genetic analysis of cavefish reveals molecular convergence in the
evolution of albinism. *Nat. Genet.* **38**, 107–111 (2006).

20. S. E. McGaugh, *et al.*, The cavefish genome reveals candidate genes for eye loss. *Nat.*
Commun. **5** (2014).

21. A. K. Powers, E. M. Davis, S. A. Kaplan, J. B. Gross, Cranial asymmetry arises later in
the life history of the blind Mexican cavefish, Astyanax mexicanus. *PLoS One* **12**
(2017).

22. H. Hinaux, *et al.*, Sensory evolution in blind cavefish is driven by early embryonic
events during gastrulation and neurulation. *Development* **143**, 4521–4532 (2016).

23. J. E. Kowalko, *et al.*, Loss of schooling behavior in cavefish through sight-dependent
and sight-independent mechanisms. *Curr. Biol.* **23**, 1874–1883 (2013).

24. M. Yoshizawa, *et al.*, Distinct genetic architecture underlies the emergence of sleep
loss and prey-seeking behavior in the Mexican cavefish. *BMC Biol.* **13**, 15 (2015).

25. J. Jaggard, *et al.*, The lateral line confers evolutionarily derived sleep loss in the
Mexican cavefish. *J. Exp. Biol.* **220**, 284–293 (2017).

26. D. Moran, R. Softley, E. J. Warrant, Eyeless Mexican cavefish save energy by
eliminating the circadian rhythm in metabolism. *PLoS One* **9**, e107877 (2014).

27. A. C. Aspiras, N. Rohner, B. Martineau, R. L. Borowsky, C. J. Tabin, Melanocortin 4
receptor mutations contribute to the adaptation of cavefish to nutrient-poor
conditions. *Proc. Natl. Acad. Sci.* **112**, 9668–9673 (2015).

28. M. R. Riddle, *et al.*, Insulin resistance in cavefish as an adaptation to a nutrient-
limited environment. *Nature* **555**, 647–651 (2018).

29. M. R. Riddle, W. Boesmans, O. Caballero, Y. Kazwiny, C. J. Tabin, Morphogenesis and
motility of the Astyanax mexicanus gastrointestinal tract. *Dev. Biol.* **441**, 285–296
(2018).

30. M. R. Riddle, *et al.*, Genetic architecture underlying changes in carotenoid
accumulation during the evolution of the Blind Mexican cavefish,
Astyanax mexicanus *bioRxiv*, 788844 (2019).

31. E. S. Demitrack, L. C. Samuelson, Notch regulation of gastrointestinal stem cells. *J.*
Physiol. **594**, 4791–4803 (2016).

32.   S. Xiong, J. Krishnan, R. Peuß, N. Rohner, Early adipogenesis contributes to excess fat
accumulation in cave populations of Astyanax mexicanus. *Dev. Biol.* **441**, 297–304
(2018).
33.   R. K. Buddington, J. M. Diamond, Aristotle revisited: the function of pyloric caeca in
fish. *Proc. Natl. Acad. Sci.* **83**, 8012–8014 (1986).
34.   N. A. Ballesteros, *et al.*, The Pyloric Caeca Area Is a Major Site for IgM+ and IgT+ B Cell
Recruitment in Response to Oral Vaccination in Rainbow Trout. *PLoS One* **8**, e66118
(2013).
35.   L. D. Townsend, Variation in the Number of Pyloric Caeca and Other Numerical
Characters in Chinook Salmon and in Trout. *Copeia* **1944**, 52 (1944).
36.   K. E. O'Quin, M. Yoshizawa, P. Doshi, W. R. Jeffery, Quantitative Genetic Analysis of
Retinal Degeneration in the Blind Cavefish Astyanax mexicanus. *PLoS One* **8**, e57281
(2013).
37.   J. A. Birchler, H. Yao, S. Chudalayandi, Unraveling the genetic basis of hybrid vigor.
Proc. Natl. Acad. Sci. **103**, 12957–12958 (2006).
38.   B. B. McConnell, *et al.*, Krüppel-Like Factor 5 Is Important for Maintenance of Crypt
Architecture and Barrier Function in Mouse Intestine. *Gastroenterology* **141**, 1302-
1313.e6 (2011).
39.   T. Nakaya, *et al.*, KLF5 Regulates the Integrity and Oncogenicity of Intestinal Stem
Cells. *Cancer Res.* **74**, 2882–2891 (2014).
40.   X. Zhang, *et al.*, Somatic Superenhancer Duplications and Hotspot Mutations Lead to
Oncogenic Activation of the KLF5 Transcription Factor. *Cancer Discov.* **8**, 108–125
(2018).
41.   T. Ishitani, K. Matsumoto, A. B. Chitnis, M. Itoh, Nrarp functions to modulate neural-
crest-cell differentiation by regulating LEF1 protein stability. *Nat. Cell Biol.* **7**, 1106–
1112 (2005).
42.   T. J. Yun, M. J. Bevan, Notch-Regulated Ankyrin-Repeat Protein Inhibits Notch1
Signaling: Multiple Notch1 Signaling Pathways Involved In T Cell Development. *J.
Immunol.* **170**, 5834–5841 (2003).
43.   P. Ornelas-García, S. Pajares, V. M. Sosa-Jiménez, S. Rétaux, R. A. Miranda-Gamboa,
Microbiome differences between river-dwelling and cave-adapted populations of the

708   fish Astyanax mexicanus (De Filippi, 1853). *PeerJ* **6**, e5906 (2018).

709   44.   J. Laskowski, J. M. Thurman, "Chapter 14 - Factor B" in *The Complement FactsBook*

710   *(Second Edition)*, Factsbook., Second Edi, S. Barnum, T. Schein, Eds. (Academic Press,

711   2018), pp. 135–146.

712   45.   Andoh, *et al.*, Detection of complement C3 and factor B gene expression in normal

713   colorectal mucosa, adenomas and carcinomas. *Clin. Exp. Immunol.* **111**, 477–483

714   (1998).

715   46.   R. Peuß, *et al.*, Single cell analysis reveals modified hematopoietic cell composition

716   affecting inflammatory and immunopathological responses in Astyanax

717   mexicanus *bioRxiv*, 647255 (2019).

718   47.   M. R. Riddle, *et al.*, Genetic architecture underlying changes in carotenoid

719   accumulation during the evolution of the Blind Mexican cavefish,

720   Astyanax mexicanus *bioRxiv*, 788844 (2019).

721   48.   Mezquita, Mezquita, Two Opposing Faces of Retinoic Acid: Induction of Stemness or

722   Induction of Differentiation Depending on Cell-Type. *Biomolecules* **9**, 567 (2019).

723   49.   P. Czarnewski, S. Das, S. Parigi, E. Villablanca, Retinoic Acid and Its Role in

724   Modulating Intestinal Innate Immunity. *Nutrients* **9**, 68 (2017).

725   50.   P. Karpowicz, Y. Zhang, J. B. Hogenesch, P. Emery, N. Perrimon, The Circadian Clock

726   Gates the Intestinal Stem Cell Regenerative State. *Cell Rep.* **3**, 996–1004 (2013).

727   51.   K. Parasram, P. Karpowicz, Time after time: circadian clock regulation of intestinal

728   stem cells. *Cell. Mol. Life Sci.* (2019) https:/doi.org/10.1007/s00018-019-03323-x.

729   52.   E. Peyric, H. A. Moore, D. Whitmore, Circadian Clock Regulation of the Cell Cycle in

730   the Zebrafish Intestine. *PLoS One* **8**, e73209 (2013).

731   53.   A. Beale, *et al.*, Circadian rhythms in Mexican blind cavefish Astyanax mexicanus in

732   the lab and in the field. *Nat. Commun.* **4**, 2769 (2013).

733   54.   I. A. Frøland Steindal, A. D. Beale, Y. Yamamoto, D. Whitmore, Development of the

734   Astyanax mexicanus circadian clock and non-visual light responses. *Dev. Biol.* **441**,

735   345–354 (2018).

736   55.   E. R. Duboué, A. C. Keene, R. L. Borowsky, Evolutionary convergence on sleep loss in

737   cavefish populations. *Curr. Biol.* **21**, 671–676 (2011).

56.   Y. Elipot, L. Legendre, S. Père, F. Sohm, S. Rétaux, *Astyanax* Transgenesis and

Husbandry: How Cavefish Enters the Laboratory. *Zebrafish* **11**, 291–299 (2014).

57.   R-core-team, R: A Language and Environment for Statistical Computing (2019).

58.   C. T. Rueden, *et al.*, ImageJ2: ImageJ for the next generation of scientific image data.
BMC Bioinformatics **18** (2017).

59.   J. M. Catchen, A. Amores, P. Hohenlohe, W. Cresko, J. H. Postlethwait, Stacks: building
and genotyping Loci de novo from short-read sequences. *G3 (Bethesda).* **1**, 171–82
(2011).

60.   B. Langmead, S. L. Salzberg, Fast gapped-read alignment with Bowtie 2. *Nat. Methods*
9, 357–9 (2012).

61.   D. Arends, P. Prins, R. C. Jansen, K. W. Broman, R/qtl: high-throughput multiple QTL
mapping: Fig. 1. *Bioinformatics* **26**, 2990–2992 (2010).

62.   K. F. Kavalco, L. F. De Almeida-Toledo, Molecular Cytogenetics of Blind Mexican Tetra
and Comments on the Karyotypic Characteristics of Genus Astyanax (Teleostei,
Characidae). *Zebrafish* **4**, 103–111 (2007).

63.   P. Danecek, *et al.*, The variant call format and VCFtools. *Bioinformatics* **27**, 2156–
2158 (2011).

64.   M. I. Fariello, S. Boitard, H. Naya, M. SanCristobal, B. Servin, Detecting Signatures of
Selection Through Haplotype Differentiation Among Hierarchically Structured
Populations. *Genetics* **193**, 929–941 (2013).

65.   A. Dobin, *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

66.   F. García-Alcalde, *et al.*, Qualimap: evaluating next-generation sequencing alignment
data. *Bioinformatics* **28**, 2678–2679 (2012).

67.   Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

68.   N. Bray, H. Pimentel, P. Melsted, L. Pachter, Near-optimal RNA-Seq quantification.
bioRxiv (2015) https:/doi.org/10.1101/021592.

69.   C. Soneson, M. I. Love, M. D. Robinson, Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. *F1000Research* **4**, 1521 (2016).

70.   C. Soneson, M. I. Love, M. D. Robinson, Differential analyses for RNA-seq: transcript-
level estimates improve gene-level inferences. *F1000Research* **4**, 1521 (2015).

71.   M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. *Genome Biol.* **15**, 550 (2014).

Tables

Table 1. Location of QTL markers defining confidence internal for relative hindgut length

marker	LOD	LG: position	Scaffold: position	Chromosome: position
r70838	1.64	10: 61.11	KB871783: 144963	14: 27137229
r172341	4.82	10: 68.28	KB882083: 490666	14: 17061465
r171715	2.82	10: 70.29	KB882083: 2674927	14: 11724563

Genome-wide maximum penalized LOD score using 1000 random permutations, position on the linkage group (LG), Pachón genome scaffold, and surface fish chromosome.

Table 2. Population metrics for genes in the hindgut QTL that exhibit evidence of non-neutral evolution comparing Río Choy with Tinaja

| | | Compared to Río Choy (n=9) | | | | | | Compared to Rascón (n=8) | | | | | |
| | | Tinaja (n=10) | | Pachón (n=10) | | Molino (n=9) | | Tinaja (n=10) | | Pachón (n=10) | | Molino (n=9) | |
gene ID	gene name, description	fixed/total alleles used	Dxy (max 21837)	fixed/total alleles used	Dxy (max 21457)	fixed/total alleles used	Dxy (Max 20517)	fixed/total alleles used	Dxy (max 20328)	fixed/total alleles used	Dxy (Max 19682)	fixed/total alleles used	Dxy (Max 18007)
ENSAMXG00000007196	cfb1, complement activation	2/66	20616	2/65	20343	0/35	19153	**3/27**	**19158**	**6/29**	**18552**	**18/41**	**16986**
ENSAMXG00000007167	cfb2, complement activation	**6/40**	**20301**	**1/34**	**17327**	0/26	13357	**1/27**	**18979**	**3/28**	**15532**	6/31	15620
ENSAMXG00000007054	rbmx2, RNA binding motif protein, X-linked 2	**1/8**	19816	0/7	18688	**2/7**	**18804**	NA	231	**1/1**	**5980**	**4/4**	**15183**
ENSAMXG00000007360	prdx5, peroxiredoxin 5	**2/66**	19746	**2/6**	**19516**	0/3	4557	NA	NA	NA	NA	3/3	15431
ENSAMXG00000007379	adssl, adenylosuccinate synthase like	**4/15**	**18587**	**5/14**	**18455**	0/9	2290	0/11	10256	**1/11**	**9232**	5/16	15065
ENSAMXG00000007093	klf5l, Kruppel-like factor 5-like	**2/8**	**17219**	0/6	3676	0/4	3755	0/6	15383	0/6	7873	0/5	4384
ENSAMXG00000007382	rce1b, Ras converting CAAX endopeptidase 1b	**1/11**	**16658**	3/12	18226	1/4	14767	0/10	15380	**1/9**	**15944**	**2/7**	**12419**
ENSAMXG00000025173	RNA-binding protein lark-like	**2/6**	**12077**	**3/6**	**15675**	**2/5**	**11921**	0/2	931	**1/2**	**4971**	**4/5**	**14059**
ENSAMXG00000007446	TENM2, teneurin transmembrane protein 2	**4/46**	**10278**	**5/44**	**10039**	**8/30**	**11529**	0/14	2628	**1/15**	**2195**	**8/21**	**3533**
ENSAMXG00000007073	slc25a14, solute carrier family	**1/5**	**10190**	0/4	6696	**0/3**	**4454**	0/3	3595	**2/5**	**15257**	1/4	7264

For each gene, the number of alleles that have fixed differences out of the total sites used, and divergence (Dxy) between the indicated surface and cave population is shown. Values in bold indicate that the genome-wide estimation of the hapFLK statistic resulted in at least one p-value less than 0.05.

Table S1. Genes within the hindgut QTL 1.5-LOD support interval that are differentially expressed in the hindgut

gene stable ID	gene name, description	Average normalized expression count			lr	pw
		Tinaja	surface	hybrid		
ENSAMXG00000006912	ubiquitin interaction motif containing 1	**641.226**	298.488	525.200	*	*
ENSAMXG00000009311	h2afy, H2A histone family member Y	**433.354**	252.691	387.330	*	*
ENSAMXG00000007120	amot, Angiomotin	**396.543**	168.257	269.468	*	*
ENSAMXG00000014597	notch1a	**106.556**	54.248	<u>40.240</u>	*	
ENSAMXG00000037919	plac8l1, PLAC8-like 1	**104.402**	44.104	70.487	*	*
ENSAMXG00000034188	NA	**93.520**	19.000	24.367	*	*
ENSAMXG00000018167	ERMP1, endoplasmic reticulum metallopeptidase 1	**77.677**	41.809	56.172	*	*
ENSAMXG00000019610	f2rl1.1, coagulation factor II (thrombin) receptor-like 1	**56.584**	14.651	31.024	*	*
ENSAMXG00000043097	NA	**30.063**	3.306	17.268	*	*
ENSAMXG00000040346	NA	**22.600**	0.968	1.520	*	
ENSAMXG00000030011	NA	**22.139**	0.220	8.351	*	*
ENSAMXG00000007196	cfb1, complement factor B-like	**60.433**	36.832	56.506		*
ENSAMXG00000009861	cingulin-like protein 1	**589.163**	1125.543	<u>461.148</u>		*
ENSAMXG00000006854	grk6, G protein-coupled receptor kinase 6	**66.554**	46.243	54.880		*
ENSAMXG00000043097	NA	**30.063**	3.306	17.268		*
ENSAMXG00000017243	ubtd2, ubiquitin domain containing 2	**27.857**	13.419	15.001		*
ENSAMXG00000040346	NA	**22.600**	0.968	1.520		*
ENSAMXG00000032351	NA	**0.869**	20.972	<u>0.418</u>		*
ENSAMXG00000000826	arf-GAP with Rho-GAP domain, ANK repeat and PH domain-containing	436.604	630.912	**738.131**	*	*
ENSAMXG00000037917	txnl1, thioredoxin-like 1	259.679	321.669	**382.788**	*	
ENSAMXG00000000836	reticulon, reticulon-3-A-like	3.765	174.850	**268.463**	*	*
ENSAMXG00000014252	chchd10, coiled-coil-helix-coiled-coil-helix domain containing 10	160.164	121.477	**231.895**	*	
ENSAMXG00000042179	cmc4, C-x(9)-C motif containing 4 homolog	26.142	**28.869**	48.263	*	
ENSAMXG00000043642	pgrmc1, progesterone receptor membrane component 1	1126.239	**2175.765**	1955.909	*	*
ENSAMXG00000014522	lrrc8aa, leucine rich repeat containing 8 VRAC subunit Aa	794.214	**1434.929**	<u>1221.255</u>	*	*
ENSAMXG00000009861	cingulin-like protein 1	589.163	**1125.543**	<u>461.148</u>	*	
ENSAMXG00000031422	thioredoxin domain containing 15	348.453	**541.287**	423.549	*	*
ENSAMXG00000033220	cox7b, cytochrome c oxidase subunit 7B	280.756	**415.961**	396.242	*	*
ENSAMXG00000041114	NA	24.135	**414.594**	48.999	*	*
ENSAMXG00000040749	bada, BCL2 associated agonist of cell death a	190.098	**317.216**	305.060	*	*
ENSAMXG00000037990	NA	38.642	**237.678**	230.671	*	*
ENSAMXG00000017290	rars, arginyl-tRNA synthetase	190.139	**236.984**	<u>153.839</u>	*	
ENSAMXG00000031420	sosondowah ankyrin repeat domain family member B	60.483	**179.482**	114.594	*	*
ENSAMXG00000034277	NLR family CARD domain-containing protein 3-like	32.671	**167.225**	48.911	*	*
ENSAMXG00000008478	kif20a, kinesin family member 20A	28.861	**146.302**	69.886	*	*
ENSAMXG00000000841	GALNT10, polypeptide N-acetylgalactosaminyltransferase 10	91.798	**138.911**	103.782	*	*
ENSAMXG00000007080	pof1b, actin binding protein	86.944	**133.568**	<u>59.637</u>	*	
ENSAMXG00000007243	rin1a, Ras and Rab interactor 1a	85.462	**129.130**	92.142		*
ENSAMXG00000018210	TMEM38B, transmembrane protein 38B	1718.867	**3347.577**	3040.203		*
ENSAMXG00000041114	NA	24.135	**414.594**	48.999		*

For each gene, values in bold are highest expression across sample types. Underlined values indicate the lowest expression in the F1 hybrid samples. Asterisks indicate significant difference by likelihood ratio test between all three sample types (lr) and pairwise comparison between surface fish and Tinaja cavefish (pw).

Table S2. Differentially expressed hindgut genes exhibiting highest expression in the F1 hybrid (Group 3, Figure S1)

gene stable ID	gene name	description
ENSAMXG00000000542	gys1	glycogen synthase 1 (muscle) [Source:ZFIN;Acc:ZDB-GENE-040426-1243]
ENSAMXG00000003217		NULL
ENSAMXG00000003528	ITLN1 (1 of many)	intelectin 1 [Source:HGNC Symbol;Acc:HGNC:18259]
ENSAMXG00000005979	timeless	timeless circadian clock [Source:ZFIN;Acc:ZDB-GENE-030131-8978]
ENSAMXG00000006267	esr1	estrogen receptor 1 [Source:ZFIN;Acc:ZDB-GENE-020806-5]
ENSAMXG00000006712	caskin1	CASK interacting protein 1 [Source:ZFIN;Acc:ZDB-GENE-071115-3]
ENSAMXG00000008593	pir	pirin [Source:ZFIN;Acc:ZDB-GENE-040718-288]
ENSAMXG00000009578	adcy8	adenylate cyclase 8 (brain) [Source:ZFIN;Acc:ZDB-GENE-070912-197]
ENSAMXG00000011826	si:ch211-132f19.7	si:ch211-132f19.7 [Source:ZFIN;Acc:ZDB-GENE-130530-619]
ENSAMXG00000011960	cyp26a1	cytochrome P450, family 26, subfamily A, polypeptide 1 [Source:ZFIN;Acc:ZDB-GENE-990415-44]
ENSAMXG00000012719		heat shock 70 kDa protein 12A-like [Source:NCBI gene;Acc:103046531]
ENSAMXG00000013859	glsa	glutaminase a [Source:ZFIN;Acc:ZDB-GENE-050204-3]
ENSAMXG00000014779	sgsm3	small G protein signaling modulator 3 [Source:ZFIN;Acc:ZDB-GENE-040426-2635]
ENSAMXG00000014901	apeh	acylaminoacyl-peptide hydrolase [Source:ZFIN;Acc:ZDB-GENE-030131-9870]
ENSAMXG00000015029	rorcb	RAR-related orphan receptor C b [Source:ZFIN;Acc:ZDB-GENE-040724-215]
ENSAMXG00000018475		NULL
ENSAMXG00000018995		peroxiredoxin-1 [Source:NCBI gene;Acc:103031813]
ENSAMXG00000019373	nudcd3	NudC domain containing 3 [Source:ZFIN;Acc:ZDB-GENE-040426-2255]
ENSAMXG00000019396	cygb1	cytoglobin 1 [Source:ZFIN;Acc:ZDB-GENE-020513-1]
ENSAMXG00000020744	c8g	complement component 8, gamma polypeptide [Source:ZFIN;Acc:ZDB-GENE-040426-1898]
ENSAMXG00000021464	si:ch211-276a23.5	si:ch211-276a23.5 [Source:ZFIN;Acc:ZDB-GENE-141215-4]
ENSAMXG00000024777	eno1a	enolase 1a, (alpha) [Source:ZFIN;Acc:ZDB-GENE-030131-6048]
ENSAMXG00000024896	nrarpa	NOTCH regulated ankyrin repeat protein a [Source:ZFIN;Acc:ZDB-GENE-030515-6]
ENSAMXG00000025981		NULL
ENSAMXG00000027124	RF00478	NULL
ENSAMXG00000029674	trh	thyrotropin-releasing hormone [Source:ZFIN;Acc:ZDB-GENE-020930-1]
ENSAMXG00000030396	mb	myoglobin [Source:ZFIN;Acc:ZDB-GENE-040426-1430]
ENSAMXG00000031413		solute carrier family 22 member 4-like [Source:NCBI gene;Acc:103029834]
ENSAMXG00000031569	vrk3	vaccinia related kinase 3 [Source:ZFIN;Acc:ZDB-GENE-030131-246]
ENSAMXG00000031818	zgc:92606	zgc:92606 [Source:ZFIN;Acc:ZDB-GENE-040718-462]
ENSAMXG00000034120	slc29a3	solute carrier family 29 (equilibrative nucleoside transporter), member 3 [Source:ZFIN;Acc:ZDB-GENE-081107-66]
ENSAMXG00000034331		NULL
ENSAMXG00000035277	zdhhc11	zinc finger, DHHC-type containing 11 [Source:ZFIN;Acc:ZDB-GENE-130530-795]
ENSAMXG00000035722		NULL
ENSAMXG00000036341	hexb	hexosaminidase B (beta polypeptide) [Source:ZFIN;Acc:ZDB-GENE-030131-2333]
ENSAMXG00000036770	fhdc4	FH2 domain containing 4 [Source:ZFIN;Acc:ZDB-GENE-131127-25]
ENSAMXG00000037811	SECISBP2L	SECIS binding protein 2 like [Source:HGNC Symbol;Acc:HGNC:28997]
ENSAMXG00000038275		NULL
ENSAMXG00000038302		NULL
ENSAMXG00000038552	CASQ2	calsequestrin 2 [Source:HGNC Symbol;Acc:HGNC:1513]
ENSAMXG00000038579		NULL
ENSAMXG00000039496		NULL
ENSAMXG00000040338		intelectin-like [Source:NCBI gene;Acc:103036009]
ENSAMXG00000041228		NULL
ENSAMXG00000041671	wipi1	WD repeat domain, phosphoinositide interacting 1 [Source:ZFIN;Acc:ZDB-GENE-040426-1404]
ENSAMXG00000041918	guca1c	guanylate cyclase activator 1C [Source:ZFIN;Acc:ZDB-GENE-030829-1]
ENSAMXG00000041920	slc1a2b	solute carrier family 1 (glial high affinity glutamate transporter), member 2b [Source:ZFIN;Acc:ZDB-GENE-030131-7779]
ENSAMXG00000042050		NULL
ENSAMXG00000042163		NULL
ENSAMXG00000042450	ankrd46a	ankyrin repeat domain 46a [Source:ZFIN;Acc:ZDB-GENE-030131-2643]
ENSAMXG00000042617	plpp1a	phospholipid phosphatase 1a [Source:ZFIN;Acc:ZDB-GENE-080225-26]
ENSAMXG00000042843	rdh1	retinol dehydrogenase 1 [Source:ZFIN;Acc:ZDB-GENE-030912-15]
ENSAMXG00000042921	PSMB11 (1 of many)	proteasome subunit beta 11 [Source:HGNC Symbol;Acc:HGNC:31963]
ENSAMXG00000043019		NULL
ENSAMXG00000044068		NULL

Table S2. Differentially expressed hindgut genes exhibiting lowest expression in the F1 hybrid (Group 4, Figure S1)

gene stable ID	gene name	description
ENSAMXG00000001449	mat2al	methionine adenosyltransferase II, alpha-like [Source:ZFIN;Acc:ZDB-GENE-060421-5255]
ENSAMXG00000002813		NULL
ENSAMXG00000004203	dock3	dedicator of cytokinesis 3 [Source:ZFIN;Acc:ZDB-GENE-100624-1]
ENSAMXG00000004229	gdpd1	glycerophosphodiester phosphodiesterase domain containing 1 [Source:ZFIN;Acc:ZDB-GENE-040822-22]
ENSAMXG00000004732	bag3	BCL2 associated athanogene 3 [Source:ZFIN;Acc:ZDB-GENE-040801-40]
ENSAMXG00000005293	evplb	envoplakin b [Source:ZFIN;Acc:ZDB-GENE-030131-212]
ENSAMXG00000007764	krt1-c5	keratin, type 1, gene c5 [Source:ZFIN;Acc:ZDB-GENE-060316-3]
ENSAMXG00000009861		cingulin-like protein 1 [Source:NCBI gene;Acc:103022577]
ENSAMXG00000010263	CASP6	caspase 6 [Source:HGNC Symbol;Acc:HGNC:1507]
ENSAMXG00000010610	ppl	periplakin [Source:ZFIN;Acc:ZDB-GENE-041008-196]
ENSAMXG00000011397	spire1a	spire-type actin nucleation factor 1a [Source:ZFIN;Acc:ZDB-GENE-061013-119]
ENSAMXG00000011672	slc9a3r1a	solute carrier family 9, subfamily A (NHE3, cation proton antiporter 3), member 3 regulator 1a [Source:ZFIN;Acc:ZDB-GENE-031006-7]
ENSAMXG00000011790	zgc:172079	zgc:172079 [Source:ZFIN;Acc:ZDB-GENE-080204-103]
ENSAMXG00000012301	slc4a4a	solute carrier family 4 (sodium bicarbonate cotransporter), member 4a [Source:ZFIN;Acc:ZDB-GENE-060526-274]
ENSAMXG00000013027	brca2	breast cancer 2, early onset [Source:ZFIN;Acc:ZDB-GENE-060510-3]
ENSAMXG00000013552	pabpc1l	poly(A) binding protein, cytoplasmic 1-like [Source:ZFIN;Acc:ZDB-GENE-030131-5836]
ENSAMXG00000013796	pak1	p21 protein (Cdc42/Rac)-activated kinase 1 [Source:ZFIN;Acc:ZDB-GENE-030826-29]
ENSAMXG00000015091		NULL
ENSAMXG00000015951	eps8l2	EPS8 like 2 [Source:ZFIN;Acc:ZDB-GENE-131121-340]
ENSAMXG00000017060	galnt7	UDP-N-acetyl-alpha-D-galactosamine: polypeptide N-acetylgalactosaminyltransferase 7 [Source:ZFIN;Acc:ZDB-GENE-050522-284]
ENSAMXG00000017336	slc5a1 (1 of many)	solute carrier family 5 (sodium/glucose cotransporter), member 1 [Source:ZFIN;Acc:ZDB-GENE-040426-1524]
ENSAMXG00000017464	lmo7a	LIM domain 7a [Source:ZFIN;Acc:ZDB-GENE-030219-74]
ENSAMXG00000017536		polypeptide N-acetylgalactosaminyltransferase 6-like [Source:NCBI gene;Acc:103034470]
ENSAMXG00000017853		transmembrane protease serine 9-like [Source:NCBI gene;Acc:103044070]
ENSAMXG00000018625	per1b	period circadian clock 1b [Source:ZFIN;Acc:ZDB-GENE-040419-1]
ENSAMXG00000021143	slc7a8a	solute carrier family 7 (amino acid transporter light chain, L system), member 8a [Source:ZFIN;Acc:ZDB-GENE-121120-2]
ENSAMXG00000025148	STX19	syntaxin 19 [Source:HGNC Symbol;Acc:HGNC:19300]
ENSAMXG00000029238	si:ch211-137i24.10	si:ch211-137i24.10 [Source:ZFIN;Acc:ZDB-GENE-060526-34]
ENSAMXG00000029616	ugt5d1 (1 of many)	UDP glucuronosyltransferase 5 family, polypeptide D1 [Source:ZFIN;Acc:ZDB-GENE-091118-36]
ENSAMXG00000029919	b3gnt3.4	UDP-GlcNAc:betaGal beta-1,3-N-acetylglucosaminyltransferase 3, tandem duplicate 4 [Source:ZFIN;Acc:ZDB-GENE-030131-8875]
ENSAMXG00000029933		annexin A2-like [Source:NCBI gene;Acc:103024979]
ENSAMXG00000030052	pcbd1	pterin-4 alpha-carbinolamine dehydratase/dimerization cofactor of hepatocyte nuclear factor 1 alpha [Source:ZFIN;Acc:ZDB-GENE-040426-1787]
ENSAMXG00000031773		NULL
ENSAMXG00000032507		zinc finger protein 800-like [Source:NCBI gene;Acc:103026180]
ENSAMXG00000033111		NULL
ENSAMXG00000033217		NULL
ENSAMXG00000033692		NULL
ENSAMXG00000034400		NULL
ENSAMXG00000036383		suppressor of cytokine signaling 2-like [Source:NCBI gene;Acc:111191435]
ENSAMXG00000036963		cerebellin-2-like [Source:NCBI gene;Acc:103040140]
ENSAMXG00000037136	zgc:103681	zgc:103681 [Source:ZFIN;Acc:ZDB-GENE-041010-17]
ENSAMXG00000037535		NULL
ENSAMXG00000038262		NULL
ENSAMXG00000040014	serpinb1l2	serpin peptidase inhibitor, clade B (ovalbumin), member 1, like 2 [Source:ZFIN;Acc:ZDB-GENE-041001-117]
ENSAMXG00000040633		NULL
ENSAMXG00000040957		NULL
ENSAMXG00000041754	rtn2a	reticulon 2a [Source:ZFIN;Acc:ZDB-GENE-060420-1]
ENSAMXG00000042112	bco1	beta-carotene oxygenase 1 [Source:ZFIN;Acc:ZDB-GENE-010508-1]
ENSAMXG00000042476	slc25a15b	solute carrier family 25 (mitochondrial carrier; ornithine transporter) member 15b [Source:ZFIN;Acc:ZDB-GENE-060526-35]
ENSAMXG00000043501		voltage-gated potassium channel subunit beta-2 [Source:NCBI gene;Acc:103043691]
ENSAMXG00000043650		NULL
ENSAMXG00000043711	pparab	peroxisome proliferator-activated receptor alpha b [Source:ZFIN;Acc:ZDB-GENE-990415-211]
ENSAMXG00000043975	slc1a1	solute carrier family 1 (neuronal/epithelial high affinity glutamate transporter, system Xag), member 1 [Source:ZFIN;Acc:ZDB-GENE-040718-414]

Figures and figure legends

Figure 1. Histology of the *Astyanax mexicanus* surface fish gastrointestinal tract as shown by hematoxylin and eosin staining. A, Transverse section of esophageal region (L: lumen, e: epithelium, m1-2: muscle layers). B, Stomach section showing epithelium (e), eosinophilic gastric glands (g), and layers of muscle (o: orthogonal, c: circumferential, l: longitudinal). C, Cross section of a pyloric caecum showing the lumen (L) and ridged epithelium (e). D, Transverse section of the proximal midgut showing lumen filled with mucus (L), epithelium (e), lamina propria (l), and circumferential inner and longitudinal outer muscle layers (m). E, Transverse section of a more posterior section of the midgut showing epithelium (e), lamina propria (l), and circumferential inner and longitudinal outer muscle layers (m). F, Transverse section of the hindgut showing epithelium (e), and muscle layers (m). A-F image magnification is 10X. D-F' 20X images (scale bar is 50µM) showing the cell types of the gut epithelium in the proximal midgut (D'), distal midgut (E'), and hindgut (F'). Black arrows highlight the differences between secretory goblet cells in the proximal (D') and distal (E') midgut. Black arrow head in E' indicates group of immune cells (pink). Red arrow head shows pyknotic nuclei. Grey arrow in F' highlights cells with hydrophobic (mostly clear) structures within the cytoplasm. D-E are images from the same slide.

Figure 2. Pyloric caeca structure and number in *Astyanax mexicanus* surface and cave morphs. A, Image of the region of gut containing pyloric caeca. The darker lines show the ridged morphology of the epithelium (black arrow, scale bar is 2 mm). B, Drawing of the pyloric caeca indicating the position and number of caeca. The caeca at position 7, 8, and 9 are variable. C, Graph comparing percentage of one-month old fish with the indicated number of total caeca for each population (n=5, 6, 8, 6).

Figure 3. Relative length of the gastrointestinal (GI) tract of *Astyanax mexicanus* surface and cave morphs. A, Images of fish from the indicated populations that were fed 6 mg of pellet food per day for greater than 8 months and their GI tracts (scale bars is 1cm). B, Boxplots showing relative midgut and hindgut length of fish from the indicated populations (n=6 for surface, Tinaja, and Pachón. n=5 for Molino). For box plots, median, 25th, 50th, and 75th percentiles are represented by horizontal bars and vertical bars represent 1.5 interquartile ranges. Asterisks indicates Pachón relative hindgut length is significantly less than surface (p=0.05), Tinaja (p=0.002), and Molino (p=0.001, one-way ANOVA with Turkey's post hoc test).

Figure 4. Diet influences *A. mexicanus* gut morphology. Relative length, circumference, and number of folds in the epithelium in the midgut (A-C) and hindgut (D-E) in fish continuously fed a moderate diet or switched to a high-nutrient or low nutrient diet as adults (n=3 fish per population and diet, and 3 tissue sections average per individual fish). In boxplots, median, 25th, 50th, and 75th percentiles are represented by horizontal bars and vertical bars represent 1.5 interquartile ranges. Significance code (*p<0.05) from one-way ANOVA with HSD post hoc test comparing between populations (black asterisks) or within population by diet (orange asterisks = Tinaja, blue asterisks = surface fish). Bar graphs show percent change in the indicated phenotype in the midgut (G) and hindgut (H).

Figure 5. Cavefish have more proliferation in the intestinal epithelium on a low-nutrient diet compared to surface fish and, reduce proliferation in response to starvation. Cross section of midgut in surface fish (A) and Tinaja cavefish (B) stained with H&E (left) and serial section showing DAPI staining and EdU positive cells; larger view of yellow boxed region shown in right panels. Yellow arrows point to EdU positive cells. Note yellow arrow near the tip of the fold shows cell in the lamina propria. C-F, Quantification of midgut length, circumference, number of folds, and EdU positive cells in fish fed *ad libitum* versus starved for two weeks. G-J, Quantification of hindgut length, circumference, number of folds, and EdU positive cells in fish fed *ad libitum* versus fasted for two weeks (n=3 fish per population and diet and 3 tissue sections averaged per individual fish). In boxplots, median, 25th, 50th, and 75th percentiles are represented by horizontal bars and vertical bars represent 1.5 interquartile ranges. Significance code (*p<0.05, **p<.005) from one-way ANOVA with HSD post hoc test. K-L, Bar graphs showing percent change in the indicated phenotypes in the midgut (K) and hindgut (L).

Figure 6. Genetic mapping of body condition and gut morphology in adult *A. mexicanus* surface/Tinaja F2 hybrids. A-E, Histograms showing distribution of the indicated phenotypes in F2 hybrids and (right) genome wide scan showing logarithm of the odds (LOD) score for the indicated phenotypes using Haley-Knott regression mapping. A, B non-parametric model, C-E normal model. Significance thresholds at p=0.05 are indicated in parenthesis following the phenotype.

Figure 7. Gut length heterosis in surface/Tinaja hybrids. A, Image of gut from surface fish, surface/Tinaja F1 hybrid, and Tinaja cavefish of similar lengths. B, Quantification of relative hindgut length in surface (n=6), F1 hybrid (n=10), Tinaja (n=6). C, Quantification of relative hindgut length of surface/Tinaja F2 hybrids with the indicated genotype at the marker with the highest LOD score (SS: homozygous surface, SC: heterozygous, CC: homozygous cave). D, Quantification of relative midgut length in in surface (n=6), F1 hybrid (n=10), Tinaja (n=6). E, Quantification of relative midgut length of surface/Tinaja F2 hybrids with the indicated genotype at the marker with the highest LOD score (SS: homozygous surface, SC: heterozygous CC: homozygous cave).

Figure 8. Population genetics and RNA sequencing analysis of candidate genes controlling hindgut length. A, B, Heat maps showing the genome-wide divergence percentile of genes within the hindgut QTL that have significant HapFLK p-values. A, Divergence of Río Choy surface fish from three cavefish populations, and B, Divergence of Rascón surface fish from three cavefish populations. C, Log2 Fold expression change compared to surface fish of a subset of genes within the hindgut QTL (blue), or that are a part of the gene cluster that shows lowest (group3, red) and highest (group4, green) expression in F1 hybrids. Grey significance codes from likelihood ratio test comparing all three sample types and black asterisks next to bars indicate significance from Wald test comparing to only surface fish (Significance code for adjusted p-value: *<.05 **<.005 ***<.0005, ns>.05). Error bars indicate standard error estimate for log2 fold change.

Supplemental Figure 1: Gene expression clusters of genes that are differentially expressed between Tinaja, Surface, and F1 hybrid hindguts (n=5 hindguts per population).