Thiamethoxam exposure deregulates short ORF gene expression in the

honey bee and compromises immune response to bacteria

Pâmela Decio[1*], Pinar Ustaoglu[2*], Kamila Derecka[3], Ian C. W. Hardy[3], Thaisa

C. Roat[1], Osmar Malaspina[1], Nigel Mongan[4], Reinhard Stöger[3, 5] and

Matthias Soller[2, 5]

[1]Universidade Estadual Paulista (UNESP), Instituto de Biociências, Campus

Rio Claro, São Paulo, Brazil.

[2]School of Biosciences, College of Life and Environmental Sciences, University

of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom

[3]School of Biosciences, University of Nottingham, LE12 5RD, United Kingdom

[4]School of School of Veterinary Medicine and Science, University of

Nottingham, LE12 5RD, United Kingdom

Running title: Thiamethoxam exposure, short ORFs and bee immunity

Key Words: Neonicotinoid, *Apis mellifera*, anti-microbial peptides,

[5]Corresponding authors: m.soller@bham.ac.uk,

reinhard.stoger@nottingham.ac.uk

*equally contributing authors

Abstract

Maximizing crop yields relies on the use of agrochemicals to control insect pests. One of the most widely used classes of insecticides are neonicotinoids that interfere with signalling of the neurotransmitter acetylcholine, but these can also disrupt crop-pollination services provided by bees. Here, we analysed whether chronic low dose long-term exposure to the neonicotinoid thiamethoxam alters gene expression and alternative splicing in brains of Africanized honey bees, *Apis mellifera,* as adaptation to altered neuronal signalling. We find differentially regulated genes that show concentration-dependent responses to thiamethoxam, but no changes in alternative splicing. Most differentially expressed genes have no annotated function but encode short Open Reading Frames (sORFs), a characteristic feature of anti-microbial peptides. As this suggested that immune responses may be compromised by thiamethoxam exposure, we tested the impact of thiamethoxam on bee immunity by injecting bacteria. We show that intrinsically sub-lethal thiamethoxam exposure makes bees more vulnerable to normally non-pathogenic bacteria. Our findings imply a synergistic mechanism for the observed bee population declines that concern agriculturists, conservation ecologists and the public.

INTRODUCTION

The western honey bee *Apis mellifera* is highly beneficial for human societies. Besides honey production, this semi-domesticated insect plays a critical role in sustaining global food security through the provision of managed pollination services that contribute to increased yields of many crops. Globally, crop productivity is also enhanced by the application of pesticides, including insecticides. Agrochemicals are, however, among the key factors implicated in contributing to declining bee health and abundance and there is a need to find a balance between the necessity of insecticide applications and the unintended effects to non-target species [1–6].

Compared to organophosphate pesticides, neonicotinoids are relatively safe for vertebrates and have been one of the most important classes of insecticides for the last three decades [7]. Neonicotinoids can be applied to seeds and they then spread systemically within growing plants, killing leaf-eating pests. This reduces non-target effects compared to spraying but still retains toxic burden for pollinators as neonicotinoids continue to be found in wildflowers adjacent to treated fields [8,9].

Neonicotinoids act as agonists, competing with the neurotransmitter acetylcholine in binding to nicotinic acetylcholine receptors (nAChR) [10–12]. The increased toxicity for insects is thought to be caused by the characteristically high nAChR density within insect nervous systems [reviewed in [13]]. Detailed knowledge of cellular and molecular effects of insecticide exposure is required to mitigate negative effects or refine target specificity. Changes in gene expression and processing of RNAs, including alternative splicing, are among the mechanisms available to an organism to adapt to environmental

perturbations [14,15]. Sub-lethal exposure to xenobiotics, such as pesticides, can

alter alternative splicing [16–18]. Since many ion channel genes and other

important neuronal genes undergo extensive alternative splicing, they are

prime targets for changes induced by xenobiotics [17,19]. Moreover, sub-lethal

uptake of some neonicotinoids affects the honey bee brain, impairing foraging

behaviour, flight, navigation, communication, learning and memory [20–25].

Neonicotinoid exposure has also been linked to a decline in bee health,

including a reduction of immune competence [26–28]. Insects do not have

antibodies and rely on the innate immune system to fight microbial infections;

both cellular and humoral responses have been identified [29,30]. The cellular

response leading to phagocytosis or encapsulation of pathogens is mediated

by three types of haematopoietic cell lineages [29,31,32]. The humoral immune

response is activated by Toll and Imd pathways [reviewed in [33]], leading to the

expression of short (10-100 amino acids) antimicrobial peptides (AMPs) which

are highly diverse among insect species [34]. The Toll-pathway is triggered by

Gram-positive bacteria and fungi, ultimately leading to expression of AMPs via

the NF_{kappa}-B transcription factor Dif, that are then secreted from the fat body

into the haemolymph [29,35,36]. The Immune deficiency (Imd) pathway leads to

expression of a different set of AMPs via the NF_{kappa}-B transcription factor relish

in response to peptidoglycans from primarily Gram-negative bacterial through

the activation of pattern recognition receptors and a complex intracellular

signalling cascade [29,30,35]. In addition to the fatbody, AMP expression activated

by the Imd pathway has also been detected in glial cells in the brain [37–40].

We have previously shown that worker-bee larvae in colonies contaminated

with the neonicotinoid imidacloprid, have altered expression of genes belonging

to lipid-carbohydrate-mitochondrial metabolic networks [41]. We have further

demonstrated that sub-lethal exposure to thiamethoxam, another

neonicotinoid, can cause impairment in the midgut and brain of the Africanized

Apis mellifera, as well as contribute to a reduction in honey bee lifespan [9,42–44].

In this study, we analysed the effects of chronic, sub-lethal, thiamethoxam

exposure on genome-wide gene expression and alternative splicing in the

brains of honey bee workers. We found 52 differentially regulated genes

showing a concentration dependent response to thiamethoxam exposure. Most

of these genes have no annotated function but the vast majority are

characterized by encoding short Open Reading Frames (sORFs), half of which

are predicted to encode antimicrobial peptides. This result suggested that

immune responses may be compromised by thiamethoxam exposure. Indeed,

we found that thiamethoxam-exposed bees that were also infected with bacteria

had greatly decreased viability compared to infected but chemically unexposed

bees. Overall, our results suggest that thiamethoxam makes bees vulnerable

to bacterial infection by compromising immune responses.

MATERIALS AND METHODS

We assessed whether the neonicotinoid thiamethoxam alters gene expression

in the brain of Africanized honey bees, *Apis mellifera,* after long-term exposure

to field-relevant concentrations.

Thiamethoxam exposures

Genetically unrelated bees from three unrelated hives from different locations

in the apiary of the Biosciences Institute *, UNESP*, Rio Claro-SP (Brazil). Brood

frames were removed from three different colonies and kept for 24 hours in

biochemical oxygen demand (BOD) at 34° C and relative humidity of 80 ± 10%.

The newly emerged bees (aged 0 to 24h) were marked with special non-toxic

pen (Posca pen) and returned to their respective colonies until they were

collected after 15 days. All experiments followed the recommendations of the

guidelines for xenobiotic assessments on bees (OECD, 1998a, 1998b).

124• Thereafter, bees were kept in groups of 20 individuals in small cages (12

cages/treatment) within an incubator at 32° C. Bees were fed *ad libitum* with

sugar solution (1:1 water and inverted sugar) from pierced 2 ml Eppendorf

tubes for the control group (dataset A, Supplemental Data 1) and, for toxin

exposure treatments, thiamethoxam (Sigma) was added at 2 ng/ml (low dose,

LD, dataset B, Supplementary Data 1) or at 50 ng/ml (high dose, HD, dataset

C, Supplementary Data 1). Sugar-feeders were re-filled before they were empty

to guarantee continuous consumption. The tested concentrations were chosen

based on the range of realistic field concentrations found in pollen and nectar

[9,45–48]. Assuming that caged worker bees consume about 22 μl of 50% sugar

per day [49] the bees would have ingested 0.44 ng and 11 ng after ten days for

the low and high dose exposure, respectively. After ten days, bees were cold-

anaesthetised and their brains were dissected for RNA extraction.

RNA extraction, Illumina sequencing, analysis of differential gene

expression and splicing

Total RNA was extracted from ten dissected brains per sample and three

replicates were done for the control, the low and the high dose treatment

(dataset A, Supplemental Data 1). To extract total RNA, brains were cracked in

liquid nitrogen with a pestle and then homogenized in 50 µl of Tri-reagent

(SIGMA). Then the volume was increased to 500 µl and proceeded following

the manufacturer's instructions. Total RNA was then stored in 70% ethanol and

transported at ambient temperature from Brazil to the UK.

Total RNA was treated with DNase I (Ambion) and stranded libraries for Illumina

sequencing were prepared after poly(A) selection from total RNA (1 µg) with

the TruSeq stranded mRNA kit (Illumina) using random primers for reverse

transcription according to the manufacturer's instructions. Pooled indexed

libraries were sequenced on an Illumina HiSeq2500 to yield 28–41 million

paired-end 125 bp reads for three control, three low dose and three high dose

samples. After demultiplexing, sequence reads were aligned to the *Apis*

mellifera genome (Amel-4.5-scaffolds) using Tophat2.0.6 [50]. Differential gene

expression was determined by Cufflinks-Cuffdiff and the FDR-correction for

multiple tests to raw P values, with p < 0.05 considered significant [51]. Illumina

sequencing and differential gene expression analysis was carried out by

Fasteris (Switzerland). Sequencing data are deposited in GEO under

GSE132858

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE132858).

Alternative splicing was analysed by rMATS [52] and manually inspected by

uploading bam files into the Integrated Genome Viewer [53] and comparing read

frequency. Comparison of gene lists was made with Venny 2.1

(https://bioinfogp.cnb.csic.es/tools/venny/).   Protein   sequences   from

differentially expressed genes of bees were obtained from ensemble

(http://metazoa.ensembl.org/Apis_mellifera/Info/Index) and blasted against

Drosophila annotated proteins using flybase (http://flybase.org) to assign gene

functions. Proteins with no assigned functions were scanned for motifs using

the Interpro server (https://www.ebi.ac.uk/interpro/) [54]. Short ORFs were

analysed for antimicrobial peptide prediction using the following server:

https://www.dveltri.com/ascan/v2/ascan.html [55].

Reverse transcription quantitative polymerase chain reaction (RT-qPCR)

Reverse transcription was carried out with Superscript II (Invitrogen) as

previously described [56]. PCR was performed as described and PCR products

were analysed on ethidium bromide stained 3% agarose gels [19]. Neuronal

genes used as reference for gene expression comparison were bee *erect wing*

(*ewg*, *GB44653)* and *Amyloid Precursor Protein-like* (*Appl, GB48454*), based

on expression analysis in *Drosophila* [57,58]. To validate cDNAs obtained from

RT, the following primers were used to amplify bee *ewg* with AM ewgG F

(CCTGATGGTACCGTATCAATTATTCAAGTTG) and AM ewgH R

(CCGTGTCCATCTTCTCCTGTGAGAATGATTTG), bee *Appl* with AM Appl F

(GCGCGATTCCAGGAAACTGTTGCTGCTC) and AM Appl R2

(CTGCTGTCCAGCAGATGTTTGTAATGAG). Additional primers were

GB45995 (GTTGCATTTTTACGCGTACAGTTACACGACAG) and GB45995 R

(GGGAAATCCCCGGGAAGAGAGCAACTGGAG), GB45995 F

(GTTGCATTTTTACGCGTACAGTTACACGACAG) and GB45995 R

(GGGAAATCCCCGGGAAGAGAGCAACTGGAG), and GB47479 F

(GGGCTATTTGCTATCTAAGTGATCCTCC) and GB47479 R

(GGGTTTAGGAGTTTTCGTTTTAGCTGCTG) and PCR amplifications were

done by 30 sec initial denaturation at 94º C, and then 40 cycles in total with 30

sec at 94º C, annealing at 48º C with 30 sec extension at 72º C for 2 cycles,

then at 50º C with 30 sec extension at 72º C for 2 cycles and then at 52º C with

and 60 sec extension at 72º C for 36 cycles with a final extension of 2 min at

72º C.

RNA quality and quantity were assessed using an Agilent 6000Nano Kit. 250

ng of DNAse-treated RNA (the same samples used for deep seq) was reversed

transcribed using a SuperScript III Invitrogen kit at 55ºC, following

manufacturer's instructions. Primers for quantitative real-time PCR (RT-qPCR)

were designed by Primer Express software (exon/exon junctions included

except for GB41813). Amplicon sizes were assessed in 15% acrylamide gel,

PCR products were then sequenced and efficiency of every pair of primers was

included in the analysis.

Duplicate samples of cDNA were amplified in Real Time PCR with SybrGreen

chemistry under following conditions: denaturation 93ºC for 30 sec, annealing

60ºC for 30 sec, elongation 72ºC for 30sec. *GB41923* was amplified with

primers GB41923 F1 (CGCGTTGATCGTCATGATATTG) and GB41923 R1

(CTATAAGGAAATTTTGAGCCTTCGA), *GB40669* with primers GB40669 F1

(GGCCGGATATCGCTTCAAA) and GB40669 R1

(GTCTCTTTTATCTTTTCCTCGGAATTC), and *GB48969* with primers

GB48969 F1 (TTGCAGCCGTAGCAAAAGGTA), GB48969 R1

(ACCGATTTGAGCACCTTGGT) and *bubblegum* (*bgm, GB51680*) with primers

bgm F1 (CATGCACAAAGAGTACAAAAATTTCA) and bgm R1

(TGGTCCCAATTCTCCAGTAACA). Analysis of CT values was performed in

Light Cycler480 software and normalization and differential expression was

determined with the $2^{-\Delta\Delta Ct}$ method [59] by normalizing to the expression of *ewg*

(GB44653), *Appl (GB48454)* and *actin* (*GB44311*) genes. *Actin* and *ewg* were

amplified with primers ewg Fw2 (CCGCGTCTCCTACAGCTCTT) and ewg Rv2

(TGTAAAACTGCCGTAGGATATTGG)   and   actF1

(TTCCCATCTATCGTCGGAAGA) and actR1 (TTTGTCCCATGCCAACCAT),

and *Appl* with primers AM appl F and AM appl R2, following [41,60].

Bacterial infection assays

Thiamethoxam-induced alteration of anti-microbial peptide gene expression

could disrupt bees' immune response. To evaluate the effect of thiamethoxam

on immunity, we adopted an assay procedure, initially developed for

Drosophila, which assesses how efficiently injected non-pathogenic bacteria

are cleared by anti-microbial peptides [61]. To assay clearance in bees we used

Bacillus badius, a normally non-pathogenic bacterium commonly found in the

environment and *Ochrobactrum anthropi*, which are Gram-positive and Gram-

negative members of the honey bee microbiome [62,63]. *O. anthropi* were isolated

from worker bee gut cultures by plating the gut content on LB agar plates

incubated at 30º C.

Bacterial species were identified by colony PCR and ribosomal 16S

sequencing: A colony was picked from an LB agar plate with a yellow tip and

placed into 10 µl TE in a PCR tube and heated to 94º C for 5 min. The PCR mix

was added adjusting the MgCl concentration to 1.5 mM. PCR was done for 20-

40 cycles with 54º C annealing for 40 sec and 1 min extension at 72º C. A 490

bp fragment of the ribosomal 16S gene was amplified with primers 16S F

(ACTGAGACACGGYCCAGACTCCTACGTC)   and   16S   R

(GCGTGGACTACCAGGGTATCTAATCC) and sequenced with primer 16S

Fseq (CTCCTACGGGAGGCAGCAGTRGGGTC). If sequences did not yield a

single species, primers 16S F2 (GTGGACTACCAGGGTATCTAATCCTG) and

16S R2 (CCTACGGTTACCTTGTTACGACTTCAC) were used for amplification

of a 733 bp fragment, which was sequenced by 16S R2seq

(CCATGGTGTGACGGGCGGTGTGTAC).

Forager honey bees for infection assays were collected from colonies of the

Winterbourne Garden of the University of Birmingham (UK). They were kept

and injected with bacteria as we described previously [19]. Bacteria for injections

were freshly plated and grown overnight on LB plates. Then a single colony

was used to inoculate a 5 ml LB in a 50 ml Falcon tube and grown overnight to

saturation: 2 µl of this culture was then injected. Bacterial titres of cultures were

determined at the time of injections by plating 100 µl of 10^5, 10^6 and 10^7

dilutions in LB and counting the colonies the next day: this showed that 2-8 x

10^6 *B. badius* or *O. anthropi* were typically injected. In some treatments the

injected bacteria were diluted 10-fold (Fig. 3). Groups of 8 to 12 bees were

exposed to combinations of bacterial and thiamethoxam doses and survival

was assessed at 24 hours and 48 hours.

Data on survival were analysed using backwards stepwise logistic ANOVAs

adopting a logit-link function and assuming quasi-binomial distributed errors.

These analyses were carried out using GenStat v.19 (VSN International, Hemel

Hempstead). Statistical analysis of viability were done with GraphPad prism

using ANOVA followed by Tukey-Kramer post-hoc tests.

RESULTS

Chronic thiamethoxam exposure effects gene expression

After low (LD) and high dose (HD) exposure to thiamethoxam, there were 222 up- and 181 down-regulated genes for LD (Fig. 1a) and 233 up- and 114 down-regulated genes for HD with a 1.5 fold difference in expression compared to the control treatment (Fig. 1b; Supplemental Table 1). From these differentially regulated genes, 37 were up-regulated and 15 were down-regulated in a dose-sensitive manner (Fig. 1; Supplemental Table 1).

To validate these results from the Illumina sequencing, we performed RT-qPCR for three of the dose-responsive genes: *GB41923*, a putative sodium-chloride co-transporter, and *GB48969, GB40669,* two genes with unknown function. We detected an expression difference for all three genes upon thiamethoxam exposure (Supplemental Fig. 1). We also validated and confirmed differential expression of *bubblegum*, encoding a very long-chain acyl-CoA synthetase, which has been found to be down-regulated in honey bee larvae exposed to the neonicotinoid imidacloprid [41] (Supplemental Fig. 1).

Alternative splicing has been suggested as a mechanism to adapt gene expression to environmental changes [17,64]. We analysed the RNA-seq data for changes in alternative splicing but found no conclusive patterns (Supplementary Information 1; Supplemental data 2).

Dose-responsive expression occurs mostly in genes encoding uncharacterized ORFs

Next, we categorized the genes dose-responsive to thiamethoxam according to their functions, taking into account known functions of orthologues in

Drosophila and functions deduced from annotated protein domains retrieved by

BLAST analysis. Amongst the dose-sensitive genes that were up-regulated

(Fig. 1A), 14 % (5/37) were assigned roles in cellular signalling (with potential

links to altered neuronal function, such as olfactory and taste perception) and

as structural components (cytoskeleton), and 8 % (3/37) were assigned

functions in transcriptional regulation of gene expression (Fig. 2A). However,

59% (22/37) of dose-sensitive up-regulated genes and 73 % (11/15) of dose-

sensitive down-regulated genes had neither clear orthologues in *Drosophila* nor

any recognizable protein domains that would indicate a biological function (Fig.

2A), which is in contrast to about 20% of genes with unknown function in gene

expression studies in *Drosophila,* [65,66].

Most thiamethoxam dose-responsive genes encode short proteins

Many of the dose-responsive, differentially expressed genes with unknown

function coded for short ORFs. For thiamethoxam-induced differentially up- and

down-regulated genes, respectively 73 % (27/37) and 87 % (13/15) encode for

genes with ORFs of 250 amino acids or shorter (Fig. 2). Using a machine

learning algorithm [55], we predicted the 40 genes coding for peptides of ≤250

amino acids, 17 (43 %), the peptides have antimicrobial function (11/17 genes

were up-regulated and 6/17 were down-regulated, Fig. 2).

Thiamethoxam makes worker bees more vulnerable to bacterial infection

When saturated liquid cultures (2-8 Mio bacteria in 2 µl) of *B. badius* and *O.*

anthropi were injected into bees that were not exposed to thiamethoxam,

viability was little affected, indicating that the bee immune system usually clears

the infection efficiently (Fig. 3, Table 1). In contrast, co-injection of normally sub-lethal doses of thiamethoxam together with either *B. badius* or *O. anthropi* resulted in bee death (Fig. 3, Table 1). In particular, at a dose of 1 µM thiamethoxam (arrow in Fig 3) bees generally survived for 48 h. In contrast, presence of either *B. badius* or *O. anthropi* together with thiamethoxam killed almost all bees within 48 h.

Using data on each bacterial species separately, along with replicates without bacterial exposure, showed that bee viability was negatively affected by increasing doses of thiamethoxam and of bacteria (Table 1: 2-way logistic ANOVAs). At 48 hours, a synergistic interaction between these main effects was detected when the bacterial species was *B. badius*: bee viability declined more rapidly in response to bacterial dose when bees were exposed to higher doses of thiamethoxam (Table 1, Fig. 3).

To test for differences between the effects of the two bacterial species, replicates without bacterial exposure were excluded and the effects of species, non-zero bacterial dose, thiamethoxam and their interactions were explored (Table 1: 3-way logistic ANOVAs). At 24 hours, bee viability declined with both thiamethoxam dose and bacterial dose and was lower when the bacterium applied was *O. anthropi*, without interactions (Table 1). At 48 hours, the same dosage effects were found but the species effect was not significant (Table 1). There was, however, a marginally non-significant interaction between bacterial species and bacterial dose: as probability estimates from logistic analyses are inexact, we explored the consequences of retaining this interaction in the model. This generated a further significant interaction between thiamethoxam

dose and the species of bacteria applied ($F_{1,21}$ = 5.85, P=0.025): *O. anthropi*

affected bees more negatively than did *B. badius* as pesticide dose increased.

We conclude that thiamethoxam suppresses the abilities of bees to cope with

natural challenges of the immune system, which would normally not be fatal.

DISCUSSION

Our study addresses the effect of thiamethoxam on gene expression at field

relevant doses of thiamethoxam found in pollen and nectar. In particular we

chose for these experiments a low and a high dose, which is about 0.04 ng and

1.1 ng per bee per day for a 10 day chronic long-term exposure. A number of

studies have used thiamethoxam doses in the same range for analysing

changes in gene expression or observing an impact on behaviour following

short term exposure [18,20–24,67,68]. In fact, it was found that chronic low-dose long-

term thiamethoxam exposure altered bee activity, motor function and

movement to light [24], but to our knowledge no previous analysis of gene

expression under such conditions has been done.

A key finding of our analysis of honey bee transcriptomes is the highly enriched

fraction of dose-responsive, uncharacterised genes encoding short open

reading frames (sORFs). Such sORFs have only recently been recognized to

encode functional peptides [69–71], some of which play important roles during

development or have a function in the immune system [72,73]. Intriguingly, the

majority of the dose-responsive sORFs we have identified are predicted to

encode peptides with antimicrobial function (anti-microbial peptides, AMPs).

The sORFs we identified have not been reported in prior whole-transcriptome

evaluations of neonicotinoid exposure in bee brains despite using similar

concentrations of neonicotinoids [18,67,68]. A possible explanation for the different gene set we obtained after thiamethoxan exposure is that we analysed changes after long-term, low dose exposure, while prior studies used shorter exposures. The finding that mostly sORFs, and among them many predicted AMPs are differentially expressed points to the immune system for being dysregulated by neonicotinoids. The best-characterized AMPs are expressed in the fat body upon pathogen exposure, but it is now recognized that glial cells in the brain also exert immune functions and express AMPs [37,39,40]. Accordingly, the sORFs that we found to be differentially expressed upon chronic low dose thiamethoxam exposure might form a basal immune defence, similar to the antimicrobial environment present in saliva in vertebrates which many AMPs. Glia are support cells of neurons and are protective in a number of ways. Dysregulated AMP's in the brain have been linked to neurodegenerative disease [37,38,40]. Hence, the dysregulation of sORFs in response to thiamethoxam exposure could indicate disturbed communication between these two cell types beyond immune functions. Likewise, antimicrobial peptides have also been identified in having a role in learning and memory in *Drosophila*, but the molecular mechanism how AMPs enhance learning and memory remains unknown [74].

We noted that other studies found an overrepresentation of down-regulated genes with known function in immune and defence processes upon exposure to different types of neonicotinoids [18,26,67,68,75]. In contrast we did not find differentially regulated immune genes including the known AMPs in the brain. AMPs are highly expressed in the fat body. For the analysis of gene expression in bee brains we dissected them from cold-anaesthetized bees and ensured

that all fat body surrounding the brain was removed. It is possible, that freezing

of brains prior to dissection [67] or residual fat body adhering to the brain

contributed considerably to a different profile of immune gene expression.

Immunosuppression has also observed been upon exposure to neonicotinoids

co-infecting bees with viruses of *Nosema, a* unicellular parasite of bees. In

particular, this affected the cellular response, but also expression of AMPs

[18,26,67,68,75]. Since we did not detect differential expression of known immune

genes, the bees in our study were not infected with these known pathogens,

but we cannot exclude an impact of other microbiota that are non-pathogenic

in healthy bees. Moreover, our analysis focused on gene expression in the

brain, where glial cells govern immune functions, which is a different humoral

response than mediated by fatbody cells secreting AMPs into the hemolymph.

Bees, including *Apis mellifera*, are characterised by their limited set of canonical

immune genes, compared to non-social insects, such as the fruit fly *D.*

melanogaster [76–78]. Currently, only six antimicrobial-peptide genes comprising

four gene families have been described in honey bees [78]. In contrast,

Drosophila has 20 antimicrobial-peptide genes comprising eight gene families

[29,78]. From these genes, only *defensin* is conserved between honey bees and

Drosophila, consistent with the idea that antimicrobial peptides evolve fast to

adapt to species-specific environmental conditions [34]. Given the low number of

known antimicrobial peptides in bees, it is conceivable that new (currently

uncharacterised) genes encoding antimicrobial peptides are evolving.

Consistent with this idea, some of the sORFs differentially regulated by

thiamethoxam are bee or hymenopteran specific.

Various agrochemicals, have been shown to alter the gut microbiome [79,80]. Our

results are consistent with previous findings where neonicotinoid exposure

adversely affects insect immunity [80,81]. Specifically, we have shown that

immune challenges from what are normally non-pathogenic bacteria become

fatal to bees when combined with thiamethoxam exposure. How foreign

bacteria enter the body of bees appears to be crucial For example, Dickel and

colleagues found that oral ingestion of *Enterococcus faecalis* bacteria,

increases the survival neonicotinoid-exposed, caged worker bees [82]. In our

study, the bacteria were injected between segments, which mimics punctures

sustained by attacks of the *Varroa* mite. We note that pathogens can enter bee

haemolymph through punctures inflicted by *Varroa destructor* mites [63] and thus

there may be considerable mortality within hives infested with *Varroa* and also

exposed to thiamethoxam.

In summary, the most prominent changes in gene expression upon long-term

low-dose thiamethoxam exposure identified mostly genes that encode short

ORFs, around half of which are predicted to code for antimicrobial peptides.

Furthermore, thiamethoxam exposure reduced the capacity of bees to

withstand microbial infection. Taken together, these findings imply that low

doses of neonicotinoids may be intrinsically sub-lethal to bees but can be

ultimately fatal via a weakened immune response to extrinsic pathogens. The

roles of the identified genes in the immune response of bees will need to be

identified to establish how bee immunity might be strengthened to resist

bacterial infections.

Authors' contributions

P.D., P.U. and K.D. performed the experiments, T.C.R, I.C.W.H., O.M., N.M.

analysed data, R.S. and M.S. supervised experiments and analysed data, R.S.

and M.S. wrote the manuscript with help from P.D., P.U. and I.C.W.H.

Acknowledgments

We thank N. Parker and the Winterbourne garden for bees, N. Parker for bee

suits, V. Soller-Haussmann and K. Nallasivan for help with bee collections. G.

Salmond for bacterial strains, D. Scocchia for culturing and genotyping bacteria,

and L. Orsini, E. Davies and I. Haussmann for comments on the manuscript.

For this work we acknowledge funding from the Foundation for Research

Support of the State of São Paulo, FAPESP (2012 / 13370-8; 2014 / 23197-7),

the Biotechnology and Biological Sciences Research Council (BBSRC), the

Nottingham-Birmingham Fund, and the Sukran Sinan Memory Fund.

References

1. Klein, S., Cabirol, A., Devaud, J.-M., Barron, A. B. & Lihoreau, M. Why Bees Are So Vulnerable to Environmental Stressors. *Trends Ecol. Evol.* **32**, 268–278 (2017).

2. Steinhauer, N. *et al.* Drivers of colony losses. *Curr. Opin. insect Sci.* **26**, 142–148 (2018).

3. Sánchez-Bayo, F. & Wyckhuys, K. A. G. Worldwide decline of the entomofauna: A review of its drivers. *Biol. Conserv.* **232**, 8–27 (2019).

4. Rundlöf, M. *et al.* Seed coating with a neonicotinoid insecticide negatively affects wild bees. *Nature* **521**, 77–80 (2015).

5. Shah, F. M. *et al.* Field evaluation of synthetic and neem-derived alternative insecticides in developing action thresholds against cauliflower pests. *Sci. Rep.* **9**, 7684 (2019).

6. Decourtye, A. & Devillers, J. Ecotoxicity of neonicotinoid insecticides to bees. *Advances in Experimental Medicine and Biology* **683**, 85–95 (2010).

7. Tomizawa, M. & Casida, J. E. Neonicotinoid insecticide toxicology: mechanisms of selective action. *Annu Rev Pharmacol Toxicol* **45**, 247–268 (2005).

8. Wood, T. J. & Goulson, D. The environmental risks of neonicotinoid pesticides: a review of the evidence post 2013. *Environ. Sci. Pollut. Res. Int.* **24**, 17285–17325 (2017).

9. Tavares, D. A., Roat, T. C., Silva-Zacarin, E. C. M., Nocelli, R. C. F. & Malaspina, O. Exposure to thiamethoxam during the larval phase affects synapsin levels in the brain of the honey bee. *Ecotoxicol.*

Environ. Saf. **169**, 523–528 (2019).

10. Matsuda, K. *et al.* Neonicotinoids: insecticides acting on insect nicotinic acetylcholine receptors. *Trends Pharmacol. Sci.* **22**, 573–80 (2001).

11. Jeschke, P. & Nauen, R. Neonicotinoids-from zero to hero in insecticide chemistry. *Pest Manag Sci* **64**, 1084–1098 (2008).

12. Crossthwaite, A. J. *et al.* The invertebrate pharmacology of insecticides acting at nicotinic acetylcholine receptors. *J. Pestic. Sci.* **42**, 67–83 (2017).

13. Gauthier, M. State of the Art on Insect Nicotinic Acetylcholine Receptor Function in Learning and Memory. in 97–115 (Springer, New York, NY, 2010). doi:10.1007/978-1-4419-6445-8_9

14. Nilsen, T. W. & Graveley, B. R. Expansion of the eukaryotic proteome by alternative splicing. *Nature* **463**, 457–63 (2010).

15. Pai, A. A. & Luca, F. Environmental influences on RNA processing: Biochemical, molecular and genetic regulators of cellular response. *Wiley Interdiscip. Rev. RNA* **10**, e1503 (2019).

16. Sumanasekera, C., Watt, D. S. & Stamm, S. Substances that can change alternative splice-site selection. *Biochem. Soc. Trans.* **36**, 483–90 (2008).

17. Zaharieva, E., Chipman, J. K. & Soller, M. Alternative splicing interference by xenobiotics. *Toxicology* **296**, 1–12 (2012).

18. Colgan, T. J. *et al.* Caste- and pesticide-specific effects of neonicotinoid pesticide exposure on gene expression in bumblebees. *Mol. Ecol.* **28**, 1964–1974 (2019).

19. Decio, P. *et al.* Acute thiamethoxam toxicity in honeybees is not

enhanced by common fungicide and herbicide and lacks stress-induced changes in mRNA splicing. *Sci. Rep.* **9**, 19196 (2019).

20. Cabirol, A. & Haase, A. The Neurophysiological Bases of the Impact of Neonicotinoid Pesticides on the Behaviour of Honeybees. *Insects* **10**, 344 (2019).

21. Fischer, J. *et al.* Neonicotinoids Interfere with Specific Components of Navigation in Honeybees. *PLoS One* **9**, e91364 (2014).

22. Tison, L. *et al.* Effects of sublethal doses of thiacloprid and its formulation Calypso ® on the learning and memory performance of honey bees. *J. Exp. Biol.* **220**, 3695–3705 (2017).

23. Williamson, S. M. & Wright, G. A. Exposure to multiple cholinergic pesticides impairs olfactory learning and memory in honeybees. *J. Exp. Biol.* **216**, 1799–807 (2013).

24. Tosi, S., Burgio, G. & Nieh, J. C. A common neonicotinoid pesticide, thiamethoxam, impairs honey bee flight ability. *Sci. Rep.* **7**, (2017).

25. Aliouane, Y. *et al.* Subchronic exposure of honeybees to sublethal doses of pesticides: Effects on behavior. *Environ. Toxicol. Chem.* **28**, 113–122 (2009).

26. Brandt, A., Gorenflo, A., Siede, R., Meixner, M. & Büchler, R. The neonicotinoids thiacloprid, imidacloprid, and clothianidin affect the immunocompetence of honey bees (*Apis mellifera* L.). *J. Insect Physiol.* **86**, 40–47 (2016).

27. Perveen, N. & Ahmad, M. Toxicity of some insecticides to the haemocytes of giant honeybee, *Apis dorsata* F. under laboratory conditions. *Saudi J. Biol. Sci.* **24**, 1016–1022 (2017).

28. Tesovnik, T. *et al.* Immune related gene expression in worker honey bee (Apis mellifera carnica) pupae exposed to neonicotinoid thiamethoxam and Varroa mites (Varroa destructor). *PLoS One* **12**, e0187079 (2017).

29. Lemaitre, B. & Hoffmann, J. The Host Defense of *Drosophila melanogaster*. *Annu. Rev. Immunol.* **25**, 697–743 (2007).

30. Buchon, N., Silverman, N. & Cherry, S. Immunity in Drosophila melanogaster--from microbial recognition to whole-organism physiology. *Nat. Rev. Immunol.* **14**, 796–810 (2014).

31. Govind, S. Innate immunity in *Drosophila*: Pathogens and pathways. *Insect Sci.* **15**, 29–43 (2008).

32. Parsons, B. & Foley, E. Cellular immune defenses of *Drosophila melanogaster*. *Dev. Comp. Immunol.* **58**, 95–101 (2016).

33. Sheehan, G., Garvey, A., Croke, M. & Kavanagh, K. Innate humoral immune defences in mammals and insects: The same, with differences ? *Virulence* **9**, 1625–1639 (2018).

34. Vilcinskas, A. Evolutionary plasticity of insect immunity. *J. Insect Physiol.* **59**, 123–129 (2013).

35. Ferrandon, D., Imler, J.-L., Hetru, C. & Hoffmann, J. A. The Drosophila systemic immune response: sensing and signalling during bacterial and fungal infections. *Nat. Rev. Immunol.* **7**, 862–874 (2007).

36. Kleino, A. & Silverman, N. The Drosophila IMD pathway in the activation of the humoral immune response. *Dev. Comp. Immunol.* **42**, 25–35 (2014).

37. Kounatidis, I. & Chtarbanova, S. Role of glial immunity in lifespan

determination: A *Drosophila* perspective. *Frontiers in Immunology* **9**, (2018).

38. Kounatidis, I. *et al.* NF-κB Immunity in the Brain Determines Fly Lifespan in Healthy Aging and Age-Related Neurodegeneration. *Cell Rep.* **19**, 836–848 (2017).

39. Petersen, A. J., Rimkus, S. A. & Wassarman, D. A. ATM kinase inhibition in glial cells activates the innate immune response and causes neurodegeneration in *Drosophila*. *Proc. Natl. Acad. Sci. U. S. A.* **109**, E656–E664 (2012).

40. Lye, S. H. & Chtarbanova, S. *Drosophila* as a model to study brain innate immunity in health and disease. *Int. J. Mol. Sci.* **19**, (2018).

41. Derecka, K. *et al.* Transient exposure to low levels of insecticide affects metabolic networks of honeybee larvae. *PLoS One* **8**, e68191 (2013).

42. Oliveira, R. A., Roat, T. C., Carvalho, S. M. & Malaspina, O. Side-effects of thiamethoxam on the brain andmidgut of the africanized honeybee *Apis mellifera* (Hymenopptera: Apidae). *Environ. Toxicol.* **29**, 1122–1133 (2014).

43. Tavares, D. A., Roat, T. C., Carvalho, S. M., Silva-Zacarin, E. C. M. & Malaspina, O. In vitro effects of thiamethoxam on larvae of Africanized honey bee Apis mellifera (Hymenoptera: Apidae). *Chemosphere* **135**, 370–378 (2015).

44. Friol, P. S., Catae, A. F., Tavares, D. A., Malaspina, O. & Roat, T. C. Can the exposure of *Apis mellifera* (Hymenoptera, Apiadae) larvae to a field concentration of thiamethoxam affect newly emerged bees? *Chemosphere* **185**, 56–66 (2017).

45. Krupke, C. H., Hunt, G. J., Eitzer, B. D., Andino, G. & Given, K. Multiple routes of pesticide exposure for honey bees living near agricultural fields. *PLoS One* **7**, (2012).

46. Mullin, C. A. *et al.* High Levels of Miticides and Agrochemicals in North American Apiaries: Implications for Honey Bee Health. *PLoS One* **5**, e9754 (2010).

47. Pilling, E., Campbell, P., Coulson, M., Ruddle, N. & Tornier, I. A Four-Year Field Program Investigating Long-Term Effects of Repeated Exposure of Honey Bee Colonies to Flowering Crops Treated with Thiamethoxam. *PLoS One* **8**, (2013).

48. Stoner, K. A. & Eitzer, B. D. Movement of soil-applied imidacloprid and thiamethoxam into nectar and pollen of squash (*Cucurbita pepo*). *PLoS One* **7**, (2012).

49. Free, J. B. & Spencer-Booth, Y. Observations on the Temperature Regulation and Food Consumption of Honeybees (*Apis mellifera*). *J. Exp. Biol.* **35**, (1958).

50. Kim, D. *et al.* TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol.* **14**, R36 (2013).

51. Trapnell, C. *et al.* Differential gene and transcript expression analysis of RNA-seq experiments with TopHat and Cufflinks. *Nat. Protoc.* **7**, 562–78 (2012).

52. Shen, S. *et al.* rMATS: Robust and flexible detection of differential alternative splicing from replicate RNA-Seq data. *Proc. Natl. Acad. Sci.* **111**, E5593–E5601 (2014).

53. Robinson, J. T. *et al.* Integrative genomics viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).

54. Jones, P. *et al.* InterProScan 5: genome-scale protein function classification. *Bioinformatics* **30**, 1236–1240 (2014).

55. Veltri, D., Kamath, U. & Shehu, A. Deep learning improves antimicrobial peptide recognition. *Bioinformatics* **34**, 2740–2747 (2018).

56. Koushika, S. P., Soller, M., DeSimone, S. M., Daub, D. M. & White, K. Differential and Inefficient Splicing of a Broadly Expressed *Drosophila erect wing* Transcript Results in Tissue-Specific Enrichment of the Vital EWG Protein Isoform. *Mol. Cell. Biol.* **19**, 3998–4007 (1999).

57. Koushika, S. P., Soller, M. & White, K. The Neuron-Enriched Splicing Pattern of Drosophila erect wing Is Dependent on the Presence of ELAV Protein. *Mol. Cell. Biol.* **20**, 1836–1845 (2000).

58. Haussmann, I. U., Li, M. & Soller, M. ELAV-Mediated 3′-End Processing of *ewg* Transcripts Is Evolutionarily Conserved Despite Sequence Degeneration of the ELAV-Binding Site. *Genetics* **189**, 97–107 (2011).

59. Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **25**, 402–408 (2001).

60. Scharlaken, B. *et al.* Reference Gene Selection for Insect Expression Studies Using Quantitative Real-Time PCR: The Head of the Honeybee, *Apis mellifera* , After a Bacterial Challenge. *J. Insect Sci.* **8**, 1–10 (2008).

61. Nehme, N. T. *et al.* Relative Roles of the Cellular and Humoral Responses in the Drosophila Host Defense against Three Gram-

Positive Bacterial Infections. *PLoS One* **6**, e14743 (2011).

62. Engel, P., Martinson, V. G. & Moran, N. A. Functional diversity within the simple gut microbiota of the honey bee. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 11002–7 (2012).

63. Raymann, K. & Moran, N. A. The role of the gut microbiome in health and disease of adult honey bee workers. *Curr. Opin. Insect Sci.* **26**, 97–104 (2018).

64. Soller, M. Pre-messenger RNA processing and its regulation: a genomic perspective. *Cell. Mol. Life Sci.* **63**, 796–819 (2006).

65. Haussmann, I. U., White, K. & Soller, M. Erect wing regulates synaptic growth in Drosophila by integration of multiple signaling pathways. *Genome Biol.* **9**, R73 (2008).

66. Haussmann, I. U. & Soller, M. Differential activity of EWG transcription factor isoforms identifies a subset of differentially regulated genes important for synaptic growth regulation. *Dev. Biol.* **348**, 224–230 (2010).

67. Christen, V., Schirrmann, M., Frey, J. E. & Fent, K. Global Transcriptomic Effects of Environmentally Relevant Concentrations of the Neonicotinoids Clothianidin, Imidacloprid, and Thiamethoxam in the Brain of Honey Bees (*Apis mellifera*). *Environ. Sci. Technol.* **52**, 7534–7544 (2018).

68. Li, Z. *et al.* Brain transcriptome of honey bees (Apis mellifera) exhibiting impaired olfactory learning induced by a sublethal dose of imidacloprid. *Pestic. Biochem. Physiol.* **156**, 36–43 (2019).

69. Couso, J.-P. & Patraquim, P. Classification and function of small open

reading frames. *Nat. Rev. Mol. Cell Biol.* **18**, 575–589 (2017).

70. Pueyo, J. I., Magny, E. G. & Couso, J. P. New Peptides Under the s(ORF)ace of the Genome. *Trends Biochem. Sci.* **41**, 665–678 (2016).

71. Andrews, S. J. & Rothnagel, J. A. Emerging evidence for functional peptides encoded by short open reading frames. *Nat. Rev. Genet.* **15**, 193–204 (2014).

72. Pueyo, J. I. *et al.* Hemotin, a Regulator of Phagocytosis Encoded by a Small ORF and Conserved across Metazoans. *PLoS Biol.* **14**, (2016).

73. Galindo, I., Pueyo, I., Fouix, S., Bishop, S. A. & Couso, J. P. Peptides Encoded by Short ORFs Control Development and Define a New Eukaryotic Gene Family. **5**, (2007).

74. Barajas-Azpeleta, R. *et al.* Antimicrobial peptides modulate long-term memory. *PLOS Genet.* **14**, e1007440 (2018).

75. De Smet, L. *et al.* Stress indicator gene expression profiles, colony dynamics and tissue development of honey bees exposed to sub-lethal doses of imidacloprid in laboratory and field experiments. *PLoS One* **12**, e0171529 (2017).

76. Barribeau, S. M. *et al.* A depauperate immune repertoire precedes evolution of sociality in bees. *Genome Biol.* **16**, 83 (2015).

77. López-Uribe, M. M., Sconiers, W. B., Frank, S. D., Dunn, R. R. & Tarpy, D. R. Reduced cellular immune response in social insect lineages. *Biol. Lett.* **12**, 20150984 (2016).

78. Evans, J. D. *et al.* Immune pathways and defence mechanisms in honey bees Apis mellifera. *Insect Mol. Biol.* **15**, 645–656 (2006).

79. Motta, E. V. S., Raymann, K. & Moran, N. A. Glyphosate perturbs the

gut microbiota of honey bees. *Proc. Natl. Acad. Sci.* **115**, 10305–10310 (2018).

80. Chmiel, J. A., Daisley, B. A., Burton, J. P. & Reid, G. Deleterious Effects of Neonicotinoid Pesticides on Drosophila melanogaster Immune Pathways. *MBio* **10**, e01395-19 (2019).

81. Di Prisco, G. *et al.* Neonicotinoid clothianidin adversely affects insect immunity and promotes replication of a viral pathogen in honey bees. *Proc. Natl. Acad. Sci.* **110**, 18466–18471 (2013).

82. Dickel, F., Münch, D., Amdam, G. V., Mappes, J. & Freitak, D. Increased survival of honeybees in the laboratory after simultaneous exposure to low doses of pesticides and bacteria. *PLoS One* **13**, e0191256 (2018).

Table 1. Effects of thiamethoxam dose, bacterial dose and species on bee viability.

Source	Survival after 24 hours				Survival after 48 hours			
	d.f.	Deviance	*F*	*P*	*d.f.*	Deviance	*F*	*P*
2-way logistic ANOVAs								
B. badius dose	1	60.69	59.26	<0.001	1	74.64	50.04	<0.001
Thiamethoxam dose	1	73.10	71.38	<0.001	1	49.92	33.47	<0.001
Bacteria × Thiamethoxam	1	0.55	0.54	0.472	1	36.79	24.66	<0.001
Residual	17	17.41			17	25.36		
Total	20	121.44			20	186.72		
O. anthropi dose	1	55.42	18.99	<0.001	1	76.28	1.76	0.003
Thiamethoxam dose	1	59.99	20.55	<0.001	1	44.51	11.12	<0.001
Bacteria × Thiamethoxam	1	0.09	0.03	0.865	1	7.06	19.06	0.199
Residual	20	58.38			20	80.04		
Total	23	156.51			23	191.16		
3-way logistic ANOVAs								
Bacterial species	1	15.41		0.021	1	6.21	0.25	0.622
Bacterial dose	1	32.67		0.002	1	14.87	6.28	0.021
Thiamethoxam dose	1	90.19		<0.001	1	85.56	36.12	<0.001
Bacterial species × Bacterial dose	1	3.01		0.281	1	8.73	3.68	0.069
Bacterial species × Thiamethoxam	1	0.13		0.204	1	6.21	2.62	0.121
Bacterial dose × Thiamethoxam	1	4.23		0.824	1	2.56	1.08	0.311
Residual	20	49.08			20	47.39		
Total	26	155.30			26	151.08		

Figure legends

Figure 1: Thiamethoxam induces differential expression in a subset of genes. Venn diagrams indicating the number of differentially expressed genes between control bees and bees exposed to a low dose (left) and high dose (right) of thiamethoxam that were up- (A) or down- (B) regulated. 37 genes were up-regulated and 15 genes were down-regulated in a dose-sensitive manner.

Figure 2

Figure 2: Classification of thiamethoxam induced differentially expressed genes according to function and size. (A) Numbers of genes are plotted according to functions for thiamethoxam induced differentially expressed genes with annotated functions for up- (top) and down-regulated (bottom) genes. (B) Pie charts indicating the fraction of thiamethoxam induced differentially expressed genes according the ORF length for up- (left) and down- (right) regulated genes.

Figure 3

Figure 3: Thiamethoxam exposure makes bees vulnerable to infection by B. badius and O. anthropi. Viability of groups of 8-12 bees 24 h (grey bars) and 48 h (black bars) after injection with *B. badius* (A) or *O. anthropi* (B) (2-8 x 10^6 and diluted 1:10) alone or together with thiamethoxam (at a range of doses, μM) shown as means and standard errors of three replicates. Arrows indicate injection of a 1 μM Thiamethoxam solution with (A right and B) or without bacteria (A left). Statistical significance is indicated by ** ($p<0.01$) and *** ($p<0.001$).

SUPPLEMENTAL INFORMATION AND DISCUSSION

Analysis of alternative splicing

As alternative splicing has been suggested as a mechanism to adapt gene expression to environmental changes [17,64], we analysed the RNA-seq data for changes in alternative splicing. We found significant differences (Supplemental Table 2). Subsequent inspection of sequencing traces using Integrated Genome Viewer (IGV) [53] found that most of these genes had complex splicing patterns that showed no obvious differences in the number of sequence reads over alternatively spliced gene sections (data not shown); therefore, we did not explore these genes further in this study.

Not to find changes in alternative splicing in response to low dose long-term exposure was unexpected, but we also did not find changes upon acute high dose xenobiotic exposure in selected genes in our earlier study on bees [19]. In other insects, alternative splicing of the *Dscam* gene, which acts as a pattern recognition receptor in the immune response, can generate many different isoforms [83,84] and the splicing patterns can change upon bacterial infection [85]. Since *Dscam* alternative splicing is robust against perturbations of splicing factors [86,87], the absence of changes might indicate that a specific immune challenge is required to change its splicing pattern [85]. We previously also analysed alternative splicing in bee *elav* and *Xbp-1* genes as potential markers for defects in synapse formation and the stress response upon high dose acute thiamethoxam exposure, but also low dose chronic exposure did not affect their splicing [19,65,66]. In contrast to most other species, bees have only a single *elav* gene making it unlikely that the lack of splicing differences is due to redundancy

among close related ELAV RNA binding proteins [88]. Potentially, alternative splicing might be changed only in a few cells in the brain, eluding detection without single cell analysis, or could in addition also be more subtle, requiring more replicates for detection of significantly altered alternative splicing changes [17].

References

83. Hemani, Y. & Soller, M. Mechanisms of Drosophila Dscam mutually exclusive splicing regulation. *Biochem. Soc. Trans.* **40**, 804–809 (2012).

84. Schmucker, D. *et al.* Drosophila Dscam is an axon guidance receptor exhibiting extraordinary molecular diversity. *Cell* **101**, 671–84 (2000).

85. Dong, Y., Taylor, H. E. & Dimopoulos, G. AgDscam, a Hypervariable Immunoglobulin Domain-Containing Receptor of the Anopheles gambiae Innate Immune System. *PLoS Biol.* **4**, e229 (2006).

86. Ustaoglu, P. *et al.* Srrm234, but not canonical SR and hnRNP proteins, drive inclusion of *Dscam* exon 9 variable exons. *RNA* **25**, 1353–1365 (2019).

87. Haussmann, I. U. *et al.* Plasmid-based gap-repair recombineered transgenes reveal a central role for introns in mutually exclusive alternative splicing in Down Syndrome Cell Adhesion Molecule exon 4. *Nucleic Acids Res.* **47**, 1389–1403 (2018).

88. Zaharieva, E., Haussmann, I. U., Bräuer, U. & Soller, M. Concentration and Localization of Coexpressed ELAV/Hu Proteins Control Specificity of mRNA Processing. *Mol. Cell. Biol.* **35**, 3104–15 (2015).

Supplemental Figure 1

Supplemental Figure legend

Supplemental Figure 1: Validation of thiamethoxam induced differentially expressed genes by RT-qPCR. Means with standard error from three experiments are represented by a log2 fold change in expression levels normalized to *ewg*, *Appl* and *actin* genes.