1 **Selection on a pleiotropic color gene block underpins early differentiation between two**
2 **warbler species**
3
4
5
6

Silu Wang[1]*, Sievert Rohwer[2], Devin R. de Zwaan[3], David P. L Toews[4], Irby J. Lovette[5], Jacqueline
Mackenzie[1], Darren E. Irwin[1]
[1] Department of Zoology, 6270 University Blvd, University of British Columbia, BC, Vancouver, BC,
V6T1Z4, Canada
[2] Department of Biology and Burke Museum, Box 353010, University of Washington, Seattle, WA,
98195, USA
[3] Department of Forest and Conservation Sciences, 2424 Main Mall, University of British Columbia, BC,
V6T1Z4, Canada
[4] Department of Biology, 619 Mueller Laboratory, Pennsylvania State University, University Park, PA,
16802, USA
[5] Fuller Evolutionary Biology Program, Cornell Lab of Ornithology, 159 Sapsucker Woods Road, Ithaca,
NY, 14850, USA
19
20 *Corresponding author
21 Email: wangsmt@zoology.ubc.ca

22
23

24   **Abstract**

25   When one species gradually splits into two, divergent selection on specific traits can cause
peaks of differentiation in the genomic regions encoding those traits. Whether speciation is
initiated by strong selection on a few genomic regions with large effects or by more diffused
selection on many regions with small effects remains controversial. Differentiated phenotypes
between differentiating lineages are commonly involved in reproductive isolation, thus their
genetic underpinnings are key to the genomics architecture of speciation. When two species
hybridize, recombination over multiple generations can help reveal the genetic regions
responsible for the differentiated phenotypes against a genomic background that has been
homogenized via backcrossing and introgression. We used admixture mapping to investigate
genomic differentiation and the genetic basis of differentiated plumage features (relative melanin
and carotenoid pigment) between hybridizing sister species in the early stage of speciation:
Townsend's (*Setophaga townsendi*) and Hermit warblers (*S. occidentalis*). We found a few
narrow and dispersed divergent regions between allopatric parental populations, consistent with
the 'divergence with gene flow' model of speciation. One of the divergent peaks involves three
genes known to affect pigmentation: ASIP, EIF2S2, and RALY (the ASIP-RALY gene block).
After controlling for population substructure, we found that a single nucleotide polymorphism
(SNP) inside the intron of RALY displays a strong pleiotropic association with cheek, crown, and
breast coloration. In addition, we detect selection on the ASIP-RALY gene block, as the
geographic cline of the RALY marker of this gene block has remained narrower than the plumage
cline, which remained narrower than expected under neutral diffusion over two decades. Despite
extensive gene flow between these species across much of the genome, the selection on ASIP-
RALY gene block maintains stable genotypic and plumage difference between species allowing
further differentiation to accumulate via linkage to its flanking genetic region or linkage-
disequilibrium genome-wide.
49
50
51   **Key words:** *speciation, admixture mapping, cline, plumage, RALY, ASIP, hybrid zone,*
52   *Setophaga*

Introduction

Examining the genomic distribution of differentiation between two populations can reveal

targets of divergent selection, advancing our understanding of the speciation process [1–6]. At the

onset of divergence between sister taxa, differentiation at narrow regions of chromosomes, or

"islands of differentiation", is expected [7–12]. Strong selection on a few genes of large

phenotypic effects can result in a small number of highly differentiated regions. Physical linkage

results in elevated divergence of the nearby neutral regions, a phenomenon known as divergence

hitchhiking [2,7,11]. More differentiation can accumulate as speciation progresses, thus

extending regions of high differentiation and forming "continents of divergence" [4].

Alternatively, speciation can be initiated via many regions of small effects on traits under

multifarious selection (one selection targeting multiple traits) and gradually accumulate genome-

wide differentiation via correlated differentiation across the genome [13–18]. A central debate in

speciation research has therefore been whether speciation occurs through gradual differentiation

throughout the genome or more rapidly at a few key regions. Genomic analyses of sister species

in the early stages of speciation uncovers the evolutionary process of speciation, thus inform this

debate.

Islands of divergence may be associated with divergent (and often diagnostic) traits that

are involved in reproductive isolation [19–21]. Divergent phenotypic traits frequently play

important roles in social interactions at geographic boundaries between closely-related species

[22,23]. These features are commonly involved in mate-choice [19,20] or aggressive signaling

[21,24] that are divergently selected [25–30] and facilitate species recognition and reproductive

isolation [31,32]. We are just beginning to understand the genetic underpinnings of traits related

to species divergence [26,27,29,33,34]. Integrating the genetic basis underlying pigmentation

with corresponding evolutionary forces will deepen our understanding of the formation and

maintenance of species barriers.

Carotenoid and melanin pigmentation are commonly involved in species-diagnostic traits

in avian signals [19,25–28,32,35]. At the boundaries between closely-related species, the

intensity of coloration might not differ, but rather the differential patterning of the colors is

involved in mate choice favoring conspecifics [32,35,36]. In such cases, the regulatory genes for

melanin and carotenoid pathways are strong candidates for differentiation between species as

these loci are likely to cause differences in plumage pattern.

Hybrid zones, locations where divergent lineages interbreed, provide opportunities to

understand the relationship between species-diagnostic features and genomic differentiation.

Social signals can be crucial for conspecific mate recognition and intrasexual competition for

mating opportunities [19,20,37–39]. When the plumage signals established in isolated

populations come into secondary contact, hybrids can be selected against due to their

intermediate or mismatched parental signals interfering with mate recognition and competition

[37,38].

The hybrid zone between *Setophaga townsendi* (abbreviated as *townsendi*) and *S.*

occidentalis (abbreviated as *occidentalis*) along the Cascade mountains harbors extensive gene

flow between these closely related species that started diverging an estimated 400,000 years ago

[40–44]. Males of this species pair differ in several plumage features related to both carotenoid

and melanin patterning on the crown, cheek, breast, and back. For example, *townsendi* has a

melanin-based cheek patch (Figure 1A right), whereas *occidentalis* displays a completely

carotenoid-based cheek (Figure 1A, left). Because apparent hybrids predominantly resemble

occidentalis in crown and cheek coloration (Figure 1A center), with intermediate breast (the

extent of yellow plumage) and back colors (green plumage in the mantle), Rohwer & Wood [43]

predicted that face coloration of Hermit Warblers and hybrids would be controlled by a single-

locus dominant allele. In addition, whether the other carotenoid and melanin patterning

differences between species are underpinned by the same genetic mechanism is an open question.

These signals can be important in male-male competition for territories, as individuals in the

hybrid zone demonstrate an aggression bias towards different plumage types [42,45].

We investigated whether speciation in this young species pair occurred through gradual

widespread differentiation across the genome or via a small number of restricted regions. If

selection for coloration phenotypes has targeted a small number of specific genomic regions,

genomic islands of differentiation should be narrow, dispersed, and associated with loci involved

in pigmentation pathways [3,7,8]. In addition, we asked the following questions regarding the

genetic basis of species-specific traits: (1) is there an island of divergence pleiotropically

underpins species-specific features (crown and cheek darkening, breast yellow, bib size, greenish

back); (2) Is cheek darkening influenced largely by an allele of dominant effect, consistent with

the prediction made two decades ago by Rohwer & Wood (1998); and (3) Is there stable selection

maintaining differentiation in the gene region underlying key species diagnostic features?

Materials and Methods

Sampling

Whole specimens of *Setophaga* warblers were collected over two historical sampling

sessions in the Washington Cascade hybrid zone [41,43]. The first sampling ($N = 314$ individuals;

35 sites) was carried out in 1987-1994 [43], while the second ($N = 127$; 11 sites) was in 2005-

2008 [41] and covered a subset of the sites from the original sampling. We accessed these

specimens at the Burke Museum of Natural History and Culture (University of Washington,

Seattle, Washington).

We carried out a third round of sampling using a catch-and-release approach during the

breeding season (early May to mid-July) in 2015-16 (Figure 1B). Upon locating a territorial male

by song, a mist net with a playback (of a locally recorded song) at the bottom was set up nearby.

After capturing an adult, photographs and a blood sample were taken for further analysis. We re-

sampled the sites that were sampled in 1986-94 [43]. For details, see previous work in this system

[46].

Plumage measurements

Melanin- and carotenoid-based plumage traits allow identification of the two species

(Figure 1B), but there is also variation within each species [43,47]. To quantify plumage variation

within and between populations, we focused on five distinct plumage traits in males: the relative

amount of black (melanin) and yellow (carotenoid) in the 1) cheek, 2) crown, and 3) breast and

the 4) throat bib, as well as, 5) the intensity of green chroma on the back (Figure 1B).

For each of the 242 warblers captured in the field, we took three pictures from different

angles: 1) frontal with head tilted up (for bib and breast measurements), 2) profile (cheek), and 3)

from above (crown) (Figure S1). Plumage color metrics were measured using Adobe Photoshop

CC in CIE (*Comission Internationale de l'Eclairage*) LAB color space. LAB color space is a 3-

dimensional space consisting of 3 distinct, perpendicular axes: 1) Luminosity (L) ranging from 0

(black) to 100 (white), 2) 'a' ranging from green (negative) to red (positive), and 3) 'b' ranging

from blue (negative) to yellow (positive; Adobe 2017). We chose this color space because it

linearizes the variables of interest along three distinct axes: black ('L'), yellow (along the 'b'

axis), and green ('a').

For the cheek and mantle, we selected a standardized area and averaged the pixels to

record the 'b' and 'a' values, respectively. For the cheek, we selected and averaged the entire area

from above the eye (but excluding the eye) to the throat badge, and from the base of the bill to the

mantle using the profile photos. Differences in ambient light conditions at the time a picture is

taken can confound comparison of color metrics among individuals. To address this, we used the

white-balance feature in Photoshop, using the white plumage of each individual's belly as a

standard, to correct for differences in ambient light among photos and standardize the color

metrics. We acknowledge that without spectral analysis, we do not incorporate UV reflectance

which is a ubiquitous aspect of signaling in avian systems [48]. However, our methods allow us

to estimate the relative intensity of melanin- and carotenoid-based plumage traits during the

breeding season.

To measure the size of the black bib, we used the program Analyzing Digital Images

(ADI;[49]). A scale was included in all photos to standardize size measurements among

individuals. We measured bib size (Figure 1A) by creating a polygon around the bib and

calculating the area ($\pm$ 0.1 mm^2).

GBS pipeline

The GBS pipeline details have been described in [46]. Following [50], we prepared

genotyping-by-sequencing (GBS) [51] libraries from individual DNA samples. In brief, genomes

were digested with restriction enzyme and ligated with barcode and adaptors, amplified with

PCR, and size selected (fragment length of 300 - 400 bp) for sequencing. Libraries were

sequenced at Genome Quebec with paired-end sequencing (read length = 125 bp) on an Illumina

HiSeq 2500 automated sequencer. The resulting sequences were processed following a GBS

pipeline [52]. All the sequence can be acquired through GenBank (accession number:

PRJNA573930, ID: 573930). We demultiplexed the reads with a custom script and trimmed them

using Trimmomatic 0.36 (Bolger et al 2014) [TRAILING:3 SLIDINGWINDOW:4:10

MINLEN:30], then we aligned reads to a *Taeniopygia guttata* reference version 3.2.4 [53] using

bwa (Li et al 2009) (default settings). We assumed synteny of *Setophaga* and *T. guttata* genomes

given the limited rearrangement in avian genomes [54,55], and the conclusions of this study

would be unlikely to be affected by a moderate number of rearrangements. We conducted SNP

calling with GATK (McKenna et al. 2010), which produced 4,097,089 SNPs. The SNP filtering

was done with VCFtools [56], which includes removing indels, genotype quality (GQ) > 20,

minor allele frequency (MAF) $\geq$ 0.05, removing loci with >30% missing data, and only including

biallelic SNPs, resulting in 21,852 SNPs remaining.

Whole Genome Sequencing (WGS)

In addition, we selected 5 samples from *occidentalis* in Pinehurst, CA, U.S.A. (UWBM

66152, 66153, 66148-66150) and 5 samples from inland *townsendi* in Tok, AK, U.S.A. (UWBM

84816-84819, 84860) from the Burke museum for whole genome re-sequencing. For DNA

extraction, we used 2 mm^3 of tissue digested in Qiagen buffers—following the tissue extraction

procedure—and separated DNA using UPrep spin columns (Genesee). We standardized DNA

concentrations with after quantifying concentrations with a Qubit fluorometer, and then generated

sequencing libraries with the llumina TruSeq Nano kit—which includes an 8-cycle PCR

enrichment—selecting 350 bp insert sizes. We individually indexed each sample, and sequenced

the combined libraries across a single lane of an Illumina NextSeq using the paired-end 150 bp

sequencing chemistry. We combined these 10 samples with other wood warblers from other

projects, but consistently included 24 individuals per sequencing lane in order to generate

comparable coverage across samples.

Genomic differentiation

To quantify the level of differentiation throughout the genome, we calculated F_{ST} [57]

with VCFtools [56] between allopatric *townsendi* (Montana and Idaho, USA; $N = 38$) and

occidentalis (Oregon and California, USA; $N = 23$) for each of the filtered SNPs.

The genomic architecture of differentiation from the GBS was compared to that of the

WGS data. The reads from the WGS data were aligned to the same *Taeniopygia guttata* reference

(version 3.2.4) [53] with bwa (Li et al 2009) (default settings). ANGSD [58] was employed to

calculate genotype frequencies accounting for genotyping uncertainty before estimating F_{ST} for

each of the non-overlapping (step size = 10kb) 10kb windows.

Admixture mapping

To identify SNPs that are associated with variation in the five plumage traits, we

conducted admixture mapping with GenABEL package in R [59] for the subset ($N = 242$) of the

GBS data produced in [46] that had pigment quantification (see above): 189 birds sampled from

hybrid zone (around the center of the hybrid zone in Washington, U.S.A.), 46 from the

occidentalis zone (Southern Oregon and California, U.S.A.), and 7 from the *townsendi* zone

(Northeast Washington & Montana, U.S.A.). The GenABEL function *ibs*, which calculates

identity-by-state relatedness among individuals, was used to control for population structure.

Genomic control of inflation factor λ (which represents the effect of genetic structure and sample

size) was conducted [59]. Briefly, λ was estimated from the genomic data assuming that

randomly-selected markers from the genome are not associated with the trait (after controlling for

population substructure). λ was then used to correct the test statistic χ^2 of each association test, so

that the test is relative to the null hypothesis of no phenotype-genotype association [59]. Because

the cheek darkening and breast yellow intensity were left-skewed, we rank order transformed

these phenotype data.

To find the number of independent hypotheses for multiple hypothesis correction, we

estimated the total number of independent linkage blocks (a proxy for the number of independent

hypotheses) in the dataset. Principle Component Analysis calculating genotypic covariance

[60,61] was run until 99.5% of the variance was explained with the R package SNPRelate [62],

producing 210 PCs. We applied a Bonferroni correction to the independent blocks, such that

significance level alpha = 0.05/210. Because we conducted admixture mapping on five plumage

traits, we further applied Bonferroni correction across the 5 plumage traits, which gave rise to the

final alpha value of $0.05/(210 \times 5) \approx 4.76 \times 10^{-5}$. Candidate SNPs (i.e., those with *p*-values below

this alpha value) were annotated in Ensembl Zebra Finch (taeGut3.2.4;[63]).

Geographical cline analysis on candidate loci

As one candidate gene, RALY (see Results), stood out in the above analysis as

particularly strongly associated with plumage variation, we investigated the spatial and temporal

variation in this locus relative to the plumage hybrid index and rest of the genome (using samples

described previously [46]). We fit the relationship between RALY allele frequency and location

using an equilibrium geographic cline model [64,65].

Geographical cline analysis followed [46]. Briefly, we collapsed the two-dimensional

sampling into a one-dimensional transect by measuring the location of each site to the 0.5 isocline

of the plumage hybrid index (HI) in the historical sampling [43], as follows. First, a Local

Polynomial Regression Fitting (LOESS) model in R [66] was used to fit variation in HI across the

hybrid zone, and this model was used to estimate the HI = 0.5 isocline in the 1987-94 sampling

[43]. Then for each site, the shortest distance to the 0.5 isocline was calculated with 'sp' package

[67]. The sites east or west of the isocline were specified as having positive versus negative

distance values, respectively. We added 1200 km to the distance score of each site so that all

distance values are above zero, while the relative distance of each site to the isocline is preserved.

Then the data was fit to the equilibrium sigmoidal cline model [64,65] $y = \frac{1}{1+e^{-\frac{4(x-c)}{w}}}$, in which

cline center (c) and width (w) was estimated (where y is HI and x is location with respect to the

HI = 0.5 isocline). To examine whether selection (i.e., divergent selection and/or selection against

hybrids) is acting on the candidate loci, we followed [46] and tested whether the increase in cline

width ($w^2_{2015\text{-}16}$ -$w^2_{1987\text{-}94}$) is significantly less than expected under the neutral diffusion model

[68]. The change of w^2 (between sampling periods) as opposed to w allows comparison of the

observed change in cline width to the neutral expectation [46]. To understand the RALY cline

width change relative to the plumage and genomic cline, we compared $w^2_{2015\text{-}16}$ -$w^2_{1987\text{-}94}$ relative

to the plumage and genomic HI cline [46], which was respectively based on the scores of the 8

plumage landmarks (with 0 representing pure *occidentalis* and 1 being pure *townsendi*), and the

scaled genomic PC1 (with 0 representing pure *occidentalis* and 1 being pure *townsendi*).

Results

Across the Cascade mountain range, the *occidentalis* plumage type was observed on the

southwest and the *townsendi* plumage type was on the northeast [43,46] (Figure 1 B). The overall

pattern of genomic differentiation was consistent with the plumage divergence: *occidentalis* and

townsendi plumage types were most different along genomic PC1. There was low genome-wide

weighted average F_{ST} [57] between allopatric *occidentalis* and *townsendi* populations (Weir and

Cockerham's F_{ST} = 0.03; Figure 2A). However, four high regions of differentiation (F_{ST} > 0.6)

were found (Figure 2A, Figure S2, gene association and functions summarized in Table S2) that

map to Zebra Finch chromosome (chr) 5 (nucleotide position 25064223-25875302 in the Zebra

Finch genome, mean F_{ST} = 0.75), chr 20 (1981369, mean F_{ST} = 0.90), and chr Z (66226657, F_{ST}

= 0.82). Such genomic architecture of divergence revealed by GBS data is consistent with the

pattern from the WGS data (Figure S3), where the genome is largely undifferentiated except a

few peaks at chr 1A, 4, 5, 20, and Z (Figure S3G).

The F_{ST} peak on chr 20 (Figure 2B) from the GBS data is within an intron of the gene

RALY, which encodes heterogeneous nuclear ribonucleoprotein, the cofactor for cholesterol

biosynthetic genes [69]. We hereafter refer to this SNP as the RALY SNP. This RALY SNP

demonstrated the highest F_{ST} (0.90) inside an "island" of relatively high F_{ST} on chr 20 between

occidentalis and *townsendi* parental populations across the genome (Figure 2B). The pure

occidentalis population mostly contains GG homozygotes (referred to hereafter as OO), while the

townsendi population contains CC homozygotes (referred to as TT). In the WGS data (Figure

S4), which has much higher density of markers across the genome than the GBS dataset, this

'island' of differentiation contains pigmentation genes ASIP, EIF2S2, and RALY (Figure S4BC),

thus we refer to it hereafter as the "ASIP-RALY" block. The RALY SNP revealed by the GBS

data (Figure 2B) conveniently becomes the marker representing this ASIP-RALY divergent gene

block surveyed across a large number of individuals (Figure S4BC).

Admixture mapping showed a strong pattern of few loci associated with divergent

plumage traits in the hybrid zone. In particular, the colors of the crown, cheek, and the breast

were each very strongly associated with the same RALY SNP (on chr 20) that was the outlier in

the F_{ST} analysis (crown: χ^2 = 57.89, r^2 = 0.57, p = 5.95 x 10^{-15}, Figure 3A, Figure 4AD; cheek: χ^2

= 44.76, r^2 = 0.55, p = 2.23 x 10^{-11}, Figure 3C, Figure 4BE; breast: χ^2 = 28.47, r^2= 0.57, p = 9.53 x

10^{-8}, Figure 3D, Figure 4CF). The RALY SNP shows a partial dominance pattern of the G allele

for the three plumage traits, with G/C heterozygotes tending to have similar phenotypes as G/G

homozygotes, although there appears to be some additivity as well (Figure 4). The RALY gene

(Figure 3B) is associated with yellow pigmentation in mice and quail [70,71], and is adjacent to

two other pigmentation genes -- ASIP (115941 bases away) and EIF2S2 (30223 bases away)

(Figure 3B, S4C).

A low-resolution sequencing approach such as GBS, which sequences less than 1% of the

genome, would be unlikely to detect a SNP that is directly causal for phenotypic variation. It is

more likely that the RALY SNP is closely physically linked to the causal DNA variant (SNP or

some other type of variant) for the phenotypic differences. It is also possible that multiple linked

SNPs in this region are responsible for the variation in different phenotypic traits, such that the

apparent pleiotropic effect (i.e., strong association with three plumage traits) of the RALY SNP

might be due to separate causal genes that are closely linked. The continuous divergence of the

ASIP-RALY gene block between allopatric species in the WGS dataset suggests that the ASIP-

RALY gene linkage block might be involved in plumage color divergence as a whole (Figure

S4).

Three additional SNPs were found to be significantly associated with the intensity of

yellow on the breast (Figure 3D). Two are on chr 4A at nucleotide location 5588139 and

5588235, both inside an ortholog of the mammal Immunoglobulin Binding Protein 1 (IGBP1)

gene. One is at nucleotide location 6623798 on chr 13, inside gene Annexin A6, which codes for

a phospholipid binding protein, Annexin VI [72]. We did not detect SNPs that significantly

explained variation in either bib size or green coloration on the back (Figure 3EF). The RALY

SNP explains 57% of variation in crown (Figure 4AD), 52% of the variation in cheek (Figure 4B,

E), and 43% variation in breast coloration (Figure 4CF).

We found evidence that the ASIP-RALY region is under divergent selection, as it is the

most extreme F_{ST} outlier, while most of the genome is not very differentiated (average = 0.03)

(Figure 5D). In addition, the RALY SNP (representing the ASIP-RALY gene block)

demonstrated a spatial cline that was stable in location over two decades (Figure 5A). The cline

center was at 1216.35 ± 4.0 (SE) km in 1987-94 (Figure 5A, blue curve), and did not significantly

shift in 2015-16 sampling (Figure 5A, yellow curve), occurring at 1213.140 ± 2.2 (SE) km.

Furthermore, the width of the RALY cline stayed narrower (95% CI of $w^2_{2015\text{-}16}$ - $w^2_{1987\text{-}94}$: -

39004.70 to 1035.13 km^2) than under a model of neutral diffusion ($w^2_{2015\text{-}16}$ - $w^2_{1987\text{-}94}$ = 7540

km^2) (Figure 5B, see [46] for explanation of method).

Discussion

We observed clear but narrow genomic 'islands of differentiation' between Townsend's and Hermit Warblers on chr 1A, 5, 20, and Z (Figure 2). This agrees with the predicted pattern of speciation in the face of gene flow: as most of the genome remains homogenized due to gene flow and lack strong selection, restricted targets of divergent selection have become strongly differentiated [3,7,11,73].

The RALY SNP (chr 20) within the ASIP-RALY gene block represents one of the 'islands of differentiation' and is strongly associated with key divergent traits: cheek, crown, and breast coloration. The consistently narrow geographical RALY cline in the hybrid zone over several decades, despite ongoing gene flow [46], further implicates selection (divergent selection and/or selection against hybrids) around the RALY SNP as contributor to reproductive isolation between these young sister species. The $w^2_{2015-16}$ -$w^2_{1987-94}$ estimate at the RALY cline was less than the plumage cline and the genomic PC1 cline (Figure 5B), suggesting that a narrow region centered at the RALY locus might be the direct target of selection maintaining the stable plumage and genomic clines [46]. Such selection may further extend genomic differentiation by elevating the differentiation at linked sites.

Although there is signature of genetic hitchhiking that underpins extended genetic divergence around the ASIP-RALY targets of selection (Figure S5), it seems that the plumage pigmentation effect is more narrowly underpinned by ASIP-RALY region. In the hybrid zone, the recombination over multiple generations can break down the hitchhiking gene blocks (that tend to co-segregate with the trait-determining loci in allopatric populations), revealing specific genetic underpinnings of the differentiated phenotypes. The genetic underpinning of the plumage trait should be very close to the narrow region around the RALY SNP.

The 'divergence with gene flow' pattern, in which closely related taxa hybridize

extensively and have little differentiation across much of the genome, along with evidence of a

narrow and highly differentiated genetic region that has large effect on divergent traits suggests a

simple genomic architecture of divergence. Such a strong target of selection could effectively

counteract gene flow and maintain a species boundary at secondary contact, allowing speciation

to progress [4,74,75]. There may also be weak multilocus divergent selection on a large set of

loci, yet such weak polygenic selection is inherently challenging to detect.

Pleiotropy

This ASIP-RALY block underlying multiple plumage patches could provide a strong

pleiotropic mechanism (one region that affects multiple phenotypes [76]), linking divergent

selection and possibly reproductive isolation. The RALY SNP is associated with a suite of

melanin- and carotenoid-related plumage patches (crown, cheek, and breast coloration) that differ

diagnostically between species (Figure 3) and are likely under some form of selection. Previous

admixture mapping in *Setophaga coronata auduboni*/*S. c. coronata* revealed SCARF2 on chr 15

that is associated with multiple carotenoid and melanin traits [77]. Intriguingly, in this sister pair

within the same genus, drastically different color regions are involved in carotenoid versus

melanin plumage divergence. Such distinct pleiotropic genetic underpinnings of plumage

divergence may facilitate rapid speciation in *Setophaga* [78].

Melanin and carotenoid plumage traits are often involved in mate-choice and male-male

competition [79–82]. If carotenoid-related plumage traits are targets of female mate-choice or

male-male competition for mating opportunities in *Setophaga* warblers, strong selection on

pleiotropic genes can play a powerful role in limiting mating between species and facilitating

speciation. Pleiotropy is a more effective mechanism linking selection to reproductive isolation

than linkage or polygenic inheritance, although genetic evidence for pleiotropic genes that

influence both selected traits and reproductive isolation has been scarce (see review [83]). Among

the limited existing examples, the classic cases involve coloration: the wingless gene affecting

reproductive isolation and wing coloration in *Heliconius* butterflies [19], and the YUP locus

affects pollinator isolation and flower coloration in monkey flowers [10]. RALY could be an

additional example of a speciation gene. The stable and narrow cline of RALY SNP and extreme

differentiation at the RALY locus imply divergent selection and/or selection against hybrids at

this locus, which suggests that RALY could be associated with reproductive isolation in this

system. Future study is needed to validate the possibility by examining the role of RALY-ASIP-

regulated plumage signals in mate preference or male-male competition.

Dominance

Our results support the prediction that Rohwer and Wood [43] made two decades ago that

the cheek coloration in hybrids is underpinned by a single dominant locus. Indeed, heterozygotes

of the RALY SNP exhibit similar yellow cheeks as the homozygous GG individuals (i.e., yellow

cheek), which were found predominantly in the *occidentalis* population (Figure 4D-F).

Dominance in the signal trait might reduce gene flow if it is in the opposite direction of

dominance in the receiver trait (i.e., heterozygous receivers are more sensitive to *townsendi*

signals), due to 'opposing genetic dominance' in receiver and signaler traits [83,84]. This is

because the F1 hybrids (heterozygous for both signaling and receiving) might for example,

demonstrate *occidentalis* signal and preference for *townsendi*, thus cannot mate with other F1

hybrids. Therefore the *occidentalis* RALY dominance effect on the cheek signal could suppress

gene flow if it is opposed by a *townsendi* dominant effect on receiving trait.

RALY SNP

The RALY SNP is at the peak of divergence at allopatry (Figure 2B) and is tightly

associated with plumage variation in the hybrid zone, although its neighboring SNPs are not

significantly associated with plumage traits (after controlling for population substructure) (Figure

3, 6C). The roughly 180kb extended divergence around RALY in WGS data (Figure S2, S4)

suggests the entire color gene block (ASIP-RALY) is involved in divergence.

The RALY gene encodes for heterogeneous nuclear ribonucleoprotein, the RNA binding

protein that is known to regulate the expression of its downstream gene that codes for agouti

signaling protein (ASIP) [70]. Deletion of RALY in Japanese quails and mice leads to novel

transcripts of ASIP, which cause a yellow skin phenotype called "lethal yellow" [70,71]. In mice,

a lethal yellow mutation (Ay) in ASIP exhibits dominant pleiotropic effects on other traits in

addition to skin coloration, which includes obesity, diabetic condition, etc., that are unrelated to

the ASIP gene [71]. ASIP itself is well known for its influence on skin pigment by binding to

melanocortin receptors and competitively excluding its agonists, preventing black/brown

pigmentation [85,86]. In birds, ASIP has been repetitively involved in genomic divergence

between closely-related species in *Vermivora* [26], *Sporophila* [87], and *Munia* [88]*,* in a

mendelian inheritance pattern with the melanic form being the recessive phenotype. Genetic

variants of RALY in warblers could result in differential expression ASIP leading to variations in

carotenoid and melanin patterning. The pleiotropic dominance effect of the RALY SNP we

identified in this natural hybrid zone is consistent with these known effects of RALY mutants in

the lab. Future study should investigate the functional interaction of RALY and ASIP in plumage

divergence, and whether there is any convergent mutation in ASIP between *Vermivora* and

Setophaga.

On top of the strong pleiotropic ASIP-RALY association with plumage, there could be

other regions of the genomes contributing to further refinement of species-specific coloration in

this system. Specifically, other significant SNPs on chr 4A and 13 are associated with breast

coloration. In addition, there might be narrow but important genetic regions that this GBS data

did not cover. However, this is unlikely because the WGS data did not reveal additional

differentiation peaks.

Conclusion

We identified a few narrow and dispersed genomic regions of differentiation between two

hybridizing *Setophaga* warblers, consistent with divergent selection in the face of gene flow. A

highly divergent peak spans among three closely linked pigmentation genes: ASIP, EIF2S2, and

RALY. Admixture mapping within a hybrid zone between these species showed that this SNP

was strongly associated with three distinct plumage traits that distinguish the species, revealing a

major-effect region related to speciation. In contrast, we did not find genetic regions significantly

associated with the other two species-diagnostic traits, implying many genes of small effects.

The extensive similarity in the rest of the genome raises a question regarding the

distinctiveness of *townsendi* and *occidentalis* and their future. Key to this question is whether

these regions of differentiation represent sufficient reproductive isolation that will maintain

separation for further differentiation to build up. The genetic divergence of the ASIP-RALY gene

region between allopatric populations and the stable narrow geographical cline of this region

suggest that selection maintains such difference between species. Future studies should track the

genomic differentiation in linkage disequilibrium with the ASIP-RALY region and investigate

the mechanism of apparent selection. Overall, these results show that avian sister species can be

very similar across the great majority of their genomes, sometimes differing primarily at narrow

genomic regions under selection maintaining phenotypic differences between species.

Acknowledgement

Ruth Midgley and Else Mikkelsen provided excellent assistance in the field and lab for this study.

We thank Chris Wood (Burke Museum) for help relocating the historical sites and accessing the

historical plumage samples, and Sharon Birks (Burke Museum) for accessing tissue samples for

sequencing. Sally Otto, Ailene MacPherson, Dolph Schluter, Loren Rieseberg, Armando

Geraldes, Tom Booker, Toby Bradshaw, Molly Schumer, and members of the Irwin Lab provided

helpful discussion. We thank research funding provided by the Natural Sciences and Engineering

Research Council of Canada (grants 311931-2012, RGPIN-2017-03919 and RGPAS-2017-

507830 to DEI; and PGS D 331015731 to SW), a Werner and Hildegard Hesse Research Award

in Ornithology and a UBC Four Year Doctoral Fellowship to SW. Finally, we thank Environment

Canada, U. S. Geological Survey, Departments of Fish & Wildlife of Washington, Oregon,

California, Idaho, and Montana, and the UBC Animal Care Committee for providing research

permits.

Reference

1.   Feder JL, Gejji R, Yeaman S, Nosil P. Establishment of new mutations under divergence
and genome hitchhiking. Philos Trans R Soc B Biol Sci. 2012;367: 461–474.
doi:10.1098/rstb.2011.0256
2.   Via S. Divergence hitchhiking and the spread of genomic isolation during ecological
speciation-with-gene-flow. Philosophical Transactions of the Royal Society B: Biological
Sciences. 2012. pp. 451–460. doi:10.1098/rstb.2011.0260
3.   Rice WR, Hostert EE. Laboratory Experiments on Speciation: What Have We Learned in
40 Years? Evolution (N Y). 1993;47: 1637. doi:10.2307/2410209
4.   Nosil P, Harmon LJ, Seehausen O. Ecological explanations for (incomplete) speciation.

454   Trends in Ecology and Evolution. 2009. pp. 145–156. doi:10.1016/j.tree.2008.10.011

455   5.   Irwin DE, Milá B, Toews DPL, Brelsford A, Kenyon HL, Porter AN, et al. A comparison
456   of genomic islands of differentiation across three young avian species pairs. Mol Ecol.
457   2018;27: 4839–4855. doi:10.1111/mec.14858

458   6.   Wolf JBW, Ellegren H. Making sense of genomic islands of differentiation in light of
459   speciation. Nature Reviews Genetics. 2017. p. :87-100. doi:10.1038/nrg.2016.133

460   7.   Nosil P, Funk DJ, Ortiz-Barrientos D. Divergent selection and heterogeneous genomic
461   divergence. Molecular Ecology. 2009. pp. 375–402. doi:10.1111/j.1365-
462   294X.2008.03946.x

463   8.   Turner TL, Hahn MW, Nuzhdin S V. Genomic islands of speciation in Anopheles
464   gambiae. PLoS Biol. 2005;3: 1572–1578. doi:10.1371/journal.pbio.0030285

465   9.   Wu CI. The genic view of the process of speciation. Journal of Evolutionary Biology.
466   2001. pp. 851–865. doi:10.1046/j.1420-9101.2001.00335.x

467   10.   Bradshaw HD, Schemske DW. Allele substitution at a flower color locus produces a
468   pollinator shift in monkeyflowers. Nature. 2003;426: 176–178. doi:10.1038/nature02106

469   11.   Via S. Natural selection in action during speciation. Proc Natl Acad Sci. 2009;106: 9939–
470   9946. doi:10.1073/pnas.0901397106

471   12.   Schluter D, Conte GL. Genetics and ecological speciation. Proc Natl Acad Sci. 2009;106:
472   9955–9962. doi:10.17226/12692

473   13.   Orr HA. The population genetics of speciation: The evolution of hybrid incompatibilities.
474   Genetics. 1995.

475   14.   Orr HA, Turelli M. The evolution of postzygotic isolation: Accumulating Dobzhansky-
476   Muller incompatibilities. Evolution (N Y). 2001. doi:10.1111/j.0014-3820.2001.tb00628.x

477   15.   Matsubayashi KW, Katakura H. Contribution of multiple isolating barriers to reproductive
478   isolation between a pair of phytophagous ladybird beetles. Evolution (N Y). 2009;63:
479   2563–2580. doi:10.1111/j.1558-5646.2009.00738.x

480   16.   Lawniczak MKN, Emrich SJ, Holloway AK, Regier AP, Olson M, White B, et al.
481   Widespread divergence between incipient Anopheles gambiae species revealed by whole
482   genome sequences. Science (80-). 2010. doi:10.1126/science.1195755

483   17.   Johnson NA, Porter AH. Rapid speciation via parallel, directional selection on regulatory
484   genetic pathways. J Theor Biol. 2000;205: 527–542. doi:10.1006/jtbi.2000.2070

485   18.   Williams LM, Oleksiak MF. Ecologically and evolutionarily important SNPs identified in
486   natural populations. Mol Biol Evol. 2011;28: 1817–1826. doi:10.1093/molbev/msr004

487   19.   Kronforst MR, Young LG, Kapan DD, McNeely C, O'Neill RJ, Gilbert LE. Linkage of
488   butterfly mate preference and wing color preference cue at the genomic location of
489   wingless. Proc Natl Acad Sci. 2006;103: 6575–6580. doi:10.1073/pnas.0509685103

490   20.   Whibley AC, Langlade NB, Andalo C, Hanna AI, Bangham A, Thébaud C, et al.
491   Evolutionary paths underlying flower color variation in Antirrhinum. Science (80-).
492   2006;313: 963–966. doi:10.1126/science.1129161

493   21.   Seehausen O, Schluter D. Male-male competition and nuptial-color displacement as a
494   diversifying force in Lake Victoria cichlid fishes. Proc R Soc B Biol Sci. 2004;271: 1345–
495   1353. doi:10.1098/rspb.2004.2737

496   22.   Anderson CN, Grether GF. Interspecific aggression and character displacement of
497   competitor recognition in *Hetaerina* damselflies. Proc R Soc London, Ser B. 2010;277:
498   549–55. doi:10.1098/rspb.2009.1371

23.   West-Eberhard M-J. Sexual selection, competitive communication, and species-specific
signals in insects. In: Lewis T, editor. Insect Communication. Academic Press; 1984. pp.
283–324.

24.   Mikami OK, Kohda M, Kawata M. A new hypothesis for species coexistence: Male-male
repulsion promotes coexistence of competing species. Population Ecology. 2004. pp. 213–
217. doi:10.1007/s10144-004-0189-5

25.   Uy JAC, Moyle RG, Filardi CE, Cheviron ZA. Difference in Plumage Color Used in
Species Recognition between Incipient Species Is Linked to a Single Amino Acid
Substitution in the Melanocortin-1 Receptor. Am Nat. 2009;174: 244–254.
doi:10.1086/600084

26.   Toews DPL, Taylor SA, Vallender R, Brelsford A, Butcher BG, Messer PW, et al.
Plumage Genes and Little Else Distinguish the Genomes of Hybridizing Warblers. Curr
Biol. 2016;26: 2313–2318. doi:10.1016/j.cub.2016.06.034

27.   Toews DPL, Brelsford A, Grossen C, Milá B, Irwin DE. Genomic variation across the
Yellow-rumped Warbler species complex. Auk. 2016;133: 698–717. doi:10.1642/AUK-
16-61.1

28.   Barrera-Guzmán AO, Aleixo A, Shawkey MD, Weir JT. Hybrid speciation leads to novel
male secondary sexual ornamentation of an Amazonian bird. Proc Natl Acad Sci.
2018;115: E218–E225. doi:10.1073/pnas.1717319115

29.   Knief U, Bossu CM, Saino N, Hansson B, Poelstra J, Vijay N, et al. Epistatic mutations
under divergent selection govern phenotypic variation in the crow hybrid zone. Nat Ecol
Evol. 2019;3: 570–576. doi:10.1038/s41559-019-0847-9

30.   Poelstra J. The Genetics of Speciation and Coloration in Carrion and Hooded Crows.
Digital Comprehensive Summaries of Uppsala Dissertations from the Faculty of Science
and Technology. 2013.

31.   Hill GE, Mcgraw KJ. Correlated changes in male plumage coloration and female mate
choice in cardueline finches. Anim Behav. 2004;67: 27–35.
doi:10.1016/j.anbehav.2003.02.002

32.   Sætre GP, Moum T, Bureš S, Král M, Adamjan M, Moreno J. A sexually selected
character displacement in flycatchers reinforces premating isolation. Nature. 1997.
doi:10.1038/42451

33.   Shriner D. Overview of admixture mapping. Curr Protoc Hum Genet. 2017;2017: 1.23.1-
1.23.8. doi:10.1002/cphg.44

34.   Crawford JE, Nielsen R. Detecting adaptive trait loci in nonmodel systems: Divergence or
admixture mapping? Mol Ecol. 2013;22: 6131–6148. doi:10.1111/mec.12562

35.   Hill GE. Sexiness, Individual Condition, and Species Identity: The Information Signaled
by Ornaments and Assessed by Choosing Females. Evol Biol. 2015;42: 251–259.
doi:10.1007/s11692-015-9331-x

36.   Uy JAC, Moyle RG, Filardi CE, Cheviron ZA. Difference in Plumage Color Used in
Species Recognition between Incipient Species Is Linked to a Single Amino Acid
Substitution in the Melanocortin-1 Receptor. Am Nat. 2009. doi:10.1086/600084

37.   Dijkstra PD, Border SE. How does male-male competition generate negative frequency-
dependent selection and disruptive selection during speciation? Curr Zool. 2018/01/02.
2018;64: 89–99. doi:10.1093/cz/zox079

38.   Pfennig KS, Pfennig DW. Character displacement: ecological and reproductive responses

to a common evolutionary problem. Q Rev Biol. 2009;84: 253–276. Available:
https://www.ncbi.nlm.nih.gov/pubmed/19764283

39.   van Doorn GS, Dieckmann U, Weissing FJ. Sympatric Speciation by Sexual Selection: A
Critical Reevaluation. Am Nat. 2004;163: 709–725. doi:10.1086/383619

40.   Krosby M, Rohwer S. A 2000 km genetic wake yields evidence for northern glacial refugia
and hybrid zone movement in a pair of songbirds. Proc R Soc B Biol Sci. 2009;276: 615–
621. doi:10.1098/rspb.2008.1310

41.   Krosby M, Rohwer S. Ongoing movement of the hermit warbler X Townsend's Warbler
Hybrid Zone. PLoS One. 2010;5: e14164. doi:10.1371/journal.pone.0014164

42.   Pearson SF, Rohwer S. Asymmetries in male aggression across an avian hybrid zone.
Behav Ecol. 2000;11: 93–101. doi:10.1093/beheco/11.1.93

43.   Rohwer S, Wood C. Three Hybrid Zones Between Hermit and Townsend ' S Warblers in
Washington and Oregon. Auk. 1998;115: 284–310.

44.   Weir JT, Schluter D. Ice sheets promote speciation in boreal birds. Proc R Soc B Biol Sci.
2004;271: 1881–1887. doi:10.1098/rspb.2004.2803

45.   Pearson SF. Behavioral asymmetries in a moving hybrid zone. Behav Ecol. 2000;11: 84–
92. doi:10.1093/beheco/11.1.84

46.   Wang S, Rohwer S, Delmore KE, Irwin DE. Cross-decades stability of an avian hybrid
zone. J Evol Biol. 2019.

47.   Owen-Ashley NT, Butler LK. Androgens, interspecific competition and species
replacement in hybridizing warblers . Proc R Soc B Biol Sci. 2004;271: S498–S500.
doi:10.1098/rsbl.2004.0230

48.   Eaton MD, Lanyon SM. The ubiquity of avian ultraviolet plumage reflectance. Proc R Soc
B Biol Sci. 2003;270: 1721–1726. doi:10.1098/rspb.2003.2431

49.   Snyder R. Analyzing Digital Images (ADI) Resources. STEM Digit. 2011;21.

50.   Alcaide M, Scordato ESC, Price TD, Irwin DE. Genomic divergence in a ring species
complex. Nature. 2014;511: 83–85. doi:10.1038/nature13285

51.   Elshire RJ, Glaubitz JC, Sun Q, Poland JA, Kawamoto K, Buckler ES, et al. A robust,
simple genotyping-by-sequencing (GBS) approach for high diversity species. PLoS One.
2011;6: e19379. doi:10.1371/journal.pone.0019379

52.   Irwin DE, Alcaide M, Delmore KE, Irwin JH, Owens GL. Recurrent selection explains
parallel evolution of genomic regions of high relative but low absolute differentiation in a
ring species. Mol Ecol. 2016;25: 4488–4507. doi:10.1111/mec.13792

53.   Warren WC, Clayton DF, Ellegren H, Arnold AP, Hillier LW, Künstner A, et al. The
genome of a songbird. Nature. 2010;464: 757–762. doi:10.1038/nature08819

54.   Zhang G, Li C, Li Q, Li B, Larkin DM, Lee C, et al. Comparative genomics reveals
insights into avian genome evolution and adaptation. Science (80-). 2014;346: 1311–1320.
doi:10.1126/science.1251385

55.   Ellegren H. Evolutionary stasis: the stable chromosomes of birds. Trends Ecol Evol.
2010;25: 283–291. doi:10.1016/j.tree.2009.12.004

56.   Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, et al. The variant
call format and VCFtools. Bioinformatics. 2011;27: 2156–2158.
doi:10.1093/bioinformatics/btr330

57.   Weir BS, Cockerham CC. Estimating F-Statistics for the Analysis of Population Structure.
Evolution (N Y). 1984;38: 1358. doi:10.2307/2408641

58.   Korneliussen TS, Albrechtsen A, Nielsen R. ANGSD: Analysis of Next Generation
590   Sequencing Data. BMC Bioinformatics. 2014. doi:10.1186/s12859-014-0356-4
591   59.   Aulchenko YS, Ripke S, Isaacs A, van Duijn CM. GenABEL: An R library for genome-
592   wide association analysis. Bioinformatics. 2007;23: 1294–1296.
593   doi:10.1093/bioinformatics/btm108
594   60.   Gao X, Starmer J, Martin ER. A multiple testing correction method for genetic association
595   studies using correlated single nucleotide polymorphisms. Genet Epidemiol. 2008;32:
596   361–369. doi:10.1002/gepi.20310
597   61.   Johnson RC, Nelson GW, Troyer JL, Lautenberger JA, Kessing BD, Winkler CA, et al.
598   Accounting for multiple comparisons in a genome-wide association study (GWAS). BMC
599   Genomics. 2010;11: 724. doi:10.1186/1471-2164-11-724
600   62.   Zheng X, Levine D, Shen J, Gogarten SM, Laurie C, Weir BS. A high-performance
601   computing toolset for relatedness and principal component analysis of SNP data.
602   Bioinformatics. 2012;28: 3326–3328. doi:10.1093/bioinformatics/bts606
603   63.   Zerbino DR, Achuthan P, Akanni W, Amode MR, Barrell D, Bhai J, et al. Ensembl 2018.
604   Nucleic Acids Res. 2018;46: D754–D761. doi:10.1093/nar/gkx1098
605   64.   Szymura JM., Barton NH. Genetic Analysis of a Hybrid Zone Between the Fire-Bellied
606   Toads , Bombina bombina and B . variegata , Near Cracow in Southern Poland. Evolution
607   (N Y). 1986;40: 1141–1159.
608   65.   Gay L, Crochet PA, Bell DA, Lenormand T. Comparing clines on molecular and
609   phenotypic traits in hybrid zones: A window on tension zone models. Evolution (N Y).
610   2008;62: 2789–2806. doi:10.1111/j.1558-5646.2008.00491.x
611   66.   R Core team. R Core Team. R: A Language and Environment for Statistical Computing. R
612   Foundation for Statistical Computing , Vienna, Austria. ISBN 3-900051-07-0, URL
613   http://www.R-project.org/. 2015. pp. 275–286.
614   67.   Pebesma EJ, Bivand RS. Classes and methods for spatial data in R. R News. 2005;5: 9–13.
615   doi:10.1159/000323281
616   68.   Barton NH, Hewitt G. M. Analysis of Hybrid Zones. Annu Rev Ecol Syst. 1985;16: 113–
617   148.
618   69.   Sallam T, Jones MC, Gilliland T, Zhang L, Wu X, Eskin A, et al. Feedback modulation of
619   cholesterol metabolism by the lipid-responsive non-coding RNA LeXis. Nature. 2016;534:
620   124–128. doi:10.1038/nature17674
621   70.   Nadeau NJ, Minvielle F, Ito S, Inoue-Murayama M, Gourichon D, Follett SA, et al.
622   Characterization of Japanese quail yellow as a genomic deletion upstream of the avian
623   homolog of the mammalian ASIP (agouti) gene. Genetics. 2008;178: 777–786.
624   doi:10.1534/genetics.107.077073
625   71.   Michaud EJ, Bultman SJ, Klebig ML, van Vugt MJ, Stubbs LJ, Russell LB, et al. A
626   molecular model for the genetic and phenotypic characteristics of the mouse lethal yellow
627   (Ay) mutation. Proc Natl Acad Sci. 1994;91: 2562–2566. doi:10.1073/pnas.91.7.2562
628   72.   Chang N, Sutherland C, Hesse E, Winkfein R, Wiehler WB, Pho M, et al. Identification of
629   a novel interaction between the Ca(2+)-binding protein S100A11 and the Ca(2+)- and
630   phospholipid-binding protein annexin A6. Am J Physiol Cell Physiol. 2007;292: C1417-
631   30. doi:10.1152/ajpcell.00439.2006
632   73.   Nosil P, Feder JL. Widespread yet heterogeneous genomic divergence. Molecular Ecology.
633   2012. pp. 2829–2832. doi:10.1111/j.1365-294X.2012.05580.x

74. Endler JA. Properties of Clines. Geographic Variation, Speciation, and Clines. Princeton,
New Jersey: Princeton University Press; 1977. pp. 62–65.
75. Gavrilets S. Fitness landscapes and the origin of species. Princeton University Press.
Princeton, New Jersey: Princeton University Press; 2004. Available:
https://books.google.co.uk/books?hl=en&lr=&id=duk9DwAAQBAJ&oi=fnd&pg=PR13&
dq=Fitness+Landscapes+and+the+Origin+of+Species&ots=qBMShqcSzt&sig=EXfli7MIc
LpN-IJqfmGZGrOyvTw#v=onepage&q=Fitness Landscapes and the Origin of
Species&f=false%0Ahttp://scholar.goog
76. Fisher RA. The genetical theory of natural selection. New York, NY: Oxford Univ. Press;
1930.
77. Brelsford A, Toews DPL, Irwin DE. Admixture mapping in a hybrid zone reveals loci
associated with avian feather coloration. Proc R Soc B Biol Sci. 2017;284: 20171106.
doi:10.1098/rspb.2017.1106
78. Lovette IJ, Bermingham E. Explosive speciation in the New World Dendroica warblers.
Proc R Soc B Biol Sci. 1999;266: 1629–1636. doi:10.1098/rspb.1999.0825
79. Kingma SA, Szentirmai I, Székely T, Bókony V, Bleeker M, Liker A, et al. Sexual
selection and the function of a melanin-based plumage ornament in polygamous penduline
tits Remiz pendulinus. Behav Ecol Sociobiol. 2008;62: 1277–1288. doi:10.1007/s00265-
008-0556-x
80. McGraw KJ. Carotenoid-based ornamentation and status signaling in the house finch.
Behav Ecol. 2000;11: 520–527. doi:10.1093/beheco/11.5.520
81. Pryke SR, Andersson S. Carotenoid-based epaulettes reveal male competitive ability:
Experiments with resident and floater red-shouldered widowbirds. Anim Behav. 2003;66:
217–224. doi:10.1006/anbe.2003.2193
82. Tarof SA, Dunn PO, Whittingham LA. Dual functions of a melanin-based ornament in the
common yellowthroat. Proc R Soc B Biol Sci. 2005;272: 1121–1127.
doi:10.1098/rspb.2005.3053
83. Nosil P. A Genetic Mechanism to Link Selection to Reproductive Isolation. Ecological
Speciation. Oxford: Oxford Univ. Press; 2012. pp. 109–138.
84. Forister ML. Independent inheritance of preference and performance in hybrids between
host races of Mitoura butterflies (Lepidoptera: Lycaenidae). Evolution (N Y). 2005;59:
1149–1155.
85. Lovett M, Cheng ZY, Lamela EM, Yokoi T, Epstein CJ. Molecular markers for the agouti
coat color locus of the mouse. Genetics. 1987;115: 747–754.
86. Wolf Horrell EM, Boulanger MC, D'Orazio JA. Melanocortin 1 receptor: Structure,
function, and regulation. Frontiers in Genetics. 2016. p. 95. doi:10.3389/fgene.2016.00095
87. Campagna L, Repenning M, Silveira LF, Fontana CS, Tubaro PL, Lovette IJ. Repeated
divergent selection on pigmentation genes in a rapid finch radiation. Sci Adv. 2017;3:
e1602404. doi:10.1126/sciadv.1602404
88. Stryjewski KF, Sorenson MD. Mosaic genome evolution in a recent and rapid avian
radiation. Nat Ecol Evol. 2017;1: 1912–1922. doi:10.1038/s41559-017-0364-7
675
676

678
Figure 1 A, illustration highlighting the key plumage difference between *occidentalis* (left,
turquoise), hybrids (center, yellow), and *townsendi* (right, magenta) (illustration by Gil Jorge
Barros Henriques). **B**, a map showing site mean plumage hybrid index based on 8 plumage traits
(0 for pure *occidentalis*, in turquoise; 1 for pure *townsendi*, in magenta). **C**, Genomic eigenvector
1 (EV1) and eigenvector 2 (EV2) with individual datapoints colored by plumage hybrid index.
The genomic EV1 reflects the variation among individuals that are *occidentalis*-like (low
genomic eigenvector EV1) versus *townsendi*-like (high genomic EV1).

Figure 2 F_{ST} (between inland *townsendi* and *occidentalis*) scan is consistent with the isolation
with gene flow model in which a few regions are under selection (divergent selection or selection
against hybrids) while the rest of the genome is similar between species (Weir Cockerham
Weighted average F_{ST} is indicated by the horizontal red dotted line) due to gene flow. A, F_{ST}
genomic scan of the genome revealed an island of differentiation around the highest peak in
chromosome 20 (B, zoom in chromosome 20).

Figure 3 Admixture mapping revealed a SNP inside the RALY gene in chromosome 20 at position 1981369 that was significantly associated with crown (**A**, **B**) and cheek darkening (**C**), as well as the intensity of yellow on the breast (**D**), but not with bib size (**E**) nor with the color of the back (**F**). **A-F**, genomic scan of inflation factor-corrected –log (*p*-value) of genotype-phenotype association tests. The horizontal red lines represent the Bonferroni-corrected critical threshold. The plumage trait tested in each scan is indicated by the red triangle in the cartoons. **A-C**, there was a strong peak at chromosome 20 position 1981369 inside the RALY gene that indicates a pleiotropic effect on crown, cheek, and breast coloration. **B**, genetic map of around position 1981369 on chromosome 20 and its nearby SNPs in this study. The green boxes represent the stretches of the protein-coding genes, EIF2S2 and RALY, with the exons depicted as the red vertical bars connected by introns (red lines). **E-F**, no significant SNP was detected that is associated with bib size or greenish back.

707
708
Figure 4 Association of the RALY SNP on crown (**A**, **D**), cheek (**B**, **E**), and breast (**C**, **F**)
coloration, in which the pure *occidentalis* genotypes is denoted as "OO", pure *townsendi*
genotype is denoted as "TT", and heterozygotes as "OT". **D-F**, The associations of the RALY
genotype and cheek darkening among individuals are significant ($p < 10^{-7}$) after accounting for
the underlying genomic ancestry. The data are consistent with a combination of dominance and
additive effects on the three plumage traits.

715
Figure 5 Evidence of selection on RALY locus or its linked region. The RALY SNP shows
stable clines (sampled in 1987-94 and 2015-16) across the hybrid zone, which are extremely
narrow (**A, B**, zoom in around 1100 to 1300 km), and has become significantly narrower in the
recent sampling (**C**). The bootstrap distribution of $w^2_{2015\text{-}16}$ -$w^2_{1987\text{-}94}$ (the change of squared cline
width between sampling periods) of the RALY cline is less than expected under neutral diffusion,
suggesting selection has been maintaining narrow clines of genomic PC1, plumage, and RALY
SNP (**C**). The selection at RALY (greater deviation from neutral expectation than plumage and
genomic PC1) might indirectly maintain the narrow plumage and genomic cline (**C**). The vertical
lines are $w^2_{2015\text{-}16}$ -$w^2_{1987\text{-}94}$ values (solid line: sample estimates; dotted line: bootstrap mean
values) respectively for RALY cline (yellow), plumage cline (turquoise), and genomic PC1 cline
(purple). (**D**), RALY is the extreme outlier in the distribution of F_{ST} between inland *townsendi*
and *occidentalis* across the genome (RALY SNP F_{ST} value depicted by the red line).
728
729

Supplementary Information

We then investigated whether selection at the RALY marker representing ASIP-RALY gene block has led genetic hitchhiking of its flanking regions facilitating further differentiation around this target of selection. We tested whether the bootstrap (for 10,000 iterations) difference of F_{ST} between the candidate genetic block (300kb flanking region of the ASIP-RALY genetic block, excluding the RALY locus itself, colored in yellow Figure S5A) versus the rest of chr20 (colored in light blue Figure S5A) is significantly greater than 0. Indeed, we found the signature of hitchhiking, as ASIP-RALY candidate genetic block demonstrated significantly greater F_{ST} than the rest of the regions in chr20 (Figure S5AB, grey distribution of difference is greater than 0, 95% CI: 0.107-0.109, $t_{9999} = 254.31$, $p < 10^{-15}$).

Figure S1 Field photos of a hybrid male showing three different angles: 1) frontal with head tilted up showing throat badge and breast measurements), 2) profile showing the cheek, and 3) from above showing the crown.

Figure S2 SNPs with $F_{ST} > 0.6$ are colored in red. High F_{ST} SNPS are distributed on chr1A, 5, 20, and Z.

749
750 **Figure S3** Weir Cockerham F_{ST} scan of with the WGS data, in which each dot represents a 10kb
751 non-overlapping window (A), where peaks were found on chromosome 1A (B), 4 (C), 5 (D), 20
752 (E), and Z (F). G, the ASIP-RALY gene block demonstrates extremely high F_{ST} relative to the rest
753 of the genome.
754

755
756
757 **Figure S4** The $\boldsymbol{F_{ST}}$ peak on chromosome 20 (A) resides in the ASIP-RALY gene block (B, C). C,
758 the position of the protein coding genes relative to the peak (bounded by the red vertical lines). C,
759 within the genes, the vertical strikes are the coding regions flanked by the non-coding regions
760 (horizontal lines).
761
762

764
Figure S5 There is signature of hitchhiking around the RALY peak (elevated F_{ST} around RALY),
while the effect on plumage coloration is more tightly associated with RALY. Thus the potential
causal locus of this trait should be very close to RALY. **A, B** signature of hitchhiking: the
candidate gene block (the pigmentation gene block and its 300kb flanking region excluding the
RALY locus) has higher F_{ST} than regions in the rest of the chr20 (B, the bootstrap difference of
F_{ST} is significantly greater than 0, indicated by the vertical dotted line, 95% CI: 0.107-0.109). **C,**
in comparison to F_{ST}, the trait effect is more specific to the RALY peak: there is a steeper decay
of scaled trait effect size (relative to RALY peak) than the scaled differentiation (relative F_{ST} to
RALY peak).

Table S1 Allele frequencies of SNPs between RALY and ASIP in hybrid zone and parental
zones. The causal genetic mechanism for plumage coloration might be narrower than the ASIP-
RALY linkage block around the RALY SNP. The RALY SNP is in physical proximity to the two
other pigmentation genes mentioned above, although 3 other SNPs (1955244, 1972476, and
1972481) that are 8888-26125 bp away from RALY SNP (Table S2; Figure 3B), physically
closer to ASIP and EIF2S2, did not show association with phenotype (Figure 3B). Two of these
SNPs showed low minor allele frequencies (0.05-0.09) in the hybrid zone, and thus are not
expected to be highly associated with trait variation. However, the fact that SNP 1973476 (closer
to the other genes than the candidate RALY SNP) demonstrated similar minor allele frequency as
the significant RALY SNP and was not significantly associated with plumage coloration
highlights the importance of the region around position 1981369 inside the RALY gene. Either
way, the RALY SNP represents the ASIP-RALY gene block and appears to have an effect on
multiple species-diagnostic coloration traits.
790

	POS	allele1:freq	allele2:freq
Hybrid zone			
	1955244	T:0.946	A:0.054
	1972476	A:0.72	C:0.28
	1972481	T:0.917	C:0.083
	1981369	C:0.375	G:0.625
townsendi zone			
	1955244	T:1	A:0
	1972476	A:0.5	C:0.5
	1972481	T:0.656	C:0.344
	1981369	C:1	G:0
occidentalis zone			
	1955244	T:0.95	A:0.05
	1972476	A:0.944	C:0.056
	1972481	T:0.972	C:0.028
	1981369	C:0.048	G:0.952

791
792
793
794

795 **Table S2** SNPs with $F_{ST} > 0.6$ and their position, association to genes and molecular functions

Chromosome	Position	F_{ST}	Relation to genes	Gene(s)	Function
1A	54442413	0.656	intergenic	GRM8, ENSTGUG00000004218	G protein-coupled receptor activity, glutamate receptor activity, telomere maintenance, DNA binding
5	25064223	0.822	intron	UBR1	ubiquitin-protein transferase activity
5	25174918	0.855	coding	ENSTGUG00000011205	DNA binding
5	25746680	0.644			
5	25783847	0.772	intergenic	DPF3, RGS6	histone acetyltransferase activity; G protein-coupled receptor signaling pathway, intracellular signal transduction
5	25875302	0.649	intron	RGS6	G protein-coupled receptor signaling pathway, intracellular signal transduction
20	1981369	0.892	intron	RALY	DNA and RNA binding, cholesterol biosynthesis
Z	66226657	0.818	intron	BBOX1, TNPO1	carnitine biosynthesis, oxidation-reduction; protein import into nucleus, intracellular protein transport