1   Spexin and a novel cichlid-specific spexin paralog both inhibit FSH and LH through a
2   specific galanin receptor (Galr2b) in tilapia

Yaron Cohen[1], Krist Hausken[1], Yoav Bonfil[1], Michael Gutnick[2], and Berta Levavi-

Sivan[1]*

[1]Department of Animal Sciences and [2]Koret School of Veterinary Medicine, The Robert

H. Smith Faculty of Agriculture, Food, and Environment, Hebrew University of

Jerusalem, Rehovot 76100, Israel

* Corresponding author:

berta.sivan@mail.huji.ac.il

ORCID Identifier: 0000-0002-0183-9524

19   Short title: Spexin inhibits tilapia gonadotropin secretion

Abstract

Spexin (SPX) is a 14 amino acid peptide hormone that has pleiotropic functions across vertebrates, one of which is involvement in the brain-pituitary-gonad axis of fish. SPX(1) has been identified in each class of vertebrates, and a second SPX (named SPX2) has been found in some non-mammalian species. We have cloned two spexin paralogs, designated as Spx1a and Spx1b, from Nile tilapia (*Oreochromis niloticus*) that have varying tissue distribution patterns. Spx1b is a novel peptide only identified in cichlid fish, and is more closely related to Spx1 than Spx2 homologs as supported by phylogenetic, synteny, and functional analyses. Kisspeptin, Spx, and galanin (Gal) peptides and their corresponding kiss receptors and Gal receptors (Galrs), respectively, are evolutionarily related. Cloning of six tilapia Galrs (Galr1a, Galr1b, Galr2a, Galr2b, Galr type 1, and Galr type 2) and subsequent *in vitro* second-messenger reporter assays for $G\alpha_s$, $G\alpha_q$, and $G\alpha_i$ suggests that Gal and Spx activate Galr1a/Galr2a and Galr2b, respectively. A decrease in plasma follicle stimulating hormone and luteinizing hormone concentrations was observed with injections of Spx1a or Spx1b *in vivo*. Additionally, application of Spx1a to pituitary slices decreased the firing rate of LH cells, suggesting direct inhibition at the pituitary level. These data collectively suggest an inhibitory mechanism of action against the secretion of gonadotropins for a traditional and a novel spexin paralog in cichlid species.

Introduction

Spexin (SPX1; also termed neuropeptide Q) was identified first by computational methods [1], and then also by chemical methods [2] in humans. These computational methods have been attempted mostly on the basis of the characteristics of the prohormones from which active neuropeptides are processed. The mature peptide sequence contains 14 amino acids that are flanked by monobasic and dibasic proteolytic cleavage sites. The mature spexin peptide was found to be identical in all tetrapods and elephant shark, and differs in only one amino acid ($A^{13}T$) in piscine species. A paralog of spexin, termed Spx2, was later identified in non-mammalian species [3]. In mammals, Spx was found to participate in inducing stomach contraction [1], inhibiting adrenocortical cell proliferation[4], postnatal hypoxia response [5], cardiovascular and renal modulation[6], nociceptive response [7], fatty acid absorption and weight regulation [8], and diabetes [9]. In teleosts, functional studies of Spx1 mainly focused on its inhibitory role in the regulation of reproduction [10] and food intake [2] [11]. However, a recent study reported that *spx1* knock-out zebrafish exhibited normal reproductive capability but higher food intake than wild type fish, an effect mediated via increased expression of the appetite stimulant, agouti-related peptide AgRP1 [12].

The galanergic neurotransmission system is one of the newest described signaling systems. Today, the galanin family consists of galanin (Gal), galanin-like peptide (GalP), galanin-message associated peptide (GMAP), and alarin, and this family has been shown to be involved in a wide variety of biological and pathological functions [13]. Three different types of galanin receptors have been described so far in mammals: galanin

receptor 1, 2, and 3 (GALR1, GALR2, and GALR3) [14]. All of them are members of the G protein-coupled receptor (GPCR) family and act through stimulation of various second messenger systems. The biological activity of GALR1 and GALR3 stimulation is linked to the activity of adenylate cyclase (AC) and cyclic AMP (cAMP) production, and stimulation of GALR2 receptor results in phospholipase C (PLC) activity [14]. Recently, it was reported that SPX is a functional agonist for GALR2 and GALR3 in humans as well as Galr2a and Galr2b in zebrafish [15].

Reproduction is regulated by the hypothalamic-pituitary-gonadal (HPG) axis in all vertebrates. Hypothalamic axons secrete neuropeptides into the gonadotroph cells of the pituitary in order to mediate the expression and secretion of the gonadotropins follicle-stimulating hormone (FSH) and luteinizing hormone (LH) [16]. The first identified neuropeptide that regulates this function was gonadotropin-releasing hormone (GnRH). The hypophysiotrophic type of GnRH regulates gonadotropins by neuroglandular and neurovascular anatomical connections in zebrafish[17; 18]. Recently, a plethora of neuropeptides involved in the control of reproduction were investigated. Most of them are stimulatory neuropeptides, such as neurokinin B [19], neuropeptide Y [20], secretoneurin [21], galanin [22], agouti-related peptide [23], and melanocortin [24]. However, the study of neuropeptides that relay their signal through inhibitory pathways is more challenging, and hence are less studied. The most known inhibitory pathway in fish reproduction is the dopaminergic system [25]. In fish, as in mammals, dopamine D2 receptors transduce their signal through an inhibitory ($G\alpha_i$) signaling pathway [26]. $G\alpha_i$ signaling is involved in a variety of physiologic processes, including chemotaxis, neuro-transmission, proliferation, hormone secretion, and analgesia[27].

Nile tilapia is one of the top principal aquaculture species, and is a suitable experimental model fish for reproductive endocrinological research on Perciformes, which is the most recently evolved and largest group of teleost fish that includes many other target aquaculture species. Our objective was to identify spexin in tilapia and clarify its role as a regulator in the HPG axis. This was accomplished by cloning two spexin and six Galr sequences, and performing *in vitro* and *in vivo* studies.

Materials and Methods

Fish husbandry and transgenic lines

Sexually mature Nile tilapia (*Oreochromis niloticus,* Lake Manzala strain) were kept and bred in the fish facility unit at the Hebrew University in 500-L tanks at 26°C and with a 14/10 hour light/dark photoperiod regime. Fish were fed daily ad libitum with commercial fish pellets (Raanan fish feed, Israel).

We previously created transgenic tilapia lines by the adoption of a tol2 transposon-mediated approach and Gateway cloning technology [28]. In the current study we used transgenic tilapia in which red fluorescent protein (RFP) expression is driven by the tilapia LHβ promoter, thus labeling LH gonadotrophs. The tagRFP-CAAX cassette,

used in the current study directs the fluorescent protein to the cell membranes. The use of
tagRFP eliminates the aggregation problems associated with mCherry and results in a
more uniform labeling of the cells.

All experimental procedures were in compliance with the Animal Care and Use
guidelines of the Hebrew University and were approved by the local Administrative
Panel on Laboratory Animal Care.

110 Cloning and tissue distribution of tilapia spexin ligands and receptors

The putative spexin 1a and 1b gene sequences were initially identified using
tblastn against the tilapia genome (O_niloticus_wgs_v1, Genbank 354508) with zebrafish
spexin (XM_005164774) as input, and custom full coding sequence primers were
designed for PCR. Primers were designed against the nucleic acid sequences of the six
galanin receptors (Galr1a, Galr1b, Galr2a Galr2b, Galr type 1 and Galr type 2) obtained
from the Ensembl Genome Browser (http://www.ensembl.org) and the GenBank database
(http://www.ncbi.nlm.nih.gov/genbank). Putative transmembrane domains, glycosylation
sites and phosphorylation sites were identified using TMHMM v.2.0, NetNGlyc 1.0 [29],
and NetPhos 2.0 [30].

Total RNA was extracted from sexually mature female tilapia brain using TRIzol
reagent (Life Technologies), and 5 µg was used as template for cDNA synthesis using
Smart MMLV reverse transcriptase (Clontech). All cloning PCRs were performed with
an initial denaturation at 94 °C for 2.5 min, followed by 30 cycles of denaturation at 94
°C for 30s, annealing at each of the primers' specific Tm (Table 1) for 30s, and extension
at 72 °C for 90s, and a final extension at 72 °C for 10 min using Advantage2 polymerase
mix (Clontech); specific primers were designed for cloning the putative spexin ligands
and receptors (Table 1). The PCR products were ligated into pCRII-TOPO vector and
cloned into competent DH5α *E. coli* cells. Plasmid DNA was isolated from overnight
cultures by miniprep columns (QIAgen) and sequenced with T7 and SP6 primers.

Cloned tilapia spexin 1a and 1b sequences were submitted to GenBank under
accession numbers MN399812 and MN399813 respectively. Cloned tilapia galanin
receptors sequences were submitted to GenBank under accession numbers MN326828,
MN326829, MN326830, MN326831, MN614146, and MN614147 for Galr1a, Galr1b,
Galr2a, Galr2b, Galr type 1, and Galr type 2, respectively.

Tissue samples were collected from three mature male and female tilapia. Total
RNA was extracted from each of the following tissues: brain, pituitary gland, spleen,
gills, kidneys, muscles, fat, ovaries/testes, retina, heart, caudal and front intestines, and
liver. We dissected the brain into three parts, of which the anterior part contains the
olfactory bulbs and preoptic area, the midbrain contains the optic tectum and
hypothalamus, and the hindbrain contains the medulla oblongata and the cerebellum.
cDNA samples were prepared from 2µg of total RNA according to [31]. The tissue
expression patterns of Spx1a, Spx1b, Galr1a, Galr1b, Galr2a, and Galr2b in various

tissues were analyzed by qPCR with the primer sets described in Table 2. The cycling
parameters consisted of preincubation at 95 °C for 10 min followed by 45 cycles of
denaturation at 95°C for 10 s, annealing at 60°C for 30 s, and extension at 72°C for 10s,
followed by a melting curve analysis (95°C for 60s, 65°C for 60s, 97°C for 1s).

Phylogenetic and synteny analyses

All sequences used for spexin and galanin receptors were identified from NCBI
and Ensembl databases (Supplementary Table 1) and aligned by MUSCLE using Mega7.
Evolutionary analyses were conducted in MEGA7 [32]. The evolutionary history of
mature spexin peptide homologs was inferred using the Maximum Likelihood method
based on the JTT matrix-based model, and an additional frequency distribution (JTT+F)
was utilized for the receptors [33]. The bootstrap consensus trees were inferred from 500
replicates to represent the evolutionary history of the taxa analyzed [34]. The percentage
of replicate trees in which the associated taxa clustered together in the bootstrap test are
shown next to the branches. Initial tree(s) for the heuristic search were obtained
automatically by applying Neighbor-Join and BioNJ algorithms to a matrix of pairwise
distances estimated using a JTT model, and then selecting the topology with superior log
likelihood value. The analysis involved 34 and 77 amino acid sequences and there were a
total of 15 and 264 positions in the final datasets for spexins and Galrs, respectively.

We examined and compared the genomic environment around spexin1 and
spexin2 in zebrafish, Nile tilapia, and Burtoni. Cichlid orthologs were identified by tblasn
query using zebrafish genes in NCBI genome browsers, and organized 5' to 3' on their
respective chromosomes/linkage groups/scaffolds. Accession numbers can be found in
Supplementary Table 2.

Peptide synthesis

Tilapia spexin 1a (NWTPQAMLYLKGTQ-NH2), tilapia spexin 1b
(NWTSQAILYLKGAQ-NH2) and human galanin
(GWTLNSAGYLLGPHAVGNHRSFSNKNGLTS-NH2) were synthesized by GL
Biochem. Peptides were synthesized by the automated solid-phase method by applying
Fmoc active-ester chemistry, purified by HPLC to >95% purity and the carboxy terminus
of each peptide was amidated. The peptides were dissolved to the desired concentration
in fish saline (0.85% NaCl in DDW) for *in vivo* experiments.

In vivo effect of Spx1a and Spx1b on FSH and LH release

Adult female tilapia (body weight (BW)= 82.4±21.7g) were injected intraperitoneally with saline, tilapia spexin 1a, or spexin 1b at 10 μg/kg fish (n=8 fish per group). The fish were bled from the caudal blood vessels into heparinized syringes at 2, 4, 8, and 24 h after injection. Blood was centrifuged (3,200 rpm for 30min at 4°C) to obtain plasma samples, which were stored at −20°C until assayed. This time course is according to standard protocols used previously to test the effect of various hypothalamic neuropeptides on circulating levels of LH and FSH in tilapia [35] [36]. Six independent experiments were carried out for the *in vivo* studies.

ELISA for the measurement of tilapia FSH and LH

Plasma LH and FSH concentrations were measured by specific competitive ELISAs developed for tilapia [37] using primary antibodies against recombinant tilapia LHβ or FSHβ, respectively, and recombinant tilapia LHβα [38] or FSHβα [37] for the standard curves. The sensitivity was 2.43 and 1.52 ng/mL for LH and FSH, respectively. The inter-assay covariance was 14.8% and 12.5%, and the intra-assay variation was 7.2% and 8% for LH and FSH, respectively.

Determination of brain Spx and Galr expression in fed vs. fasted adult tilapia by qPCR

An *in vivo* experiment where sexually mature, male tilapia (n = 24, BW = 41.7 ± 1.64) were divided into two groups (one fed ad libitum and the other fasted for 26 days) was previously performed [39]. Total RNA from the brains were taken for real-time PCR expression analyses from the fed and starved groups. The RNA generated from this experiment was used as template for generated cDNA and performing qPCR analysis of brain Spx1a, Spx1b, Galr2a, and Galr2b in this study.

The qPCR cycling parameters consisted of preincubation at 95 °C for 10 min followed by 45 cycles of denaturation at 95°C for 10s, annealing at 60°C for 30s, and extension at 72°C for 10s, followed by a melting curve analysis (95°C for 60s, 65°C for 60s, 97°C for 1s). Reaction conditions were according to [40], and amplification was carried out in Lightcycler 96® (Roche Diagnostic International). Serial dilutions were prepared from a cDNA pool, and the efficiencies of the specific gene amplifications were compared by plotting Ct vs log (template concentration). A dissociation-curve analysis was run after each real-time experiment to confirm the presence of a single product. To control for false-positives, a no reverse transcriptase negative control was run for each template and primer pair. Data were interpreted by the comparative cycle threshold method using 18S as a reference gene [40].

.

Second messenger reporter assays

Transient transfection, cell procedures, and stimulation protocols were generally performed as described previously [26] [57]. Briefly, the entire coding sequence of tilapia

Galr1a, Galr1b, Galr2a, Galr2b, Galr type 1, and Galr type 2 were inserted into
pcDNA3.1 (Invitrogen). Co-transfection of the receptors (3 µg/plate for each Galr) and a
cAMP-response element-luciferase (CRE-Luc), serum response element (SRE-Luc;
Invitrogen), or Gqi5 reporter plasmid (3µg/plate) was carried out with TransIT-X2®
System (Mirus). The cells were serum-starved for 16 h, stimulated with various
stimulants for 6 h, and then harvested and analyzed according to [26] [57]. Experiments
were repeated a minimum of three times from independent transfections and each
treatment was performed in triplicate wells. EC_{50} values were calculated from dose
response curves by means of computerized non-linear curve fitting on baseline-corrected
(control) values using Prism version 6 software (GraphPad).

To confirm which of the novel receptors that were cloned in this study transduce
effects through the inhibitory Gi signaling pathway, we used a plasmid that contains a
chimeric G-protein, Gqi5, in which the five C-terminal amino acids of Gq were changed
to those of Gi1,2, (obtained from Addgene, Inc. Cambridge, MA, USA) and has been
described previously [41]. Gqi5 is activated by Gi-linked GPCRs, but couples to the
effector protein normally activated by Gq, phospholipase C. Since phospholipase C
produces IP_3 and actuates Ca^{2+} release, the inhibitory nature of Gi-linked receptors can be
observed as stimulation via the SRE-luc reporter [42].

Preparation of pituitary slices and electrophysiology

Pituitary slices were prepared according to [43] [44]. Tilapia of both sexes were
anesthetized with 2-phenoxyethanol (1 ml/ liter; Sigma–Aldrich Corp., St. Louis, MO)
and decapitated. Pituitaries were removed and kept at room temperature in gassed (5%
CO_2) Ringer's saline containing (in mM): 124 NaCl, 3 KCl, 2 $CaCl_2$, 2 $MgSO_4$, 1.25
NaH_2PO_4, 26 $NaHCO_3$, and 10 glucose (pH 7.35) up to 5 min before being transferred to
2% agarose low gelling temperature (Sigma) at 32°C. The agarose with the pituitary was
placed in the -20°C for a short time until the agarose solidified. The block was glued to
the stage of a vibratome (series 1000, Vibratome Co., St. Louis, MO), and 150 µM slices
were transferred to cooled gassed Ringer's saline.

For electrophysiological recordings, slices were gently transferred to a chamber
attached to the stage of an upright microscope (Axioskop FS, Zeiss, Oberkochen,
Germany) continuously superfused with Ringer's saline at room temperature. Endocrine
cells were viewed with a 40x, 0.8 numerical aperture, water immersion objective lens
(Olympus, Munich, Germany). Patch pipettes were pulled from borosilicate glass
capillaries (Hilgenberg, Maisfield, Germany) on a Narishige PP83 puller. Membrane
currents were recorded using an on-cell patch technique - cell-attached recording in
which a patch electrode is attached to the cell, but the membrane is not broken. The
standard pipette solution contained (in mM): 135 potassium gluconate, 2 $MgCl_2$, 1 $CaCl_2$,
11 EGTA, 3 ATP (magnesium salt), and 10 HEPES (potassium salt), pH 7.25. Only
fluorescently labeled cells from the pituitary were recorded. An Axoclamp-2B amplifier
(Axon Instruments, Union City, CA) was used in Bridge mode. Experiments were
controlled by USB-6341 data acquisition board (National Instruments, USA) and

WinWCP V5.3.7 (Strathclyde Electrophysiology Software, UK). Data were analyzed
using PClamp 10 (Axon Instruments) and Microcal Origin 6.0 software.

Statistical analyses

Data are presented as means ± SEM. All samples had equal variance, as
determined by an unequal variance test performed using JMP 7.0 software. The
significance of differences between group means of plasma gonadotropin levels were
determined by ANOVA, followed by Tukey's test using Graph-Pad Prism 5.01 software
(San Diego, CA). EC_{50} values of the receptors assays were calculated using log treatment
vs. luciferase intensity on a nonlinear regression curve using Prism.

Results

Cloning and tissue distribution of spexin1a, 1b, and gal receptors in tilapia

The cloned full coding sequence of tilapia spexin 1a and 1b were 363 and 315
base pairs and coded preprohormones of 120 and 104 amino acids, respectively. Both
mature tilapia spexin peptides were presumed to be 14 aa based on conserved peptide-
flanking monobasic and dibasic cleavage sites (Figure 1). The amino acid sequence of
Spx1b differs from that of orthologous piscine Spx1a at positions 4 (Pro to Ser), 7 (Met
to Ile), and 13 (Thr to Ala). Both Spx1a and Spx1b are likely amidated at their C-termini
due to the GRR motif. The amino acid sequence of Spx2 differs from that of Spx1 at
positions 3 (Gly vs Thr), 6 (Ser vs Ala), 13 (Arg vs Thr (piscine species) or Ala (shark
and tetrapods)), and 14 (Tyr or His vs Gln) (Figure 1).

Six different Galrs were found in the tilapia genome and their corresponding
coding regions were cloned and sequenced. According to their positions in the
phylogenetic tree, they were named Galr1a, Galr1b, Galr2a Galr2b, Galr type 1 and Galr
type 2. They encoded proteins of 350, 357, 325, 379, 344 and 379 amino acids for
Galr1a, Galr1b, Galr2a Galr2b, Galr type 1, and Galr type 2, respectively. All seven-
transmembrane receptors contain 1-2 potential sites for N-linked glycosylation (NXT/S)
in the N-terminal region, and 1-3 serine residues for potential protein kinase C
phosphorylation (SSR/SKK/SKR) (Supplementary Figure 3).

To shed light on the potential physiological roles of spexins/Galrs in tilapia, we
examined their mRNA tissue distribution by real-time PCR (Supplementary Figure 1).
The two spexins and their related receptors exhibit overlapping and distinctive patterns of
expression in various tissues. Spx1a expression was detected primarily in the midbrain
and ovary, whereas Spx1b was detected in pituitary, kidney, and all brain parts, but
mostly in the anterior brain. Spx1b expression levels were a few orders of magnitude
higher than that of Spx1a relative to 18S ribosomal subunit expression. The expression
patterns of Galr1a and Galr2a were similar, where both were detected in most of the
tissues with the highest expression observed in the kidney and head kidney. However,

Galr1b was detected mostly in the anterior gut and anterior brain, with lower expression found in all other tissues except the gills. High expression of Galr2b was detected in the anterior and midbrain relative to the other tissues, where virtually no expression was detected.

Evolutionary history of spexins and gal receptors

The phylogenetic analysis showed that the vertebrate SPX sequences fall into two distinct clades, with both cichlid spexins grouped with Spx1 (Figure 2A) and all Spx2 orthologs grouped together in a distinct clade. However, the second tilapia spexin paralog was more phylogenetically related with SPX1 than with SPX2 homologs. We also found that other cichlids, like zebra mbuna and Burtoni, contain two spexin paralogs that are more similar to orthologous spexin 1 sequences. Thus, we named the two cichlid spexins Spx1a and Spx1b.

The genomic organization of spexin paralogs between zebrafish, Nile tilapia, and Burtoni was determined. The genomic environment for *spx1a* was identified on zebrafish chromosome 4, Nile tilapia LG15, and Burtoni on an unplaced scaffold, and was highly conserved (Figure 2B). *Spx1b* was identified on Nile tilapia LG7 and an unplaced scaffold in Burtoni between *golt1bb* and *ldhbb*. *Spx2* was identified on zebrafish chromosome 25 between *gal* and *ldhd* (Figure 2C).

To determine the phylogenetic identity of the cloned Gal receptors we performed a phylogenetic analysis consisting of vertebrate Galr and Kissr sequences (Figure 3). The Galr and Kissr sequences formed two distinct clades. The Galr clade was subdivided into Galr1 and Galr2/3 clades, which, respectively, further formed distinct groups of Galr1a and Galr1b, and Galr3, Galr2a, and Galr2b. Tilapia Galr type 1 and 2 and medaka Galr type 1 belong to a sister group of Galr1.

Tilapia spexins suppress FSH and LH release *in vivo*

We next aimed to evaluate the biological effects of the two spexin paralogs on the release of tilapia gonadotropins. After a single intraperitoneal administration of 10 µg/kg fish Spx1a, both FSH and LH plasma levels significantly decreased after 120 min, compared with the levels observed at 0 min, and stayed at this lower level even after 24 h (Figure 4A, C). Injection of Spx1b caused a gradual decrease in plasma levels of FSH and LH, with lower levels shown after 4 h (Figure 4B, D). In the control groups, plasma gonadotropin levels did not change for the whole experimental period.

Fasting lowers Spexin and Galr2 mRNA expression in adult tilapia

Most of the documented knowledge relates spexin to metabolic processes or feeding behaviors. Even though our study primarily focused on reproduction, we aimed

335 to determine differences in spexin expression in adult tilapia that were fasted or fed for
336 26 days. Spx1a and Spx1b mRNA expression was significantly lower in the fasted vs. fed
337 fish. Galr2a expression was significantly lower in fasted fish than in fed fish, and Galr2b
338 expression not affected (Figure 5).

Activation and signaling of tilapia galanin receptors

All six cloned, full-length tilapia Galrs were sub-cloned into the pcDNA3.1
expression vector and co-transfected into COS7 cells with a luciferase plasmid containing
either cAMP response element (CRE), serum responsive element (SRE), or Gqi5 to
evaluate their signal transduction pathway(s) upon treatment with tilapia Spx1a, Spx1b,
or human galanin (Figure 6). Since each Galr subtype is coupled to different G proteins
($G\alpha_i$ for GalR1 and GalR3; $G\alpha_{q/11}$ for GalR2) [14], we tested all the receptors using three
different signal transduction pathways: CRE-luc for the activation of PKA/cAMP, SRE-
luc for the activation of PKC/Ca^{2+}, and Gqi5/SRE-luc for further verification of
inhibition through $G\alpha_i$. The 5 aa at the carboxyl-terminus of Gqi5 are sufficient to enable
interaction with $G\alpha_i$-coupled receptors [41]. Ligand stimulation of a $G\alpha_i$-coupled receptor
is expected to trigger the following cascade: Gqi5, phospholipase C, protein kinase C,
SRE-luciferase transcription, luciferase activity, light emission. For verification of the
activation of Gqi5 trough $G\alpha_i$, we used the tilapia dopamine D2 receptor [26][57]. A
dose-dependent increase in luciferase activity confirmed that tilapia D2 receptor relayed
its activity trough the inhibitory Gi (Supplementary Figure 4).

Tilapia Spxs or hGAL did not increase CRE-luc or SRE-luc levels in cells
expressing Galr1a or Galr1b, but hGAL activated Gqi5 signaling via Galr1a. Increasing
concentrations of Gal resulted in significant increases in CRE-luc, SRE-luc, and Gqi5
activity in cells expressing Galr2a. Stimulation by high concentrations of both tilapia Spx
paralogs resulted in an increase in Gqi5 in cells expressing Galr2a. Exposure to
increasing concentrations of either Spx resulted in a dose-dependent increase in SRE-luc,
CRE-luc, and Gqi5 activity in cells expressing Galr2b. Both Spx1a and Spx1b were also
very efficient in transcription of SRE-luc in cells expressing Gqi5, suggesting that Galr2b
relays its signal through Gi ($EC_{50}=1nM$). The EC_{50} values of Spxs and Gal for each
receptor are summarized in Supplementary Table 3.

Effect of spexin1a on the electrophysiological activity of pituitary LH cells

Following the results that spexins activate Galr2a and 2b, of which the former was
also found to be expressed in the pituitary of tilapia, we aimed to examine whether spexin
affects LH gonadotroph electrical activity. To address this question, we used an
electrophysiological method on disconnected pituitary slices to keep the cells in their
natural organization, but without signals from the brain. We measured the frequency of
action potentials before, during, and after application of 100nm Spx1a. Exposure to
Spx1a decreased the spike rate, which slowly increased back to basal frequencies when
spexin was washed out from the recording chamber (Figure 7).

Discussion

Spexin is a highly conserved peptide hormone that may have pleiotropic functions in different vertebrate species. Cloning of two spexin paralogs and six galanin receptors permitted phylogenetic, synteny, and functional analyses in Nile tilapia, supporting the involvement of the spexin/galanin receptor system in fish reproduction and metabolism. Evolutionary analyses support that cichlid species possess a novel form of Spx1, named Spx1b, but not Spx2 like in some other vertebrates. Demonstration of Spx1a and Spx1b involvement in reproduction was shown *in vivo* and *in vitro*, and starvation limited the expression of both paralogs. Despite their differential tissue distribution and expression profiles, both Spx1a and Spx1b had similar biological effects, which are likely mediated through galanin receptor 2b.

Phylogenetic and synteny analyses coupled with vertebrate ancestral chromosome (VAC) reconstruction supported that *kiss*, *gal*, and *spx* genes were located adjacent to each other on VAC D and individually arose by local tandem duplication prior to the first whole genome duplication event [15]. This genomic proximity was conserved after two rounds of whole genome duplication (1R, 2R) and is observed in modern vertebrate genomes, where *spx1* is found near *kiss2*, and *spx2* is found near *gal*. Our synteny analysis supported this organization for Nile tilapia and Burtoni *spx1*, but not *spx2*. We initially had thought that the spexin-like gene flanked by *golt1bb* and *ldhbb* in tilapia was *spx2*, but this did not correspond with the genomic organization of zebrafish *spx2*, which is flanked by *gal* and *ldhd*. The genomic distance between Nile tilapia *gal* and *ldhd* is >34 million base pairs, and *spx1b* lies outside of this syntenic block between *golt1bb* and *ldhbb*. No zebrafish spexin-like gene was found between *golt1bb* and *ldhbb*, suggesting that this tilapia and Burtoni spexin paralog is not *spx2*. Additionally, Spx1b has 85-93% (12-13/14aa) sequence conservation across all vertebrate Spx1 peptides, but only 57-71% (8-10/14aa) conservation with the variable and meager Spx2 sequences. Phylogenetic analysis of mature spexin peptide sequences shows that cichlid Spx1b forms a sister clade to vertebrate Spx1, and is not placed with other teleost Spx1 homologs. Due to the highly conserved nature of short peptide sequences and a 20% divergence in amino acid sequence identity from Spx1, we propose that Spx1b is a novel spexin peptide found only in cichlid species.

Spexin has been implicated in regulating the hypothalamic-pituitary-gonad (HPG) axis in some fish species. Our *in vivo* experiments support that not only Spx1a, but also the novel form, Spx1b, inhibits LH and FSH release in tilapia. The first report on fish spexin in zebrafish and goldfish identified high and dynamic expression in the brain throughout the reproductive stages [10]. Spexin inhibited LH release in goldfish pituitary cultures and *in vivo*, and an estradiol feedback mechanism on hypothalamic spexin expression was observed. Ovariectomized goldfish had increased hypothalamic spexin expression, and estrogen replacement returned expression to basal levels [10]. A similar effect was seen in the spotted scat, where estradiol decreased spexin expression in a dose-dependent manner *in vitro* [45]. However, zebrafish $spx^{-/-}$ knockouts displayed normal reproductive phenotypes, so spexin may only play a direct role in regulating the reproductive axis in certain fish species [12]. With that being said, knockout models of reproductive hormones in fish species tend to reproduce normally (reviewed in [46]).

There are few reports on the role of spexin in reproduction, all of which have been carried out in fish; additional studies in non-piscine species are required to determine if spexin has a functionally conserved role along the HPG axis.

A distinctive feature of the teleost pituitary gland is that LH and FSH are synthesized by different gonadotrope cells, which suggests a functional and differential regulation of gonadotropin release mechanisms. The cells of the teleost pituitary are distinctly organized and receive hypothalamic signals directly by nerve innervation as well as via neurovasculature [18]. Teleost LH cells form dense networks throughout the pars distalis of the adenohypophysis, and FSH cells form small clusters that are more broadly distributed throughout the pars distalis [17]. LH cells, but not FSH cells, are functionally coupled, as shown by perfusion of pituitary fragments exposed to GnRH with gap-junction blockers and by a patch-clamp technique [17]. These organizational characteristics of the teleost pituitary complicate our understanding about the various ways by which the gonadotropes decode stimulatory and inhibitory inputs. The principal stimulatory and inhibitory factors for LH release are gonadotropin-releasing hormone (GnRH) and dopamine (DA), respectively, but less is known about FSH release. LH and FSH cells are regulated by more than 20 neurohormones, but only a small portion of them function as inhibitory factors [46]. In fish, the dopamine receptor (D2-R) is expressed on LH cells [47] and on GnRH neurons [25], suggesting that direct and indirect mechanisms of LH release inhibition exist. Gonadotrophs are excitable cells, and action potential frequency of LH and FSH cells were shown to be influenced by GnRH exposure [48]. We performed electrophysiological measurements on LH cells in pituitary slices and observed that Spx1a decreases the action potential frequency, suggesting that spexin decouples the LH gonadotrope functional network directly.

Spexin mRNA expression levels may also be correlated with reproductive development. Increases in LH and FSH mRNA expression is seasonal and corresponds to gonadal development [49] [50]. In goldfish, hypothalamic Spx1 expression significantly decreased as the breeding season progressed and GSI increased [10], potentially contributing to progressive LH release. A similar decrease was seen in the orange-spotted grouper over the course of oogenesis, where the highest hypothalamic Spx1 expression was observed prior to oocyte primary growth [11], and in the spotted scat, expression significantly decreased over the course of vitellogenesis [45]. In zebrafish, however, Spx1 expression in the brain gradually increased from the ovarian primary growth stage, peaked in early-mid vitellogenesis, then returned to primary growth levels during final maturation [10]. Given that Spx1 is negatively regulated by estrogen [10], which increases in circulation until final oocyte maturation, it seems reasonable that Spx expression correlates with reproductive development. Additionally, Spx1 treatment affects the expression of other reproductive factors, such as increasing gonadotropin inhibitory hormone (GnIH) and GnRH-III expression and decreasing GpHα and FSHβ expression in a sole [51].

Spexin has wide tissue distribution patterns in tetrapods and fish, suggesting that spexin has pleiotropic biological consequences in feeding behavior, metabolism, nociception, cardiovascular function, muscle motility, stress, depression, and anxiety [3]. The localization of spexins at the cellular level inform these functions. In tilapia, the two

spexin paralogs exhibited wide differential tissue distribution patterns, where Spx1a was primarily expressed in the midbrain (containing the hypothalamus in our preparations), and Spx1b was primarily expressed in the anterior brain, which contains the preoptic area. Similarly, in a detailed report of spexin neuron circuitry in zebrafish, Spx1 expression was restricted to the midbrain tegmentum and the hindbrain and Spx2 expression was found in the preoptic area, however neither was detected in the hypothalamus [52]. In goldfish, Spx1 was identified in the hypothalamus, thalamus, and medial longitudinal fasciculus, however immunoreactivity was detected using heterologous antibodies that may have recognized both Spx1 and Spx2 [10]. Even though our tissue distribution analysis is crude and unable to implicate any detailed biological function from expression, these data support the conservation of Spx1 expression, which differs from Spx1b localization. In order to determine if the neuronal circuits of spexins are conserved in fish, methodologies for determining specific spexin localization (IHC with custom antibodies, fluorescent *in situ* hybridization, etc.) are needed from additional species.

Spexin plays a role in feeding behavior and metabolism in fish and mammals (reviewed in [3]). In the orange-spotted grouper [11], half-smooth tongue sole [45], and spotted scat [45], starvation increased Spx1 expression. We observed a decrease in brain Spx1a, Spx1b, and Galr2a expression in fasted adult tilapia. Brain Spx1 expression was also decreased in unfed Ya-fish [53]. In goldfish, Spx1 injections decreased surface feeding and increased food rejection, and increased circulating Spx1 and brain and liver Spx1 mRNA expression after feeding [2] [54]. This effect was also observed in $spx1^{-/-}$ zebrafish, which had higher food intake compared to wildtype fish, and Spx1 administration suppressed agouti-related peptide (AgRP) expression [12]. In support of this mechanism of satiety control, central administration of Spx1 in goldfish caused a decrease in expression of appetite stimulants (NPY, AgRP, and apelin) and increased expression of anorexigenic factors (proopiomelanocortin (POMC), cocaine- and amphetamine-regulated transcript (CART), cholecystokinin (CCK), melanin-concentrating hormone (MCH), and corticotropin-releasing hormone (CRH)) in different parts of the brain [2]. Furthermore, intracerebroventricular injection of Spx1 inhibited feeding behavior induced by NPY and orexin. In addition to appetite factors, insulin was shown to increase circulating levels of Spx1 and its expression in the telencephalon, hypothalamus, and optic tectum [54]. Therefore, local and central Spx1 appears to be an integral part of appetite control and energy homeostasis in fish.

Spexin mediates its effects by activating an inhibitory G-protein ($G\alpha_i$) via galanin receptor (GalR) 2/3. Like the kisspeptin, galanin, and spexin peptides, the kisspeptin and galanin receptors are related. Ancestral forms of KissR, GalR1, and GalR2/3 were identified on different VACs with 1R and 2R expanding each clade, giving rise to four KissR (KissR1, 3, 2, 4), two GalR1 (1a, 1b), two GalR2 (2a, 2b) and GalR3 paralogs [15]. Our phylogenetic analysis supports that of Kim *et al.* (2014), showing that GalRs are divided into two major clades, namely GalR1 and GalR2/3, with KissRs forming a distinctive sister clade. We cloned two additional Galrs based on predicted NCBI sequences named Galr type 1 and type 2, which formed a sister group to GalR1 sequences, but were not activated by galanin or spexin (Supplementary Figure 2). It has been previously shown that the cognate receptor(s) for Spx1/Spx2 are GalR2/3, whereas

GalR1 is the cognate receptor for galanin; teleosts do not possess GalR3 [15]. Our second
messenger reporter assays revealed that tilapia Spx1a and Spx1b activated $G\alpha_s$ (via
CRE), $G\alpha_q$ (via SRE), and $G\alpha_i$ (via Gqi5) signaling pathways through Galr2b, but not
Galr1a or Galr1b. Galanin was more efficacious than Spx1a or Spx1b in activating $G\alpha_s$
and $G\alpha_i$ via Galr2a, but did not activate $G\alpha_q$ (Figure 8). Determining inhibitory effects via
$G\alpha_i$ is challenging, so we utilized a reporter plasmid that permits $G\alpha_i$-coupled receptors to
stimulate PLC [41]. Human, Xenopus, and zebrafish GalRs were activated in a similar
manner using an alternative expression system [15]. GalR2 displays differential signaling
preferences to either Gal or Spx, which induce different conformational changes in the
receptor and bias intracellular signaling cascades [55]. Spx quickly dissociated after G-
protein signaling, whereas Gal binding was more stable and promoted arrestin-dependent
receptor internalization. It has been shown that galanin and spexin have an inverse
relationship in regards to LH release (reviewed in [15]), perhaps due in part to the
receptor displaying biased agonism in order to decipher endogenous hormone signaling.

We have shown evolutionary and functional evidence that cichlid fish possess two
paralogs of Spx1 but not Spx2, and that tilapia Spx1a and Spx1b can inhibit LH and FSH
secretion via Galr2b. Why cichlids have evolved a second form of Spx1 and lost Spx2
remains unknown. One in ten teleosts are cichlids, making them the most species-rich
group of vertebrates and excellent genetic models of adaptive radiation [56].
Additionally, Nile tilapia is a major globally aquacultured species. Further research into
the potential for manipulation of growth and reproductive processes with spexin or
galanin receptor agonists/antagonists is warranted.

Acknowledgements

532 The research was funded by the Israel Science Foundation (ISF) no. 1540/17.
533 KNH was supported by BARD, the United States - Israel Binational Agricultural
534 Research and Development Fund, Vaadia-BARD Postdoctoral Fellowship Award No.
535 FU-561-2017.

536 We would like to thank Dr. Michal Shpilman for her technical assistance.

Author Contributions Statement

539
540 YC, KH, and BS prepared and wrote the manuscript. YC, KH, and YB curated
541 and analyzed all data. MG and BS acquired funding, and supervised and conceptualized
542 the manuscript. All authors contributed to manuscript revision, read and approved the
543 submitted version.
544

Conflict of Interest Statement

546   The authors declare no conflicts of interest.
547
548   **Contribution to the Field Statement**
549
Reproduction is regulated by a hierarchical hormonal signaling cascade via the
brain-pituitary-gonadal axis in all vertebrates. Many neurohormones release from the
brain have been identified to stimulate the release of pituitary gonadotropins (follicle
stimulating hormone and luteinizing hormone), which stimulate reproductive
developmental processes (I.E. gametogenesis) in the gonads. A plethora of neuropeptides
involved in the control of reproduction have been investigated over the last decades, and
most of them are stimulatory neuropeptides. However, the study of neuropeptides that
relay their signal through inhibitory pathways is more challenging. Spexin is a highly
conserved 14 amino acid peptide hormone that may have pleiotropic functions in
different vertebrate species, but was shown to inhibit pituitary gonadotropin release in
fish. Our work supports that the conserved spexin peptide, as well as a novel cichlid-
specific spexin, directly inhibits gonadotropin release in tilapia. Understanding the
stimulatory and inhibitory factors that regulate fish reproduction is critical to aquaculture
success, as well as our basic understanding of comparative reproductive endocrinology in
vertebrates.

Figure legends

Figure 1. Multiple sequence alignment of mature vertebrate spexin1, spexin2, and
galanin peptides with flanking monobasic or dibasic cleavage motifs. The last three
amino acids of spexin sequences (GRR) were aligned to the end of the galanin sequences,
but manually moved for brevity. Cichlid spexin 1a and 1b sequences are labeled to
differentiate them, and spexin1b are boxed.

Figure 2. Evolutionary history of spexin sequences. A. The phylogenetic tree of
mature vertebrate spexin paralogs was generated by the maximum likelihood method and
the percentage of replicate trees in which the associated taxa clustered together in the
bootstrap test (500 replicates) are shown next to the branches; bootstrap scores lower than
50% are condensed. **B.** The genomic environment of *spx1* and **C.** *spx2* in zebrafish, Nile
tilapia, and Burtoni. Spx1a and spx1b are surrounded in red boxes (broken to emphasize
lack of a spx-related gene in zebrafish) and Spx2 has a bolded black box. Chromosomes
(Ch)/ linkage groups (LG)/ scaffolds (scaf.) are organized 5'-3' and genomic location (in
million (M) or thousand (K) base pairs) is indicated above and below each column. Note
that the range of zebrafish Ch25 and tilapia LG7 are broken for brevity.

Figure 3. Phylogenetic analysis of vertebrate galanin and kisspeptin receptors. The
phylogenetic tree was generated by the maximum likelihood method and the percentage
of replicate trees in which the associated taxa clustered together in the bootstrap test (500
replicates) are shown next to the branches. This cladogram was rooted with human

melanocortin receptors (MCR). A solid line subdivides the Galr and KissR clades, and subsequent subdivisions within these clades are separated by dashed lines.

Figure 4. Spexins inhibit LH and FSH *in vivo*. Adult female tilapia were injected with Spx1a, Spx1b, or saline and blood was sampled up to 24 hours. Plasma FSH (**A,B**) and LH (**C,D**) were determined by ELISA. Control values are shown in black, and spexin treatment values are shown in blue. N=8 fish per group. Data are represented as mean FSH or LH (ng/ml) ± SEM.

Figure 5. Brain spexin and Galr mRNA expression in response to fasting. Sexually mature tilapia were fed *ad libitum* (Fed), or fasted (Fast) for 26 days. Results were normalized to the amount of *18S* by the comparative cycle threshold method. N = 24. Values are given as mean relative expression ± SEM. Different letters indicate statistically significant differences between groups ($p < 0.05$).

Figure 6. *In vitro* reporter assays for tilapia Galanin receptors. Tilapia Galrs were cotransfected with either CRE-luc (**A-D**), SRE-Luc (**E-H**), or Gqi5 (**I-L**) into Cos7 cells and treated with either Spx1a (black squares), Spx1b (red triangles), or human galanin (Gal; blue upside-down triangles). Data are represented as mean ± SEM fold-change in luciferase activity over basal (no treatment).

Figure 7. Effect of spexin1a on firing rate of LH cells in mature tilapia. Transgenic tilapia (LH-RFP) pituitaries were dissected for electrophysiological recording, and a cell-attached configuration was used to monitor action potentials without breaking the cell membrane. The pituitary slice was briefly exposed to Spx1a during the recording (blue bar). **A.** A graphic representation of firing-rate before (**B**), during (**C**), and after spexin application. Each dot represents a action potential along the time scale (X-axis) and the instantaneous frequency from the action potential before (Y-axis). Note that the firing-rate decreased after Spx1a was applied, and after a few minutes the spike rate slowly rises back to the control level. **D.** Two spikes with an example of the firing-rate measurement.

Figure 8. A schematic representation of the Spx/Galr system in tilapia.

Tables

Table 1. Gene-specific primers used for the identification of tilapia Spx peptides and Galrs. The primers sequences and their melting temperatures used for the PCR are presented.

Primer	5' to 3' sequence	Direction	Gene	Tm (°C)

1	ATGAACTTGACAGACTCTGTTTGGCT	Forward	Galr1a	64
2	TCAAACATTTGTGCAGTTAGTTGAGG	Reverse		
3	ATGGCGCTGTCGGAAAAC	Forward	Galr1b	63
4	TCATATGTTGGTGGTGGAGTGT	Reverse		
5	ATGAATGACTCTGTGTCTAACTCCTCC	Forward	Galr2a	62
6	TTAATTGGACACATCGATGCTG	Reverse		
7	ATGTCAGACTTTGAGGATTTCACCT	Forward	Galr2b	64
8	TCAAGTCTGCTGCTGCTGAAAG	Reverse		
9	ATGGAGAGTTCGGGGCAG	Forward	Galr type 1	65
10	TCAGTTTTGGCTTCTCCCTGAG	Reverse		
11	ATGGCTGTTCCCAACACCTATG	Forward	Galr type 2	66
12	CTAAAGGCTGTCTGTGGTAACGCT	Reverse		
13	ATGAAGGGTTTGAAGACCGTCACA	Forward	Spx1a	60

14	TTAGAAATATTCCCTCTTCCACACTGGTA	Reverse		59
15	GCAGGAGAGCACAACGCTTAAAG	Forward	Spx1b	65
16	GCTCTGATGAATGTTCACAAGTGG	Reverse		65

621 **Table 2. Gene-specific primers used for real-time PCR**. The primer sequences and
622 PCR efficiency ($10^{-1/slope}$) are shown.

Primer	5' to 3' sequence	Direction	Gene	Efficiency
1	GAAAAAGACTGCTCAGACGGTG	Forward	Galr 1a	1.82
2	GTTTAGCGGGAAGGTCCCAA	Reverse		
3	TGTTGGGGTACCTGCTACCG	Forward	Galr 1b	1.94
4	GAGGACCGTCTGAGCCGTT	Reverse		
5	TGGACAGGTATTTAGCCATTTGC	Forward	Galr 2a	2.15
6	ACAACAGTGCCAGCCAGGT	Reverse		
7	GTGTCTGTTGACAGGTATCT	Forward	Galr 2b	1.89

8	GTCATAGTAGCTGAGATATGG	Reverse		
9	TCTCCCAGTCATGGAGCACG	Forward	Spx1a	1.86
10	TTCTTCCCTGGGTG	Reverse		
11	TCCACAGAGGAGAAACTGG	Forward	Spx1b	1.96
12	CTGTTCTGCTTGTGAGTCCC	Reverse		
13	GCACCACCACCCACAGAATC	Forward	18S	1.97
14	CGACCATAAACGATGCCAACTAG	Reverse		

REFERENCES

[1] O. Mirabeau, E. Perlas, C. Severini, E. Audero, O. Gascuel, R. Possenti, E. Birney, N. Rosenthal, and C. Gross, Identification of novel peptide hormones in the human proteome by hidden Markov model screening. Genome Res 17 (2007) 320-7.

[2] M.K. Wong, K.H. Sze, T. Chen, C.K. Cho, H.C. Law, I.K. Chu, and A.O. Wong, Goldfish spexin: solution structure and novel function as a satiety factor in feeding control. Am J Physiol Endocrinol Metab 305 (2013) E348-66.

[3] C.H. Lim, M.Y.M. Lee, T. Soga, and I. Parhar, Evolution of Structural and Functional Diversity of Spexin in Mammalian and Non-mammalian Vertebrate Species. Front Endocrinol (Lausanne) 10 (2019) 379.

[4] M. Rucinski, A. Porzionato, A. Ziolkowska, M. Szyszka, V. Macchi, R. De Caro, and L.K. Malendowicz, Expression of the spexin gene in the rat adrenal gland and evidences suggesting that spexin inhibits adrenocortical cell proliferation. Peptides 31 (2010) 676-82.

[5] A. Porzionato, M. Rucinski, V. Macchi, C. Stecco, G. Sarasin, M.M. Sfriso, C. Di Giulio, L.K. Malendowicz, and R. De Caro, Spexin is expressed in the carotid body and is upregulated by postnatal hyperoxia exposure. Adv Exp Med Biol 758 (2012) 207-13.

[6] L. Toll, T.V. Khroyan, K. Sonmez, A. Ozawa, I. Lindberg, J.P. McLaughlin, S.O. Eans, A.A. Shahien, and D.R. Kapusta, Peptides derived from the prohormone proNPQ/spexin are

potent central modulators of cardiovascular and renal function and nociception. FASEB J
26 (2012) 947-54.

[7] T. Pirzeh L, and a. M., Effect of Intra Hippocampal CA1 Injection of Spexin on Pain Sensitivity
in Female Rat. . Bull. Env. Pharmacol. Life Sci. 3 (2014) 71-74.

[8] J.L. Walewski, F. Ge, H.t. Lobdell, N. Levin, G.J. Schwartz, J.R. Vasselli, A. Pomp, G. Dakin, and
P.D. Berk, Spexin is a novel human peptide that reduces adipocyte uptake of long chain
fatty acids and causes weight loss in rodents with diet-induced obesity. Obesity (Silver
Spring) 22 (2014) 1643-52.

[9] A. Karaca, F. Bakar-Ates, and N. Ersoz-Gulcelik, Decreased Spexin Levels in Patients with Type
1 and Type 2 Diabetes. Med Princ Pract 27 (2018) 549-554.

[10] Y. Liu, S. Li, X. Qi, W. Zhou, X. Liu, H. Lin, Y. Zhang, and C.H. Cheng, A novel neuropeptide in
suppressing luteinizing hormone release in goldfish, Carassius auratus. Mol Cell
Endocrinol 374 (2013) 65-72.

[11] S. Li, Q. Liu, L. Xiao, H. Chen, G. Li, Y. Zhang, and H. Lin, Molecular cloning and functional
characterization of spexin in orange-spotted grouper (Epinephelus coioides). Comp
Biochem Physiol B Biochem Mol Biol 196-197 (2016) 85-91.

[12] B. Zheng, S. Li, Y. Liu, Y. Li, H. Chen, H. Tang, X. Liu, H. Lin, Y. Zhang, and C.H.K. Cheng, Spexin
Suppress Food Intake in Zebrafish: Evidence from Gene Knockout Study. Sci Rep 7 (2017)
14643.

[13] K.E. Webling, J. Runesson, T. Bartfai, and U. Langel, Galanin receptors and ligands. Front
Endocrinol (Lausanne) 3 (2012) 146.

[14] J. Sipkova, P. Sida, N. Kasprikova, I. Kramarikova, S. Hynie, and V. Klenerova, Effect of Stress
on the Expression of Galanin Receptors in Rat Heart. Folia Biol (Praha) 63 (2017) 98-104.

[15] D.K. Kim, S. Yun, G.H. Son, J.I. Hwang, C.R. Park, J.I. Kim, K. Kim, H. Vaudry, and J.Y. Seong,
Coevolution of the spexin/galanin/kisspeptin family: Spexin activates galanin receptor
type II and III. Endocrinology 155 (2014) 1864-73.

[16] B. Levavi-Sivan, J. Bogerd, E.L. Mananos, A. Gomez, and J.J. Lareyre, Perspectives on fish
gonadotropins and their receptors. Gen Comp Endocrinol 165 (2010) 412-437.

[17] M. Golan, A.O. Martin, P. Mollard, and B. Levavi-Sivan, Anatomical and functional
gonadotrope networks in the teleost pituitary. Scientific Reports 6 (2016) 23777-Article
No.: 23777.

[18] M. Golan, E. Zelinger, Y. Zohar, and B. Levavi-Sivan, Architecture of GnRH-Gonadotrope-
Vasculature Reveals a Dual Mode of Gonadotropin Regulation in Fish. Endocrinology 156
(2015) 4163-4173.

[19] J. Biran, O. Palevitch, S. Ben-Dor, and B. Levavi Sivan, Neurokinin B and neurokinin B
receptor: a novel system involved in controlling fish reproduction. PNAS 109 (2012)
10269-10274.

[20] Y. Zohar, J.A. Munoz-Cueto, A. Elizur, and O. Kah, Neuroendocrinology of reproduction in
teleost fish. General and Comparative Endocrinology 165 (2010) 438-455.

[21] E. Zhao, J.R. McNeilly, A.S. McNeilly, R. Fischer-Colbrie, A. Basak, J.Y. Seong, and V.L.
Trudeau, Secretoneurin stimulates the production and release of luteinizing hormone in
mouse LβT2 gonadotropin cells. Am J Physiol Endocrinol Metab 301 (2011) E288-97.

[22] P. Pinto, Z. Velez, C. Sousa, S. Santos, A. Andrade, M.V. Alvarado, A. Felip, S. Zanuy, and
A.V.M. Canario, Responsiveness of pituitary to galanin throughout the reproductive
cycle of male European sea bass (Dicentrarchus labrax). Gen Comp Endocrinol 250
(2017) 152-161.

[23] C. Zhang, P.M. Forlano, and R.D. Cone, AgRP and POMC neurons are hypophysiotropic and
coordinately regulate multiple endocrine axes in a larval teleost. Cell Metab 15 (2012)
256-64.
[24] D.N. Jiang, J.T. Li, Y.X. Tao, H.P. Chen, S.P. Deng, C.H. Zhu, and G.L. Li, Effects of
melanocortin-4 receptor agonists and antagonists on expression of genes related to
reproduction in spotted scat, Scatophagus argus. J Comp Physiol B 187 (2017) 603-612.
[25] S. Dufour, M.E. Sebert, F.A. Weltzien, K. Rousseau, and C. Pasqualini, Neuroendocrine
control by dopamine of teleost reproduction. Journal of Fish Biology 76 (2010) 129-160.
[26] B. Levavi-Sivan, J. Aizen, and A. Avitan, Cloning, characterization and expression of the D2
dopamine receptor from the tilapia pituitary. Molecular and Cellular Endocrinology 236
(2005) 17-30.
[27] S. Gaillard, L. Lo Re, A. Mantilleri, R. Hepp, L. Urien, P. Malapert, S. Alonso, M. Deage, C.
Kambrun, M. Landry, S.A. Low, A. Alloui, B. Lambolez, G. Scherrer, Y. Le Feuvre, E.
Bourinet, and A. Moqrich, GINIP, a Galphai-interacting protein, functions as a key
modulator of peripheral GABAB receptor-mediated analgesia. Neuron 84 (2014) 123-
136.
[28] M. Golan, and B. Levavi-Sivan, Social Dominance in Tilapia is Associated with Gonadotroph
Hyperplasia. General and Comparative Endocrinology 192 (2013) 126-135.
[29] N. Blom, T. Sicheritz-Pontén, R. Gupta, S. Gammeltoft, and S. Brunak, Prediction of post-
translational glycosylation and phosphorylation of proteins from the amino acid
sequence. Proteomics 4 (2004) 1633-1649.
[30] N. Blom, S. Gammeltoft, and S. Brunak, Sequence and structure-based prediction of
eukaryotic protein phosphorylation sites1 1Edited by F. E. Cohen. Journal of Molecular
Biology 294 (1999) 1351-1362.
[31] J. Biran, S. Ben-Dor, and B. Levavi-Sivan, Molecular identification and functional
characterization of the kisspeptin/kisspeptin receptor system in lower vertebrates. Biol
Reprod 79 (2008) 776-786.
[32] S. Kumar, G. Stecher, and K. Tamura, MEGA7: Molecular Evolutionary Genetics Analysis
Version 7.0 for Bigger Datasets. Mol Biol Evol 33 (2016) 1870-4.
[33] D.T. Jones, W.R. Taylor, and J.M. Thornton, The rapid generation of mutation data matrices
from protein sequences. Comput Appl Biosci 8 (1992) 275-82.
[34] J. Felsenstein, Confidence Limits on Phylogenies: An Approach Using the Bootstrap.
Evolution 39 (1985) 783-791.
[35] N. Mizrahi, C. Gilon, I. Atre, S. Ogawa, I.S. Parhar, and B. Levavi-Sivan, Deciphering Direct
and Indirect Effects of Neurokinin B and GnRH in the Brain-Pituitary Axis of Tilapia.
Frontiers in Endocrinology 10 (2019).
[36] J. Biran, M. Golan, N. Mizrahi, S. Ogawa, I.S. Parhar, and B. Levavi Sivan, LPXRFa, the piscine
prtholog of GnIH, and LPXRF-receptor positively regulate gonadotropin secretion in
tilapia (Oreochromis niloticus). Endocrinology 155 (2014) 4391-401.
[37] J. Aizen, H. Kasuto, and B. Levavi-Sivan, Development of specific enzyme-linked
immunosorbent assay for determining LH and FSH levels in tilapia, using recombinant
gonadotropins. Gen Comp Endocrinol 153 (2007) 323-332.
[38] H. Kasuto, and B. Levavi-Sivan, Production of biologically active tethered tilapia LHβα by the
methylotrophic yeast Pichia pastoris. Gen Comp Endocrinol 140 (2005).
[39] M. Shpilman, L. Hollander-Cohen, T. Ventura, A. Gertler, and B. Levavi-Sivan, Production,
gene structure and characterization of two orthologs of leptin and a leptin receptor in
tilapia. Gen Comp Endocrinol 207 (2014) 74-85.

[40] J. Biran, O. Palevitch, S. Ben-Dor, and B. Levavi-Sivan, Neurokinin Bs and neurokinin B
receptors in zebrafish-potential role in controlling fish reproduction. Proceedings of the
National Academy of Sciences of the United States of America 109 (2012) 10269-74.
[41] B.R. Conklin, Z. Farfel, K.D. Lustig, D. Julius, and H.R. Bourne, Substitution of three amino
acids switches receptor specificity of Gq alpha to that of Gi alpha. Nature 363 (1993)
274-6.
[42] K. Coward, and N.R. Bromage, Spawning frequency, fecundity, egg size and ovarian
histology in groups of Tilapia zillii maintained upon two distinct food ration sizes from
first-feeding to sexual maturity. Aquatic Living Resources 12 (1999) 11-22.
[43] B. Levavi-Sivan, C.L. Bloch, M.J. Gutnick, and I.A. Fleidervish, Electrotonic coupling in the
anterior pituitary of a teleost fish. Endocrinology 146 (2005) 1048-1052.
[44] C.L. Bloch, N. Kedar, M. Golan, M.J. Gutnick, I.A. Fleidervish, and B. Levavi-Sivan, Long-term
GnRH-induced gonadotropin secretion in a novel hypothalamo-pituitary slice culture
from tilapia brain. General and Comparative Endocrinology 207 (2014) 21-27.
[45] S.P. Deng, H.P. Chen, Y. Zhai, L.Y. Jia, J.Y. Liu, M. Wang, D.N. Jiang, T.L. Wu, C.H. Zhu, and
G.L. Li, Molecular cloning, characterization and expression analysis of spexin in spotted
scat (Scatophagus argus). Gen Comp Endocrinol 266 (2018) 60-66.
[46] V.L. Trudeau, Facing the Challenges of Neuropeptide Gene Knockouts: Why Do They Not
Inhibit Reproduction in Adult Teleost Fish? Front Neurosci 12 (2018) 302.
[47] R. Fontaine, P. Affaticati, K. Yamamoto, C. Jolly, C. Bureau, S. Baloche, F. Gonnet, P. Vernier,
S. Dufour, and C. Pasqualini, Dopamine Inhibits Reproduction in Female Zebrafish (Danio
rerio) via Three Pituitary D2 Receptor Subtypes. Endocrinology 154 (2013) 807-818.
[48] R.A.U. Strandabø, K. Hodne, E. Ager-Wick, O. Sand, F.-A. Weltzien, and T.M. Haug, Signal
transduction involved in GnRH2-stimulation of identified LH-producing gonadotropes
from lhb-GFP transgenic medaka (Oryzias latipes). Molecular and Cellular Endocrinology
372 (2013) 128-139.
[49] A. Hellqvist, M. Schmitz, I. Mayer, and B. Borg, Seasonal changes in expression of LHβ and
FSHβ in male and female three-spined stickleback, *Gasterosteus aculeatus*. General and
Comparative Endocrinology 145 (2006) 263-269.
[50] C.J. Martyniuk, K.J. Kroll, W.F. Porak, C. Steward, H.J. Grier, and N.D. Denslow, Seasonal
relationship between gonadotropin, growth hormone, and estrogen receptor mRNA
expression in the pituitary gland of largemouth bass. Gen Comp Endocrinol 163 (2009)
306-17.
[51] S. Wang, B. Wang, and S. Chen, Spexin in the half-smooth tongue sole (Cynoglossus
semilaevis): molecular cloning, expression profiles, and physiological effects. Fish Physiol
Biochem 44 (2018) 829-839.
[52] E. Kim, I. Jeong, A.Y. Chung, S. Kim, S.H. Kwon, J.Y. Seong, and H.C. Park, Distribution and
neuronal circuit of spexin 1/2 neurons in the zebrafish CNS. Sci Rep 9 (2019) 5025.
[53] H. Wu, F. Lin, H. Chen, J. Liu, Y. Gao, X. Zhang, J. Hao, D. Chen, D. Yuan, T. Wang, and Z. Li,
Ya-fish (Schizothorax prenanti) spexin: identification, tissue distribution and mRNA
expression responses to periprandial and fasting. Fish Physiol Biochem 42 (2016) 39-49.
[54] A. Ma, M. He, J. Bai, M.K. Wong, W.K. Ko, and A.O. Wong, Dual Role of Insulin in Spexin
Regulation: Functional Link Between Food Intake and Spexin Expression in a Fish Model.
Endocrinology 158 (2017) 560-577.
[55] A. Reyes-Alcaraz, Y.N. Lee, S. Yun, J.I. Hwang, and J.Y. Seong, Conformational signatures in
beta-arrestin2 reveal natural biased agonism at a G-protein-coupled receptor. Commun
Biol 1 (2018) 128.

[56] W. Salzburger, Understanding explosive diversification through cichlid fish genomics. Nat
Rev Genet 19 (2018) 705-717.

[57] E. Ciani, R. Fontaine, G. Maugars, N. Mizrahi, I. Mayer, B. Levavi-Sivan, & F.-A. Weltzien.
Melatonin receptors in Atlantic salmon stimulate cAMP levels in heterologous cell lines
and show season-dependent daily variations in pituitary expression levels. Journal of
Pineal Research, 67(3),(2019) e12590.

A

Spx1a

64
90
54

Spx1

Spx1b
86

Spx2

68
61

Tilapia
Zebra mbuna
Burtoni
Medaka
Goldfish
Fugu
Platyfish
Zebrafish
Cave fish
Human
Macaque
Mouse
Rat
Rabbit
Sheep
Cow
Chicken
Turkey
Turtle
Anole
Gar
Coelacanth
Elephant shark
P.nyererei
Zebra mbuna
Tilapia
Burtoni
Zebrafish
Xenopus SPX 2a
Xenopus SPX 2b
Medaka
Chicken
Coelacanth
Anole

B

Zebrafish *Ch4*	Nile tilapia *LG15*	Burtoni *Unplaced scaf.*
16.7M	28.04M	159.8K
golt1b	slc25a3a	slc25a3a
spx1	ldhba	ldhba
gys2	kiss2	kiss2
kiss2	gys2	gys2
ldhba	spx1a	spx1a
slc25a3a	golt1b	golt1b
16.85M	28.09M	237.9K

C

Zebrafish *Ch25*	Nile tilapia *LG7*	Burtoni *Unplaced scaf.*
3.25M	4.06M	31.3K
GALR type 2	ldhd	slc35b4
pthlhb	syt9b	golt1bb
ldhbb	5.95M	spx1b
	39.28M	ldhbb
golt1bb	lrp5	pthlhb
slc35b4	ppp6r3	GALR type 2
3.34M	gal	ldhd
8.06M	GALR type 2	lrp5
kcnc1	pthlhb	ppp6r3
lrp5	ldhbb	kcnc1
ppp6r3	spx1b	gal
gal	golt1bb	syt9b
spx2	slc35b4	650K
ldhd	kcnc1	
syt9b	50.48M	
15.90M		