Distinct neural substrates for opening and closing the gate from

perception to working memory

NEURAL CORRELATES OF WORKING MEMORY GATING

Gal Nir-Cohen[1,3], Yoav Kessler[2,3], and Tobias Egner[4,5]

[1] Department of Cognitive and Brain Sciences, Ben-Gurion University of the Negev.

[2] Department of Psychology, Ben-Gurion University of the Negev.

[3] Zlotowski Center for Neuroscience, Ben-Gurion University of the Negev.

[4] Center for Cognitive Neuroscience, Duke University

[5] Department of Psychology and Neuroscience, Duke University.

Corresponding author e-mail: nga@post.bgu.ac.il

19 **Conflict of interest statement**: The authors declare no competing financial interests.

20 **Acknowledgments**: This study was funded by US-Israel Binational Science Foundation

21 (BSF) grant awarded to YK and TE.

Abstract

Working memory (WM) needs to protect current content from interference and simultaneously be amenable to rapid updating with newly relevant information. An influential model suggests these opposing requirements are met via a basal ganglia (BG) - thalamus gating mechanism that allows for selective updating of prefrontal cortex (PFC) WM representations. A large neuroimaging literature supports the general involvement of the PFC, BG, and thalamus, as well as posterior parietal cortex (PPC), in WM. However, the specific functional contributions of these regions to key sub-process of WM updating, namely gate-opening, content substitution, and gate closing, are still unknown, as common WM tasks conflate these processes. We therefore combined functional MRI (19 male and 29 female participants) with the reference-back task, specifically designed to tease apart these sub-processes. Participants compared externally presented face stimuli to a reference face held in WM, while alternating between updating and maintaining this reference, resulting in opening vs. closing the gate to WM. Gate opening and substitution processes were associated with strong BG, thalamic and fronto-parietal activation, but – intriguingly - the same activity profile was observed for sensory cortex supporting task stimulus processing (i.e., the fusiform face area). In contrast, gate closing was associated near-exclusively with PPC activation. These novel findings provide support for the gate opening process proposed by the BG gating model, but qualify the model's assumptions by demonstrating that gate closing does not depend on the BG but the PPC, and that gate opening also involves task-relevant sensory cortex.

Significance statement

The ability to selectively "gate" task-relevant information into working memory (WM) is crucial for the efficient use of its limited capacity. Whereas previous studies documented the involvement of multiple brain regions in WM updating, how these regions mediate specific sub-processes of WM updating has remained unknown. In this neuroimaging study we utilized a task specifically designed to dissociate gate-opening, content substitution, and gate-closing processes. Our results demonstrate the involvement of the basal ganglia, frontoparietal, and task-relevant sensory cortex in opening the gate from perception to WM. Importantly, we also document that gate-opening and gate-closing are mediated by different brain regions, with gate-closing being supported by the parietal cortex. These results inform and extend current models of WM.

Introduction

Working memory (WM) refers to our ability to temporarily hold information in mind, manipulate it, and update it in the service of goal-directed behavior (Cowan, 2017; Oberauer et al., 2018) . Models of WM have long emphasized the tension between its maintenance and updating functions (Frank et al., 2001; Miller and Cohen, 2001; O'Reilly, 2006; Badre, 2012; Fallon et al., 2017): current WM content has to be shielded from interference by irrelevant information, while at the same time being amenable to updating when new goal-relevant information appears.

An influential neurocognitive theory addressing this dilemma is the PBWM (prefrontal cortex, basal ganglia, and WM) network model, which postulates a selective input-gate for WM (Frank et al., 2001; O'Reilly and Frank, 2006). Specifically, the model proposes a basal ganglia (BG) gating mechanism that separates perceptual input, represented in sensory cortex, from WM representations, maintained in (or via) dorsolateral prefrontal cortex (dlPFC). By default, the gate is closed, thus enabling robust maintenance/shielding of WM content. However, in response to salient signals, like task-relevant stimuli or reward cues, the BG gate opens (based on phasic dopaminergic input from the midbrain), allowing for the inflow of new information into WM via a thalamus-PFC pathway.

There is a vast neuroimaging literature supporting the assumption that the dlPFC – usually in conjunction with the medial prefrontal cortex (mPFC) and posterior parietal cortex (PPC) – contributes to the maintenance of WM content (e.g., (Roth et al., 2006; Feredoes et al., 2011; Nee et al., 2013; D'Esposito and Postle, 2015). The same broad set of regions, often referred to as the frontoparietal network (FPN), has also been implicated in updating WM, as inferred from N-back (e.g.,

Owen et al., 2005), AX-CPT (e.g., Lopez-Garcia et al., 2016) and task switching

studies (e.g., Kim et al., 2012). A smaller set of studies has also provided evidence to

support the involvement of the BG (Cools et al., 2007; McNab and Klingberg, 2008;

Murty et al., 2011) and/or the midbrain (Murty et al., 2011; D'Ardenne et al., 2012) in

WM updating.

There is thus broad agreement on the FPN, BG, and thalamus being the key

players in WM input gating. However, the precise roles these regions play is not

presently clear, because the tasks typically employed in this literature do not allow

one to tease apart key sub-processes that comprise WM updating. For instance,

standard N-back, AX-CPT, and task switching protocols conflate item encoding,

updating, substitution, and other processes, and do not provide a means to

differentiate gate opening, gate closing, substitution and item removal processes

(discussed in Ecker et al., 2010; Rac-Lubashevsky and Kessler, 2016a, 2016b; Kessler

et al., 2017; Lewis-Peacock et al., 2018).

The goal of the present study was therefore to examine potential functional

specialization in the above WM network with respect to sub-processes involved in

WM updating. To this end, we paired functional magnetic resonance imaging (fMRI)

with the recently developed "reference-back" task (Rac-Lubashevsky and Kessler,

2016a, 2016b, 2018), which has been shown to successfully disentangle processing

costs associated with four key WM updating operations: (1) opening the gate to WM,

(2) updating information in WM – which may either take the form of reinforcing

current content or (3) substituting old with new information, and (4) closing the gate

in order to enable robust maintenance of the newly updated information.

By interrogating neural responses in the BG, FPN, and thalamus, as well as in

visual regions with known sensitivity to our task stimuli (see below), we observed

distinct patterns of neural substrates supporting the different WM updating processes.

Whereas dlPFC, BG, and thalamus were preferentially involved in the gate opening

process, parietal cortex also contributed to this process, but additionally displayed a

stronger contribution to substitution, and a near-exclusive involvement in the gate

closing process.

Materials and Methods

Participants

To mitigate the dangers of false-positive and –negative findings, we based our

sample size on effect size estimation (Button et al., 2013). Specifically, a recent meta-

analysis of a large fMRI data set indicated a moderate effect size for WM task

contrasts (Poldrack et al., 2017), For a desired power of 0.8 to detect this size of effect

in within-subjects contrasts, under assumption of a conservative (low) level of

correlation between paired observations (r=0.3), we required a minimum sample size

of N=45 (based on GPOWER, Erdfelder et al., 1996).

61 healthy students from Ben-Gurion University of the Negev participated in

the experiment in exchange for monetary compensation. 13 participants were

excluded from the analysis due to technical problems with the MRI during the scan

(6), extensive head movements (2) or a low accuracy rate (5; <80%).   The final

sample included 48 participants (29 females; age M=25.5, SD=2 years). All

participants were right-handed and reported normal or corrected-to-normal vision.

None of the participants had any history of neurological or psychiatric problems. The

experiment was approved by the Helsinki committee of the Soroka Medical Center,

Beer Sheva, Israel.

Stimuli

The reference-back task used 8 face images of neutral facial expression (4

males, 4 female; two faces per block) from FEI faces database

(http://fei.edu.br/~cet/facedatabase.html). The faces were displayed inside blue (RGB

values: 0,0,255) and red (RGB values: 255,0,0) colored frames (see Figure 1). The

faces' diameter was approximately 180 pixels (4.76 centimeters, subtending a visual

angle of 2.7° from a 100 centimeters viewing distance). The frame's dimensions were

380x380 pixels (10x10 centimeters), subtending a visual angle of 5.7°. The use of

face stimuli (in combination with an independent "localizer" scan) enabled us to

identify the fusiform area (FFA; Kanwisher et al., 1997) in order to assess neural task

stimulus processing in visual cortex as a function of WM updating operations. The

localizer task employed gray scale images of famous familiar faces, unfamiliar faces,

buildings, objects and scrambled objects. Those images were presented within an

elliptical shape (14.5x8 centimeters, 8.3°x4.5°) against a black background. All

stimuli were projected on a screen at the back of the scanner bore, and viewed via a

mirror affixed to the headcoil.

Procedure

The scanning session took about an hour, in the following sequence:

anatomical structural scan (10 min); 1-back task with face stimuli, serving as a face

localizer task (10 min); and four 80-trial blocks of the reference-back task (30 min).

The participants completed a behavioral practice session (2 blocks) of the reference-

back task a day or two prior to the experimental session in the scanner.

The reference-back task

We employed the reference-back task (Rac-Lubashevsky and Kessler, 2016a, 2016b), which is designed to disentangle WM updating sub-processes (i.e., gate opening, gate closing, and substitution). This task is based on the N-back task. In the classic N-back task the participant is presented with a sequence of stimuli, and is asked to decide whether the current stimulus is identical or not to the stimulus presented N trials before (Jonides et al., 1997; Owen et al., 2005). Since updating, including all of its sub-processes, takes places in each trial of the n-back task, it is difficult to isolate the sub-process and identify their distinctive neural markers. In order to overcome this limitation, the reference-back paradigm was developed.

This task is composed of two trial types: reference and comparison trials. Specifically, in each trial of the current task (see Figure 1), a face stimulus was presented inside a red or a blue frame, and the participant was required to indicate whether or not the stimulus was identical to the one presented in the *most recent red frame*. Trials involving a red frame are denoted *reference trials*. In these trials, the participant must first compare the presented stimulus to the one held in WM, i.e., the face that appeared in the previous red frame (making a same/different decision); the participant then has to update his/her WM with the stimulus that appears in the present trial, which serves as the reference for future trials. Trials involving a blue frame are denoted *comparison trials*. Like in red frame trials, the participant is required to make a same/different decision between the currently presented face and the reference held in WM; however, unlike in reference trials, WM does not have to be updated, because blue-framed faces do not serve as a reference for future trials.

Accordingly, both reference and comparison trials involve a same/different decision against a WM referent, but only the former require WM updating. This

means that the gate to WM should be open in reference trials but kept closed in comparison trials. By considering the state of the gate on the previous trial, this protocol further allows one to distinguish between trials where the gate needs to be opened and trials where the gate needs to be closed. Specifically, trials in which the previous trial-type is repeated (e.g., two reference trials in a row) do not entail a change in the state of the gate: the gate remains open for successive reference trials and it remains closed for successive comparison trials. However, switching from a comparison trial to a reference trial requires gate opening, while switching from a reference to a comparison trial requires gate closing (see Figure 1 for a trial-by-trial example).

Subsequentially, the reference-back task enables one to distinguish among WM updating sub-processes using three orthogonal contrasts. These contrasts were utilized in previous studies, showing robust behavioral effects (Rac-Lubashevsky and Kessler, 2016a, 2016b; Kessler, 2017) as well as EEG correlates (Rac-Lubashevsky and Kessler, 2018), and association with spontaneous eye blink rates, an index of central phasic dopaminergic activity (Rac-Lubashevsky et al., 2017). Moreover, reaction time costs for these contrasts demonstrate split-half reliabilities of .85-.86, and are correlated with performance in the standard N-back task (Rac-Lubashevsky and Kessler, 2016b).

The contrasts were defined as follows. (1) *Gate opening* is the difference between reference-switch and reference-repeat trials. All reference trials require the gate to WM to be open but only the former, where participants are switching from a comparison to a reference trial, involves the process of gate opening. Using a similar logic, (2) *Gate closing* was defined as the difference between comparison-switch and comparison-repeat trials – all comparison trials require a closed gate, but only on

trials where participants switch from a reference to a comparison trial does the process

of gate closing take place. Lastly, (3) *Substitution* refers to replacing old with new

information in WM, which occurs on those reference trials where the current stimulus

does not match the previous reference (see also Ecker et al., 2010). It is important to

de-confound substitution from the difference between making a "same" versus a

"different" response, and this can be achieved by using the difference between "same"

and "different" responses in comparison trials as a baseline. Accordingly, substitution

is calculated as an interaction contrast, reflecting a larger difference between "same"

and "different" responses in reference trials than in comparison trials: ("different"

$_{\text{reference}}$ − "same" $_{\text{reference}}$) − ("different" $_{\text{comparison}}$ − "same" $_{\text{comparison}}$). In addition, the

reference-back design enabled us to examine the differential neural activity of being

in an open-gate state ("update mode", see Kessler and Oberauer, 2014, 2015)

compared to a closed-gate state. Accordingly, the *updating mode* is defined by the

overall difference between reference trials, where the WM referent has to be updated,

and comparison trials, where the referent does not have to be updated. Thus, while

substitution refers to the situation where updating involves replacing of the old

referent with a new one (and possibly also includes removing the now-irrelevant item,

see Lewis-Peacock et al., 2018), *updating* refers to the more general situation of being

in an open-gate state, regardless of whether the referent has to be replaced ("different"

trials) or not (see Table 1). The updating mode contrast only involved trial-type

repetition trials, in order not to confound it with gate-switching.

223   These process designations are shown in Figure 1, and the specific contrasts

224 isolating the different updating operations are laid out below in the Statistical Data

225 Analysis section.

Trial Type	reference	comparison	reference	reference	comparison
Gate State	open	closed	open	open	closed
Gate Switch		switch (closing)	switch (opening)	repeat	switch (closing)
Response		"different"	"same"	"different"	"same"

227   Figure 1: An example sequence of reference (red frame) and comparison (blue frame)

228   trials in the reference-back task (top), along with an illustration of the putative states

229   of the WM gating process and responses required (bottom).

231   Each block of the reference-back task started with a reference trial, to which

232 the participants did not respond. Then, in each subsequent trial, a framed face was

233 presented for 2 seconds, followed by a blank inter-trial interval for 2, 4, 6 or 8

234 seconds. Each of the eight conditions (Trial-Type * Gate-Switching * Response) was

235 presented 10 times in each block, resulting in a total of 320 trials (40 trials per

236 conditions). The order of trials, as well as the duration of the inter-trial interval jitter,

237 were determined using Optseq (Free-Surfer analysis tools; Greve, 2002). Each of the

238 four experimental blocks involved two face stimuli from the same gender. The stimuli

239 were changed from one block to another and were counterbalanced between

240 participants.

FFA Localizer Task

As an FFA localizer task, we employed a block-design 1-back task (taken from Avidan et al., 2014). Different stimulus categories (familiar faces, unfamiliar faces, buildings, objects, and scrambled objects) were presented in 10 second blocks, with 6 seconds intervals between blocks. Within each block, ten images were presented, each for 800ms followed by 200ms inter-trial interval. Within each block, nine images were unique whereas one image was presented twice in a row. The participants were asked keep track of the stimuli, and to press a key each time an image was presented twice in a row (1-back). There were seven repetition of each block type.

fMRI data acquisition and preprocessing

All fMRI data were collected at the Brain Imaging Research Center, Soroka Medical Center, Beer-Sheva, using a 3-T Philips ingenia MRI scanner. The scanning of each participant started with a 3D structural scan, acquired by a T1-weighted sequence that yielded high resolution images of 1^3 mm voxel size with matrix of 256X256 for 170 slices. Functional data were collected by a T2*-weighted sequence (TR = 2000 msec, TE = 35 msec, flip angle = 90°). 35 slices were scanned in ascending order with a 96x96 matrix size, 2.61x2.61 mm voxel resolution with 3 mm thickness. A total of 215 volumes were acquired. Behavioral responses were recorded using a two-keys box the participants held in their right hand and pressed with their index or middle finger for different and same responses, respectively. Imaging data were preprocessed and analyzed using Statistical Parametric Mapping 12 (Welcome Trust Centre for Neuroimaging, London, UK; http://www.fil.ion.ucl.ac.uk/spm). Each participant's functional images were re-aligned, co-registered to the anatomical image and slice-time corrected. Then, the images were normalized into MNI space (with a 2^3

mm voxel size interpolation) and smoothed using a 6 mm gaussian kernel to full

width at half maximum (FWHM).

Statistical data analysis

Four contrasts were defined for gate opening, gate closing, substitution, and

updating mode, as explained in the task description above (See Table 1). These

contrasts were used to analyze both the behavioral and the neural data. For the FFA

localizer task, we contrasted blocks of face stimuli over blocks where non-face stimuli

were shown.

Trial type	reference				comparison			
Gate Switch	repeat		switch		repeat		Switch	
Response	same	different	same	different	same	different	same	different
Gate opening	-	-	+	+				
Gate closing					-	-	+	+
Substitution	-	+			+	-		
Updating mode	+	+			-	-		

Table 1: Contrast weights for defining distinct WM updating sub-processes.

All the participants' functional images for each of the contrasts were submitted

to a one-sample t-test group-level analysis, where participants were treated as random

effects. We pursued two broad sets of fMRI analyses, the first being an exploratory

whole-brain analysis, and the second being a region-of-interest (ROI) analysis

grounded in our *a priori* interest of closely interrogating the role of different nodes of

the WM network (the constituent parts of the FPN, the BG nuclei, and the thalamus)

and the FFA in distinct aspects of the WM updating process. Accordingly, we took a

very conservative approach for guarding against false-positives in the exploratory

analysis, using a voxel-based family wise error *(*FWE*)* with a threshold of $p < 0.05$

and a minimum cluster size (K_E) of 10 significant voxels for the whole-brain analysis,

and employed a less conservative approach for the *a priori* ROI analysis by using a

voxel-based false discovery rate (FDR) with a threshold of $p < 0.05$ and $K_E > 10$. Note

that we use voxel-based rather than cluster-based thresholding throughout to bypass

recent concerns about common cluster-based correction approaches (Eklund et al.,

2016; Cox et al., 2017). The WM network ROIs were defined anatomically, using the

WFU PickAtlas toolbox (Maldjian et al., 2003) masks of the BG and thalamus (using

AAL labels; Tzourio-Mazoyer et al., 2002), and assembling an FPN mask, using

Brodmann areas, by combining masks of the dorsolateral PFC (BA8, BA9, BA46),

medial PFC/ACC (BA24, BA32), and the posterior parietal cortex (BA7, BA40). We

specified a functional ROI of the FFA via the localizer scan. Finally, in addition to

searching for significant clusters of activation within each ROI, we also ran the

above-mentioned contrasts on mean activity (beta estimates) extracted from each ROI.

Results

Behavioral results

All the conditions, along with the four *a priori* contrasts of interest were

tested on both response time (RT) and accuracy; mean RTs for the key conditions are

shown in Figure 2, and descriptive and inferential statistics are presented in Tables 2

and 3. The results fully replicated those of the original studies on the reference-back

paradigm (Rac-Lubashevsky and Kessler, 2016b, 2016a): As can be seen in Figure 2,

mean RT for reference trials was substantially slower than for comparison trials,

reflecting the cost of WM updating mode (54ms, $p < 0.001$). As shown in Figure 2a,

switch trials were found to be significantly slower than repeat trials, both in reference

trials and in comparison trials, reflecting the costs of gate-opening (72ms, $p < 0.001$)

and gate-closing (52ms, $p < 0.001$), respectively. Finally, the interaction contrast

comparing "same" and "different" response conditions between reference and

comparison trials revealed a robust substitution cost (92ms, $p < 0.001$). In summary,

the behavioral results showed that our adaptation of the reference-back protocol was

successful in revealing the behavioral signatures of updating, gate opening and

closing, and substitution processes in WM, thus providing a solid basis for

interrogating the fMRI data for neural substrates of these processes.

	Conditions		Response Time		Accuracy	
Trial Type	Gate switch	Response	Mean(ms)	SD	Mean(%)	SD
Reference	Repeat	Same	694	113	98	3.02
		Different	882	158	97	3.28
	Switch	Same	773	160	98	2.62
		Different	945	199	96	4.48
Comparison	Repeat	Same	688	116	98	2.47
		Different	782	145	98	3.15
	Switch	Same	716	114	98	2.93
		Different	857	165	97	2.67

315 Table 2: Descriptive statistics of the RT and accuracy data.

	Response Time				Accuracy			
Contrast	Mean difference (ms)	$F_{(1,47)}$	p	η_p^2	Mean difference (%)	$F_{(1,47)}$	p	η_p^2
Gate-opening	72	71.12	<.001	.60	-.4	1.02	.31	.02
Gate-closing	52	78.33	<.001	.62	-.6	3.04	.08	.06
Substitution	92	162.45	<.001	.77	-.1	0.71	.40	.01
Updating mode	54	57.88	<.001	.47	-.1	2.36	.13	.05

Table 3: Summary of the inferential statistics of the RT and accuracy data analysis

Figure 2: Mean response time (and standard error of the mean) for reference and

comparison trials are displayed (A) as a function of whether the condition was

repeated or switched to, and (B) of whether the stimulus/response was the same or

different to the reference stimulus held in WM. The figure also highlights the four

key contrasts defining gate-opening, gate-closing, substitution and updating mode.

Imaging results

Whole-brain analysis

Exploratory whole brain analyses were conducted for each contrast of interest

using a conservative correction threshold (voxel-wise FWE $p < 0.05$, $K_E > 10$). Dorsal

views of cortical activations revealed by each contrast are presented in Figure 3 (for

full list of activated clusters, see Table 4). The process of *gate opening* (required in

reference trials that follow a comparison trial) was associated with increased

activation in dorsal and dorsomedial frontal and parietal regions, with particularly

large clusters of activity observed in the posterior and medial aspects of the PPC,

including the precuneus. Additionally, gate-opening was associated with activity in

the thalamus, as well as an extensive posterior cluster stretching from the cuneus into

parts of visual cortex, including the fusiform gyrus (for full list of activated clusters,

see Table 4). Given that this contrasts controls for basic visual input (which is equated

between reference and comparison trials), the latter data suggest that the process of

gating visual information into WM may be directly reflected in enhanced activity in

relevant visual regions (see also FFA ROI analyses, below).

The analysis of the gate-closing process (required on comparison trials that

follow reference trials) did not yield significant activations with the conservative

FWE whole-brain correction. To probe further for potential neural substrates of gate-

closing, we applied a more lenient form of whole-brain correction (voxelwise FDR

$p < 0.05$, $K_E > 10$) to this contrast, which revealed primarily activity in bilateral PPC,

specifically in the superior parietal lobule (SPL)/intraparietal sulcus (IPS), along with

smaller clusters of dorsal frontal activation (for full listing of active clusters, see Table

4). When WM did not only have to be updated but information in WM had to be

replaced (*substitution*, required on reference trials where the current stimulus

mismatched the WM referent), activity was enhanced in left dlPFC (middle frontal

gyrus) and inferior parietal lobule (for full listing of active clusters, see Table 4).

Finally, being in an *Updating* mode was also associated with increased activation of

dorsal frontal and parietal regions, including most prominently the left PPC.

In sum, in line with expectations, the exploratory whole-brain analysis

identified core components of the FPN as supporting the regulation of WM

updating/protection processes. However, these contrasts also documented some clear

regional differences, suggesting a relatively greater involvement of medial posterior

parietal (and visual) cortex in gate opening, of more lateral posterior parietal regions

in gate closing operations, and relatively stronger prefrontal involvement in the

substitution process. To get a better grasp on regional activation differences and

commonalities between processes, we next conducted a set of conjunction analyses,

361   before turning to the ROI based analysis of the key WM network components to gain

362   a clearer quantitative picture of a possible functional fractionation of this network.

363   Figure 3: Whole-brain group search results for neural substrates of the different WM
364   updating sub-processes/states are displayed as rendered dorsal 3D views.

Conjunction analysis

We next conducted whole-brain conjunction analyses between pairs of

contrasts in order to identify regions that are significantly activated exclusively in

relation to one process, or shared among several sub-processes. This analysis was

again conducted using a conservative voxelwise threshold of FWE $p < 0.05$ paired

with a minimum cluster of $K_E > 10$, except for conjunctions involving the gate-closing

contrast, with the latter being thresholded at FDR $p < 0.05$ with minimum cluster of

$K_E > 10$ (see above). Axial view slices presenting some key conjunction results are

shown in Figure 4. (For full listing of active clusters, see Table 5).

We started by examining the conjunction between updating mode and each of

the other three sub-processes (i.e., gate opening, gate closing and substitution).

Updating shared activation of the supplementary motor area (SMA) with substitution.

Additionally, updating and gate closing, as well as updating and gate-opening, shared

neural activity in the left PPC (i.e., BA 7, BA 40). The only shared activation between

gate opening and substitution was found in the SMA. No overlapping activity was

found between gate closing and gate opening or substitution, implying that the gate

closing relies on different neural substrates than the other two processes.

Figure 4: Results of the whole-brain group search conjunction analysis across the

different WM updating sub-processes/states are displayed on axial brain slices (left

to right, slices: Z = 42, 44, 46, 48, 50, 52, 54, 56).

ROI analysis

Activity related to the different WM updating operations in *a priori* ROIs was

examined with ROI-wide FDR correction, using a voxelwise threshold of FDR $p <$

0.05 paired with a minimum cluster extent of $K_E > 10$. Dorsal views/axial slices

illustrating key findings are shown in Figure 5. A list of peak coordinates is shown in

Table 6. Additionally, we extracted and analyzed mean activity estimates from each

of the ROIs. While this analysis is necessarily less sensitive, as it averages activity

over entire anatomical regions, it allowed us to further quantify the potential regional

functional specializations with respect to WM sub-processes, and to ensure that the

inferences derived from the ROI-based search do not simply reflect (quantitative)

thresholding effects, but genuinely (qualitatively) different activity patterns. Figures 6

and 7 present mean beta values extracted for each ROI (broken down into nuclei, in

the case of the BG).

BG   In the ROI-based search, we observed enhanced activity in BG nuclei

for gate opening and substitution, but not for gate closing or being in an updating

mode (see Figures 5 and 6). More specifically, opening of the gate to WM was

associated with the most widespread increase in activity, involving all of the BG

nuclei bilaterally (Figure 5). In contrast, no activation increase was detected in any

part of the BG during WM gate closing or updating mode. Finally, the substitution of

old WM content with new information was associated with increased activation in the

caudate, left putamen, and left pallidum. This pattern of results was largely replicated

in the analysis of mean activation estimates across the BG, as well as for individual

nuclei (Figure 6). Mean activation of the caudate was significantly higher during

switch than repeat trials in reference trials (gate opening: $F_{(1,47)} = 9.66$, $p < 0.05$), but

not in comparison trials (gate closing: $F_{(1,47)} < 1$, $p = .41$). This pattern also held for

mean activation in the putamen (gate opening: $F_{(1,47)} = 5.28$, $p < 0.05$, gate closing:

$F_{(1,47)} = 1.25$, $p = .26$) and the pallidum (gate opening: $F_{(1,47)} = 13.41$, $p < 0.001$, gate

closing: $F_{(1,47)} < 1$, $p = .45$). At the level of mean ROI activity, however, substitution

cost was not significant for any contrast (caudate: $F_{(1,47)} = 2.80$, $p = .10$; putamen:

$F_{(1,47)} < 1$, $p = .72$; pallidum: $F_{(1,47)} < 1$, $p = .35$), and the same was true for updating

mode related activity (caudate: $F_{(1,47)} = 2.09$, $p = .15$; putamen: $F_{(1,47)} < 1$, $p = .53$;

pallidum: $F_{(1,47)} < 1$, $p = .53$). In sum, the present data support the longstanding

proposal of BG involvement in input-gating of WM content, by showing that the BG

are robustly associated with the process of opening the gate to WM. We also observed

some evidence for substitution related activity, but – importantly – the BG nuclei

seem to play no active role in closing the gate to WM.

Thalamus Similar to the BG, the thalamus was found to display activity

increases during WM gate opening and substitution, but displayed no detectable

increase in activity during the gate closing operation or with respect to being in an

updating mode. This was born out both by the search for significant clusters within

the thalamus ROI (Figure 5), as well as by the analyses run on mean thalamus

activation (Figure 6): Switch trials evoked higher mean activation than repeat trials in

reference trials (reflecting gate-opening: $F_{(1,47)} < 20.99$, $p < .001$), but not in

comparison trials (reflecting gate-closing: $F_{(1,47)} < 1$, $p = .52$). Moreover, at the mean

ROI activity level, neither the substitution cost ($F_{(1,47)} < 1$, $p = .48$) nor updating mode

contrasts ($F_{(1,47)} < 1$, $p = .89$) were significant. In sum, as in the BG, the most

pronounced activity increase in the thalamus was observed when the gate to WM had

to be opened, where we observed no evidence for an involvement in gate closing.

FPN As already suggested by the whole-brain analysis above, we found that

components of the FPN were activated by all WM updating sub-processes, but the

pattern of activation revealed a functional fractionation of the different FPN nodes

(see Figures 5 and 7). Specifically, the search for significant activated clusters within

the ROIs showed that, whereas the entire FPN was robustly and bilaterally activated

during the gate opening operation and by being engaged in an updating mode, the

process of WM content substitution produced much more lateralized activity in the

left parietal and left lateral frontal cortex, and gate closing was associated almost

exclusively with enhanced left parietal activation along the IPS (Figure 5).

The analysis of mean ROI activity confirmed this picture: As shown in Figure

7, the gate-opening contrast was significant for mean activation in all FPN

components (dlPFC: $F_{(1,47)} = 22.93$, $p < .001$; mPFC\ACC: $F_{(1,47)} = 12.41$, $p < .001$;

PPC: $F_{(1,47)} = 19.24$, $p < .001$). In contrast, the gate closing contrast at the level of

mean ROI activity was not found to be significant in any of the FPN components,

though in line with the ROI-based search results above, the PPC mean activity

displayed a trend in this direction (dlPFC: $F_{(1,47)} < 1$, $p = .26$; mPFC\ACC: $F_{(1,47)} < 1$,

$p = .91$; PPC: $F_{(1,47)} = 2.34$, $p = 0.13$). The substitution cost contrast was significant

for the PPC ($F_{(1,47)} = 5.83$, $p < .05$) but not the frontal FPN components (dlPFC: $F_{(1,47)}$

$= 2.38$, $p = 0.12$; mPFC\ACC: $F_{(1,47)} < 1$, $p = .40$). Finally, the updating mode contrast

was significant for mean activity in the PPC ($F_{(1,47)} = 4.26$, $p < .05$) and marginally

significant for the frontal regions (dlPFC: $F_{(1,47)} = 3.75$, $p = .06$; mPFC\ACC: $F_{(1,47)} =$

3.81, $p = .06$). Thus, in line with the results of the whole-brain analysis, the ROI-

based analysis provided additional evidence for a functional dissociation of FPN

components' roles in WM updating, with the frontal and parietal nodes being

concerned with gate opening and substitution processes (and being in an updating

mode), but posterior parietal cortex additionally serving a more exclusive role in the

gate closing process.

FFA   The (functionally defined) FFA ROI displayed a similar activity

pattern to that observed in the BG, thalamus, and frontal (but not parietal) FPN

components. In the search for active clusters, we observed loci in the FFA that

showed significant activation increases during substitution, but the most pronounced

activity was observed during gate opening (Figure 5). In contrast, no activity increase

was detected during gate closing or in relation to the updating mode. The same pattern

was confirmed in the analysis of mean FFA activation (Figure 7). Specifically, there

was greater mean activation during switch trials than repeat trials in reference trials

(gate opening: $F_{(1,46)} = 24.95$, $p < .001$) but not in comparison trials (gate closing:

$F_{(1,46)} < 1$, $p = .55$). Substitution cost and updating mode contrasts were not significant

at the mean ROI activity level, though both showed trending patterns for these effects

($F_{(1,46)} = 2.90$, $p =.09$; $F_{(1,46)} = 2.56$, $p =.11$, respectively). Thus, intriguingly,

processes related to allowing sensory content to enter WM (gate opening) or to

replace WM representations (substitution) seem to have a direct impact on activity in

the sensory regions that are involved in processing and/or representing the relevant

stimulus material.

Figure 5: Results of the ROI analyses are displayed as a function of WM updating

sub-process (rows) and ROI (columns) on dorsal-view rendered 3D brains (left

column) and axial slices (other columns). FPN = frontoparietal network, BG = basal

ganglia, FFA = fusiform face area.

Figure 6: Mean activity estimates (and standard error of the mean) for the BG nuclei

and thalamus are shown for reference and comparison trials as a function of whether

the condition was repeated or switched to (upper panel) and of whether the

stimulus/response was the same or different to the reference stimulus held in WM

(lower panel). The figure also highlights the four key contrasts defining gate-opening,

gate-closing, substitution and updating mode († denotes $p < .01$; * denotes $p < .05$; **

denotes $p < .001$).

Figure 7: Mean activity estimates (and standard error of the mean) for the FPN components and the FFA are shown for reference and comparison trials as a function of whether the condition was repeated or switched to (upper panel) and of whether the stimulus/response was the same or different to the reference stimulus held in WM (lower panel). The figure also highlights the four key contrasts defining gate-opening, gate-closing, substitution and updating mode († denotes $p < .01$; * denotes $p < .05$; ** denotes $p < .001$).

Region	HM	K_E	MNI coordinates			Z
			x	y	z	
Gate opening (reference switch > reference repeat)						
Precuneus		4,647	0	-70	34	7.33
Fusiform gyrus	R	133	30	-74	-10	7.05
SMA	L	43	-2	-8	52	5.56
MFG	L	20	-22	14	58	5.41
IFG	L	37	-50	24	22	5.39
Thalamus	L	18	-8	-18	10	5.34
Insula	L	17	-28	28	0	5.25
Gate closing (comparison switch > comparison repeat)						
Parietal lobule	L	943	-36	-66	50	5.05
Parietal lobule	R	276	40	-52	-28	5.18
BA46	L	42	-46	28	24	4.23
Middle frontal gyrus	R	50	34	2	58	4.10
Middle frontal gyrus	L	46	-30	-2	66	3.97
SMA	L	57	-2	14	48	4.10
BA10	L	30	-42	44	0	4.10
Substitution (different-same reference > different-same comparison)						
SMA	L	259	-2	8	56	6.06
Middle frontal gyrus	L	83	-50	-44	50	5.65
IPL	L	37	-44	2	50	6.53
Updating (repeat reference > repeat comparison)						
IPL	L	302	-48	-42	50	6.17
IPL	R	82	34	-44	38	5.61
MFG	R	49	38	4	58	5.54
SPL	L	22	-34	-64	52	5.42
MFG	L	10	-28	0	66	5.21

501 Table 4: Whole-brain analysis' list of peak activation in MNI coordinates; Z refer to
502 z-score at peak activated voxel

Region	HM	K_E	MNI coordinates			Z
			x	y	z	
Updating ∩ Gate opening						
IPL	L	16	-30	-50	38	4.82
Updating ∩ Gate closing						
IPL	L	51	-48	-44	50	5.25
Updating ∩ Substitution						
SMA	L	25	-2	8	58	5.30
Middle frontal gyrus	L	13	-46	2	50	4.96
Gate opening ∩ Substitution						
SMA	L	33	-4	10	52	5.33

504 Table 5: Conjunction analysis' list of peak activation in MNI coordinates; Z refer to z-
505 score at peak activated voxel

Region		HM	K_E	MNI coordinates			Z
				x	y	z	
Gate opening (reference switch > reference repeat)							
BG	Caudate	L	477	-8	22	2	4.66
		R	450	10	14	-2	4.90
	Putamen	L	608	-28	12	-4	3.76
		R	254	36	-8	-2	3.48
	Pallidum	L	206	-14	0	6	4.98
		R	108	16	4	2	4.09
Thalamus		L	1,031	-8	-18	10	5.34
		R	865	8	-14	8	4.71
FPN	lPFC	R	567	26	20	44	5.58
		L	428	-46	20	26	4.97
	mPFC\ACC	R	578	4	-8	50	4.67
		L	661	-2	-8	50	5.54
	PPC	L	2,286	-2	-72	34	6.83
		R	1,697	4	-68	32	7.10
FFA		L	208	-38	-62	-18	4.93
		R	276	36	-52	-16	5.88
Gate closing (comparison switch > comparison repeat)							
FPN	lPFC	L	10	-46	28	24	4.23
	mPFC\ACC	L	15	-4	14	46	3.99
	PPC	L	425	-46	-40	50	4.96
		R	25	34	-68	50	4.05
Substitution (different-same reference > different-same comparison)							
BG	Caudate	L	24	-8	22	2	3.60
		R	68	16	4	20	3.46
	Putamen	L	42	-16	6	6	3.61
	Pallidum	L	35	-14	6	2	3.89
Thalamus		L	13	-2	-10	12	3.21
		R	15	14	-30	10	3.26
FPN	lPFC	L	129	-50	6	40	5.03
		R	24	4	14	56	4.56
	mPFC\ACC	L	162	-2	8	52	5.43
		R	51	4	16	46	3.98
	PPC	L	1,289	-52	-40	50	5.57
		R	161	14	-66	60	4.42
FFA		R	14	44	-64	-20	3.90
		L	14	-40	-62	-20	3.52
Updating (repeat reference > repeat comparison)							
FPN	lPFC	L	122	-50	8	42	4.26
		R	81	38	40	36	5.42
	mPFC\ACC	L	33	-4	12	46	4.25
	PPC	L	1218	-50	-40	48	6.11
		R	830	40	-42	40	5.07

507 Table 6: ROI analysis peak activation in MNI coordinates; Z refer to z-score at peak

508 activated voxel

Discussion

There is copious evidence for the involvement of the FPN, BG, and thalamus in maintaining and updating WM content, but how the regions may differentially contribute to different sub-processes of WM updating is not well understood. To address this important question, we combined fMRI with the recently developed reference-back task, which enabled us – for the first time - to tease apart neural substrates of gate-opening, gate-closing, and substitution processes, as well as an updating mode of operation. Moreover, we used face stimuli in order to examine updating-related activity in sensory cortex specialized for processing the WM items, namely the FFA. Our behavioral results fully replicated previous studies using this protocol (Rac-Lubashevsky and Kessler, 2016a, 2016b; Kessler, 2017). Moreover, the imaging data indicate that the FPN, BG, and thalamus all contribute to gate-opening and substitution processes, with the FPN, but not the subcortical components, also being associated with being in an updating mode. Most notably, only the PPC was additionally found to be reliably associated with the closing of the gate to WM, and we found that FFA activity displayed robust effects of WM gate opening and substitution processes. We discuss the implications of these results in turn.

Gate opening, substitution, and updating mode in the WM network

The use of the reference-back paradigm allowed us to isolate different sub-processes involved in WM updating processes, thus enabling more precise process-to-brain region attribution and, accordingly, a stronger test of the PBWM model (Frank et al., 2001; O'Reilly and Frank, 2006). In general support of that model's proposal of BG-thalamus-PFC circuits supporting WM, we observed involvement of all components of this network (as well as PPC) in the processes of opening the gate to

WM, and in subsequently replacing the current content of WM with new perceptual
information (substitution). The PBWM model posits that it is specifically the BG (and
not the FPN) that are implementing the gating operation. However, under the
assumptions of the model it is nevertheless plausible that one would also observe gate
opening and substitution related activity more broadly throughout the WM network,
as the opening of the gate, and in particular the substitution process, would be
expected to have knock-on effects in the thalamus and FPN. Moreover, in support of
the model's differentiation between the gating mechanism in the BG and the
representation of WM content in FPN, we found that an "updating mode", which
refers to an open-gate state that does not involve any change in gating status, was
associated exclusively with FPN, and not with BG or thalamic activity.

Evidence for distinct neural substrates of gate-opening vs. gate-closing

Arguably the most novel and intriguing findings of the present study concern
the gate closing process. Previous behavioral studies demonstrated that both opening
and closing the gate to WM involves a reaction time cost. This was observed both
using the reference-bask task (Rac-Lubashevsky and Kessler, 2016a, 2016b, 2018;
Rac-Lubashevsky et al., 2017) and in a sequence updating task (Kessler and Oberauer,
2014, 2015). Here we show that, under roughly equivalent behavioral costs of opening
and closing the gate, the neural mechanisms involved in the two operations are
distinct. Notably, the BG and thalamus were only involved in gate opening, but not in
gate closing. The region most reliably associated with closing the gate to WM was the
PPC, in particular the (left) intraparietal sulcus (IPS). While the PPC in general, and
the IPS in particular, has previously been implicated in WM maintenance (e.g.,
Majerus et al., 2016; Quentin et al., 2019), to our knowledge the present study

represents the first evidence to suggest a specialized role for closing the gate to WM

in the service of shielding to-be-maintained WM items.

These findings seem to be at odds with the PBWM model, which does not

specify an active gate closing mechanism originating outside of the BG (Frank et al.,

2001; O'Reilly and Frank, 2006). The lack of a BG activation increase in relation to

gate closing *per se* may plausibly reflect the fact that the closed gate (i.e., tonic

inhibition of the thalamus) represents a BG default state (e.g., Chevalier and Deniau,

1990), returning to which might not impose additional local metabolic demands on the

BG. However, the observation of robust behavioral gate closing costs (both in the

current and prior studies), combined with the increased activation we detected in the

PPC in relation to the gate closing process, suggest that closing the gate to WM

nevertheless involves the contribution of an active control process. Based on the

present results, we propose that this process originates in the PPC. Given its ample

direct projections to the BG (e.g., Cavada and Goldman-Rakic, 1991; Jarbo and

Verstynen, 2015) we speculate that gate closing involves the PPC signaling to the BG

the need to close the gate to WM. Clearly, future studies, perhaps involving targeted

neuro-stimulation of the PPC, are needed to rigorously test this new proposal.

Another possible account for the lack of active BG involvement in the gate

closing process could be that closing the gate to WM relies on direct projections from

the VTA to frontal regions, and hence does not activate the BG (cf. Braver and

Cohen, 2000; D'Ardenne et al., 2012). Specifically, while gate opening should be

selective by nature, and hence involve activating specific BG-PFC loops, gate closing

may be non-selective and hence could operate through the (non-specific) VTA-PFC

pathway. Future studies could potentially probe whether a different neural signature

emerges under conditions where gate-closing has to be selective, too, such as when

items from multiple categories have to be updated independently.

The role of posterior cortex in WM updating

Many prior studies have established that the PFC is a crucial component of

short-term memory (Fuster and Alexander, 1971), cognitive control (Miller and Cohen,

2001) and WM updating and maintenance (e.g., Narayanan et al., 2005). However, the

specific role it plays in supporting these abilities is still under debate. One traditional

view, which is also implemented in PBWM, holds that the PFC plays a key role in

encoding and storing goal-directed representation within WM (Goldman-Rakic et al.,

1996; Courtney et al., 1998; O'Reilly and Frank, 2006). In contrast, a more recent view

(the sensory recruitment hypotheses) does not identify the PFC with temporary storage

per se, but rather characterizes the lateral PFC as responsible for directing selective

attention towards memory representations, with the actual representations being held in

the posterior cortex, that is, in areas that also serve the perception and long-term storage

of the memoranda (D'Esposito et al., 2000; Postle, 2006; Feredoes et al., 2011; Lara

and Wallis, 2015; Serences, 2016).

Relevant to this debate, our findings demonstrate the involvement of the FFA

in WM updating sub-processes. In particular, gate opening and substitution processes,

but not gate-closing, were associated with elevated neural activity in the FFA. This

finding supports the notion that posterior, "perceptual" regions play a major role not

only in WM maintenance (as already posited by the sensory recruitment account), but

also in WM updating. We offer two tentative accounts for this finding. The first is that

the posterior cortex is more heavily recruited for WM maintenance in situations that

call for substitution of the information, or for switching from "passive maintenance"

(as reflected in comparison trials) to updating. A second, not mutually exclusive

interpretation is that the same signaling cascade that leads to gating information into

WM also serves to concurrently boost attention to the to-be-encoded items represented

in posterior perceptual regions. While input gating involves a feed-forward flow of

information from perception to WM, representations held in WM may in turn lead to

directing attention toward perceptual regions, in a concurrent feed-backward fashion.

These possibilities could plausibly be evaluated by combining the reference-back task

with more time-sensitive measures of neural activity in future studies.

To conclude, the present study revealed distinct neural activity related to gate

opening, gate closing, substitution, and updating mode, by combining the reference-

back protocol with fMRI. While gate opening was associated with activation of the BG-

PFC loop, in accordance with the PBWM model, the same was true for FFA activity,

while gate closing was associated predominantly with PPC activity. These results

support a novel functional fractionation of the WM network, with distinct neural

substrates mediating the opening versus closing of the gate between perception and

WM.

References

Avidan G, Tanzer M, Hadj-Bouziane F, Liu N, Ungerleider LG, Behrmann M (2014)
Selective Dissociation Between Core and Extended Regions of the Face Processing
Network in Congenital Prosopagnosia. Cereb Cortex 24:1565–1578.

Badre D (2012) Opening the gate to working memory. Proc Natl Acad Sci USA 109:19878–
19879.

Braver TS, Cohen JD (2000) On the Control of Control: The Role of Dopamine in Regulating
Prefrontal Function and Working Memory. In: Attention and performance, Vol
XVIII, Control of cognitive processes (Monsell S, Driver J, eds), pp 713–737.
Cambridge, MA: MIT.

Button KS, Ioannidis JPA, Mokrysz C, Nosek BA, Flint J, Robinson ESJ, Munafò MR (2013)
Power failure: why small sample size undermines the reliability of neuroscience. Nat
Rev Neurosci 14:365–376.

Cavada C, Goldman-Rakic PS (1991) Topographic segregation of corticostriatal projections
from posterior parietal subdivisions in the macaque monkey. Neuroscience 42:683–
696.

Chevalier G, Deniau JM (1990) Disinhibition as a basic process in the expression of striatal
functions. Trends Neurosci 13:277–280.

Cools R, Sheridan M, Jacobs E, D'Esposito M (2007) Impulsive Personality Predicts
Dopamine-Dependent Changes in Frontostriatal Activity during Component
Processes of Working Memory. J Neurosci 27:5506–5514.

Courtney SM, Petit L, Haxby JV, Ungerleider LG (1998) The role of prefrontal cortex in
working memory: examining the contents of consciousness. Philos Trans R Soc Lond
B Biol Sci 353:1819–1828.

Cowan N (2017) The many faces of working memory and short-term storage. Psychon Bull

Rev 24:1158–1170.

Cox RW, Chen G, Glen DR, Reynolds RC, Taylor PA (2017) FMRI Clustering in AFNI:

False-Positive Rates Redux. Brain Connectivity 7:152–171.

D'Ardenne K, Eshel N, Luka J, Lenartowicz A, Nystrom LE, Cohen JD (2012) Role of

prefrontal cortex and the midbrain dopamine system in working memory updating.

Proc Natl Acad Sci USA 109:19900–19909.

D'Esposito M, Postle BR (2015) The Cognitive Neuroscience of Working Memory. Ann Rev

Psych 66:115–142.

D'Esposito M, Postle BR, Rypma B (2000) Prefrontal cortical contributions to working

memory: evidence from event-related fMRI studies. Exp Brain Res 133:3–11.

Ecker UKH, Lewandowsky S, Oberauer K, Chee AEH (2010) The components of working

memory updating: An experimental decomposition and individual differences. J Exp

Psychol Learn Mem Cogn 36:170–189.

Eklund A, Nichols TE, Knutsson H (2016) Cluster failure: Why fMRI inferences for spatial

extent have inflated false-positive rates. Proc Natl Acad Sci U S A 113:7900–7905.

Erdfelder E, Faul F, Buchner A (1996) GPOWER: A general power analysis program. Behav

Res Methods Instrum Comput 28:1–11.

Fallon SJ, van der Schaaf ME, ter Huurne N, Cools R (2017) The Neurocognitive Cost of

Enhancing Cognition with Methylphenidate: Improved Distractor Resistance but

Impaired Updating. J Cogn Neurosci 29:652–663.

Feredoes E, Heinen K, Weiskopf N, Ruff C, Driver J (2011) Causal evidence for frontal

involvement in memory target maintenance by posterior brain areas during distracter

interference of visual working memory. PNAS 108:17510–17515.

Frank MJ, Loughry B, O'Reilly RC (2001) Interactions between frontal cortex and basal

ganglia in working memory: A computational model. Cogn Affect Behav Neurosci

1:137–160.

Fuster JM, Alexander GE (1971) Neuron Activity Related to Short-Term Memory. Science

173:652–654.

Goldman-Rakic PS (1996) The prefrontal landscape: implications of functional architecture

for understanding human mentation and the central executive. Philos Trans R Soc

Lond B Biol Sci 351:1445–1453.

Greve DN (2002) Optseq Home Page Available at http://surfer.nmr.mgh.harvard.edu/optseq

Jarbo K, Verstynen TD (2015) Converging Structural and Functional Connectivity of

Orbitofrontal, Dorsolateral Prefrontal, and Posterior Parietal Cortex in the Human

Striatum. J Neurosci 35:3865–3878.

Jonides J, Schumacher EH, Smith EE, Lauber EJ, Awh E, Minoshima S, Koeppe RA (1997)

Verbal Working Memory Load Affects Regional Brain Activation as Measured by

PET. J Cogn Neurosci 9:462–475.

Kanwisher N, McDermott J, Chun MM (1997) The Fusiform Face Area: A Module in Human

Extrastriate Cortex Specialized for Face Perception. J Neurosci 17:4302–4311.

Kessler Y (2017) The Role of Working Memory Gating in Task Switching: A Procedural

Version of the Reference-Back Paradigm. Front Psychol.

Kessler Y, Oberauer K (2014) Working memory updating latency reflects the cost of

switching between maintenance and updating modes of operation. J Exp Psychol

Learn Mem Cogn 40:738–754.

Kessler Y, Oberauer K (2015) Forward scanning in verbal working memory updating.

Psychon Bull Rev 22:1770–1776.

Kim C, Cilles SE, Johnson NF, Gold BT (2012) Domain general and domain preferential

brain regions associated with different types of task switching: A Meta-Analysis.

Hum Brain Mapp 33:130–142.

Lara AH, Wallis JD (2015) The Role of Prefrontal Cortex in Working Memory: A Mini

Review. Front Syst Neurosci.

Lewis-Peacock JA, Kessler Y, Oberauer K (2018) The removal of information from working

memory: The removal of information from working memory. Ann NY Acad Sci

1424:33–44.

Lopez-Garcia P, Lesh TA, Salo T, Barch DM, MacDonald AW, Gold JM, Ragland JD,

Strauss M, Silverstein SM, Carter CS (2016) The neural circuitry supporting goal

maintenance during cognitive control: a comparison of expectancy AX-CPT and dot

probe expectancy paradigms. Cogn Affect Behav Neurosci 16:164–175.

Majerus S, Cowan N, Péters F, Van Calster L, Phillips C, Schrouff J (2016) Cross-Modal

Decoding of Neural Patterns Associated with Working Memory: Evidence for

Attention-Based Accounts of Working Memory. Cereb Cortex 26:166–179.

Maldjian JA, Laurienti PJ, Kraft RA, Burdette JH (2003) An automated method for

neuroanatomic and cytoarchitectonic atlas-based interrogation of fMRI data sets.

NeuroImage 19:1233–1239.

McNab F, Klingberg T (2008) Prefrontal cortex and basal ganglia control access to working

memory. Nat Neurosci 11:103–107.

Miller EK, Cohen JD (2001) An Integrative Theory of Prefrontal Cortex Function. Annu Rev

Neurosci 24:167–202.

Murty VP, Sambataro F, Radulescu E, Altamura M, Iudicello J, Zoltick B, Weinberger DR,

Goldberg TE, Mattay VS (2011) Selective updating of working memory content

modulates meso-cortico-striatal activity. NeuroImage 57:1264–1272.

Narayanan NS, Prabhakaran V, Bunge SA, Christoff K, Fine EM, Gabrieli JDE (2005) The

Role of the Prefrontal Cortex in the Maintenance of Verbal Working Memory: An

Event-Related fMRI Analysis. Neuropsychology 19:223–232.

Nee DE, Brown JW, Askren MK, Berman MG, Demiralp E, Krawitz A, Jonides J (2013) A

Meta-analysis of Executive Components of Working Memory. Cereb Cortex 23:264–

282.

Oberauer K, Lewandowsky S, Awh E, Brown GDA, Conway A, Cowan N, Donkin C, Farrell

S, Hitch GJ, Hurlstone MJ, Ma WJ, Morey CC, Nee DE, Schweppe J, Vergauwe E,

Ward G (2018) Benchmarks for models of short-term and working memory. Psychol

Bull 144:885–958.

O'Reilly RC (2006) Biologically Based Computational Models of High-Level Cognition.

Science 314:91–94.

O'Reilly RC, Frank MJ (2006) Making Working Memory Work: A Computational Model of

Learning in the Prefrontal Cortex and Basal Ganglia. Neural Comput 18:283–328.

Owen AM, McMillan KM, Laird AR, Bullmore E (2005) N-back working memory paradigm:

A meta-analysis of normative functional neuroimaging studies. Hum Brain Mapp

25:46–59.

Poldrack RA, Baker CI, Durnez J, Gorgolewski KJ, Matthews PM, Munafò MR, Nichols TE,

Poline J-B, Vul E, Yarkoni T (2017) Scanning the horizon: towards transparent and

reproducible neuroimaging research. Nat Rev Neurosci 18:115–126.

Postle BR (2006) Working memory as an emergent property of the mind and brain.

Neuroscience 139:23–38.

Quentin R, King J-R, Sallard E, Fishman N, Thompson R, Buch ER, Cohen LG (2019)

Differential Brain Mechanisms of Selection and Maintenance of Information during

Working Memory. J Neurosci 39:3728–3740.

Rac-Lubashevsky R, Kessler Y (2016a) Dissociating working memory updating and

automatic updating: The reference-back paradigm. J Exp Psychol Learn Mem Cogn

42:951–969.

Rac-Lubashevsky R, Kessler Y (2016b) Decomposing the n-back task: An individual

differences study using the reference-back paradigm. Neuropsychologia 90:190–199.

Rac-Lubashevsky R, Kessler Y (2018) Oscillatory Correlates of Control over Working

Memory Gating and Updating: An EEG Study Using the Reference-back Paradigm. J

Cogn Neurosci 30:1870–1882.

Rac-Lubashevsky R, Slagter HA, Kessler Y (2017) Tracking Real-Time Changes in Working

Memory Updating and Gating with the Event-Based Eye-Blink Rate. Sci Rep 7:2547.

Roth JK, Serences JT, Courtney SM (2006) Neural System for Controlling the Contents of

Object Working Memory in Humans. Cereb Cortex 16:1595–1603.

Serences JT (2016) Neural mechanisms of information storage in visual short-term memory.

Vision Res 128:53–67.

Tzourio-Mazoyer N, Landeau B, Papathanassiou D, Crivello F, Etard O, Delcroix N, Mazoyer

B, Joliot M (2002) Automated Anatomical Labeling of Activations in SPM Using a

Macroscopic Anatomical Parcellation of the MNI MRI Single-Subject Brain.

NeuroImage 15:273–289.