Kinship does not predict the structure of a shark social network

Johann Mourier[1,2,3], Serge Planes[1,2]

[1] PSL Research University, EPHE-UPVD-CNRS, USR 3278, Université de Perpignan,

Perpignan, France

[2] Laboratoire d'Excellence "CORAIL," EPHE, PSL Research University, UPVD, CNRS,

USR 3278 CRIOBE, Papetoai, Moorea, French Polynesia

[3] UMR MARBEC (IRD, Ifremer, Univ. Montpellier, CNRS), Sète, France

ORCID: JM, 0000-0001-9019-1717; SP, 0000-0002-5689-5371

Corresponding author:

Johann Mourier

Email: johann.mourier@gmail.com

ABSTRACT:

Genetic relatedness in animal societies is often a factor that drives the structure of social

groups. In the marine world, most studies which have investigated this question have focused

on marine mammals such as whales and dolphins. For sharks, recent studies have

demonstrated preferential associations among individuals from which social communities

emerge. Assortment patterns have been found according to phenotypic or behavioural traits

but the role of genetic relatedness in shaping the social structure of adult shark populations

has, to the best our knowledge, never been investigated. Here, we used a social network

analysis crossed with DNA microsatellite genotyping to investigate the role of the genetic

relatedness in the social structure of a blacktip reef shark (*Carcharhinus melanopterus*)

population. Based on data from 156 groups of sharks, we used generalized affiliation indices

to isolate social preferences from non-social associations, controlling for the contribution of

sex, size, gregariousness, spatial and temporal overlap on social associations, to test for the

influence of genetic relatedness on social structure. Kinship was not a predictor of

associations and affiliations among sharks at the dyad or community levels as individuals

tended to associate independently of the genetic relatedness among them. The lack of

parental care in this species may contribute to the breakdown of family links in the

population early in life, thereby preventing the formation of kin-based social networks.

Keywords: genetic relatedness, assortment, social organisation, *Carcharhinus melanopterus*,

elasmobranch

INTRODUCTION

Group formation is an adaptive strategy, widespread across the animal kingdom, that can take various forms, from temporary unstable associations to long-term stable groups in complex societies (Krause and Ruxton 2002). Understanding the factors that influence the formation and evolution of social groups is important in order to understand the evolution of animal societies as well as to gain insight into population dynamics and to inform conservation strategy (Snijders et al. 2017). Associations among individuals can provide benefits to improve individual fitness, for example, by reducing predation risk or improving foraging efficiency (Krause and Ruxton 2002). While individuals can benefit by simply associating with other conspecifics (Kerth et al. 2011), the benefit of grouping can be enhanced by associating with similar individuals, also called social assortativity. By associating with individuals of the same size or the same sex, individuals are more likely to avoid conflict or harassment (Dadda et al. 2005) and their risk of predation is reduced via the confusion effect (Landeau and Terborgh 1986). Further, assorting with kin can also provide indirect fitness benefits (Hamilton 1964). Kin assortment has been shown to provide benefits in reducing aggression (Olsén and JäUrvi), increasing growth rate (Brown and Brown 1993) or allowing cooperative behaviour such as predator inspection (Milinski 1987).

Kin structuring has received extensive attention in many animal societies, especially where animals form stable breeding groups or where groups arise from the retention of offspring and delayed dispersal that facilitates the development of interactions with relative and kin-based groups (Wolf and Trillmich 2008; Hatchwell Ben J. 2010; Wiszniewski et al. 2010). In groups composed of relatives, kin selection should play a role in determining cooperation among group members (Hamilton 1964), although cooperation can arise also between non-kin (Clutton-Brock 2009). The role of relatedness in structuring animal societies that are characterised by a dynamic fission-fusion social system has been well studied in

species with parental care such as dolphins, giraffes, elephants or bats (Wittemyer et al. 2009; Wiszniewski et al. 2010; Kerth et al. 2011; Carter et al. 2013), but much less is known for species without parental care, as is the case for many species of fish (but see Croft et al. 2012). While the link between social networks and kinship has been extensively studied in terrestrial animals (Holekamp et al. 2012; Carter et al. 2013; Arnberg et al. 2015), kinship structure in social networks of marine and freshwater organisms has been primarily limited to marine mammals (Wiszniewski et al. 2010; Mann et al. 2012; Reisinger et al. 2017). Several cetacean societies show strong kin-based social network structures. However, in fishes, kin structure is less clear. Work on shoaling fish, for example, did not find kin assortment, even in species that are capable of kin discrimination (Croft et al. 2012). While sharks have recently been shown to be able to develop preferred associations and organise into structured social networks (Mourier et al. 2018), kinship has only been explored in one case study that focused on juvenile sharks (Guttridge et al. 2011) but did not find any clear influence of kinship in association patterns even for juvenile sharks, highlighting a lack of information on the potential for kin-based associations to arise in shark populations. Another study on spotted eagle rays did not find any evidence of relatedness in the formation of groups (Newby et al. 2014), although association strength was not quantified using association indices.

Overall, most studies that have explored the relationship between genetic relatedness and social interactions have focused on highly social species and in particular, on species that exhibit parental care (Wolf and Trillmich 2008; Wiszniewski et al. 2010; Kerth et al. 2011). Studying less social vertebrates should significantly improve our understanding of how social and genetic structure interact to shape the evolution of sociality in the animal kingdom.

In this study, we investigate the interaction between socio-spatial patterns and genetic relatedness in a population of blacktip reef sharks (*Carcharhinus melanopterus*) monitored over a 3-year period on the north shore of Moorea Island (French Polynesia). Sharks

represent an interesting and unique model to explore the extent to which individuals interact

with genetically related associates due to ecological traits that differ from most social

vertebrates. Like most social animals, sharks are now increasingly recognised as being

capable of complex social interactions, developing preferred social associations (Guttridge et

al. 2009; Jacoby et al. 2010; Mourier et al. 2012), showing unexpected learning abilities

(Guttridge et al. 2013; Mourier et al. 2017), developing foraging strategies by associating

with other species to improve predation success (Labourgade et al. 2020) and developing

patterns of leadership and dominance hierarchy (Guttridge et al. 2011; Jacoby et al. 2016;

Brena et al. 2018). However, contrary to many social organisms, reef sharks do not show

parental care and almost all shark species drop their progeny in specific nurseries outside

adult habitats and leave them to interact by themselves (Mourier and Planes 2013). These

discrete nurseries are chosen to potentially provide the neonates with a safe environment

where they will spend their first months of life. Recent studies suggested that females show

reproductive and even natal philopatry to these particular birthing grounds (Mourier and

Planes 2013; Feldheim et al. 2014), suggesting that newborn sharks may have the opportunity

to develop strong relationships with close kin. When juvenile sharks reach a certain size or

age, they leave their nursery to explore a wider home range (Chapman et al. 2009); they then

integrate within the adult population and start interacting with older individuals, but it is not

known whether they coexisted with other newborn during their juvenile stage or disperse

alone. Therefore, while aggregations of kin are possible during the early stages, it is currently

unknown if they persist through adulthood after dispersal. In shark populations, interactions

between kin are also diluted by the presence of numerous neighbours and average relatedness

quickly drops with increasing group size. In some shark species, the likelihood of associating

with a related peer is reduced due to small litter size and a high mortality rate at the juvenile

stage, leading to a lack of first order relatives to reach adulthood. However, in a closed

system, such as an isolated island, and in the case of blacktip reef sharks which spend their

entire life cycle within Moorea (Mourier and Planes 2013), relatives will have more chances

to encounter each other and to interact in social groups. Thus, in these conditions, the limited

number of related pairs might decrease the risk of inbreeding.

To understand the assortative forces which underpin the structural properties of the

system is challenging for elusive underwater animals. As the blacktip reef shark displays a

high degree of site fidelity (Papastamatiou et al. 2009) and shares some of its areas with

many conspecifics (Mourier et al. 2012), exploring this network holds the potential to work

out the relationship between spatial, social and genetic structure in a reef shark. Size, sex and

gregariousness of sharks have been shown to influence assortment at the population and

community levels (Mourier et al. 2012; Mourier et al. 2017). However, whether genetic

relatedness plays a role in structuring the network at both the individual and community

levels remain unknown. In particular, whether sharks benefit from associating with kin

remains unknown as cooperation has not been proven and social foraging may not require

associations with kin to improve predation success (Labourgade et al. 2020).

We aim to describe the social structure and to determine whether sharks had genuine

social preferences (caused by active choice of individuals to interact) by controlling for non-

social structural factors, including space use, time, phenotype, and individual gregariousness.

We then tested whether the social structure at different scales can be explained by the genetic

relatedness between individuals.

MATERIAL AND METHODS

Field observations and data collection

Between 2008 and 2010, observation surveys were conducted along approximately 10km

of coastline of the Northern reef of Moorea Island (French Polynesia) (Figure 1). The surveys

consisted of 40 min dives at 7 sites along a 10 km portion of reef (total = 180 dives, site 1 =

20, site 2 = 50, site 3 = 8, site 4 = 33, site 5 = 30, site 6 = 34 and site 7 = 7). Individual

blacktip reef sharks were identified by photo-identification, using unique, lifelong colour-

shape of the dorsal fin (Mourier et al. 2012).

Associations between individuals were defined using the "Gambit of the Group"

(Whitehead and Dufault 1999) assuming that all individuals observed together are then

considered as "associated". This approach is appropriate when individuals move between

groups and direct interactions are difficult to observe, but where groups can be easily defined

(Franks et al. 2010; Farine and Whitehead 2015). An experienced diver conducted a

stationary visual census at each site monitored, moving and identifying sharks within a ~100

m radius area (made possible by the high visibility conditions in these tropical waters). All

individuals observed during a dive were considered as part of the same group if they were

encountered within 10 min periods. We are confident that observed associations represented

true grouping structure, because groups were spatio-temporally well-defined and sharks were

engaged into specific social behaviour (e.g. following, parallel swimming or milling; Mourier

et al. 2012). To avoid the potential for weak and non-relevant associations between pairs of

individuals with very low number of sightings, we used a restrictive observation threshold to

include only individuals observed more than the median number of sightings (median = 14;

mean ± sd = 14.92 ± 8.04; Supplementary Figure S1). Thus, all individuals seen less than 15

times were removed from the analyses to ensure that associations were estimated with high

accuracy and precision.

DNA sampling and laboratory procedures

Shark fishing sessions using rod and reel and barbless hooks were conducted to obtain tissue

samples for genetic analysis. Once hooked, sharks were brought alongside the boat where

they were inverted and placed in tonic immobility while biological data and tissue samples

were collected. Each shark was identified by photo-identification of the dorsal fin, sexed and

measured to the nearest centimeter (Mourier et al. 2012; Mourier, Mills, et al. 2013). Fishing

sessions were conducted directly after underwater surveys to avoid perturbations of the

experimental setup (Mourier et al. 2017) and to increase the chance of getting DNA samples

from sharks that were part of the social network. Fishing effort was maintained until sharks

failed to respond to the bait (generally <30 min and after catching 2-3 individuals). A fin clip

was collected from the second dorsal fin or anal fin and samples were individually preserved

in 95% ethanol and returned to the laboratory for genotyping (Mourier and Planes 2013).

DNA was extracted using the QIAGEN DX Universal Tissue Sample DNA Extraction

protocol. PCR amplification and the microsatellite loci used are described in detail in

previous studies (Mourier and Planes 2013; Vignaud et al. 2013; Vignaud et al. 2014). The

software MICROCHECKER (Van Oosterhout Cock et al. 2004) was used to test for null

alleles and other genotyping errors.

We compared the suitability of seven pairwise relatedness estimators: five non-likelihood

estimators (Queller and Goodnight 1989; Li et al. 1993; Ritland 1996; Lynch and Ritland

1999; Wang 2002) and two maximum-likelihood estimators (Milligan 2003; Wang 2007) in

the R package *related* (Pew et al. 2015) and determined that the triadic maximum-likelihood

estimator (TrioML; Wang 2007) was best suited to our microsatellite panel (Supplementary

materials, Supplementary Figure S2) as it showed the highest correlation (i.e. 0.831) with the

true values and the smallest variation around the mean for every relationship (except for full-

sibs). This analysis generates simulated individuals of known relatedness based on the

observed allele frequencies and calculates the genetic relatedness using the different

estimators. The correlation between observed and expected genetic relatedness was obtained

for each estimator, and the one with the highest correlation coefficient was selected for

further analysis.

Defining associations

Using R package *asnipe* (Farine 2013), we calculated dyadic association strengths (i.e.

associations among pairs of individuals) among photo-identified individual sharks seen in

groups from the spatio-temporal co-occurrences, and the proportion of time two individuals

were observed together at the same site given that at least one was observed, using the

simple-ratio association index (SRI) (Cairns and Schwager 1987). The SRI is the

recommended association index when calibration data are unavailable (Hoppitt and Farine

2018).

To measure the diversity of associations, we calculated the social differentiation (S) in

the network that is the estimated coefficient of variation (standard deviation divided by mean)

of the true association indices. If the social differentiation of the network is 0, then

relationships among members are completely homogeneous. Conversely, if the social

differentiation is above 1.0, there is considerable diversity in the relationships between pairs

of individuals within the network (Whitehead 2008). For our data, the standard error of S was

generated by bootstrapping (1 000 replications).

Potential structural factors of social associations

We quantified five structural factors that could affect shark association patterns: spatial

overlap, temporal overlap, gregariousness, and size and sex similarity for each pair of

individuals. Genetic relatedness was not included as a structural factor as it will be tested

independently when other factors are extracted.

For each individual, an encounter rate (i.e., no. sightings of individual at site, divided by

no. sampling occasions at site) was calculated by site to define individual spatial utilization

(Supplementary Figure S3). We then generated a Bray-Curtis similarity matrix of space use

to construct a matrix of spatial overlap between individuals using R package "vegan" (Dixon

2003).

Individuals using an area at the same time are more likely to be associated with each

other. The study period corresponds to a total of 28 months between February 2008 and June

2010. The temporal overlap was calculated as the custom SRI calculated on whether pairs

were observed in the study area within sampling periods of 60 days.

Gregariousness was calculated following Whitehead and James's (2015) correction,

where the gregariousness predictor between two individuals (a and b) is the log of the sum of

the association indices involving a (except the ab index) multiplied by the sum of those

involving b (except the ba index): $G_{ab} = \log(\Sigma SRI_a \Sigma SRI_b)$ where ΣSRI_a and ΣSRI_b are the

sums of all the SRIs for individuals a and b, respectively.

Shark length was classified into size classes ranging from 1 to 6 (1: TL < 110 cm; 2:

110-119 cm; 3: 120-129 cm; 4: 130-139 cm; 5: 140-150 cm; 6: TL > 150 cm). For sex and

size similarity, we constructed a binary matrix in which elements $a_{ij} = 1$ when individuals i

and j were of the same class and $a_{ij} = 0$ otherwise (sex class, 1 if same sex, 0 if not; size class,

1 if same size class, 0 if not).

Influence of structural factors on social associations

We quantified the contribution of all five structural factors in driving social patterns with

a multiple regression quadratic assignment procedure (MRQAP) modified by Farine (2013)

that enables null models built from pre-network data permutations to be used in conjunction

with a MRQAP regression. This approach was shown to be more accurate than classic

MRQAP procedures (Farine 2017). We assessed possible linear relationships between the

social associations and the structural factors using the SRI association matrix as the

dependent variables and the matrices representing pairwise similarity of each of the five

structural factors as independent variables. We used 20,000 permutations to build randomized

distributions to compare with the empirical coefficient. The *P*-values were the proportion of

the estimated coefficient regression which were smaller or greater than what would have been

expected by chance. We used the *mrqap.custom.null* function from *asnipe* R package (Farine

2013) to run MRQAP tests in R v. 3.3.0 (R Core Team 2019).

Removing the effects of structural factors from associations

We developed generalized affiliation indices (GAI, Whitehead and James 2015) to

remove the effects of the significant structural factors from the associations and test the

existence of true affiliations between dyads (i.e. active association preferences). For this, we

fitted a binomial generalized linear model (GLM) with the unfolded SRI matrix as the

dependent variable, and the significant structural factors selected from the MRQAP as

independent variables. GAI represents the assortment of individuals not explained by the

significant structural factors and corresponds to the deviance residuals of the model. The

model was: SRI ~ TO + SO, where SRI is the association matrix, TO is the temporal overlap

matrix and SO is the spatial overlap matrix (as only TO and SO were significant factors in the

MRQAP, Table 1).

Social preferences and null models

We used a null model to test both for social preferences and the significance of the

observed network modularity. We generated 20,000 randomized association and affiliation

networks based on 25,000 data-stream permutations of the raw observation data with a

swapping algorithm (Bejder et al. 1998). We permuted the empirical group-by-individual

matrix constraining the number of groups, individuals and occurrences (matrix dimension

and fill), group size (row totals) and individual frequency of observation (column totals). To

minimize the effect of initial values potentially correlated to the empirical data, we removed

the first 5,000 randomized matrices. From the randomised group-by-individual matrix, we

calculated a simple-ratio index association matrix, with which we built a generalized

affiliation index using the same predictors selected via MRQAP for the empirical data. We

used a modified version of R codes available from Machado et al. (2019) to build null models

and to calculate SRI, GAI and modularity.

We compared the standard deviation (SD) of the observed simple-ratio index (SRI) and

the SD of the observed generalized affiliation index (GAI) with the distribution of the SD of

corresponding randomized SRI and GAI matrices generated by the null models detailed

above. An observed SD significantly higher than the null expectation indicates the presence

of preferred and avoided associations and affiliations. We also tested for strongly connected

social communities by comparing the empirical modularity (Q) (Newman 2006) of SRI and

positive GAI matrices with that of the randomized matrices. Empirical SD and Q values were

considered statistically significant if they fell outside the 95% confidence interval of their

randomized distributions.

Genetic relatedness, social structure and sex differences

To assess whether relatedness differs for same-sex dyads, we constructed three binary

matrices (0,1), each encoding the presence of a certain dyad type (female–female, male–male

or female–male). We then tested for a correlation with the relatedness matrix using three

Mantel tests (20,000 permutations), via the vegan R package (Dixon 2003).

For each type of dyad (female–female, male–male or female–male), we then tested for a

correlation between the SRI and GAI matrices and the pairwise genetic relatedness among

sharks using Mantel tests and compared the test statistics to those of the 20,000 permuted

networks.

We also compared the gregariousness of individual sharks between the sexes. For this,

we used two measures of gregariousness: node degree (or binary degree) which is the number

of direct neighbours each individual is connected to in the network and node strength (or

weighted degree) that is the sum of associations of an individual. We then used these network

metrics in order to determine whether males and females differed in their gregariousness. We

constructed generalized linear models (GLMs) to test how sex affected the observed network

degree (degree ~ sex) and strength (strength ~ sex). We ran these same models with

randomized permutations of the network data to evaluate statistical significance (Farine and

Whitehead 2015; Farine 2017).

To determine whether individuals within groups (size class and communities) were more

or less closely related than expected, we compared the observed values for each group against

a distribution of expected relatedness values generated by randomly shuffling individuals

between groups for 1000 permutations, where size was kept constant, using the R

package *related* (Pew et al. 2015). If the observed mean relatedness was greater than that of

the permuted data, then the null hypothesis which predicted that the mean within-community

relatedness is random, was rejected.

If only a few closely related individuals were present, then it is possible that their within-

community overabundance compared to between social communities might not be detected

using mean coefficient of relatedness (Buston et al. 2009). In turn, we verified whether the

proportion of closely related pairs was higher within than between social communities using

a chi-squared test following the same approach as the preceding analysis with mean

relatedness. We compared the χ^2 statistics of the observed difference in proportions of

relatedness values above a certain threshold between within- and between-communities to

that of expected relatedness values generated by randomly shuffled individuals between

community groups for 1000 permutations and keeping size constant. We tested with a

threshold relatedness value of 0.25 corresponding to the theoretical relatedness of half-sibs.

RESULTS

Data summary

Of 241 catalogued sharks (150 males, 91 females; Mourier et al. 2012), 49 (36 males, 13

females) were observed on more than 15 occasions (mean resightings = 14.92 ± 8.04 SD,

Supplementary Figure S1). A total of 225 adult sharks were genotyped from the studied area.

From the 49 sharks included in our social network analyse, 87% (43) were genotyped.

Therefore, 43 individuals (30 males, 13 females) were included in the remaining analyses.

This resulted in 156 observed groups (mean group size = 8.60 ± 4.92 SD). From the 17

microsatellite markers taken from our previous study (Mourier and Planes 2013), the

presence of null alleles was detected at Cli12 which was then removed from our dataset for

further genetic analyses. We conducted the genetic analyses with 16 loci (Supplementary

Table S1).

Social structure

The social differentiation of the population was higher than 1 (S ± SE = 1.474 ± 0.037),

revealing a diverse range of associations and a well-differentiated society. The most

significant predictors of shark associations were the temporal and spatial overlaps, which

explained 94% of the total variance in SRI (MRQAP results, Table 1).

We rejected the null hypothesis that sharks associate randomly, as the observed SD of

SRI was higher than the random SD. When GAI removed the influence of temporal and

spatial overlaps from SRI, we also rejected the null hypothesis of random affiliations (Figure

2a) demonstrating the presence of preferred social affiliations. At the population level, the

modularity (Q) of the association (SRI) and affiliation (GAI) networks were higher than

expected by chance (Figure 2b). While the three communities from the SRI network had

distinct use of space, some communities from the GAI network had similar spatial

distributions (e.g., communities yellow and purple, Figure 2c).

Crossing of genetic relatedness and association patterns

When testing for kin-biased relatedness, adult male-male (MM), female–female (FF) and

male-female (MF) dyads did not have clear higher or lower genetic relatedness (Mantel test:

MM, r = -0.012, n = 31, P = 0.061; FF, r = -0.019, n = 12, P = 0.685; MF, r = 0.023, n = 43, P

= 0.259). In addition, mean genetic relatedness was not higher within than between size

classes (mean r = 0.043, random 95% CI = 0.058-0.065, P = 0.88; Supplementary materials,

Table S2). While individuals were relatively spatially clustered, genetic relatedness appeared

much more homogeneously distributed across individuals and space (Mantel test between

matrices of spatial overlap and genetic relatedness: r = 0.011, n = 43, P = 0.351;

Supplementary Figure S4).

Average pairwise relatedness among individuals was 0.062 ± 0.001 (mean ± SE) ranging

from 0 to 0.774. Associations were only significantly positively correlated with genetic

relatedness between males (Mantel test: r= 0.103, P = 0.026) but no significant correlation

was found between GAI and genetic relatedness for any sex dyad (Table 2). Males were

generally more gregarious than females, as they significantly interacted with more

individuals (higher degree) but did not have stronger relationships (higher strength) (Table 3,

Figure 3).

Within-community relatedness estimate was inferred for each community and index (SRI

and GAI, Table 4). Relatedness within all communities was not higher than expected if

communities were randomly organized (SRI network: within mean ± SE = 0.071 ± 0.005,

between mean ± SE = 0.058 ± 0.003, P = 0.093; GAI network: within mean ± SE = 0.060 ±

0.005, between mean ± SE = 0.063 ± 0.005, P = 0.722) (Table 4).

Among the 903 potential pairs, 39 (4.31%) had values higher than 0.25. In addition, there

was no higher proportion of close relatives within than between communities for relatedness

value r > 0.25 for SRI (chi-squared test: $n_{within/between}$ = 17/ 22, d.f. = 1, χ^2 = 0.928, P = 0.119)

and GAI (chi-squared test: $n_{within/between}$ = 8/35, d.f. = 1, χ^2 = 0.386, P = 0.326) (Figure 4).

Together, these results suggest that no differences exist for within- and between-

community membership with respect to the genetic relatedness of their members.

DISCUSSION

We found a fine-scale social structure in blacktip reef sharks in Moorea. Taking into

account the confounding effects of 5 structural variables (spatial and temporal overlap,

gregariousness, size and sex), which are known to influence association patterns (e.g. Godde

et al. 2013; Diaz-Aguirre et al. 2019; Machado et al. 2019; Perryman et al. 2019), we found

that blacktip reef sharks in Moorea had both preferred associations and affiliations. However,

social proximity was not predicted by the genetic relatedness between sharks both at the

association, affiliation and community levels. At the dyad level, only male-male associations,

but not affiliations, were slightly correlated with genetic relatedness. In addition, individuals

had low probabilities of interacting with a close-kin which could explain the lack of influence

of kinship in structuring the social network in this population. These results therefore suggest

that genetic relatedness does not drive the structure of the social network in this shark

population.

Compared to previous work conducted on this population (Mourier et al. 2012) which

only analysed social structure through associations among sharks, the present study also

considered the effects of two structural variables to estimate affiliation indices. Affiliations

are an increasingly used method to investigate the true social interactions experienced by

animals (Whitehead and James 2015), in particular by considering the strong correlation that

can exist between space use overlap and association indices in a variety of taxa (e.g., Mourier

et al. 2012; Carter et al. 2013; Best et al. 2014). In our study, the network built from

associations using the simple ratio index (SRI) was composed of three main communities

relatively spatially separated and only low overlap (Figure 2). When removing the influence

of spatial and temporal overlap from association patterns, the network built from the

generalized affiliation indices (GAI) revealed five communities that were less spatially

separated. This means that associations between sharks were the result of more than just

similarities in habitat use or overlaps in time, indicating that individuals actively chose to

group with preferred social partners. The differences between the three SRI communities in

the present study and the four communities found in Mourier et al. (2012) can be due to the

high threshold we used that may highlight only string relationships and the use of SRI instead

of HWI. To our knowledge, only one study on elasmobranchs has investigated social

structure using GAI, demonstrating that manta rays also preferred affiliations (Perryman et al.

2019). Individuals' site preferences and being present in the study at the same time was a

strong predictor of association between pairs. Site fidelity is often a prerequisite for sociality,

creating an environment for social relationships to develop and the emergence of social

preferences. However, the presence of preferred social affiliations demonstrates that sharks

show active social preferrences that do not rely on preferrences for sites and periods. Our

study confirms that the observed shark social structure resembles that of a fission-fusion

society characterized by an open and fluid social structure, long-term social recognition and a

high number of potential affiliates, which is flexible depending on environemental

conditions.

Among the adult sharks in our population, there was a generally low level of

relatedness, and only a small number of dyads had close familial relatives. Interactions

frequently occurred between distant kin and non-kin. This implies that the social structure

among adult blacktip reef sharks was not based on associations between close kin as

demonstrated by our analyses which compared association and affiliation patterns with

genetic relatedness among dyads at the pairwise or community levels. This is confirmed by

the low number of close kin available for each shark in the population (< 6 % pairs with r >

0.25), thereby limiting the probability of an individual to encounter a family member and to

develop strong associations with them. The lack of genetic relatedness structure within size

classes and the lack of decreasing genetic relatedness as sharks grow also suggests that

juveniles are unlikely to leave their nursery ground with other kin. If young sharks were

developing and maintaining strong bonds with their littermates throughout their entire life,

we would have expected to find high mean relatedness and high proportion of close kin

across all size classes. The low relatedness we found within each size class indicates that

sharks favoured associations with non-kin. These results can be explained in part by the life

history and life cycle of blacktip reef sharks. In fact, in contrast to most social animals that

show some forms of family structure and parental care, female reef-associated sharks such as

blacktip reef shark, leave their pups in their nursery after birth (Mourier and Planes 2013).

Moreover, litter size in this species does not exceed five pups (Mourier, Mills, et al. 2013)

while litter size in Moorea was limited to a maximum of two pups (Mourier and Planes

2013). In addition, blacktip reef sharks follow a yearly breeding cycle with females giving

birth every year and potentially being fertilized by multiple males within or across years,

which increases the probability of having maternal and paternal half-siblings. Our ongoing

long-term nursery monitoring shows that capture probabilities rapidly decline after March

(unpublished data), 2 to 3 months after parturition, which suggests a dramatic mortality rate

within the nursery areas during the first months of life (i.e. survival rate expected to be

inferior to 50% during the first year of life). Together with a small litter size and absence of

parental care, this high mortality rate, which is common in many shark species, is likely to

limit the opportunity to find family members and develop strong affiliations with close

relatives at adulthood. Even in nurseries, juvenile lemon sharks did not clearly assort by

relatedness (Guttridge et al. 2011), even if the probability of finding a relative is higher for

this species with a larger litter size. When juvenile sharks grow, they progressively explore

their environment and increase their home range (Chin A et al. 2013), creating an opportunity

to find related individuals such as parents or maternal half-siblings from previous

reproductive seasons. At adulthood, our results confirm that preferred associations and

affiliations are not driven by genetic relatedness as sharks are associating with conspecifics of

variable genetic distances. This suggests that sharks might not have the ability for kin

recognition simply based on visual or olfactory cues and that kin-based preferred associations

and affiliations may only develop within nursery areas from increased familiarity with

littermates, or that they are not seeking for associations with related individuals. Through

investigation of social groups of spotted eagle rays *Aetobatus narinari* in Florida, Newby et

al. (2014) found no kin-structure in the social organization, although the analysis was based

on group composition rather that quantitatively inferred using association indices. However,

our results revealed that males sighly preferred to associate with other related males but this

tendency was not confirmed for affiliations (accounting for spatial and temporal structural

components). This suggests that genetic relatedness among males was spatially structured

and that males may disperse less than females. The lack of differences in relatedness between

males and females suggests that the risk of inbreeding might be low if these interactions

represented potential mating pairs and not only social bonds.

The emerging literature suggests that genetic structure of animal social networks can

vary dramatically, from highly cohesive kin-based groups like African elephants (*Loxodonta*

africana) (Archie et al. 2006) or spotted hyenas (*Crocuta crocuta*) (Holekamp et al. 2012), to

groups with moderate levels of genetic relatedness due to limited dispersal like the Galápagos

sea lion (*Zalophus wollebaeki*) (Wolf and Trillmich 2008) or the eastern grey kangaroo

(*Macropus giganteus*) (Best et al. 2014), or to groups with little to no genetic relatedness like

guppies (*Poecilia reticulata*) (Croft et al. 2012), the common raccoon (*Procyon lotor*)

(Hirsch et al. 2013) or migratory golden-crowned sparrows (*Zonotrichia atricapilla*)

(Arnberg et al. 2015). These patterns of variation provide opportunities to explore how

ecological factors interact with kinship to produce variations in the structures of animal

societies. Kinship is expected to promote the evolution of cooperation and sociality in

animals (Hamilton 1964). However, our understanding of the evolution of sociality results to

a great extent from the study of closed societies, in which interactions mainly involve

relatives and can hence be explained by kin selection (Hamilton 1964). However, the kin

selection theory has recently been challenged by results from studies showing that fitness

benefit can emerge in social groups composed mainly of non-relatives (e.g., Cameron et al.

2009; Riehl 2011; Wilkinson et al. 2016). In many natural populations, dispersal tends to be

limited, favouring local competition between neighbours and the emergence of a social

component, whether it be selfish, aggressive, cooperative or altruistic (Lehmann and Rousset

2010). But how social behaviours translate into fitness costs and benefits depends

considerably on life-history features, as well as on local demographic and ecological

conditions. The fission – fusion social dynamics lead to unstable group membership, and

dispersal and occasional recruitment of unrelated individuals lead to low average relatedness

in groups. Then under such conditions, selection is not expected to favour kin recognition

mechanisms based on familiarity alone.

Therefore, contrary to the kin selection hypothesis which predicts stronger associations

among kin, sharks tended to assort randomly according to relatedness. As kinship does not

explain the strength of social affiliations in blacktip reef sharks, the question remains as to

how and why sharks form preferred associations and affiliations organised in social

communities (Mourier et al. 2012). Although cooperation has been mainly explained in the

context of kin selection, there might be potential benefits of non-kin sociality in blacktip reef

sharks such as for other animals in which association with non-kin emerges via reciprocal

altruism (Carter and Wilkinson 2013; Wilkinson et al. 2016). While evidence of shark

cooperation has not been confirmed, gregarious behaviour can have several benefits in sharks

(Jacoby et al. 2012), including increased foraging success by hunting in groups (Weideli et al.

2015; Mourier et al. 2016), protection from predators (Mourier, Planes, et al. 2013), or

increased tolerance relationships and reduced aggression rate (Brena et al. 2018).

Heterospecific foraging associations have been found to develop and increase predation

success (Labourgade et al. 2020), which suggests that sharks can benefit from hunting

associations without associating with kin. These benefits do not necessarily imply kin

selection and can simply build on the development of familiarity from repeated interactions.

Social structure in reef sharks can arise from multiple simple ecological factors such as the

distribution of resources in space and time leading to aggregations of individuals even in the

absence of benefits of direct social affiliation (Ramos-Fernández et al. 2006) or mitigation of

the cost of unnecessary aggression when competing for resources mediated by individual

recognition (Brena et al. 2018). Regardless of the exact cause of social preferences in reef

sharks, the absence of kinship as an important factor in structuring association patterns

suggests that there are important benefits of sociality in sharks that we still need to uncover.

With an increasing use of social network analyses applied to shark populations (Mourier et al.

2018), future work on social networks and genetic relatedness in different populations or

species is necessary to confirm our results and to improve our understanding of population

dynamics in sharks and the evolution of sociality.

SUPPLEMENTARY MATERIAL

Supplementary information including two tables and four figures can be found online.

REFERENCES

Archie EA, Moss CJ, Alberts SC. 2006. The ties that bind: genetic relatedness predicts the
fission and fusion of social groups in wild African elephants. Proc R Soc Lond B Biol Sci.
273(1586):513–522. doi:10.1098/rspb.2005.3361.

Arnberg NN, Shizuka D, Chaine AS, Lyon BE. 2015. Social network structure in wintering
golden-crowned sparrows is not correlated with kinship. Mol Ecol. 24(19):5034–5044.

Bejder L, Fletcher D, Bräger S. 1998. A method for testing association patterns of social
animals. Anim Behav. 56(3):719–725. doi:10.1006/anbe.1998.0802.

Best EC, Dwyer RG, Seddon JM, Goldizen AW. 2014. Associations are more strongly
correlated with space use than kinship in female eastern grey kangaroos. Anim Behav. 89:1–
10. doi:10.1016/j.anbehav.2013.12.011.

Brena PF, Mourier J, Planes S, Clua EE. 2018. Concede or clash? Solitary sharks competing
for food assess rivals to decide. Proc R Soc B. 285(1875):20180006.
doi:10.1098/rspb.2018.0006.

Brown GE, Brown JA. 1993. Social dynamics in salmonid fishes: do kin make better
neighbours? Anim Behav. 45(5):863–871. doi:10.1006/anbe.1993.1107.

Buston PM, Fauvelot C, Wong MYL, Planes S. 2009. Genetic relatedness in groups of the
humbug damselfish Dascyllus aruanus: small, similar-sized individuals may be close kin.
Mol Ecol. 18(22):4707–4715. doi:10.1111/j.1365-294X.2009.04383.x.

Cairns SJ, Schwager SJ. 1987. A comparison of association indices. Anim Behav.
35(5):1454–1469. doi:10.1016/S0003-3472(87)80018-0.

Cameron EZ, Setsaas TH, Linklater WL. 2009. Social bonds between unrelated females
increase reproductive success in feral horses. Proc Natl Acad Sci. 106(33):13850–13853.
doi:10.1073/pnas.0900639106.

Carter GG, Wilkinson GS. 2013. Food sharing in vampire bats: reciprocal help predicts
donations more than relatedness or harassment. Proc R Soc B Biol Sci. 280(1753):20122573.
doi:10.1098/rspb.2012.2573.

Carter KD, Seddon JM, Frère CH, Carter JK, Goldizen AW. 2013. Fission–fusion dynamics
in wild giraffes may be driven by kinship, spatial overlap and individual social preferences.
Anim Behav. 85(2):385–394. doi:10.1016/j.anbehav.2012.11.011.

Chapman DD, Babcock EA, Gruber SH, Dibattista JD, Franks BR, Kessel SA, Guttridge T,
Pikitch EK, Feldheim KA. 2009. Long-term natal site-fidelity by immature lemon sharks
(Negaprion brevirostris) at a subtropical island. Mol Ecol. 18(16):3500–3507.
doi:10.1111/j.1365-294X.2009.04289.x.

Chin A, Heupel MR, Simpfendorfer CA, Tobin AJ. 2013. Ontogenetic movements of
juvenile blacktip reef sharks: evidence of dispersal and connectivity between coastal habitats
and coral reefs. Aquat Conserv Mar Freshw Ecosyst. 23(3):468–474. doi:10.1002/aqc.2349.

Clutton-Brock T. 2009. Cooperation between non-kin in animal societies. Nature.
462(7269):51–57. doi:10.1038/nature08366.

Croft DP, Hamilton PB, Darden SK, Jacoby DMP, James R, Bettaney EM, Tyler CR. 2012.
The role of relatedness in structuring the social network of a wild guppy population.
Oecologia. 170(4):955–963. doi:10.1007/s00442-012-2379-8.

Dadda M, Pilastro A, Bisazza A. 2005. Male sexual harassment and female schooling
behaviour in the eastern mosquitofish. Anim Behav. 70(2):463–471.
doi:10.1016/j.anbehav.2004.12.010.

Diaz-Aguirre F, Parra GJ, Passadore C, Möller L. 2019. Genetic relatedness delineates the
social structure of southern Australian bottlenose dolphins. Behav Ecol. 30(4):948–959.
doi:10.1093/beheco/arz033.

Dixon P. 2003. VEGAN, a package of R functions for community ecology. J Veg Sci.
14(6):927–930. doi:10.1111/j.1654-1103.2003.tb02228.x.

Farine DR. 2013. Animal social network inference and permutations for ecologists in R using
asnipe. Methods Ecol Evol. 4(12):1187–1194. doi:10.1111/2041-210X.12121.

Farine DR. 2017. A guide to null models for animal social network analysis. Methods Ecol
Evol.:n/a-n/a. doi:10.1111/2041-210X.12772.

Farine DR, Whitehead H. 2015. Constructing, conducting and interpreting animal social
network analysis. J Anim Ecol. 84(5):1144–1163. doi:10.1111/1365-2656.12418.

Feldheim KA, Gruber SH, DiBattista JD, Babcock EA, Kessel ST, Hendry AP, Pikitch EK,
Ashley MV, Chapman DD. 2014. Two decades of genetic profiling yields first evidence of
natal philopatry and long-term fidelity to parturition sites in sharks. Mol Ecol. 23(1):110–
117. doi:10.1111/mec.12583.

Franks DW, Ruxton GD, James R. 2010. Sampling Animal Association Networks with the
Gambit of the Group. Behav Ecol Sociobiol. 64(3):493–503.

Godde S, Humbert L, Côté SD, Réale D, Whitehead H. 2013. Correcting for the impact of
gregariousness in social network analyses. Anim Behav. 85(3):553–558.
doi:10.1016/j.anbehav.2012.12.010.

Guttridge TL, Dijk S van, Stamhuis EJ, Krause J, Gruber SH, Brown C. 2013. Social learning
in juvenile lemon sharks, Negaprion brevirostris. Anim Cogn. 16(1):55–64.
doi:10.1007/s10071-012-0550-6.

Guttridge TL, Gruber SH, DiBattista JD, Feldheim KA, Croft DP, Krause S, Krause J. 2011.
Assortative interactions and leadership in a free-ranging population of juvenile lemon shark
Negaprion brevirostris. Mar Ecol Prog Ser. 423:235–245. doi:10.3354/meps08929.

Guttridge TL, Gruber SH, Gledhill KS, Croft DP, Sims DW, Krause J. 2009. Social
preferences of juvenile lemon sharks, Negaprion brevirostris. Anim Behav. 78(2):543–548.
doi:10.1016/j.anbehav.2009.06.009.

Hamilton WD. 1964. The genetical evolution of social behaviour. I. J Theor Biol. 7(1):1–16.
doi:10.1016/0022-5193(64)90038-4.

Hatchwell Ben J. 2010. Cryptic Kin Selection: Kin Structure in Vertebrate Populations and
Opportunities for Kin-Directed Cooperation. Ethology. 116(3):203–216. doi:10.1111/j.1439-
0310.2009.01732.x.

Hirsch BT, Prange S, Hauver SA, Gehrt SD. 2013. Genetic relatedness does not predict
racoon social network structure. Anim Behav. 85(2):463–470.
doi:10.1016/j.anbehav.2012.12.011.

Holekamp KE, Smith JE, Strelioff CC, Van Horn RC, Watts HE. 2012. Society, demography
and genetic structure in the spotted hyena. Mol Ecol. 21(3):613–632. doi:10.1111/j.1365-
294X.2011.05240.x.

Hoppitt WJE, Farine DR. 2018. Association indices for quantifying social relationships: how
to deal with missing observations of individuals or groups. Anim Behav. 136:227–238.
doi:10.1016/j.anbehav.2017.08.029.

Jacoby DMP, Busawon DS, Sims DW. 2010. Sex and social networking: the influence of
male presence on social structure of female shark groups. Behav Ecol. 21(4).
doi:10.1093/beheco/arq061. [accessed 2017 Jan 4].
http://beheco.oxfordjournals.org/content/early/2010/05/09/beheco.arq061.

Jacoby DMP, Croft DP, Sims DW. 2012. Social behaviour in sharks and rays: analysis,
patterns and implications for conservation. Fish Fish. 13(4):399–417. doi:10.1111/j.1467-
2979.2011.00436.x.

Jacoby DMP, Papastamatiou YP, Freeman R. 2016. Inferring animal social networks and
leadership: applications for passive monitoring arrays. J R Soc Interface. 13(124):20160676.
doi:10.1098/rsif.2016.0676.

Kerth G, Perony N, Schweitzer F. 2011. Bats are able to maintain long-term social
relationships despite the high fission–fusion dynamics of their groups. Proc R Soc Lond B
Biol Sci. 278(1719):2761–2767. doi:10.1098/rspb.2010.2718.

Krause J, Ruxton GD. 2002. Living in Groups. OUP Oxford.

Labourgade P, Ballesta L, Huveneers C, Papastamatiou Y, Mourier J. 2020. Heterospecific
foraging associations between reef-associated sharks: first evidence of kleptoparasitism in
sharks. Ecology.

Landeau L, Terborgh J. 1986. Oddity and the 'confusion effect' in predation. Anim Behav.
34(5):1372–1380. doi:10.1016/S0003-3472(86)80208-1.

Lehmann L, Rousset F. 2010. How life history and demography promote or inhibit the
evolution of helping behaviours. Philos Trans R Soc Lond B Biol Sci. 365(1553):2599–2617.
doi:10.1098/rstb.2010.0138.

Li CC, Weeks DE, Chakravarti A. 1993. Similarity of DNA fingerprints due to chance and
relatedness. Hum Hered. 43(1):45–52.

Lynch M, Ritland K. 1999. Estimation of Pairwise Relatedness With Molecular Markers.
Genetics. 152:1753–1766.

Machado AMS, Cantor M, Costa APB, Righetti BPH, Bezamat C, Valle-Pereira JVS,
Simões-Lopes PC, Castilho PV, Daura-Jorge FG. 2019. Homophily around specialized
foraging underlies dolphin social preferences. Biol Lett. 15(4):20180909.
doi:10.1098/rsbl.2018.0909.

Mann J, Stanton MA, Patterson EM, Bienenstock EJ, Singh LO. 2012. Social networks reveal
cultural behaviour in tool-using dolphins. Nat Commun. 3:980. doi:10.1038/ncomms1983.

Milinski M. 1987. TIT FOR TAT in sticklebacks and the evolution of cooperation. Nature.
325(6103):433–435. doi:10.1038/325433a0.

Milligan BG. 2003. Maximum-Likelihood Estimation of Relatedness. Genetics.
163(3):1153–1167.

Mourier J, Brown C, Planes S. 2017. Learning and robustness to catch-and-release fishing in
a shark social network. Biol Lett. 13(3):20160824. doi:10.1098/rsbl.2016.0824.

Mourier J, Lédée EJI, Guttridge TL, Jacoby DMP. 2018. Network analysis and theory in
shark ecology - methods and applications. In: Shark Research: Emerging Technologies and
Application for the Study of Shark Biology. Carrier J, Heithaus M and Simpfendorfer C.
CRC Press. p. 337–356.

Mourier J, Maynard J, Parravicini V, Ballesta L, Clua E, Domeier ML, Planes S. 2016.
Extreme Inverted Trophic Pyramid of Reef Sharks Supported by Spawning Groupers. Curr
Biol. 26(15):2011–2016. doi:10.1016/j.cub.2016.05.058.

Mourier J, Mills SC, Planes S. 2013. Population structure, spatial distribution and life-history
traits of blacktip reef sharks Carcharhinus melanopterus. J Fish Biol. 82(3):979–993.
doi:10.1111/jfb.12039.

Mourier J, Planes S. 2013. Direct genetic evidence for reproductive philopatry and associated
fine-scale migrations in female blacktip reef sharks (Carcharhinus melanopterus) in French
Polynesia. Mol Ecol. 22(1):201–214. doi:10.1111/mec.12103.

Mourier J, Planes S, Buray N. 2013. Trophic interactions at the top of the coral reef food
chain. Coral Reefs. 32(1):285–285. doi:10.1007/s00338-012-0976-y.

Mourier J, Vercelloni J, Planes S. 2012. Evidence of social communities in a spatially
structured network of a free-ranging shark species. Anim Behav. 83(2):389–401.
doi:10.1016/j.anbehav.2011.11.008.

Newby J, Darden T, Bassos-Hull K, Shedlock AM. 2014. Kin structure and social
organization in the spotted eagle ray, Aetobatus narinari, off coastal Sarasota, FL. Environ
Biol Fishes. 97(9):1057–1065. doi:10.1007/s10641-014-0289-9.

Newman MEJ. 2006. Finding community structure in networks using the eigenvectors of
matrices. Phys Rev E. 74(3):036104. doi:10.1103/PhysRevE.74.036104.

Olsén KH, JäUrvi T. Effects of kinship on aggression and RNA content in juvenile Arctic
charr. J Fish Biol. 51(2):422–435. doi:10.1111/j.1095-8649.1997.tb01676.x.

Papastamatiou YP, Lowe CG, Caselle JE, Friedlander AM. 2009. Scale-dependent effects of
habitat on movements and path structure of reef sharks at a predator-dominated atoll.
Ecology. 90(4):996–1008. doi:10.1890/08-0491.1.

Perryman RJY, Venables SK, Tapilatu RF, Marshall AD, Brown C, Franks DW. 2019. Social
preferences and network structure in a population of reef manta rays. Behav Ecol Sociobiol.
73(8):114. doi:10.1007/s00265-019-2720-x.

Pew J, Muir PH, Wang J, Frasier TR. 2015. related: an R package for analysing pairwise
relatedness from codominant molecular markers. Mol Ecol Resour. 15(3):557–561.
doi:10.1111/1755-0998.12323.

Queller DC, Goodnight KF. 1989. Estimating Relatedness Using Genetic Markers. Evolution.
43(2):258–275. doi:10.1111/j.1558-5646.1989.tb04226.x.

R Core Team. 2015. R: a language and environment for statistical computing. Vienna,
Austria: R Computing Foundation for Science.

Ramos-Fernández G, Boyer D, Gómez VP. 2006. A complex social structure with fission–
fusion properties can emerge from a simple foraging model. Behav Ecol Sociobiol.
60(4):536–549. doi:10.1007/s00265-006-0197-x.

Reisinger RR, Beukes (née Janse van Rensburg) C, Hoelzel AR, de Bruyn PJN. 2017.
Kinship and association in a highly social apex predator population, killer whales at Marion
Island. Behav Ecol. 28(3):750–759. doi:10.1093/beheco/arx034.

Riehl C. 2011. Living with strangers: direct benefits favour non-kin cooperation in a
communally nesting bird. Proc R Soc Lond B Biol Sci. 278(1712):1728–1735.
doi:10.1098/rspb.2010.1752.

Ritland K. 1996. Estimators for pairwise relatedness and individual inbreeding coefficients.
Genet Res. 67(2):175–185. doi:10.1017/S0016672300033620.

Snijders L, Blumstein DT, Stanley CR, Franks DW. 2017. Animal Social Network Theory
Can Help Wildlife Conservation. Trends Ecol Evol. 32(8):567–577.
doi:10.1016/j.tree.2017.05.005.

Van Oosterhout Cock, Hutchinson William F., Wills Derek P. M., Shipley Peter. 2004.
micro-checker: software for identifying and correcting genotyping errors in microsatellite
data. Mol Ecol Notes. 4(3):535–538. doi:10.1111/j.1471-8286.2004.00684.x.

Vignaud T, Clua E, Mourier J, Maynard J, Planes S. 2013. Microsatellite Analyses of
Blacktip Reef Sharks (Carcharhinus melanopterus) in a Fragmented Environment Show
Structured Clusters. PLOS ONE. 8(4):e61067. doi:10.1371/journal.pone.0061067.

Vignaud TM, Mourier J, Maynard JA, Leblois R, Spaet JLY, Clua E, Neglia V, Planes S.
2014. Blacktip reef sharks, Carcharhinus melanopterus, have high genetic structure and
varying demographic histories in their Indo-Pacific range. Mol Ecol. 23(21):5193–5207.
doi:10.1111/mec.12936.

Wang J. 2002. An Estimator for Pairwise Relatedness Using Molecular Markers. Genetics.
160(3):1203–1215.

Wang J. 2007. Triadic IBD coefficients and applications to estimating pairwise relatedness.
Genet Res. 89(3):135–153. doi:10.1017/S0016672307008798.

Weideli OC, Mourier J, Planes S. 2015. A massive surgeonfish aggregation creates a unique
opportunity for reef sharks. Coral Reefs. 34(3):835–835. doi:10.1007/s00338-015-1290-2.

Whitehead H. 2008. Analyzing Animal Societies: Quantitative Methods for Vertebrate Social
Analysis. Chicago: University of Chicago Press.

Whitehead H, Dufault S. 1999. Techniques for Analyzing Vertebrate Social Structure Using
Identified Individuals: Review and Recommendations. In: Slater PJB, Rosenblat JS, Snowden
CT, Roper TJ, editors. Advances in the Study of Behavior. Vol. 28. Academic Press. p. 33–
74. [accessed 2020 Apr 20].
http://www.sciencedirect.com/science/article/pii/S0065345408602156.

Whitehead H, James R. 2015. Generalized affiliation indices extract affiliations from social
network data. Methods Ecol Evol. 6(7):836–844. doi:10.1111/2041-210X.12383.

Wilkinson GS, Carter GG, Bohn KM, Adams DM. 2016. Non-kin cooperation in bats. Phil
Trans R Soc B. 371(1687):20150095. doi:10.1098/rstb.2015.0095.

Wiszniewski J, Lusseau D, Möller LM. 2010. Female bisexual kinship ties maintain social
cohesion in a dolphin network. Anim Behav. 80(5):895–904.
doi:10.1016/j.anbehav.2010.08.013.

Wittemyer G, Okello JBA, Rasmussen HB, Arctander P, Nyakaana S, Douglas-Hamilton I,
Siegismund HR. 2009. Where sociality and relatedness diverge: the genetic basis for
hierarchical social organization in African elephants. Proc R Soc B Biol Sci.
276(1672):3513–3521. doi:10.1098/rspb.2009.0941.

Wolf JBW, Trillmich F. 2008. Kin in space: social viscosity in a spatially and genetically
substructured network. Proc R Soc Lond B Biol Sci. 275(1647):2063–2069.
doi:10.1098/rspb.2008.0356.

Tables

Table 1: Multiple Regression Quadratic Assignment Procedure and the influence of all structural variables on shark social associations. Matrices representing structural variables (predictors) were regressed against the association matrix (SRI) using a subset of the individuals in the population ($n = 43$) to which genetic relatedness was available. TO: temporal overlap; SO: spatial overlap; Gregariousness: typical number of associates based on (Godde et al. 2013); Size and Sex: binary matrices where individuals of the same size/sex classes are represented by 1, and different classes by 0. Adjusted R^2 indicates how much of the variation on association indices was explained by the predictors. Significant predictors in which *P*-values are given by the proportion of times the empirical regression coefficient was smaller or greater than the null expectancy from 20,000 randomisations (*P*–values are complementary, totalling 1) are indicated in bold. We considered all predictors were significant when $\beta \leq r$, thus $P > 0.95$ (**).

Period (SRI)	Predictors	Regression Coefficient (β)	*P ($\beta \geq r$)*	*P ($\beta \leq r$)*	Adjusted R^2
All	**TO**	0.5534	1.000**	<0.001*	94%
	SO	0.1069	1.000**	<0.001*	
	Gregariousness	-0.0018	<0.001	1.000	
	Size	0.0057	0.9101	0.0895	
	Sex	-0.0025	0.0003	0.9996	

753   **Table 2** Results of Mantel test for the correlation between the simple-ratio association (SRI)

754   and generalized affiliations (GAI) matrices and the pairwise genetic relatedness among

755   individual sharks for each period and each sex relationship. Observed estimates were

756   compared to those of 20,000 estimates from the randomized networks and significant p-

757   values are indicated in bold.

Index	SRI		GAI	
	Estimate (rho)	p-value	Estimate (rho)	p-value
Male-male	0.1028	**0.0267**	0.0404	0.4187
Female-female	-0.0653	0.5767	0.0361	0.7644
Male-female	0.0136	0.8291	-0.0240	0.7050

761   **Table 3** Effects of sex on shark gregariousness (degree and strength) in the social network.

		Coefficient	SE	t statictic	P_{rand}
Degree	*Intercept*	17.308	1.800	9.613	
	Sex (male)	4.726	2.156	2.192	**< 0.001**
Strength	*Intercept*	4.493	0.468	9.596	
	Sex (male)	0.369	0.560	0.660	0.548

770 **Table 4** Community level information on structure, association and affiliation indices, and

771 genetic relatedness. For each social index (SRI and GAI) and each community, are reported

772 number of community members (no. of individuals), mean index value, mean genetic

773 relatedness and p-values of the tests for within-between differences in relatedness.

Index	Communities	Modularity Q	No. of individuals	Mean index (SD)	Mean r (SE)	*p-values*
SRI	Overall	0.493	43	0.133 (0.210)	0.071 (0.005)	0.093
	Red		14	0.487 (0.178)	0.071 (0.009)	0.267
	Yellow		9	0.256 (0.307)	0.073 (0.016)	0.24
	Green		20	0.282 (0.194)	0.069 (0.006)	0.11
GAI	Overall	0.312	43	-0.028 (0.126)	0.060 (0.006)	0.722
	Red		1	/	/	/
	Yellow		12	0.078 (0.147)	0.056 (0.010)	0.707
	Green		6	-0.003 (0.155)	0.043 (0.015)	0.735
	Cyan		13	0.022 (0.127)	0.066 (0.009)	0.415
	Purple		11	0.053 (0.140)	0.061 (0.013)	0.408

   Figure legends:

Figure 1 Map of the study location indicating the monitored sites along the 10 km reef edge

of the North coast of Moorea.

Figure 2 Shark social preferences at the individual and population levels. Nodes (N = 43;

male:female = 30:13) representing photo-identified individuals are proportional to their size

and colour-coded by social modules; individuals are connected by links whose thicknesses

are proportional to SRI in the association networks (left panel), and to GAI removing

confounding factors in the affiliation networks where only positive GAIs are plotted (right

panel). In the density plots, red dots denote statistically significant observed values, shaded

distributions indicate null expectancy and blue whiskers indicate 95% confidence intervals.

The shark social network is characterized by (a) significant standard deviations (SD) of SRI

and GAI indicating social preferences and (b) significant modularity (Q) indicating social

division. In (c) are represented the density distribution of sightings of social community

members.

Figure 3 (a) Genetic network using the same layout as in Figure 2 where nodes (N = 43;

male:female = 30:13) are proportional to individual size and colour-coded by sex (males in

grey and females in white) and edges are proportional to genetic relatedness in the genetic

network (only close kin are plotted, i.e. r >0.25). (b) Differences in gregariousness (degree

and strength) of individual sharks between males and females (colours: pink indicates

females and blue indicates males). Comparison of the coefficient from the GLM based on the

observed data (red vertical line) and the frequency distribution of coefficients from the same

model based on the randomized data are indicated over each box plot to report significance of

the difference. (c) Relationship between association and affiliation indices and the triadic

likelihood estimator (TrioML) of genetic relatedness in male-male, female-female and male-

female pairs.

806   **Figure 4** Numbers of close kin (r > 0.25, the expected value of half-sibs) within and between

807   social communities for association (SRI) and affiliation (GAI) indices.