Inter-domain dynamics and fibrinolytic activity of matrix metalloprotease-1

Lokender Kumar[1,#], Joan Planas-Iglesias[2,3#], Chase Harms[1], Sumaer Kamboj[1], Derek Wright[1], Judith Klein-Seetharaman[2,4,*], and Susanta K. Sarkar [1, *]

[1]*Department of Physics, Colorado School of Mines, 1500 Illinois Street, CO 80401, U.S.A.*
[2]*Warwick Medical School, University of Warwick, Coventry CV4 7AL, U.K.*
[3]*Loschmidt Laboratories, Department of Experimental Biology, Faculty of Science, Masaryk University, Kamenice 5/A13, 625 00 Brno, Czech Republic.*
[4]*Department of Chemistry, Colorado School of Mines, 1500 Illinois Street, CO 80401, U.S.A.*

[#]*These authors contributed equally*

Corresponding author:
[*]ssarkar@mines.edu

Abstract

The roles of protein conformational dynamics and allostery in function are well-known. However, the roles that inter-domain dynamics have in function are not entirely understood. We used matrix metalloprotease-1 (MMP1) as a model system to study the relationship between inter-domain dynamics and activity because MMP1 has diverse substrates. Here we focus on fibrin, the primary component of a blood clot. Water-soluble fibrinogen, following cleavage by thrombin, self-polymerize to form water-insoluble fibrin. We studied the inter-domain dynamics of MMP1 on fibrin without crosslinks using single-molecule Forster Resonance Energy Transfer (smFRET). We observed that the distance between the catalytic and hemopexin domains of MMP1 increases or decreases as the MMP1 activity increases or decreases, respectively. We modulated the activity using 1) an active site mutant (E219Q) of MMP1, 2) MMP9, another member of the MMP family that increases the activity of MMP1, and 3) tetracycline, an inhibitor of MMP1. We fitted the histograms of smFRET values to a sum of two Gaussians and the autocorrelations to an exponential and power law. We modeled the dynamics as a two-state Poisson process and calculated the kinetic rates from the histograms and autocorrelations. Activity-dependent inter-domain dynamics may enable allosteric control of the MMP1 function.

Keywords: Inter-domain dynamics and activity, MMP1 conformational dynamics on fibrin, Single-molecule FRET.

Introduction

An understanding of the relationship between the structure and dynamics of a protein is crucial for understanding its function[1]. Researchers have shown the correlation between the function and intra-domain dynamics within a protein domain at different hierarchies of timescales[2-4]. However, the roles of inter-domain dynamics in function are not entirely understood[5,6]. Matrix metalloproteases (MMPs) are suitable for investigating the functional relationship between inter-domain dynamics and activity because the sequence of the catalytic domain remains conserved across the 23-member family[7]. The differences in activities among MMPs likely originate from the hemopexin domain with significant variations in the sequence[7]. The hemopexin domain seems to influence substrate/ligand specificity and activation/inhibition of various MMPs[8]. Also, prior research has reported the regulation of MMP1 catalytic activity by the hemopexin domain[9]. MMPs are capable of degrading numerous proteins[10], including collagen, the primary component of the extracellular matrix (ECM) that provides a scaffold for cells to maintain tissue integrity[11]. MMPs are calcium- and zinc-dependent enzymes that are produced by cells in pro forms, i.e., they need to be activated by cleaving off the pro domain for activity[11,12]. MMP1, a collagenase that degrades triple-helical type-1 collagen, stands out in the 23-member MMP family because it has crystal structures available[13]. Also, MMP1 itself is a broad-spectrum protease, an attribute that we previously used to degrade *E. coli* proteins and purify recombinant MMP1 using a protease-based method[14]. Extensive studies of MMP1 interacting with collagen monomers have revealed significant insights into the MMP1 activity, including the roles of conformational dynamics[9,15-22]. MMP1 consists of a catalytic domain that degrades substrates, a hemopexin domain that helps MMPs bind to the substrates, and a linker that mediates communications between the two domains (**Figure 1**).

Recently, we showed that inter-domain dynamics of MMP1 on type-1 collagen fibrils correlate with activity, and the two domains can allosterically communicate via the linker region as well as the substrate[23]. Since MMP1 is promiscuous, we investigated whether the insights into collagen fibril degradation applies to other substrates. In this paper, we focus on MMP1 inter-domain dynamics and activity on fibrin, the primary component of a blood clot. Fibrin is a critical physiological substrate because ~900,000 patients suffer from complications from clot formation leading to ~300,000 deaths[24] and ~$28 billion annual cost[25] in the US alone. The number of adults with clot-related health issues is estimated to reach ~1.82 million by 2050[26]. Blood clot formation through the coagulation cascade is a part of the natural response to injury and cut and prevents loss of blood from severed blood vessels[27]. The life-saving process of clot formation can become life-threatening if a clot blocks a blood vessel. Indeed, pathologies of

Figure 1. MMP1 structure (PDB ID: ISU3). The hemopexin domain is connected to the catalytic domain by a flexible linker. The pro domain is cleaved off to activate MMPs for catalysis. Yellow and purple spheres represent the van der Waals radii of the calcium and zinc atoms, respectively. The pro domain, the catalytic domain, the linker, and the hemopexin domains are roughly defined by the ranges of residues D32-Q99, F100-Y260, G261-C278, and D279-C466, respectively.

clot formation include heart attack, stroke, and pulmonary embolism, accounting for ~50% of all
hospital deaths.

Reperfusion, the act of restoring blood flow, with tissue plasminogen activator (tPA) remains the
standard treatment for ischemic stroke[28]. Plasminogen becomes plasmin after cleavage by tPA and
is able to degrade blood clots[29]. However, initiating tPA treatment after 3-4.5 h of stroke onset can
lead to adverse side effects, notably, hemorrhagic transformation[30]. Therefore, it is important to
explore alternative approaches that could either replace tPA treatment or extend the time window
for tPA perfusion beyond the current time frame[28]. For example, staphopain A and staphopain B,
cysteine proteases released by pathogenic Gram-positive bacteria *Staphylococcus aureus*, impair
clotting by degrading fibrinogen[31]. Also, the observation of clot degradation in plasminogen-
deleted animals suggests alternative fibrinolytic pathways[32]. Many proteases other than plasmin
have shown fibrinolytic activity. MMPs are potential alternatives in the fibrinolytic system because
MMPs are in the blood both in pro[33,34] and activated[35-38] forms. There are reports of fibrinolytic
activity of MMP3[39], MMP9[40], and MMP14[41], but the field is relatively unexplored[42]. Prior research
has reported that MMP1 does not have significant fibrinolytic activity[39], which is in contrast to our
observations presented in this paper.

Here we show that MMP1 inter-domain dynamics on fibrin are activity-dependent and modulated
by an enhancer or an inhibitor of MMP activity. We calculated the distance between S142 in the
catalytic domain and S366 in the hemopexin domain of MMP1 using smFRET measurements. The
area-normalized histograms of smFRET values represent the distributions of conformations,
whereas the normalized autocorrelations represent correlations between structures at different time
points. From the histograms, we defined two MMP1 structures with 1) inter-domain distance of
~4.5 nm and 2) inter-domain distance of ~5.4 nm as the closed and open states, respectively. A
comparison of active and active site mutant suggests that the open conformation of MMP1 is
functionally essential. The open conformations with well-separated domains appear more in the
presence of an activity enhancer (MMP9) and less in the presence of an inhibitor (tetracycline).
From autocorrelations, we learned that inter-domain dynamics are not entirely random and have
exponentially-distributed correlations. We fitted a sum of two Gaussians to the histograms and an
exponential to the autocorrelations. We modeled the dynamics as a two-state process, where the
locations of two states are the centers of Gaussian fits, and the sum of two kinetic rates between
the two states is the decay rate of exponential fits. Anisotropic Network Modeling (ANM) of
MMP1 dynamics revealed that a larger separation between the two domains (open conformation)
often accompanies a larger catalytic pocket opening between N171 and T230. A larger catalytic
pocket opening, in turn, enables closer proximity to the three chains of fibrin. ANM simulations
further revealed that the α-chain of fibrin is closest to the MMP1 active site, whereas the γ-chain
is furthest. SDS PAGE of ensemble fibrin degradation confirmed that MMP1 cleaves the α-chain
first, followed by the β- and γ- chains. The synergistic combination of smFRET, stochastic
simulations, ANM simulations, and ensemble assays provides an integrative approach to
investigate the functional roles of inter-domain dynamics. This approach applies to other
biochemical processes where water-soluble enzymes interact with water-insoluble substrates.

Results and discussion

Single-molecule measurement of MMP1 inter-domain dynamics on fibrin without crosslinks.
Water-soluble fibrinogen, upon limited cleavage by thrombin, self-polymerize into water-insoluble
fibrin without crosslinks. The addition of factor XIII and $CaCl_2$ creates fibrin with crosslinks. For
single molecule measurements, we used fibrin without crosslinks because we used a crystal
structure of fibrin without crosslinks for simulations. Besides, fibrin without crosslinks is optically
clearer than fibrin with crosslinks, which facilitates imaging with less background. To measure
MMP1 inter-domain dynamics at the single molecule level, we measured smFRET between two
dyes attached to the catalytic and hemopexin domains of MMP1. We mutated S142 in the catalytic
domain and S366 in the hemopexin domain of MMP1 to cysteines for labeling. The distances
between S142 and S366 are ~5.4 nm and ~4.5 nm for the open and closed conformations of MMP1,
respectively. We selected a pair of dyes, Alexa555 and Alexa647, because the Forster radius
between the two fluorophores is 5.1 nm. The FRET efficiency is 50% when the distance between
the fluorophores is equal to the Forster radius. More importantly, FRET is most sensitive to any
distance change around the Forster radius.

Figure 2. Single-molecule measurement of MMP1 dynamics on fibrin without crosslinks. (**A**)
Crystal structure of MMP1 (PDB ID 1SU3). Mutations of S142 and S366 to cysteines enables attaching
Alexa555 and Alexa647 dyes. (**B**) Scanning Electron Microscope (SEM) images of fibrin without
crosslinks. (**C**) Schematics of the TIRF microscope used for measuring MMP1 inter-domain dynamics
on fibrin. (**D**) Emission intensities of the two dyes. (top panel) Low FRET conformations lead to high
Alexa555 emission, whereas High FRET conformations lead to Low Alexa555 emission. Anticorrelated
Alexa647 and Alexa555 emissions, I_A and I_D, respectively; (bottom panel) Calculated smFRET trajectory
to show MMP1 inter-domain dynamics as a function of time.

Single molecule measurements suffer[43] from labeling[44], solution conditions[45], and properties of
fluorophores[46-49]. The stochastic nature of labeling CYS142 and CYS366 can also be problem,
which we discussed in our previous publication[23]. Since active MMP1 and active site mutant of
MMP1 would be affected equally by these complications, we can distinguish the effects due to
activity. Note that the amino acid E219[14] is the same as E200[13], which differs because we included
19 residues of the pro domain. In a previous publication, we showed that the specific activities of
labeled and unlabeled MMP1 were not affected by labeling[50]. A flexible linker connects the
catalytic and hemopexin domains of MMP1 (**Figure 2A**). **Figure 2B** shows fibrin without
crosslinks. We measured MMP1 inter-domain dynamics on a thin layer of water-insoluble fibrin
in a flow cell using a Total Internal Reflection Fluorescence (TIRF) microscope (**Figure 2C**).
Figure 2D shows one smFRET trajectory. For each condition, we collected more than 300,000
smFRET values for analysis of MMP1 inter-domain dynamics on fibrin (**Figure 3**).

Figure 3. Inter-domain dynamics of MMP1 on fibrin correlates with activity. More than 300,000
smFRET values at 22° C with a 100 ms time resolution for each condition are used to create area-
normalized histograms of MMP1 inter-domain distance (bin size=0.005). (**A**) Histograms without ligand,
(**B**) Histograms in the presence of MMP9 (an enhancer), and (**C**) Histograms in the presence of
tetracycline (an inhibitor) for active (blue) and active site mutant (orange) MMP1. Histograms are fitted
to a sum of two Gaussians (active: solid blue line; active site mutant: solid red line). Autocorrelations of
MMP1 inter-domain distance are calculated from the time series of smFRET values. (**D**)
Autocorrelations without ligand, (**E**) Autocorrelations in the presence of MMP9, and (**F**)
Autocorrelations in the presence of tetracycline for active MMP1 (blue) and active site mutant of MMP1
(orange). Autocorrelations are fitted to exponentials and power laws (exponential fit to active: dashed
black line; power law fit to active: dashed red line; exponential fit to active site mutant: solid black line;
power law fit to active site mutant: solid green line). The error bars in the histograms and autocorrelations
represent the square roots of the bin counts and the standard errors of the mean (sem) and are too small
to be seen. The supplementary information contains the fit equations and the best-fit parameters for
histograms and autocorrelations (**Table S1**). We approximated smFRET efficiency by $I_A/(I_A + I_D)$[51] and
has no unit.

MMP1 inter-domain dynamics on fibrin correlate with activity. Each value of smFRET
represents a conformation of MMP1. The histograms of smFRET values measuring inter-domain
distances (**Figure 3A**) for active MMP1 and active site mutant of MMP1 suggest that low FRET
conformations, where the two domains are well-separated, occur more for active MMP1 (blue
histogram) compared to active site mutant (orange histogram). The prevalence of low FRET
conformations for active MMP1 suggests that the two domains need to move away from each other
for performing catalytic activity on fibrin. Previously, we observed the same effects for MMP1
function on type-1 collagen[23]. Collagen degradation by MMP1 is enhanced by MMP9 and
inhibited by tetracycline. For collagen, we found that low FRET conformations appear more for
MMP9 and less for tetracycline consistent with the enhancement and inhibition of MMP1 activity.
To test whether or not MMP1 follows similar inter-domain dynamics when bound to fibrin, we
performed smFRET measurements in the presence of MMP9 and tetracycline. In the presence of
tetracycline, low FRET conformations of MMP1 on fibrin significantly disappear on fibrin (**Figure
3C**), as observed on type-1 collagen. Surprisingly, in the presence of MMP9, the MMP1 inter-
domain dynamics on fibrin (**Figure 3B**) show more high FRET states in contrast to the occurrence

FIGURE 4. MMP1 inter-domain dynamics as a two-state system. (**A**) Examples of simulated
smFRET trajectories with noise for active MMP1 (blue) and active site mutant of MMP1 (orange) using
experimentally-determined parameters for MMP1 without tetracycline. (**B**) Area-normalized histograms
of simulated smFRET values with best fits to a sum of two Gaussians (solid black line). (**C**)
Autocorrelations of simulated smFRET trajectories with best fits to exponentials (active: dashed black
line; active site mutant: solid black line). As expected, power law did not fit autocorrelations (active:
dashed red line; active site mutant: solid green line). k1+k2 was recovered from exponential fits with and
without noise. The error bars are the standard errors of mean for histograms and autocorrelations
and are too small to be seen.

of more low FRET states on collagen. To determine how a conformation at one time point
correlates with a conformation at another time point, we calculated the autocorrelations of

conformations (**Figure 3D-F**). Without any ligand, the correlation of dynamics on fibrin (**Figure**
3D) is higher for active MMP1 at shorter times, similar to previously reported dynamics on
collagen. However, the correlations in the presence of MMP9 and tetracycline show opposite
orders on fibrin (**Figure 3E** and **3F**) and collagen (previously reported)[23]. The correlated motions
indicate a decrease in conformational entropy. They can affect kinetics and thermodynamics of
biological processes, including catalysis[52]. The observed modulations of correlations under
different conditions suggest allosterically controlled inter-domain communications. The
quantitative comparisons of best-fit parameters (**Table S1**) strongly indicate that inter-domain
dynamics and activity of MMP1 are functionally related and allosteric. Further, a comparison with
inter-domain dynamics on fibrin (this paper) and collagen[23] suggest that MMP1 undergoes
substrate-dependent and allosterically-controlled dynamics.

A two-state Poisson process describes MMP1 inter-domain dynamics on fibrin. While the
histograms reveal distributions of conformations, the autocorrelations reveal whether or not
conformations at different time points are related. We found that a sum of two Gaussians fits the
histograms of smFRET values. Power law fits the autocorrelations for two-state simulations
without noise at millisecond timescales and molecular dynamics simulations at picosecond
timescales for collagen[23]. As such, we tried to fit both power law and exponential to the
autocorrelations for fibrin. The fit equations and best-fit parameters are in the supplementary
information. The exponential fits to autocorrelations enable a more straightforward interpretation
of the decay rates of correlations if we approximate the conformations of MMP1 as a two-state
Poisson process[23]. As a result, we can establish a quantitative connection between the histograms
and autocorrelations to calculate the kinetic rates between the two states. To test this, we
considered the two centers of Gaussian fits to histograms for MMP1 without ligands (**Figure 3A**)
as the two states, S1 (low FRET) and S2 (high FRET). We defined the two kinetic rates as k1
(S1→S2) and k2 (S2→S1) for interconversion between S1 and S2. For a two-state system, we
calculated k1/k2 from the ratio of Gaussian area(S2)/area(S1) and k1+k2 from the decay rates of
autocorrelations. We solved the two equations to calculate k1 and k2 (**Table S1C**) for both active
MMP1 and active site mutant of MMP1. With experimentally-determined S1, S2, k1, k2, and noise
(widths of the histograms), we simulated smFRET trajectories for active MMP1 and active site
mutant of MMP1 and analyzed the same way as we did experimental smFRET trajectories (**Figure**
4). For comparison, experimentally-determined inputs and recovered parameters from two-state
simulations are in **Table S2**. In summary, we show that a two-state system describes MMP1 inter-
domain dynamics.

For active MMP1 without ligands, the two states are S1=0.42 and S2=0.51 on fibrin (**Table S1A**),
which are similar to the two states S1=0.44 and S2=0.55 on collagen. However, the correlation
decay rate of 0.08 s^{-1} on fibrin (**Table S1C**) is lower than the rate of 0.13 s^{-1} on collagen. In the
presence of MMP9 and tetracycline, the two states and kinetic rates change even though a two-
state description still applies for active MMP1 and active site mutant of MMP1 on fibrin (**Figure**
3 and **Table S1**) and collagen. In other words, MMP1 undergoes substrate-dependent inter-domain
dynamics that follow a two-state Poisson process.

The two-state description also reveals the importance of noise in autocorrelations. Without noise,
both power law and exponential fit the autocorrelations of simulated smFRET trajectories (**Figure**
S1). With noise, however, only exponential fits the autocorrelations. Just as noise can convert a
Lorentzian line shape into Gaussian line shape[53], noise seems to turn a power law correlation into

an exponential one. Note that the exponential fits recover the underlying sum of the simulated
kinetic rates with and without noise.

Figure 5. Correlations of MMP1 inter-domain distance with catalytic pocket opening when MMP1 is not bound to a substrate. Examples of (**A**) open and (**B**) closed conformations of MMP1. The inter-domain distances between S142 and S366 and corresponding catalytic pocketing openings between N171 and T230 have been noted. Spheres and cages represent the van der Waals radii. Yellow cage: E219; Wheat sphere: calcium; Mauve sphere: zinc. (**C**) and (**D**) are scatter plots of inter-domain distance and catalytic pocket opening for the open and closed conformations, respectively. Two distances are calculated using Anisotropic Network Modeling (ANM) simulations (see methods).

Larger inter-domain distance can accompany larger catalytic pocket opening. The catalytic
cleft of MMP1 (~0.5 nm) is narrow[17], and as such, cleavage of substrates such as fibrin monomer
(diameter ~2-5 nm)[54] and collagen monomer (diameter ~1.5 nm)[55] needs more opening of the
catalytic pocket for easier access to the catalytic cleft. To test this, we used ANM simulations[56] to
calculate normal modes of MMP1 dynamics. ANM models proteins as a system of beads (amino
acids) connected by springs (bonds) and calculates the preferred/normal modes of protein motion
and corresponding eigenfrequencies of movement. Single molecule experiments (**Figure 3**)
suggest that both active MMP1 and active site mutant of MMP1 has an open conformation (lower
FRET) and a closed conformation (higher FRET). Therefore, we chose two MMP1 conformations
(**Figure 5A** and **5B**) for ANM simulations (see Methods for details). To select the open
conformation, we performed coarse-grained simulations of free MMP1 using ANM and decided
on the MMP1 conformation with the highest inter-domain distance (**Figure 5A**). For closed
conformation, we selected the conformation in the crystal structure (PDB ID: 1SU3) (**Figure 5B**).
We considered 20 modes (20 frames for each mode) with the lowest frequencies (slower motion)

to calculate the inter-domain distance (S142-S366) and catalytic pocket opening (N171-T230). A
comparison of **Figure 5C** and **5D** shows that MMP1 in the open conformation has an overall
catalytic pocket opening (~2.7 nm), whereas the closed structure has an opening (~2.6 nm). Since
a larger catalytic pocket opening enables access to the catalytic cleft, we can infer that larger inter-
domain distances (lower FRET) correlate with MMP1 activity.

Figure 6. Correlations of MMP1 inter-domain distance with catalytic pocket opening when MMP1 is bound to the reconstructed fibrin model combining 3GHG and 1FZC. Examples of (**A**) open and (**B**) closed conformations of MMP1 bound to the reconstructed model of fibrin. Three dimensional scatter plots (blue circle) of inter-domain distance (S142-S366), catalytic pocket opening (N171-T230), and rms proximity between the MMP1 catalytic site and the three fibrin chains for (**C**) open and (**D**) closed MMP1 conformations. Two-dimensional projections of the scatter plots are in gray. The open structure shows larger catalytic pocket openings.

To investigate whether or not a larger inter-domain distance correlates with a larger catalytic
pocket opening when MMP1 is bound to fibrin, we needed to determine the docking poses of
MMP1 with fibrin because there is no crystal structure of fibrin-bound MMP1. To this end, open
and closed conformations of MMP1 (**Figure 5A** and **5B**) were docked to a reconstructed model of
fibrin using ClusPro[57], a protein-protein docking server. The reconstructed model (see methods for
detailed procedure) is a combination of the crystal structures of fibrinogen (PDB ID: 3GHG) and
fibrin (PDB ID: 1FZC). The docking poses were sorted based on the distance from the α-carbon
of E219 at the MMP1 catalytic site to the potential cleaving sites on fibrin. Since the cleavage sites
on fibrin for MMP1 are unknown, we considered the cleavage sites for MMP3, MMP7, and
MMP14 as substitutes: the α-chain at Asp97-Phe98, and Asn102-Asn103; the β-chain at Asp123-
Leu124, Ans137-Val138, and Glu141-Tyr142; and the γ-chain at Thr83-Leu84[58] (**Figure S2**). For
each docking pose, we calculated the minimum distance possible for the three chains and
considered the average of three lengths as the metric for the catalytic domain-fibrin proximity. In
addition, we assumed that the hemopexin domain remains bound to the substrate, and thus, we
considered the distance from the center of mass (CoM) of the hemopexin domain to any atom of
the substrate as the metric for the hemopexin domain-fibrin proximity. We sorted the docking
poses according to the two measures of proximity to the MMP1 domains. We selected the docking
pose with minimum scores for the two criteria for ANM simulations (**Figure 6A** and **6B**). We
considered 20 modes (20 frames for each mode) with the lowest frequencies (slower motion) to
calculate inter-domain distance (S142-S366), catalytic pocket opening (N171-T230), and rms
proximity between the MMP1 catalytic site and the three fibrin chains (**Figure 6C** and **6D**). The
inter-domain distance reduces after binding fibrin for both the open and closed conformations of
MMP1. As is the case of free MMP1 (**Figure 5C** and **5D**), fibrin-bound MMP1 also showed that
larger inter-domain distance for the open conformation accompany larger catalytic pocket
openings (**Figure 6C** and **6D**). In contrast, fibrinogen-bound MMP1 shows larger catalytic pocket
opening for the closed conformation (**Figure S3**). The importance of open conformations on fibrin
and closed conformations on fibrinogen is an experimentally testable insight from ANM
simulations. At the single molecule level, we could test with fibrinogen attached to a quartz slide
for TIRF imaging. Note that the overall charge is negative for both fibrinogen and a clean quartz
slide, making it challenging to attach fibrinogen on a quartz slide. Nevertheless, histograms of
smFRET values would be peaked at a higher FRET value on fibrinogen as compared to fibrin. One
could also try to crystalize MMP1 with fibrinogen and fibrin, and check the conformations.
Nevertheless, both fibrinogen- and fibrin-bound MMP1 show closer proximity to the three chains
for the open conformation. The proximity suggests that a larger inter-domain distance is relevant
for MMP1 activity.

The proximity of MMP1 to the three chains correlates with the sequence of degradation.
Since MMP1 needs to come in the proximity before cleavage, we performed molecular docking
of MMP1 with fibrin. **Figure 7A** shows MMP1 molecular docking poses superimposed on the
reconstructed model of fibrin. MMP1 binds to fibrin at specific places in both open and closed
conformations. To gain further insights, we calculated the distance between every atom in MMP1
and the three fibrin chains. We counted the number of distances below 0.5 nm for the top 30
docking poses and calculated the mean and standard deviation (**Figure 7B**). On average, MMP1
in open conformation showed the closest proximity to the α-chain (**Figure 7B**, left panel). We
quantified the statistical significance by p-value (**Figure 7B**, left panel). In contrast, the closed
conformation of MMP1 did not show any significant difference in proximity to the three chains
(**Figure 7B**, right panel).

To test whether or not the sequence of MMP1 proximity to the three chains has any implication in
fibrinolytic activity, we performed the time-dependent degradation of fibrinogen. **Figure S4B**
shows that the overall catalytic activity is fastest on the α-chain and slowest on the γ-chain.
Fibrinogen, the basic building block of fibrin, has three pairs of amino acid chains: 1) the α-chain
at ~63 kDa, 2) the β-chain at ~56 kDa, and 3) the γ-chain at ~47 kDa. SDS PAGE of fibrinogen
control approximately confirms molecular weights of the three chains (**Figure S4A**, lane 2 from
left). These three chains are connected by a dimeric disulfide knot (DSK) at the N-termini and
many other disulfide bonds along the length of fibrinogen[59,60]. Treatment of 1 mg/mL of fibrinogen
with 0.1 mg/mL of active MMP1 at 37° C for 24 h resulted in cleavage of all three chains and a
final prominent fragment at ~40 kDa (**Figure S4A**, lane 3 from left). MMP1 completely degraded
the α-chain and partially degraded the β-chain within 2 h; however, the γ-chain took nearly 6 h to
be degraded (**Figure S4B**). After 8 h, a prominent band at ~40 kDa remained. In other words, the
computationally determined sequence of degradation in the open conformation agrees with the
experimentally observed sequence (**Figure S4B**). The agreement between the computational and
experimental sequence of degradation further confirms that the open conformations observed in
smFRET experiments are functionally relevant.

Figure 7. Proximity of MMP1 to the chains of reconstructed fibrin model. (**A**) The clustered 30 docking poses obtained from ClusPro using fibrin as the ligand and MMP1 as the receptor. The catalytic domain of MMP1 is red, whereas the hemopexin domain is gray. (**B**) We measured the distance between all possible pairs of atoms between the fibrin chains and MMP1 for 30 docking poses obtained from ClusPro. We counted any distance less than 5 Å and plotted the distributions of total count (y-axis) for each chain for the open and closed conformations. p-value<0.05:*; p-value<0.01:**; p-value<0.001:***.

We also imaged fibrin without crosslinks after treatment with MMP1 using SEM (**Figure S4C**),
which shows increased porosity of fibrin for active MMP1 consistent with the degradation. We
used fibrin without crosslinks because it is easier to form a thin layer of fibrin on a quartz slide for
smFRET measurements, and the crystal structure without crosslinks (PDB ID: 1FZC) is available.
However, fibrin with crosslinks is more physiologically relevant (see methods for making fibrin
with crosslinks). As such, we quantified the degradation rate of fibrin with crosslinks using a
weight-based assay (Figure S5). We also performed gel electrophoresis of degradation product
(**Figure S6A**), imaged the reaction volume to show the degradation within 5 h (**Figure S6B**), and
imaged the surface morphology using SEM (**Figure S6C**).

Figure 8. Effects of the linker and reconstructed fibrin on the catalytic pocket opening. The catalytic pocket openings on fibrin represented by the blue line connecting N171 and T230 for (**A**) the full-length MMP1 (open conformation: 2.68±0.01 nm; closed conformation: 2.61±0.01 nm); (**B**) the two MMP1 domains without the linker (open conformation: 2.68±0.01 nm; closed conformation: 2.61±0.01 nm); and (**C**) the catalytic domain alone (open conformation: 2.68±0.01 nm; closed conformation: 2.68±0.01 nm). The error bars represent the standard deviation of 60 measurements of the catalytic pocket opening obtained from 20 frames each for the three slowest normal modes.

Unlike collagen, MMP1 domains do not communicate via fibrin. Previously, we performed
ANM simulations of MMP1 interacting with collagen using the crystal structure of MMP1 bound
to a collagen model (4AUO)[23]. We reported that the catalytic and hemopexin domains could
communicate via collagen even if the physical linker is removed[23], which explained the
experimental observation that a mixture of the two domains purified separately could degrade

triple-helical collagen[18]. We wanted to investigate whether or not the two domains communicate
via fibrin. We created three forms of MMP1 using the crystal structure (PDB ID: 1SU3): 1) full-
length MMP1, 2) MMP1 with the linker domain removed, and 3) MMP1 with the linker and
hemopexin domain removed. We docked both the open and closed conformations with the
reconstructed fibrin using ClusPro and chose the binding pose shown in **Figure 8**. Three forms of
MMP1 interacting with fibrin resulted in similar mean and standard deviation of the catalytic
pocket opening (**Figure 8**). In contrast, the catalytic pocket opening on collagen changed even
without the physical linker[23]. In other words, fibrin interacts with MMP1 passively, whereas
collagen interacts with MMP1 actively. Using single molecule tracking of labeled MMP1 on type-
1 collagen fibrils, we also showed that fibrils play a role in MMP1 activity due to vulnerable sites
on fibrils created during fibril assembly[61].

Both fibrin and collagen are extracellular and triple-helical but have microstructural and
mechanical differences[62]. Fibrin monomers are 45 nm long and have three right-handed chains
forming a left-handed structure in between two globular regions[54]. In contrast, collagen monomers
are 300 nm long and have three left-handed chains forming a right-handed structure[63]. The
difference in handedness between fibrin and collagen may lead to in-phase and out-of-phase
motion of MMP1 domains. To understand why fibrin and collagen interact differently, however,
we need further studies since fibrin and collagen have different sequences and structures despite
the triple-helical similarities.

In summary, we have shown that the inter-domain dynamics of MMP1 on fibrin correlates with
its fibrinolytic activity. We measured smFRET between two fluorophores attached to the catalytic
and hemopexin domains of MMP1. A comparison of distributions of smFRET values between
active MMP1 and active site mutant of MMP1 suggests that MMP1 needs to have the two domains
well-separated for function. To investigate the roles of open conformations, we measured smFRET
in the presence of MMP9 and tetracycline. MMP9 is an enhancer, whereas tetracycline is an
inhibitor of MMP1 activity on collagen and a synthetic substrate. In the presence of tetracycline,
the inter-domain distance on fibrin becomes smaller consistent with the trend on collagen. MMP9
causes the inter-domain distance of MMP1 to be shorter on fibrin, which is opposite to the
observation on collagen. One possible reason is that MMP9 by itself can degrade fibrin but cannot
degrade triple-helical collagen. As a result, fibrin degradation by MMP1 in the presence of MMP9
is more complicated. A two-state Poisson process quantitatively describes the inter-domain
dynamics. We fitted the histograms of smFRET values to a sum of two Gaussians. The best-fit
parameters for the centers are the two states used for simulations. The ratio of kinetic rates equals
to the ratio of areas of the two Gaussians fitted to a histogram. Additionally, the sum of kinetic
rates equals to the decay rate of autocorrelations of smFRET values. We simulated smFRET
trajectories with experimentally determined parameters and noise levels. We recovered the
underlying parameters to validate our analyses. The presence of noise in the signal appears to
convert a power law autocorrelations into an exponential one.

We performed molecular docking of the open and closed conformations of MMP1 with fibrin. In
the closed conformation, MMP1 approaches the three chains equally. In the open conformation,
however, MMP1 approaches the α-chain first and the γ-chain last, which we confirmed using SDS
PAGE of fibrinogen degraded by MMP1. As such, the open conformation appears to be more
functionally relevant. One or both domains of MMP1 binds to fibrin at many places along the
length of fibrin. However, the primary location on fibrin is near the globular regions where both
the domains bind. We, therefore, selected the docking pose near the globular part for ANM

simulations. ANM simulations of MMP1-fibrin interactions revealed that the open conformation
with well-separated domains has a larger catalytic pocket opening. A larger opening enables the
fibrin chains to approach closer to the MMP1 active site. In the closed conformation, the fibrin
chains get closer to the catalytic site only when the catalytic domain is alone. We performed limited
digestion of fibrin by MMP1 with and without crosslinks and observed increased porosity due to
degradation. Also, MMP1 degrades and dissolves crosslinked fibrin within 5 hr, suggesting a role
of MMP1 in the fibrinolytic pathway. We quantified the degradation using a weight-based assay
because interactions of water-soluble MMP1 with water-insoluble fibrin are challenging to
quantify using solution-based biochemical assays.

We discussed the results of the MMP1-fibrin system in light of our previous studies on the MMP1-
collagen system. For both fibrin and collagen, the open conformation of MMP1 is functionally
relevant and responds to tetracycline similarly. The open conformation enables larger catalytic
pocket opening and closer proximity to the three strands of fibrin and collagen. A two-state Poisson
process describes MMP1 dynamics on both fibrin and collagen. However, MMP1 dynamics are
opposite to each other on fibrin and collagen in the presence of MMP9. Note that MMP9 can
degrade fibrin but cannot degrade triple-helical collagen and, as such, can influence MMP1
dynamics differently. Collagen plays an active role in mediating allosteric communications
between the two domains and opening the catalytic pocket. In contrast, fibrin plays a passive role
and cannot mediate communications between the two domains. Further studies are needed to
investigate the detailed mechanisms of different MMP1 interactions on fibrin and collagen, which
may enable allosteric control of MMP1 activity in a substrate-dependent manner for exploring
MMPs as drug targets.

375
Methods

Purification and labeling of MMP1. Full-length recombinant MMPs were expressed in *E. coli*
and activated using 0.1 mg/mL trypsin. Activated MMP1 and trypsin together create a chain
reaction of further MMP1 activation and broad-spectrum proteolytic cleavage of *E. coli* proteins.
Molecular weights of activated MMP1 and trypsin are ~43 kDa and ~23 kDa, respectively. We
used a 30 kDa cut-off Amicon filter to centrifuge and purify MMP1 in the activated form. For
further details, see the previous publication on protease-based purification method[14]. We labeled
purified MMP1 with AlexaFluor555 (ThermoFisher Scientific, Cat# A20346, donor dye) and
AlexaFluor647 (ThermoFisher Scientific, Cat# A20347, acceptor dye) in a ratio of ~1:1 using
maleimide chemistry. 1 mL of MMP1 at 1 mg/mL concentration was incubated with 20 µL of 1
mg/mL AlexaFluor555 and AlexaFluor647 for 1 h in a 5 mL glass vial in a nitrogen environment
to avoid oxidation of the dyes. After incubation, we used a 30 kDa cut-off Amicon filter to remove
free dyes from the solution. We compared the specific activities of labeled and unlabeled MMP1
on the synthetic substrate, MCA-Lys-Pro-Leu-Gly-Leu-DPA-Ala-Arg-NH2, (R&D Systems, Cat#
ES010) as described before[14]. Labeling MMP1 does not affect its specific activity[23].

Single-molecule measurements. We prepared 200 µL of reaction volume in 10 mM Phosphate
Buffer Saline (PBS) (Sigma, Cat# P3813, pH 7.4) by mixing 10 units of thrombin (Cayman
chemical, Cat# 13188) and 50 µg of fibrinogen (Cayman chemical, Cat# 16088). A thin layer of
fibrin was created on a quartz slide similar to the blood smear protocol used in diagnostics[64] and
incubated at 37º C for 18 h without shaking. We made a flow cell of thickness 120 µm using a
piece of double-sided adhesive tape, a clean quartz slide, and a glass coverslip. The quartz slides
had two holes to connect the input and output tubes for exchanging buffers and solutions. We

created an evanescent wave at the interface of the quartz slide and sample in a Total Internal
Reflection Fluorescence (TIRF) Microscope as described before[65-68]. We incubated 50 µL of 0.1
mg/mL labeled MMP1 with 50 µL of protein buffer (50 mM Tris, 100 mM NaCl, pH 8.0) and 100
µg/mL tetracycline for 30 min at 22° C. The labeled MMP1 was serially diluted to prepare a
working concentration of ~100 pM and flowed into the flow cell. Alexa555 dyes (donor dyes)
attached to MMP1 was excited using a 532 nm laser and imaged using the TIRF microscope at
22° C at 100 ms time resolution. We imaged Alexa647 and Alexa555 emissions using an EMCCD
camera (Andor iXon). The two emission channels were superimposed using a pairwise stitching
plugin of ImageJ, and the intensities from the two dyes were extracted and analyzed using Matlab.

Reconstruction of the extended fibrin molecule. We superimposed the crystal structure of
fibrinogen (PDB ID: 3GHG)[69] with the crystal structure of D-D fibrin fragment (PDB ID: 1FZC)[70]
using the Combinatorial Extension (CE) algorithm[71] implemented in PyMOL[72] to obtain the
reconstructed model of fibrin. The fibrinogen asymmetric unit presents two replicas (either the
chains A-F+M-P or G-L+Q-T). Each replica has two ends, which are represented by chains A-C
and chains D-F in one and by chains G-I and J-L in the other. Each of such fibrinogen ends could
be superimposed to any of the ends of the fibrin crystal (either the chains A-C or chains D-F),
yielding a total of 16 different possibilities to reconstruct the fibrin molecule. For each of the
superimposed assembly, we superimposed a new fibrin crystal to the fibrinogen biological
assembly not used for the initial superimposition. Thus, we superimposed two extra fibrin crystals
(Fa and Fb, one per superimposed fibrinogen biological assembly) on each of the 16
different reconstruction attempts. A reconstruction compatible with the original fibrinogen
packing would show no differences in the positioning of Fa and Fb. Thus, to select the best
superimposition combination to reconstruct the fibrin molecule, the one that minimized the
RMSD in between Fa and Fb was selected.

Selection of closed and open conformations of MMP1. The activation peptide (residues 32 to
98) was removed from the first biological assembly (chain A) of the crystal structure of MMP1
(PDB ID: 1SU3), leaving only the catalytic (residue numbers 107 to 260), linker (261-277) and
hemopexin (278-466) domains of the enzyme, which we selected as the closed conformation. To
obtain the open conformation, we submitted the coordinates of the closed conformation to the
ANM web server 2.1[56] using default parameters. We extracted the coordinates of the three-
dimensional fluctuations for 20 frames of the 20 slowest modes. For each frame on each model,
we calculated distances using PyMOL. We used the distance between S142 and S366 to analyze
the fluctuations between the catalytic and hemopexin domains. We selected the frame that
maximized this distance (second mode, second frame) as the MMP1 open conformation.

Molecular docking of MMP1 with fibrin. The closed and open conformations of MMP1 were
docked individually with the reconstructed fibrin molecule using the default parameters of ClusPro
2.0 web server[57]. We used the reconstructed fibrin as the receptor and the MMP1 conformations
as the ligands. We considered the centroid structure of each of the resulting clusters (maximum
30) as the representative docking solution for a cluster, and we obtained the corresponding
weighted scores for the best coefficient for each binding pose. We adjusted different poses to the
same Cartesian space by aligning the reconstructed fibrin from each pose to a template one using
PyMOL. To analyze the accessibility of MMP1 to the three chains, we counted the number of
times an atom in MMP1 was within 5 Angstrom from an atom in the chains.

Preparation of fibrin without and with crosslinks. We prepared fibrin without crosslinks by
adding 10 units of thrombin (Cayman chemical, Cat# 13188) to 50 µg of fibrinogen (Cayman

chemical, Cat# 16088) in 10 mM PBS to a total reaction volume of 200 μL and incubating at 37º
C for 18 h with 250 rpm shaking inside an incubator (Thermo Fisher Scientific, MaxQ600).
Preparation of crosslinked fibrin involved two additional reagents: factor XIII (Abcam, Cat #
ab62427) and $CaCl_2$. Thrombin activates factor XIII to factor XIIIa, which in the presence of Ca^{++}
induces crosslinks in fibrin. We added 10 units of thrombin to 2.5 μg of factor XIII in a 0.5 mL
PCR tube, diluted with 10 mM PBS to a total reaction volume of 100 μL, and incubated at 37º C
for 10 min. After incubation, we added 50 μg of fibrinogen and 10 μL of 5 mM $CaCl_2$, diluted
with PBS buffer to a total reaction volume of 200 μL, and incubated at 37º C for 15 min to obtain
gel-like crosslinked fibrin. Note that fibrin without crosslinks requires a longer incubation time
and shaking, but fibrin with crosslinks forms within 15 min without shaking. For every reaction,
we put a part of the sample on a sample mount before drying and imaging the surface morphology
of fibrin using a Phenom Pro Scanning Electron microscope.

Sodium Dodecyl Sulphate-Polyacrylamide Gel Electrophoresis (SDS PAGE). The degradation
profile after fibrinolysis was analyzed using SDS-PAGE. We mixed SDS PAGE loading buffer
containing BME (Biorad, Cat# 610737) with each sample in a 1:1 (v/v) ratio and incubated at 95º
C for 10 min. We loaded the samples onto the wells of 8% Tris.glycine gels. We ran electrophoresis
in Tris.glycine buffer (Biorad, Cat# 1610732) for 30 min at 50 V, followed by 1 h at 120V. We
used the precision plus protein kaleidoscope pre-stained protein standards (Biorad, Cat# 1610375)
as molecular weight markers. The SDS-PAGE gel was stained with Coomassie Brilliant Blue R-
250 (Biorad, Cat# 161-0436) and imaged using a digital camera after destaining.
463

464 **Acknowledgments**

465 One grant to S.K.S. and J.K.S. from the National Institutes of Health (RGM137295A) partially
466 supported this work. This project was also partially funded by the professional development funds
467 given to S.K.S and J.K.S. at the Colorado School of Mines. JPI was supported by the
468 grant MSCAfellow@MUNI.
469

470 **Author contributions**

471 S.K.S. conceived and designed the overall project. L.K. and S.K.S. designed experiments, L.K.
472 performed experiments, J.P. performed docking and ANM analysis. L.K., J.P., J.K., C.H., J.K.,
473 D.W., and S.K.S. analyzed data. S.K.S. wrote the manuscript. All authors edited the manuscript.
474
475 **Competing financial interests**

476 The authors declare no competing financial interests.
477
478 **Data availability**

479 The datasets generated during and/or analyzed during the current study are available from the
480 corresponding author on reasonable request.

References

1.   Karplus, M. & Kuriyan, J. Molecular dynamics and protein function. *Proceedings of the National Academy of Sciences* **102**, 6679-6685 (2005).

2.   Henzler-Wildman, K. & Kern, D. Dynamic personalities of proteins. *Nature* **450**, 964-972 (2007).

3.   Henzler-Wildman, K. A. *et al.* A hierarchy of timescales in protein dynamics is linked to enzyme catalysis. *Nature* **450**, 913-916 (2007).

4.   Lewandowski, J. R., Halse, M. E., Blackledge, M. & Emsley, L. Direct observation of hierarchical protein dynamics. *Science* **348**, 578-581 (2015).

5.   Bhaskara, R. M., de Brevern, A. G. & Srinivasan, N. Understanding the role of domain–domain linkers in the spatial orientation of domains in multi-domain proteins. *Journal of Biomolecular Structure and Dynamics* **31**, 1467-1480 (2013).

6.   Esteban-Martín, S. *et al.* Correlated inter-domain motions in adenylate kinase. *PLoS computational biology* **10** (2014).

7.   Ratnikov, B. I. *et al.* Basis for substrate recognition and distinction by matrix metalloproteinases. *Proceedings of the National Academy of Sciences* **111**, E4148-E4155 (2014).

8.   Piccard, H., Van den Steen, P. E. & Opdenakker, G. Hemopexin domains as multifunctional liganding modules in matrix metalloproteinases and other proteins. *Journal of leukocyte biology* **81**, 870-892 (2007).

9.   Singh, W., Fields, G. B., Christov, C. Z. & Karabencheva-Christova, T. G. Importance of the linker region in matrix metalloproteinase-1 domain interactions. *RSC advances* **6**, 23223-23232 (2016).

10.   Rodríguez, D., Morrison, C. J. & Overall, C. M. Matrix metalloproteinases: what do they not do? New substrates and biological roles identified by murine models and proteomics. *Biochimica et Biophysica Acta (BBA)-Molecular Cell Research* **1803**, 39-54 (2010).

11.   Page-McCaw, A., Ewald, A. J. & Werb, Z. Matrix metalloproteinases and the regulation of tissue remodelling. *Nature reviews. Molecular cell biology* **8**, 221-233, doi:10.1038/nrm2125 (2007).

12.   Jackson, B. C., Nebert, D. W. & Vasiliou, V. Update of human and mouse matrix metalloproteinase families. *Human genomics* **4**, 194 (2010).

13.   Manka, S. W. *et al.* Structural insights into triple-helical collagen cleavage by matrix metalloproteinase 1. *Proceedings of the National Academy of Sciences* **109**, 12461-12466 (2012).

14.   Kumar, L., Colomb, W., Czerski, J., Cox, C. R. & Sarkar, S. K. Efficient protease based purification of recombinant matrix metalloprotease-1 in E. coli. *Protein expression and purification* **148**, 59-67 (2018).

15.   Karabencheva-Christova, T. G., Christov, C. Z. & Fields, G. B. Conformational Dynamics of Matrix Metalloproteinase-1· Triple-Helical Peptide Complexes. *The Journal of Physical Chemistry B*, 5316-5326 (2017).

16.   Fields, G. B. Interstitial collagen catabolism. *J Biol Chem* **288**, 8785-8793, doi:10.1074/jbc.R113.451211 (2013).

17.   Bertini, I. *et al.* Structural basis for matrix metalloproteinase 1-catalyzed collagenolysis. *Journal of the American Chemical Society* **134**, 2100-2110 (2012).

Chung, L. D. *et al.* Collagenase unwinds triple-helical collagen prior to peptide bond hydrolysis. *Embo Journal* **23**, 3020-3030 (2004).

Lauer-Fields, J. L., Juska, D. & Fields, G. B. Matrix metalloproteinases and collagen catabolism. *Biopolymers* **66**, 19-32 (2002).

Salsas-Escat, R., Nerenberg, P. S. & Stultz, C. M. Cleavage site specificity and conformational selection in type I collagen degradation. *Biochemistry* **49**, 4147-4158 (2010).

Nerenberg, P. S. & Stultz, C. M. Differential unfolding of alpha 1 and alpha 2 chains in type I collagen and collagenolysis. *Journal of Molecular Biology* **382**, 246-256 (2008).

Nerenberg, P. S., Salsas-Escat, R. & Stultz, C. M. Do collagenases unwind triple-helical collagen before peptide bond hydrolysis? Reinterpreting experimental observations with mathematical models. *Proteins: Structure, Function, and Bioinformatics* **70**, 1154-1161 (2008).

Kumar, L. *et al.* Allosteric communications between domains modulate the activity of matrix metalloprotease-1. *bioRxiv*, 804559, doi:10.1101/804559 (2020).

Heit, J. A., Cohen, A. T. & Anderson, F. A. Estimated annual number of incident and recurrent, non-fatal and fatal venous thromboembolism (VTE) events in the US. (2005).

Mahan, C. E. *et al.* Venous thromboembolism: annualised United States models for total, hospital-acquired and preventable costs utilising long-term attack rates. *Thrombosis and haemostasis* **107**, 291-302 (2012).

Deitelzweig, S., Johnson, B., Lin, J. & Schulman, K. Prevalence of clinical venous thromboembolism in the USA: current trends and future projections. *American journal of hematology* **86**, 217-220 (2011).

Collen, D. The plasminogen (fibrinolytic) system. *Thrombosis and haemostasis* **82**, 259-270 (1999).

dela Peña, I., Borlongan, C., Shen, G. & Davis, W. Strategies to extend thrombolytic time window for ischemic stroke treatment: an unmet clinical need. *Journal of stroke* **19**, 50 (2017).

Rijken, D. & Lijnen, H. New insights into the molecular mechanisms of the fibrinolytic system. *Journal of thrombosis and haemostasis* **7**, 4-13 (2009).

Agnelli, G., Buchanan, M. R., Fernandez, F. & Hirsh, J. The thrombolytic and hemorrhagic effects of tissue type plasminogen activator: Influence of dosage regimens in rabrits. *Thrombosis research* **40**, 769-777 (1985).

Ohbayashi, T. *et al.* Degradation of fibrinogen and collagen by staphopains, cysteine proteases released from Staphylococcus aureus. *Microbiology* **157**, 786-792 (2011).

Lund, L. R. *et al.* Functional overlap between two classes of matrix-degrading proteases in wound healing. *The EMBO Journal* **18**, 4645-4656 (1999).

Saito, S. *et al.* Role of matrix metalloproteinases 1, 2, and 9 and tissue inhibitor of matrix metalloproteinase-1 in chronic venous insufficiency. *Journal of Vascular Surgery* **34**, 930-937 (2001).

Asahi, M. *et al.* Role for matrix metalloproteinase 9 after focal cerebral ischemia, effects of gene knockout and enzyme inhibition with BB-94. *Journal of Cerebral Blood Flow and Metabolism* **20**, 1681-1689 (2000).

Fujimura, M. *et al.* Early appearance of activated matrix metalloproteinase-9 and blood-brain barrier disruption in mice after focal cerebral ischemia and reperfusion. *Brain Research* **842**, 92-100 (1999).

36 Koshiba, T. *et al.* Detection of matrix metalloproteinase activity in human pancreatic
cancer. *Surgery Today-the Japanese Journal of Surgery* **27**, 302-304 (1997).
37 Hanemaaijer, R., Visser, H., Konttinen, Y. T., Koolwijk, P. & Verheijen, J. H. A novel and
simple immunocapture assay for determination of gelatinase-B (MMP-9) activities in
biological fluids: Saliva from patients with Sjogren's syndrome contain increased latent
and active gelatinase-B levels. *Matrix Biology* **17**, 657-665 (1998).
38 Sakalihasan, N., Delvenne, P., Nusgens, B. V., Limet, R. & Lapiere, C. M. Activated forms
of MMP(2) and MMP(9) in abdominal aortic aneurysms. *Journal of Vascular Surgery* **24**,
127-133 (1996).
39 Bini, A., Itoh, Y., Kudryk, B. J. & Nagase, H. Degradation of Cross-Linked Fibrin by
Matrix Metalloproteinase 3 (Stromelysin 1): Hydrolysis of the γ Gly 404− Ala 405 Peptide
Bond. *Biochemistry* **35**, 13056-13063 (1996).
40 Lelongt, B. *et al.* Matrix Metalloproteinase 9 Protects Mice from Anti–Glomerular
Basement Membrane Nephritis through Its Fibrinolytic Activity. *Journal of Experimental*
Medicine **193**, 793-802 (2001).
41 Hotary, K. B. *et al.* Matrix metalloproteinases (MMPs) regulate fibrin-invasive activity via
MT1-MMP–dependent and–independent processes. *Journal of Experimental Medicine*
195, 295-308 (2002).
42 Lijnen, H. Matrix metalloproteinases and cellular fibrinolytic activity. *Biochemistry*
(Moscow) **67**, 92-98 (2002).
43 Merchant, K. A., Best, R. B., Louis, J. M., Gopich, I. V. & Eaton, W. A. Characterizing
the unfolded states of proteins using single-molecule FRET spectroscopy and molecular
simulations. *Proceedings of the National Academy of Sciences* **104**, 1528-1533 (2007).
44 Best, R. B. *et al.* Effect of flexibility and cis residues in single-molecule FRET studies of
polyproline. *Proceedings of the National Academy of Sciences* **104**, 18964-18969 (2007).
45 Stennett, E. M., Ciuba, M. A. & Levitus, M. Photophysical processes in single molecule
organic fluorescent probes. *Chemical Society Reviews* **43**, 1057-1075 (2014).
46 Cordes, T., Vogelsang, J. & Tinnefeld, P. On the mechanism of Trolox as antiblinking and
antibleaching reagent. *Journal of the American Chemical Society* **131**, 5018-5019 (2009).
47 Rasnik, I., McKinney, S. A. & Ha, T. Nonblinking and long-lasting single-molecule
fluorescence imaging. *Nature methods* **3**, 891 (2006).
48 Kochevar, I. E. & Redmond, R. W. in *Methods in enzymology* Vol. 319   20-28 (Elsevier,
2000).
49 Levitus, M. & Ranjit, S. Cyanine dyes in biophysical research: the photophysics of
polymethine fluorescent dyes in biomolecular environments. *Quarterly reviews of*
biophysics **44**, 123-151 (2011).
50 Kumar, L. *et al.* Allosteric communications between domains modulate activity of matrix
metalloprotease-1. *bioRxiv*, 804559 (2019).
51 Zhuang, X. *et al.* A single-molecule study of RNA catalysis and folding. *Science* **288**,
2048-2051 (2000).
52 Jarymowycz, V. A. & Stone, M. J. Fast time scale dynamics of protein backbones: NMR
relaxation methods, applications, and functional consequences. *Chemical reviews* **106**,
1624-1671 (2006).
53 Wertheim, G., Butler, M., West, K. & Buchanan, D. Determination of the Gaussian and
Lorentzian content of experimental line shapes. *Review of Scientific Instruments* **45**, 1369-
1371 (1974).

54 Weisel, J. W. & Litvinov, R. I. in *Fibrous proteins: structures and mechanisms* 405-456
(Springer, 2017).
55 Shoulders, M. D. & Raines, R. T. Collagen structure and stability. *Annual review of*
biochemistry **78**, 929-958 (2009).
56 Eyal, E., Lum, G. & Bahar, I. The anisotropic network model web server at 2015 (ANM
2.0). *Bioinformatics* **31**, 1487-1489 (2015).
57 Kozakov, D. *et al.* The ClusPro web server for protein–protein docking. *Nature protocols*
12, 255-278 (2017).
58 Bini, A., Wu, D., Schnuer, J. & Kudryk, B. J. Characterization of stromelysin 1 (MMP-3),
matrilysin (MMP-7), and membrane type 1 matrix metalloproteinase (MT1-MMP) derived
fibrin (ogen) fragments D-dimer and D-like monomer: NH2-terminal sequences of late-
stage digest fragments. *Biochemistry* **38**, 13928-13936 (1999).
59 GÅRDLUND, B., HESSEL, B., BLOMBÄCK, B., MARGUERIE, G. & MURANO, G.
Primary Structure of Human Fibrinogen: Characterization of Disulfide-Containing
Cyanogen-Bromide Fragments. *European journal of biochemistry* **77**, 595-610 (1977).
60 Blombäck, B. & Blombäck, M. The molecular structure of fibrinogen. *Annals of the New*
York Academy of Sciences **202**, 77-97 (1972).
61 Dittmore, A. *et al.* Internal strain drives spontaneous periodic buckling in collagen and
regulates remodeling. *Proceedings of the National Academy of Sciences*, 201523228
(2016).
62 Lai, V. K. *et al.* Microstructural and mechanical differences between digested collagen–
fibrin co-gels and pure collagen and fibrin gels. *Acta biomaterialia* **8**, 4031-4042 (2012).
63 Gelse, K., Pöschl, E. & Aigner, T. Collagens—structure, function, and biosynthesis.
Advanced drug delivery reviews **55**, 1531-1546 (2003).
64 Bain, B. J. Diagnosis from the blood smear. *New England Journal of Medicine* **353**, 498-
507 (2005).
65 Dittmore, A. *et al.* Internal strain drives spontaneous periodic buckling in collagen and
regulates remodeling. *Proceedings of the National Academy of Sciences*, 8436-8441
(2016).
66 Sarkar, S. K., Marmer, B., Goldberg, G. & Neuman, K. C. Single-molecule tracking of
collagenase on native type I collagen fibrils reveals degradation mechanism. *Current*
Biology **22**, 1047-1056 (2012).
67 Colomb, W., Czerski, J., Sau, J. & Sarkar, S. Estimation of microscope drift using
fluorescent nanodiamonds as fiducial markers. *Journal of Microscopy* **266**, 298-306
(2017).
68 Czerski, J., Colomb, W., Cannataro, F. & Sarkar, S. Spectroscopic identification of
individual fluorophores using photoluminescence excitation spectra. *Journal of*
microscopy **270**, 261-271 (2018).
69 Kollman, J. M., Pandi, L., Sawaya, M. R., Riley, M. & Doolittle, R. F. Crystal structure of
human fibrinogen. *Biochemistry* **48**, 3877-3886 (2009).
70 Everse, S. J., Spraggon, G., Veerapandian, L., Riley, M. & Doolittle, R. F. Crystal structure
of fragment double-D from human fibrin with two different bound ligands. *Biochemistry*
37, 8637-8642 (1998).
71 Shindyalov, I. N. & Bourne, P. E. Protein structure alignment by incremental combinatorial
extension (CE) of the optimal path. *Protein engineering* **11**, 739-747 (1998).

72   Schrodinger, L. The PyMOL molecular graphics system, version 1.8. *Schrodinger LLC,*
664   *New York, NY* (2015).

Supplementary Information

Inter-domain dynamics and fibrinolytic activity of matrix
metalloprotease-1

Lokender Kumar[1,#], Joan Planas-Iglesias[2,3#], Chase Harms[1], Sumaer Kamboj[1], Derek
Wright[1], Judith Klein-Seetharaman[2,4,*], and Susanta K. Sarkar [1, *]

[1]*Department of Physics, Colorado School of Mines, 1500 Illinois Street, CO 80401, USA*
[2]*Warwick Medical School, University of Warwick, Coventry CV4 7AL, UK*
[3]*Loschmidt Laboratories, Department of Experimental Biology, Faculty of Science, Masaryk*
University, Kamenice 5/A13, 625 00 Brno, Czech Republic
[4]*Department of Chemistry, Colorado School of Mines, 1500 Illinois Street, CO 80401, USA*

14
15 [#]*These authors contributed equally*

16
17 *Corresponding author:*

18 [*]*ssarkar@mines.edu*

22
23

24   We fitted a sum of two Gaussians to the experimental histograms using the following equation:

$$y = a_1 \times e^{-\frac{(x-b_1)^2}{c_1^2}} + a_2 \times e^{-\frac{(x-b_2)^2}{c_2^2}}$$

26   where a, b, and c are amplitude, center, and width of the Gaussian, respectively. The parameters b1
27   and b2 are the two states, S1 and S2, respectively.

Table S1. Best-fit parameters for experimental histograms and autocorrelations. (A) A sum of two Gaussians fits the experimental histograms in **Figure 3**. (B) An exponential distribution fits the experimental autocorrelations in **Figure 3.** Power law distribution does not fit the experimental autocorrelations. (C) Kinetic rates of interconversion between the two states, S1 and S2, from the histograms and autocorrelations. Yellow highlighted numbers are (A) the centers of Gaussian fits and (B) the decay rates of exponential fits, e. The error bars represent the standard errors of mean.

A Gaussian fit parameters for experimental histograms

	MMP1 without ligands		MMP1 with MMP9		MMP1 with tetracycline	
	Active	Inactive	Active	Inactive	Active	Inactive
a1	1.99±0.13	0.90±0.17	2.14±0.06	1.18±3.21	4.38±0.05	0.60±0.79
b1/S1	0.42±0.01	0.52±0.01	0.46±0.01	0.46±0.18	0.61±0.01	0.49±0.13
c1	0.08±0.01	0.10±0.01	0.11±0.01	0.11±0.06	0.10±0.01	0.11±0.06
a2	5.52±0.15	8.12±0.17	5.58±0.07	4.61±4.12	2.10±0.12	5.6±1.21
b2/S2	0.51±0.01	0.55±0.01	0.51±0.01	0.53±0.01	0.70±0.01	0.58±0.01
c2	0.07±0.01	0.06±0.01	0.06±0.01	0.09±0.01	0.06±0.01	0.09±0.01

B Exponential fit parameters for correlations

	MMP1 without ligands		MMP1 with MMP9		MMP1 with tetracycline	
	Active	Inactive	Active	Inactive	Active	Inactive
d	0.37±0.01	0.15±0.01	0.21±0.01	0.45±0.01	0.62±0.01	0.36±0.01
e	0.08±0.01	0.07±0.01	0.09±0.01	0.05±0.01	0.05±0.01	0.06±0.01
f	-0.03±0.01	-0.01±0.01	-0.01±0.01	-0.03±0.01	-0.05±0.01	-0.004±0.001

C Kinetic rates calculated from histograms and correlations

	MMP1 without ligands		MMP1 with MMP9		MMP1 with tetracycline	
	Active	Inactive	Active	Inactive	Active	Inactive
k1 (s^{-1})	0.0567	0.0591	0.0528	0.0381	0.0112	0.0531
k2 (s^{-1})	0.0233	0.0109	0.0372	0.0119	0.0388	0.0069

30   We subtracted the average FRET value from each FRET trajectory and used the following equation
31   to calculate autocorrelations:

$$C_\tau = \frac{1}{N-\tau} \sum_{t=0}^{N-\tau} \left\{ I(t) - \frac{1}{N-\tau} \sum_{t'=0}^{N-\tau} I(t) \right\} \times \left\{ I(t+\tau) - \frac{1}{N-\tau} \sum_{t'=\tau}^{N} I(t) \right\}$$

where C_τ is the autocorrelation at lag number τ, N is the number of points in a FRET trajectory,
and $I(t)$ is the FRET value at t.

We normalized autocorrelations by dividing autocorrelations at each lag by $C_{\tau=0}$. We fitted
autocorrelations between $\tau=1$ and $\tau=1000$ to both power law and exponential distributions. For
power law, we used a form of Pareto distribution[1] that satisfies the boundary conditions of our
calculated autocorrelations, i.e., $C_{\tau=0}=1$ at $t=0$ and $C_{\tau=\infty}=0$ at $t=\infty$. We fitted the
following equations of power law and exponential functions:

$$C_{\tau}=(a\times\tau+1)^{-b}$$

$$C_{\tau}=d\times\exp^{-e\times\tau}+f$$

Best-fit parameters for two-state simulations. We simulated smFRET trajectories assuming that
MMP1 undergoes interconversion between two states, S1 and S2, which are the centers of
Gaussian fits to the histograms. We considered active MMP1 and active site mutant of MMP1
without ligands (**Figure 3A**). We simulated 350 smFRET trajectories, each 1000 s long, with the
input parameters in **Table S2**. We analyzed the simulated and experimental trajectories similarly.
The recovered parameters (**Table S2**, right side) agree well with the input parameters.

Table S2. MMP1 inter-domain dynamics as a two-state Poisson process. Analysis of FRET trajectories for MMP1 without ligands recovered the centers and decay rates used as inputs. The agreement between the input and retrieved values suggests that a two-state Poisson process describes the MMP1 dynamics.

Input parameters

Active MMP1
S1 = 0.42, S2 = 0.51
k1 = 0.06 s⁻¹, k2 = 0.02 s⁻¹
k1 + k2 = 0.08 s⁻¹
σ1 = 0.08, σ2 = 0.07

Inactive MMP1
S1 = 0.52, S2 = 0.55
k1 = 0.06 s⁻¹, k2 = 0.01 s⁻¹
k1 + k2 = 0.07 s⁻¹
σ1 = 0.10, σ2 = 0.06

Recovered parameters
Gaussian fit parameters for simulated histograms

	Without noise		With noise	
	Active	Inactive	Active	Inactive
a1			2.06±0.01	0.91±0.02
b1/S1	0.42	0.52	0.42±0.01	0.52±0.01
c1			0.08±0.01	0.10±0.01
a2			5.70±0.02	7.91±0.02
b2/S2	0.51	0.55	0.51±0.01	0.55±0.01
c2			0.07±0.01	0.06±0.01

Power law and exponential fit parameters for correlations

	Without noise		With noise	
	Active	Inactive	Active	Inactive
a	0.001±0.001	0.001±0.001	NA	NA
b	1882±1354	2080±1723	NA	NA

	Without noise		With noise	
	Active	Inactive	Active	Inactive
d	1.03±0.01	1.03±0.01	0.39±0.01	0.05±0.01
e	0.08±0.01	0.07±0.01	0.08±0.01	0.07±0.01
f	-0.03±0.01	-0.04±0.01	-0.01±0.01	-0.01±0.01

48
49

Figure S1. MMP1 inter-domain dynamics as a Poisson process without noise. (**A**) An example of simulated two-state FRET trajectory without noise for active MMP1 (blue) and active site mutant of MMP1 (orange). (**B**) Histograms of the recovered FRET values with bin size=0.005. (**C**) Autocorrelations of simulated trajectories recover the sum, k1+k2, from exponential fits (active MMP1: dashed black line; active site mutant of MMP1: solid black line). Note that power law does not fit the autocorrelations with noise (**Fig. 4**). However, power law fits the autocorrelations without noise (active MMP1: dashed red line; active site mutant of MMP1: solid green line).The error bars are the sems for histograms and autocorrelations and are too small to be seen.

51   **Potential sites of interactions between MMP1 and fibrin.** MMP3, MMP7, and MMP14 cleave
52   the α-chain at Asp97-Phe98 and Asn102-Asn103; the β-chain at Asp123-Leu124, Asn137-
53   Val138, and Glu141-Tyr142; and the γ-chain at Thr83-Leu84[2] as shown in **Figure S2**.

Figure S2. Potential MMP1 cleavage sites on the three fibrin chains. The balls are potential cleavage sites on fibrin that we considered to select the best docking pose between MMP1 and fibrin.

54 **Inter-domain distance correlates with the catalytic pocket opening of MMP1 on fibrinogen.**
55 In contrast to free MMP1 (**Figure 5**) and fibrin-bound MMP1 (**Figure 6**), fibrinogen-bound

Figure S3. Correlations of MMP1 inter-domain distance with catalytic pocket opening when MMP1 is bound to fibrinogen. Examples of (**A**) open and (**B**) closed conformations of MMP1 (PDB ID: 1SU3) bound to fibrinogen (PDB ID: 3GHG). Three dimensional scatter plots (blue circle) of inter-domain distance (S142-S366), catalytic pocket opening (N171-T230), and rms proximity between the MMP1 catalytic site and the three fibrinogen chains for (**C**) open and (**D**) closed MMP1 conformations. Two-dimensional projections of the scatter plots are in gray.

MMP1 (**Figure S3**) shows smaller catalytic pocket openings for the open MMP1 conformations.
However, fibrinogen-bound MMP1 shows closer proximity to the fibrinogen chains in agreement
with free MMP1 and fibrin-bound MMP1. The smaller catalytic pocket opening suggests a
different mechanism of fibrinogen and fibrin degradation by MMP1, which needs further studies
to confirm.

MMP1 activity on fibrinogen. The point mutation E219Q renders MMP1 catalytically inactive
on collagen. For fibrinogen degradation, however, the E219Q mutant degraded the α- and β-
chains, but did not degrade the γ-chain (**Figure S4A**, lane 4 from left). We used trypsin during
MMP1 purification, and trypsin is known to degrade fibrinogen[3]; therefore, any residual trypsin
could potentially interfere with the results. To address this possibility, we added 0.5 mg/mL trypsin
inhibitor to all the reactions, which was sufficient to inhibit 0.1 mg/mL trypsin (**Figure S4A**, lane
5 from left).

Figure S4. Ensemble activity of MMP1 on fibrinogen and fibrin without crosslinks. (**A**) SDS PAGE
of fibrinogen treated with different enzymes. Since we used trypsin to activate and purify MMP1, we
added 0.5 mg/mL trypsin inhibitor to lane 2-6 to control for any residual effect of trypsin. Trypsin
inhibitor (Worthington, Lima Bean, Cat# LS002829) at 0.5 mg/mL inhibited the fibrinolytic effect of 0.1
mg/mL trypsin used in lane 5. Note that the difference between trypsin inhibitor and MMP1 inhibitor
(tetracycline). (**B**) Time-dependent fibrinogen degradation by active MMP1. (**C**) SEM images of fibrin
after MMP1 treatment. Note that the active site mutant of MMP1 is catalytically inactive on collagen[4].
However, as evident from **Figure S4A**, active site mutant of MMP1 is partially active on fibrinogen. In
contrast, active site mutant of MMP1 is catalytically inactive on crosslinked fibrin, as shown in **Figure
S5** and **Figure S6**.

Even though limited digestion of fibrinogen by thrombin leads to fibrin, the longer digestion for
24 h led to further degradation of fibrinogen (**Figure S4A**, lane 6 from left). **Figure S4A** does not

reveal the sequence in which MMP1 degrades the three chains of fibrinogen. **Figure S4B** shows
the time-dependent degradation of fibrinogen. MMP1 degrades the α-chain first and the γ-chain
last. A comparison with prior research shows that plasmin, the well-known fibrinolytic agent, also
degrades the α-chain first and the γ-chain last[5]. Next, we tested the effects of MMP1 on fibrin
morphology using SEM (**Figure S4C**), where the destruction of fibrin structures is visible for
active MMP1. We created similar thin layers of fibrin morphologies for smFRET experiments.
Overall, the ensemble experiments in **Figure S4** show that MMP1 has fibrinolytic activity on
fibrinogen and fibrin, and there is a specific sequence of degradation of the three chains. Both
thrombin and trypsin cleave the same Arg-Gly bonds in fibrinogen[6]. However, thrombin is highly
specific, and only four bonds linking the fibrinopeptides to the rest of the fibrinogen molecule are
rapidly hydrolyzed by thrombin, whereas trypsin cleaves all peptide bonds in fibrinogen that
follow arginine or lysine[6].

In comparison, degradation of fibrinogen and
crosslinked fibrin by MMPs do not have
specific cleavage sites across the MMP-family.
For example, MMP3, MMP7, and MMP14
cleave the α-chain at Asp97-Phe98 and
Asn102-Asn103; the β-chain at Asp123-
Leu124, Asn137-Val138, and Glu141-Tyr142;
and the γ-chain at Thr83-Leu84[2]. Cleavage
sites on both fibrinogen and crosslinked fibrin
for these three MMPs are close to those for
plasmin. Additionally, MMP3 solubilizes
crosslinked fibrin by cleaving the Gly404-
Ala405 bond in the γ-chain[7], resulting in a D-
like monomer fragment at ~94 kDa similar to
fibrinogen degradation by plasmin. In contrast,
MMP7 and MMP14 solubilize crosslinked
fibrin and produces D-like dimer fragments at
~186 kDa similar to crosslinked fibrin
degradation by plasmin[2]. However, MMP1,
MMP2, MMP9, and MMP15 do not show

Figure S5. Weight-based fibrin degradation assay. Time-dependent weights of crosslinked fibrin after treatment with active MMP1, active site mutant of MMP1, trypsin, thrombin, and PBS buffer as the control.

similar activity[2]. Fibrinogen degradation by catalytic domains of MMP8, MMP12, MMP13, and
MMP14 show differences in the cleavage sites[8]. Overall, MMPs show differences in fibrinolytic
activity, which is intriguing because the catalytic domain has a largely conserved sequence among
the MMP-family members[9]. Interestingly, prior research reported insignificant fibrinolytic activity
of MMP1[7], in contrast to our findings.

Weight-based degradation assay for water-insoluble fibrin. We used a weight-based
degradation assay to quantify the activity of MMP1 because fibrin is water-insoluble and as such,
solution-based biochemical assays are appropriate. We prepared fibrin with crosslinks by mixing
5 µg of human factor XIII, 20 units of thrombin (Cayman chemical, Cat# 13188), and 40 µL of
10 mM PBS (pH 7.4) in a 0.5 mL PCR tube at 22º C. We incubated the mixture for 10 min at 37º
C without shaking. We added 100 µg of fibrinogen (Cayman chemical, Cat# 16088) and 20 µL of
5 mM CaCl2 to the mixture and incubated for an additional 15 min at 37º C without shaking. After
incubation, the solution becomes turbid, indicating the formation of crosslinked fibrin. We

prepared five reactions for crosslinked fibrin and added 1) 100 µL of 10 mM PBS (pH 7.4), 2) 100
µL of 1 mg/mL active MMP1, 3) 100 µL of 1 mg/mL active MMP1, 4) 100 µL of 1 mg/mL trypsin,
and 5) 100 units of thrombin. We made the final volume of each reaction to be 200 µL by diluting
with 10 mM PBS (pH 7.4). At 0 hr, we centrifuged the reactions at 10000 rpm for 10 min using a
tabletop centrifuge, discarded the supernatant, and weighed the tubes. We subtracted the weights
of the PCR tubes for each specific reaction before preparing the reactions. For five conditions, we
measured weight of ~120 mg. After weighing, we again added 1) 100 µL of 10 mM PBS (pH 7.4),
2) 100 µL of 1 mg/mL active MMP1, 3) 100 µL of 1 mg/mL active MMP1, 4) 100 µL of 1 mg/mL
trypsin, and 5) 100 units of thrombin to a final volume of 200 µL by diluting with 10 mM PBS
(pH 7.4). We incubated the reactions at 37º C without shaking. At 1 h, we centrifuged the tubes
again, weighed, and subtracted the weight of empty tubes. We diluted again as above and incubated
for another hour. We repeated the process at 0 h, 1 h, 2 h, 3 h, 4 h, and 5 h. We performed three
biological repeats to calculate the mean and standard deviations. **Figure S5** shows that active
MMP1 and trypsin degrade crosslinked fibrin at a similar rate of ~30 mg/h. Active site mutant of
MMP1, thrombin, and control show similar changes in weights at a rate of ~7 mg/h. After
subtracting the control, the rate of crosslinked fibrin degradation by MMP1 is ~0.23 mg/h/µg. Note
that the weight includes water and, as such, has a higher value.

Figure S6. Fibrinolytic activity of MMP1 on crosslinked fibrin. (**A**) SDS PAGE of crosslinked fibrin
with active MMP1 and active site mutant of MMP1. The control uses the protein buffer (50 mM Tris,
100 mM NaCl, pH 8.0). (**B**) 100 mg of wet crosslinked fibrin treated with 0.1 mg/mL active MMP1 and
active site mutant of MMP1 at 37º C. (**C**) SEM images of crosslinked fibrin with and without MMP1
treatment.

MMP1 activity on crosslinked fibrin. Water-soluble fibrinogen becomes water-insoluble crosslinked fibrin in the presence of factor XIII, thrombin, and $CaCl_2$[10]. Thrombin converts fibrinogen into fibrin monomer by cleaving fibrinopeptide A and fibrinopeptide B[11,12]. Fibrin monomers self-assemble into protofibrils. The polymerization sites noncovalently attach to the D regions of two other fibrin(ogen) molecules. Fibrin monomers in each strand assemble end-to-end, whereas monomers across the strands arrange in a half-staggered overlap. The γ-chains of adjacent D regions in each strand is covalently attached by isopeptide bonds, which are formed due to the activity of factor XIIIa and appear as a ~94 kDa band in SDS PAGE under reducing conditions (**Figure S6A**). Since factor XIIIa does not crosslink the β-chains[13,14], the band at ~52 kDa due to the β-chains remains unchanged in SDS PAGE of fibrinogen (**Figure S6A**) and crosslinked fibrin (**Figure S6A**). As shown in **Figure S6B**, active MMP1 degrades and dissolves water-insoluble crosslinked fibrin, but active site mutant of MMP1 does not dissolve crosslinked fibrin. **Figure S6C** shows the surface morphology of crosslinked fibrin imaged using SEM. Treatment with active MMP1 resulted in a more porous structure of crosslinked fibrin.

References

Arnold, B. C. Pareto distribution. *Wiley StatsRef: Statistics Reference Online*, 1-10 (2014).

Bini, A., Wu, D., Schnuer, J. & Kudryk, B. J. Characterization of stromelysin 1 (MMP-3), matrilysin (MMP-7), and membrane type 1 matrix metalloproteinase (MT1-MMP) derived fibrin (ogen) fragments D-dimer and D-like monomer: NH2-terminal sequences of late-stage digest fragments. *Biochemistry* **38**, 13928-13936 (1999).

Astrup, T. & Sterndorff, I. Fibrinolytic activity of tissue extracts and of trypsin. *Nature* **170**, 981-981 (1952).

Sarkar, S. K., Marmer, B., Goldberg, G. & Neuman, K. C. Single-molecule tracking of collagenase on native type I collagen fibrils reveals degradation mechanism. *Current Biology* **22**, 1047-1056 (2012).

Pizzo, S. V., Schwartz, M. L., Hill, R. L. & McKee, P. A. The effect of plasmin on the subunit structure of human fibrinogen. *Journal of biological chemistry* **247**, 636-645 (1972).

Andreatta, R. H., Liem, R. K. & Scheraga, H. A. Mechanism of action of thrombin on fibrinogen, I. Synthesis of fibrinogen-like peptides, and their proteolysis by thrombin and trypsin. *Proceedings of the National Academy of Sciences* **68**, 253-256 (1971).

Bini, A., Itoh, Y., Kudryk, B. J. & Nagase, H. Degradation of Cross-Linked Fibrin by Matrix Metalloproteinase 3 (Stromelysin 1): Hydrolysis of the γ Gly 404− Ala 405 Peptide Bond. *Biochemistry* **35**, 13056-13063 (1996).

Hiller, O., Lichte, A., Oberpichler, A., Kocourek, A. & Tschesche, H. Matrix metalloproteinases collagenase-2, macrophage elastase, collagenase-3, and membrane type 1-matrix metalloproteinase impair clotting by degradation of fibrinogen and factor XII. *Journal of Biological Chemistry* **275**, 33008-33013 (2000).

Ratnikov, B. I. *et al.* Basis for substrate recognition and distinction by matrix metalloproteinases. *Proceedings of the National Academy of Sciences* **111**, E4148-E4155 (2014).

Weisel, J. W. & Litvinov, R. I. in *Fibrous proteins: structures and mechanisms*   405-456 (Springer, 2017).

Medved, L. & Weisel, J. Recommendations for nomenclature on fibrinogen and fibrin. *Journal of Thrombosis and Haemostasis* **7**, 355-359 (2009).

12   Weisel, J. W. & Litvinov, R. I. Mechanisms of fibrin polymerization and clinical
implications. *Blood, The Journal of the American Society of Hematology* **121**, 1712-1719
(2013).
13   McKee, P. A., Mattock, P. & Hill, R. L. Subunit structure of human fibrinogen, soluble
fibrin, and cross-linked insoluble fibrin. *Proceedings of the National Academy of Sciences*
66, 738-744 (1970).
14   Sobel, J. H. & Gawinowicz, M. A. Identification of the α chain lysine donor sites involved
in factor XIIIa fibrin cross-linking. *Journal of Biological Chemistry* **271**, 19288-19297
(1996).

187