Towards the molecular architecture of the peroxisomal receptor docking complex

Pascal Lill[1a], Tobias Hansen[2a], Daniel Wendscheck[3a], Bjoern Udo Klink[1], Tomasz Jeziorek[2], Julian Bender[3,5], Friedel Drepper[3,4], Wolfgang Girzalsky[2], Bettina Warscheid[3,4b], Ralf Erdmann[2b], Christos Gatsogiannis[1*]

[1]Department of Structural Biochemistry, Max Planck Institute of Molecular Physiology, Dortmund, Germany

[2]Institute of Biochemistry and Pathobiochemistry, Faculty of Medicine, System Biochemistry, Ruhr-University Bochum, Bochum, Germany

[3]Biochemistry and Functional Proteomics, Institute of Biology II, Faculty of Biology, University of Freiburg, 79104 Freiburg, Germany

[4]Signalling Research Centres BIOSS and CIBSS, University of Freiburg

[5]Current address: Interdisciplinary Research Center HALOmem, Institute for Biochemistry and Biotechnology, Martin Luther University Halle-Wittenberg, Halle (Saale), Germany.

[a]These authors contributed equally to this work

[b]These authors contributed equally to this work

*For correspondence: christos.gatsogiannis@mpi-dortmund.mpg.de

Abstract

Import of yeast peroxisomal matrix proteins is initiated by cytosolic receptors, which specifically recognize and bind the respective cargo proteins. At the peroxisomal membrane, the cargo-loaded receptor interacts with the membrane docking complex, composed of Pex14p, Pex17p and Pex13p. Previous data suggest that this interaction triggers the formation of an import pore for further translocation of the cargo. The mechanistic principles are however unclear, mainly because structures of higher order assemblies are still lacking. Here, using an integrative approach, we provide the first structural characterization of the major components of the peroxisomal docking complex Pex14p/Pex17p, in a native bilayer environment and reveal its subunit organization. Our data show that three copies of Pex14p and a single copy of Pex17p assemble to form a 20 nm rod-like particle. The different subunits are arranged in a parallel manner, showing interactions along their complete sequences and providing receptor binding-sites on both membrane sides. The long rod facing the cytosol is mainly formed by the coiled-coil domains of Pex14p and Pex17p, possibly providing the necessary structural support for the formation of the import pore. Further implications of Pex14p/Pex17p for formation of the peroxisomal translocon are discussed.

Introduction

Peroxisomes are organelles present nearly ubiquitously in eukaryotic cells, ranging from unicellular yeasts to multicellular organisms, such as plants and humans. Beside ß-oxidation of fatty acids as a main conserved function of peroxisomes, a broad range of additional metabolic functions is linked to this organelle, underscored by severe and frequently lethal phenotypes of human disorders[1,2]. These organelles do not contain DNA and thus all peroxisomal matrix proteins are encoded in the nucleus and synthesized on free polyribosomes in the cytosol. Subsequently, matrix proteins are targeted to the organelle by peroxisomal import receptors[3]. A remarkable feature of peroxisomes is that unlike the transport of unfolded polypeptides across the membranes of the endoplasmic reticulum, mitochondria, and chloroplast, they can import already folded, cofactor-bound, and even oligomeric proteins[4,5]. This transport is highly selective and mediated by specific import sequences known as peroxisomal targeting signals (PTSs)[6,7]. Peroxisomal matrix proteins equipped with either a carboxy-terminal PTS1 or an amino-terminal PTS2, are recognized and bound in the cytosol by the import receptor Pex5p or Pex7p, respectively[8,9]. A peroxisomal membrane associated complex consisting of Pex13p, Pex14p and Pex17p in yeast allows docking of the cargo-loaded receptor[10-14]. This primary interaction of the cargo-loaded receptor with the docking complex induces the formation of a transient and highly dynamic import pore, necessary for the translocation of the cargo across the peroxisomal membrane[15-17]. How translocation and release of the cargo are realized in detail still remains enigmatic but it has been previously shown that the receptor is exported from the peroxisomal membrane in an

ubiquitin- and ATP-dependent manner, a process that is discussed to provide the driving force for cargo-import according to the export-driven-import model[18-20].

The receptor-docking complex is of major importance for peroxisomal matrix protein import, as it provides a binding platform for newly formed receptor-cargo complexes at the peroxisomal membrane. Both Pex13p and Pex14p are integral peroxisomal membrane proteins providing several binding sites for the import receptors Pex5p and Pex7p[16]. Pex17p is tightly associated with Pex14p and thus with the docking-complex[21], but its precise function remains unknown. Although Pex17p is part of the docking complex in yeast, it does not significantly contribute to the assembly of the Pex13p/Pex14p subcomplex[15,21,22], and its counterpart in higher eukaryotes has not yet been identified. However, Pex17p is essential for peroxisomal import of both PTS1 and PTS2 proteins[14]. Strikingly, both import receptors, Pex5p and Pex7p, associate with the docking-complex (Pex13p, Pex14p) in absence of Pex17p, but with decreased efficiency[22]. Pex17p is apparently non-essential for receptor docking, but it increases the efficiency of the docking event.

Furthermore, albeit a close association between the core components of the docking complex (Pex13p, Pex14p) is indispensable for matrix protein import [23], an assembly between the receptor Pex5p and the docking component Pex14p in absence of Pex13p is capable per se to form a large transient channel at the peroxisome membrane[15].

However, little is known about the molecular mechanism underlying the primary docking and subsequent translocation events, largely because structures of the higher-order assemblies are not available. Here, using cryo-

electron microscopy single particle analysis (cryo-EM SPA) and cryo-electron tomography (cryo-ET) combined with crosslinking and native mass spectrometry (MS), we set out to characterize the overall architecture of the yeast Pex14p/Pex17p complex.

Results

Sample optimization for structural studies

Yeast Pex14p consists of three structural regions: a highly conserved N-terminal region containing a putative transmembrane domain, a long and a short coiled-coil domain in the middle region and an unstructured C-terminal domain (Figure 1a). Yeast Pex14p contains two separated binding sites for the PTS1 import receptor Pex5p, localized at its N- and C-termini (Figure 1a) [24]. Mammalian PEX14 contains only one Pex5p binding site and only a single, shortened central coiled-coil domain[25], which has been shown to mediate dimerization, whereas the transmembrane domain is responsible for formation of higher order assemblies[26]. The topology of Pex14p at the peroxisomal membrane has been controversial, but recent results indicate that the N-terminus faces the matrix of the organelle[27]. Pex17p contains a putative transmembrane domain near the N-terminus and two short coiled-coil domains (Figure 1a). Pex14p and Pex17p associate together to form a tight core complex, however the oligomeric composition of the complex is yet unknown. Interestingly in rice blast fungus *Mangaporthe*, Pex14p and Pex17p fuse to form a unique Pex14p/Pex17p (or Pex33) peroxin[28]. Here, we co-expressed yeast Strep$_{II}$-Pex14p and His$_6$-Pex17p in *E. coli* (see Material and Methods). Initial experiments revealed a dominant Pex14p-degradation

product giving rise to a band in SDS-PAGE at about 35 kDa, which was most likely a consequence of a site-specific cleavage around the methionine at position 51, as identified by MS (Supplementary Figure 1a-c). Hence, we replaced methionine 51 by leucine and co-expressed this Pex14p-variant (Strep$_{II}$-Pex14p(M51L)), for which the additional band at 35 kDa almost disappeared (Supplementary Figure 1d and e). The complex was purified by a two-step affinity chromatography (Strep-Tactin and Ni-NTA affinity chromatography) and size exclusion chromatography in presence of DDM (Figure 1b, Supplementary Figure 2a). Analysis by negative stain EM revealed elongated flexible pin-like particles, which however displayed limited contrast and were partially disintegrated and/or formed aggregates, most probably due to hydrophobic interactions (Supplementary Figure 2a).

To optimize the sample and increase contrast, we replaced DDM (n-Dodecyl-β-D-maltoside) with amphipol A8-35[29]. However, the tight adsorption of amphipols to the hydrophobic transmembrane domains of Pex14p/Pex17p surprisingly resulted in dissociation of the complex and separation of Pex14p and Pex17p (Supplementary Figure 3). This might suggest that the transmembrane domains of Pex14p and Pex17p or regions in close proximity to them might be involved in crucial interactions for Pex14p/Pex17p complex formation. To overcome this issue, we exploited the possibility to reconstitute the Pex14p/Pex17p complex into lipid nanodiscs. Both Pex14p and Pex17p contain a single transmembrane helix, thus depending on the oligomeric state of the complex, only few membrane-spanning helices are expected (Figure 1a). Therefore, we used the membrane scaffold protein (MSP)1D1dH5 with a rather narrow predicted scaffold diameter of 8-9 nm for nanodisc

reconstitution. Despite extensive optimization efforts, the resulting proteo-nanodiscs were heterogeneous and incorporated multiple copies of the Pex14p/17p complex, usually from both sides (Supplementary Figure 2b). We then repeated the reconstitution experiments using the recently reported circular MSP1D1Δ4-6 nanodiscs[30] (Figure 1b). Δ4-6 circularized nanodisc with a diameter of 7 nm are according to our knowledge the smallest available nanodiscs to date and, in addition, they show superior stability and homogeneity in comparison to typical linear MSPs. Indeed, subsequent negative stain EM analysis revealed high-contrast monodispersed particles, with most nanodiscs trapping only a single copy of the complex that could be easily identified and automatically picked (Supplementary Figure 2c).

Pex14p/Pex17p is an elongated rod-like complex

To select the structure of the Pex14p/Pex17p complex, we recorded high-contrast images using Volta Phase-Plate (VPP) cryo-EM (Figure 1c-d). The digital micrographs show an even spread of elongated rod-like particles (Figure 1c-d), suggesting that most particles adopt a side-view orientation on the grid. Particles were selected automatically using crYOLO[31] and further processed with SPHIRE[32]. Reference-free 2D class averages revealed a single copy of Pex14p/Pex17p within the nanodisc (Figure 1e). In contrast to other membrane proteins reconstituted in larger nanodiscs and analyzed by cryoEM[33,34], the protein in this case is not "floating" within the nanodisc, but rather inserted peripherally at close proximity to the MSP (Figure 1e). This might indicate possible interactions between the transmembrane domain of Pex14p and Pex17p and the scaffold helices. Transmembrane helices are

however not resolved within the disc, suggesting either inaccurate alignment of the small transmembrane domain due to the compact density of the disc or flexibility of the transmembrane helices relative to the cytoplasmic rod. In contrast, the cytoplasmic domains are clearly visible and appear as five sequentially arranged globular densities and are stiff, mostly straight or slightly curved (Figure 1e). The class averages show an average length of ~220 Å (Figure 1f) and reveal additional flexibility at the end of the rod (Supplementary Video 1). We finally computed a 10 Å 3D reconstruction from ~82600 particles selected from the most homogeneous 2D class averages (Supplementary Figure 4).

The final density map revealed a 22 nm elongated rod inserted at the periphery of the nanodisc (Figure 2a). The rod appears hollow with an inner diameter of 7 Å. We further performed focused refinement of the rod after signal subtraction of the nanodisc, which significantly improved the respective density (Supplementary Figure 4). The resulting structure can be dissembled in two structural regions, showing a kinked connection (Figure 2b). Further weak connections are located along the lower density; they do not however involve a change in the direction of the overall rod axis. At several segments, the resolution is now significant, so that up to four α-helices are resolved, running approximately parallel to the rod axis (Figure 2b,c, Supplementary Video 1). The segments are however weakly connected and the density is locally fragmented, thereby not allowing to trace the inter-connections of the helices along the rod. Interestingly, the observed helical arrangement resembles a typical coiled-coil structure (Figure 2c). We therefore concluded that the elongated extra-membrane rod-like domain is mainly formed by the

predicted coiled-coil domains of Pex14p and Pex17p, which are located directly after the predicted transmembrane domains (Figure 1a).

Molecular composition of the Pex14p/Pex17p complex

To investigate the composition of the complex, we used native MS and gas phase collisional activation. Figure 3a shows native MS spectra for the co-expressed Pex14p(M51L)/Pex17p complex in DDM. The dominant peaks are assigned to a charge state series in the lower *m/z* region indicating masses of monomeric Pex14p and *E. coli* DnaK. Above *m/z* 4700, three partly resolved series were observed, which were assigned to molecular masses of 144 kDa, 263 kDa and 408 kDa. Isolation of ions within these series and subsequent collisional activation resulted in MS^2 spectra showing charge state series indicative of masses of monomeric Pex14p and Pex17p (see Supplementary Figure 5). Possible stoichiometries to account for these three masses are therefore 3:1, 6:1 and 9:2 of Pex14p:Pex17p as schematically depicted above the spectrum in Figure 3a. Of note, upon collisional activation, first monomeric Pex17p appeared as a main product, whereas at higher collisional energies Pex14p was predominant (Supplementary Figure 5).

In accordance with the native MS data, negative stain analysis of Pex14p(M51L)/Pex17p in DDM also revealed three distinct populations of particles: single rods, rod-dimers and rod-trimers (Figure 3b). The higher-order assemblies (rod-dimers and -trimers) can be explained by tail-to-tail hydrophobic interactions when the complex is solubilized in DDM. Indeed, these populations could not be observed after reconstitution of the complex in MSP1D1Δ4-6 nanodiscs (Figure 1e).

We further used size-exclusion chromatography combined with multi-angle light scattering (SEC-MALS) to verify the molecular weight of Pex14p(M51L)/Pex17p. The complex indeed ran at 143 kDa (±0.02%) consistent with the 3:1 stoichiometry (Supplementary Figure 6). Based on these results, we conclude that the Pex14p(M51L)/Pex17p complex consists of three copies of Pex14p(M51L) and one copy of Pex17p, with a molecular weight of ~143-144 kDa. This is also in accordance with the tetrameric arrangement of parallel α-helices observed locally at several segments of the cryoEM structure (Figure 2b-c).

To further verify the presence of Pex14p and Pex17p within the rod complexes observed by cryoEM, we co-expressed (Pex14p/His$_6$-Pex17p) and (His$_6$-Pex14p/Pex17p), reconstituted the respective complex in MSP1D1Δ4-6 nanodiscs, labelled with Ni-NTA-gold and imaged by negative stain EM (Figure 3c-d). In both cases, the gold particles bound below the nanodiscs, suggesting co-localization of the N-termini of Pex14p and Pex17p.

In order to challenge the oligomerization and overall structure of the Pex14p/Pex17p complex in a more close-to-native lipid environment and membrane curvature, we reconstituted the complex in liposomes and visualized the liposomes by negative stain EM (Figure 3e). After successful reconstitution, the liposomes are shaped like sea urchins, with multiple copies of rods (spikes) covering isotropically their outer surface. Further analysis of the liposomes by cryo-ET confirmed that Pex14p/Pex17p do not form larger super-complexes upon liposome incorporation (e.g. 6:1, 9:2 Pex14p:Pex17p) (Figure 3f-g, Supplementary Video 3). Mostly 20 nm single rods were visible

around the circumference of the liposomes, clearly resembling the cryo-EM structure in the nanodisc.

Interactions within *in vitro* and native Pex14p/Pex17p complexes

To obtain more detailed structural information on protein interactions within the Pex14p/Pex17p complex, the SEC-purified complex was subjected to chemical crosslinking combined with MS analysis (XL-MS). Figure 4a gives a schematic overview of the connected residues identified. Several linkages indicate major contacts between the C-terminal end of Pex17p (aa177-189) and the second predicted coiled-coil of Pex14p. This is in agreement with previous results identifying the extreme C-terminus of Pex17p (corresponding to aa167-199) as the smallest fragment that is sufficient for Pex14p-interaction[35]. Additional interfaces were identified between the first coiled-coil of Pex14p and the first coiled-coil of Pex17p as well as between the N-terminal domains of Pex14p (Lys67) and Pex17p (aa20-21), indicating further secondary contacts between both proteins.

Blue lines in Figure 4a represent identified homomultimeric linkages that are crosslinks involving two overlapping peptides, and thus, originating from two molecules of Pex14p. These linkages span the full region of predicted coiled-coils and are further localized within and close to the N-terminal three-helix domain of Pex14p (Figure 4a). Remaining linkages in Figure 4a shown as grey lines represent intra- or intermolecular contacts and most of them span the N-terminal domain of Pex14p.

To further characterize the Pex14p-Pex17p interactions, we performed domain-mapping experiments based on the yeast two-hybrid system with

Pex17p$_{167-199}$ fused to the Gal4p-DNA-binding domain (DB) and Gal4p-DNA-activation domain fusions of progressive carboxyl- and/or amino-terminal truncations of Pex14p. In line with previous findings, co-expression of Pex17p$_{167-199}$ with full-length Pex14p resulted in considerable ß-galactosidase, demonstrating the interaction of both proteins[35] (Figure 4b). However, Pex14p lacking N- or C-terminal regions did still interact with Pex17p$_{167-199}$, pointing to an internal binding region. In fact, we identified the second putative coiled-coil domain of Pex14p plus a C-terminal portion as sufficient to mediate interaction to Pex17p$_{167-199}$ (Figure 4b). This is in line with the XL-MS data, which also revealed the second putative coiled-coil domain of Pex14p as a major interaction site with Pex17p (Figure 4a).

Of note, Pex17p lacking its C-terminus did not interact with Pex14p in yeast two-hybrid assays, although further N-terminal contacts were detected by XL-MS (Figure 4a). Thus, in order to characterize possible secondary binding sites, Pex17p lacking its C-terminus (Pex17p$_{1-167}$) was analyzed in its endogenous environment, the peroxisomal membrane, instead of the nucleus, as it is the case in yeast two-hybrid assays. To this end, a wild-type strain expressing either Pex17p or Pex17p$_{1-167}$ genomically tagged with protein A (Pex17pTPA, Pex17p$_{1-167}$TPA) with a TEV cleavage site between both fusion partners was used. In line with published data and the function of Pex17p as a constituent of the receptor docking complex, Pex13p, Pex14p as well as the PTS1-receptor Pex5p were co-isolated when full-length Pex17p served as bait (Figure 4c)[36]. As originally described[21], other components of the importomer, represented by the two RING-finger proteins Pex10p and Pex12p, were also part of the Pex17p-complex. Pex17p lacking its C-terminal

Pex14p-binding module displayed a significant reduction in its steady-state level when compared with wild-type protein and the constituents of the importomer that were present in the Pex17p-complex, were also associated with Pex17p$_{1-167}$TPA, albeit in a much lower amount (Figure 4c). Thus, the results demonstrate that Pex17p$_{1-167}$ is still associated with the importomer, although it is lacking its C-terminal Pex14p-binding region. This is again in line with the XL-MS results, corroborating that interactions between the first putative coiled-coils and between the N-termini of co-expressed Pex14p and Pex17p (Figure 4a) promote the insertion of Pex17p$_{1-167}$ into Pex14p membrane complexes (Figure 4c). Interestingly, growth on oleic acid of cells deficient in Pex17p was restored upon expression of Pex17p$_{1-167}$, indicating that the truncated Pex17p$_{1-167}$ is biologically active (Figure 4d).

Intermolecular interactions within the predicted coiled-coil domains of Pex14p are essential for its oligomerization.

To assess which of the Pex14p-Pex14p interactions are essential for oligomerization, we used recombinantly expressed truncated Pex14p-variants and monitored formation of oligomers by chemical crosslinking and native MS. One N-terminal variant, Pex14p$_{1-95}$, and two C-terminal variants comprising solely the second coiled-coil domain (Pex14p$_{213-341}$) or both coiled-coil domains (Pex14p$_{119-341}$) were incubated with the crosslinker BS3 and products of the reactions separated by SDS-PAGE (Figure 5a). Whereas the N-terminal and the shorter C-terminal variant remained in the monomeric form, the larger C-terminal variant showed additional bands indicative of the formation of dimers and trimers (Figure 5a). With increasing concentrations of

the crosslinker, the trimer band became stronger than the dimer band, but no higher molecular weight bands of tetrameric or higher oligomeric species appeared (Supplementary Figure 7). The same Pex14p variants were studied by native MS preserving non-covalent interactions during transfer into the gas phase and subsequent analysis (Figure 5b). In line with the crosslinking results, the two shorter variants were observed solely monomeric, whereas the longer variant comprising both coiled-coil domains predominantly formed trimers and, to a considerably lesser extent, dimers and monomers.

From the crosslinking experiment, gel-separated monomeric and oligomeric forms of Pex14p$_{119-341}$ were subjected to digestion using trypsin and LC-MS/MS analysis for peptide identification. In addition, a quantitative MS analysis was performed, enabling a direct comparison of the abundance of individual cross-linked peptides identified in the homo-oligomer to those in the monomer (Supplementary Figure 8). The map of interactions resulting from this analysis is summarized in Figure 5c. We identified a set of 12 residue pairs specifically enriched with the homo-oligomer representing intra-molecular interactions along the full region of both predicted coiled-coil domains (Supplementary Figure 8c). Only four residue pairs had ratios close to 1:1 and, thus, were present in almost equal abundance in monomers and oligomers. Of note, one residue pair of high abundance (199/233) and two pairs of lower abundance (162/190 and 161/190) were monomer-specific, suggesting that these linkages may prevent assembly into oligomeric forms.

In conclusion, *in vitro* formation of Pex14p oligomers required the presence of its first coiled-coil domain. The identified intermolecular interactions along the full length of both predicted coiled-coil domains of Pex14p homo-oligomers

indicate a parallel assembly of the individual molecules. Connections between the two coiled-coil domains within Pex14p monomers interfere with the formation of higher oligomers, possibly by forcing bended conformations not compatible with the parallel geometry.

Discussion

It is well established that Pex13p, Pex14p, and Pex17p mediate the docking of cytosolic receptor-cargo complexes to the peroxisomal membrane[10-14,37-39]. Pex13p and Pex14p recognize and physically bind both of the import receptors, Pex5p and Pex7p. This, and the fact that Pex17p interacts with Pex14p[14,39, reviewed in 40] led on the one hand to the conclusion that these three peroxins form the receptor-docking complex. On the other hand, transport of the peroxin Pex8p into peroxisomes, requires only the presence of Pex14p and the import receptor Pex5p[41]. Interestingly, *in vitro*, the receptor and Pex14p alone are capable to form a gated ion-conducting channel[15]. These data together suggest that Pex14p oligomers constitute minimal receptor-docking complexes. In our study, we characterize the overall architecture of Pex14p in complex with Pex17p. Pex17p is present only in yeast and in this case, co-expression of Pex14p with Pex17p in *E. coli* was crucial to obtain samples of sufficient quality and amount for further structural studies.

The present study provides the first insights into the architecture of this central component of the peroxisomal import machinery in a membrane-mimicking environment, which we summarize in Figure 6a. We show that Pex14p and Pex17p together form a ~140 kDa complex (Figure 3a), containing three copies of Pex14p and one copy of Pex17p. The complex resembles a rod-like

particle with one end inserted in the lipid nanodisc (Figure 1e, Figure 2a). The length of the rod-part (~18 nm) corresponds to the predicted length of the two elongated Pex14p coiled-coil domains, thus suggesting that the extramembrane region of the particle is mainly formed by these long coiled-coils. We attempted to solve the structure of the complex by single particle cryo-EM, but visualization of a thin helical bundle inserted into, according to our knowledge, the smallest available lipid nanodisc, was not possible for conventional defocus-based imaging. Therefore, we used the Volta Phase Plate[42] in order to enhance the contrast of this weak phase object. We were able to obtain a low resolution reconstruction of the complex (Figure 2a) and further focused refinement of the rod after signal subtraction allowed us to reach sub-nanometer resolution, revealing a bundle of four helices at different heights of the rod (Figure 2b,c). This is in line with the 3:1 stoichiometry of the complex, suggesting an elongated parallel assembly of the helical regions of three copies of Pex14p and a single copy of Pex17p (Figure 6a). The rod-density shows a kink after 12.5 nm (Figure 2b,c), most probably resembling the short connection between the long (~85 aa) and short (~18 aa) coiled-coil domains of Pex14p. The XL-MS data identified intermolecular interactions along the full length of Pex14p and Pex17p (Figure 4a). Mapping of the Pex14p-Pex17p interactions was further confirmed by yeast two-hybrid assays and functional expression of a Pex17p variant (Figure 4b,c). The data clearly suggest a parallel arrangement of both molecules along their complete sequence (Figure 6a). Both coiled-coil domains of Pex14p are required for homooligomerization (Figure 5) and importantly, the three coiled-coil domains of Pex14p show a parallel and not an anti-parallel arrangement relative to

each other. The C-terminal domain of Pex14p (~100 aa) is not resolved in our cryo-EM structure, however, according to secondary and 3D structure predictions (Supplementary Figure 9), this domain is almost completely unstructured and apparently flexible (Figure 6a). The transmembrane domains of Pex14p and Pex17p could not be resolved, even after signal subtraction and focused refinement on the nanodisc density, suggesting a flexible arrangement of four α-helices spanning the membrane. The conserved N-terminal domain of Pex14p is connected to the transmembrane domain by a flexible linker (Supplementary Figure 9, Figure 6a) and is apparently flexible. Indeed, two-dimensional class averages of Pex14p/Pex17p indicate that this part of the complex is highly flexible, only appearing as blurry density below the nanodisc (Supplementary Video 1). Nevertheless, gold labeling experiments allowed us to detect the N-termini of both Pex14p and Pex17p below the nanodisc, suggesting an intra-peroxisomal localization of the respective domains (Figure 3c, Figure 6a). This is in agreement with recent studies, suggesting a N_{in}-C_{out} topology of Pex14p[27]. Cryo tomograms of reconstituted Pex14p/Pex17p in liposomes revealed isotropic decoration of the liposomes by the complex resulting in a sea urchin-like proteo-liposome.

At this point, it is important to emphasize that yeast Pex14p contains two binding sites for the PTS1 receptor Pex5p, which according to our results are located on both sides of the peroxisomal membrane: the first at the N-terminal domain (intra-peroxisomal) and the second at the C-terminal domain (facing the cytosol) (Figure 6a). The C-terminal domain of Pex14p (~100 aa)

containing the cytosolic binding-domain is predicted to be intrinsically disordered and, thus, highly flexible (Figure 6a).

Our data thus strongly indicate that the peroxisomal docking complex, mainly formed by Pex14p, resembles a rod-like structure that emanates into the cytosol and uses highly flexible terminal peptides to recruit the cargo-loaded receptor complexes (Figure 6b). The further steps of pore formation and cargo translocation can only be speculated at this stage. However, the overall Pex14p architecture presented in this study renders the possibility of dramatic Pex14p conformational changes and entry of further Pex14p domains into the membrane upon receptor binding rather unlikely. In particular, the 20 nm long and compact assembly of its central coiled-coil domain is not consistent with large structural rearrangements of this domain. A cluster of charged residues in-between the transmembrane helix and the first coiled-coil helix (Supplementary Figure 9) would prevent insertion of the coiled-coil helices into the membrane bilayer. The possible distance of the C-terminal receptor binding site to the membrane is unlikely to allow entry of this unstructured domain into the membrane without dramatic conformational changes of the coiled-coil domain. Furthermore, the α-helices of the conserved intra-peroxisomal flexible N-terminal domain containing the second receptor binding site are not amphipathic, and thus not capable to enter the membrane (Figure 6b).

The overall architecture of the complex suggests a pivotal role of Pex14p in recruitment of Pex5p/cargo complexes and providing the structural support necessary for further formation of the translocation pore. This would imply that the pore may be exclusively formed by the receptor Pex5p, which is in line

with the transient pore model[43] suggesting that the receptor Pex5p is functioning like a pore-forming toxin (Figure 6b). Indeed, import-deficient *pex14Δ* yeast mutants are capable to import matrix proteins after Pex5p over-expression[44] and *in vitro,* Pex5p is capable to enter the membrane without any help of the docking complex, when present at high concentrations[45]. The interaction of the N-terminal domain of Pex14p to membrane-embedded Pex5p might then further stabilize the Pex5p transient pore assembly (Figure 6b).

Our results provide the first mechanistic insight into the yeast peroxisomal docking complex and thus into major aspects of the peroxisomal import machinery in general. Achieving atomic resolution is beyond our grasp now, but we are confident that our study will provide a strong foundation and trigger future *in vivo* and *in vitro* structural and integrative studies of higher order assemblies, towards understanding key events of peroxisomal translocation.

Figures

a

b

c

d

e

f

Figure 1. Pex14p/Pex17p reconstituted in circular MSP1D1Δ4-6 nanodiscs a) Primary domain structure of yeast Pex14p and Pex17p. Red: alpha-helix; green: beta-sheet; orange: predicted membrane domain, blue: predicted coiled-coil domain; pale green: Pex13p binding domain. Binding sites to the peroxisomal receptor Pex5p are also indicated. **b)** Size exclusion chromatography of Pex14p/Pex17p in DDM (blue) and cMSP1D1Δ4-6 nanodiscs (orange). Fractions of the Pex14p/Pex17p-peak used for subsequent EM studies are indicated by arrows and the respective SDS-PAGE is shown. **c)** Subarea of a typical low-dose VPP cryo-EM micrograph of Pex14p/Pex17p reconstituted in cMSP1D1Δ4-6. Some particles are highlighted with white circles. Scale bar, 50 nm **d)** For better visualization we also denoised this particular micrograph using JANNI. It should be noted that denoised micrographs were not used during image processing. **e)** Representative reference-free 2D class averages of Pex14p/Pex17p. Scale bar, 10 nm. The inset shows one of the class averages further magnified. Densities of the nanodisc (green) and the cytosolic domains (yellow) are highlighted. **f)** End-to-end distance of the 2D class-averages.

Figure 2. Architecture of the Pex14p/Pex17p complex a) Pex14p/Pex17p density map in nanodisc. Shown are four different side views, after horizontal rotation of 90° around their longest axis, the bottom and top view. The rod and nanodisc density, are highlighted in yellow and green, respectively. **b)** Resulting map from focused 3D local refinement on the rod region after signal subtraction of the nanodisc. Insets show slices through the rod density at different heights. The two structural regions of the rod are highlighted in gray and green. The green arrowhead indicates the kinked connection between these two structural regions. Yellow arrowheads indicate further weak connections along the rod. **c)** Manual fit of identified α-helices into the electron density.

Figure 3. Molecular composition of Pex14p/Pex17p and incorporation into liposomes. a) Native MS spectra of recombinant Pex14p(M51L)/His$_6$-Pex17p. Proteins were transferred into 200 mM ammonium acetate, pH 6.7 prior to native MS analysis. Peak series assigned to series of charge states are annotated by symbols as shown in the legend. Charge state series in the lower *m/z* region indicated masses of monomeric Pex14p and *E. coli* DnaK. Above *m/z* 4700 three non-resolved series were observed, best described by species of MW 144 kDa, 263 kDa and 408 kDa. Possible stoichiometries for the three masses are 3:1, 6:1 and 9:2 of Pex14p:Pex17p as schematically depicted above the spectrum. **b)** Representative subarea of a negative stain EM micrograph of recombinant Pex14p(M51L)/His$_6$-Pex17p in DDM. White, orange and yellow dashed circles indicate monomers, dimers and trimers of the complex, respectively, explaining the different stoichiometries observed by native MS. **c-d)** Localization of the N-terminal His$_6$-tag in the recombinant His$_6$-Pex14p(M51L)/Pex17p (c) and Pex14p(M51L)/His$_6$-Pex17p (d) with

Ni-NTA-nano-gold. The topology of the gold label is also shown graphically mapped on the model of Pex14p/Pex17p. The N-termini of both proteins are clearly located closely to the nanodisc. **e)** Representative negative stain image of Pex14p/Pex17p incorporated into liposomes. **f-g**) Slices through the tomographic cryo-EM volume of a single Pex14p/Pex17p-liposome (f) and corresponding 3D segmentation (g), showing the bilayer (gray) and the characteristic Pex14p/Pex17p rods (yellow).

Figure 4. Characterization of the interactions between Pex14p and Pex17p a) Map of connections between side chains of Pex14p and Pex17p identified by XL-MS. Purified Strep$_{II}$-Pex14p/His$_6$-Pex17p complexes were cross-linked using BS3 and subjected to in-solution digestion using trypsin. **b)** Gal4p-binding domain (DB)/Pex14p fusions with progressive N- and/or C-terminal deletions of Pex14p were assayed for their interaction with Pex17p$_{167-199}$. Double transformants of PCY2 *pex14Δ* cells expressing the indicated fusion proteins were selected, and ß-galactosidase activity was determined by a filter assay using X-gal as substrate.

Three representative independent double transformants are shown. The ß-galactosidase activity shown on the right is the average of triplicate measurements for three independent transformants harboring each set of plasmids. SEM: Standard Error of Mean. U: Units **c)** Native complexes were isolated from solubilized membrane fractions from wild-type cells expressing either Pex17pTPA or Pex17p$_{1-167}$TPA eluted from IgG-Sepharose with TEV protease. An eluate of wild-type cells served as control. Equal portion of the solubilisate (left) and eluate (right) were subjected to immunoblot analysis with antibodies as indicated. **d)** Cells of wild-type-, *pex17Δ*- or *pex17Δ*-cells expressing plasmid-based Pex17p-variants as indicated were grown overnight on glucose minimal media. Subsequently,10-fold dilutions were prepared, and 2 µl of each dilution were spotted onto oleate plates, which were scored for the appearance of colonies and halo-formation. In contrast to wild-type cells, *pex17Δ* mutant was not able to utilize oleate. Expression of wild-type Pex17p as well as Pex17p$_{1-167}$ restored the growth defect of *pex17Δ* cells, whereas Pex17p$_{1-125}$ lost its ability for functional complementation.

Figure 5. Distinguishing intra- and intermolecular cross-links of Pex14p homooligomers a) Recombinant variants of Pex14p comprising its N-terminal domain (Pex14p$_{1-95}$; circle), a shorter (Pex14p$_{213-341}$; triangle) and a longer C-terminal part (Pex14p$_{119-341}$; square) were cross-linked using BS3 and separated by SDS-PAGE. Bands assigned to monomeric, dimeric and trimeric forms are marked by one, two or three symbols. **b)** Native MS of Pex14p$_{1-95}$ (top), Pex14p$_{213-341}$ (middle) and Pex14p$_{119-341}$ (bottom spectrum). Proteins were transferred into 200 mM ammonium acetate, pH 6.7, and analyzed by nano-ESI-QTOF-MS. Main charge states are indicated based on assignments of peaks series to monomeric and oligomeric complexes as indicated. The respective charge-state deconvoluted spectra are shown on the right. **c)** Schematic view of intermolecular (straight red lines) and intramolecular (curved lines) crosslinks as well as peptide modifications by loop-links (angled lines) and mono-links (flags) identified and quantified by LC-MS in monomeric (top) and oligomeric (bottom) Pex14p$_{119-341}$. Quantitative analysis further

distinguished linkages recovered equally from monomeric and oligomeric forms (grey) and monomer-specific linkages (purple).

Figure 6. Organization of the Pex14p/Pex17p complex. a) Model of the Pex14p/Pex17p complex. The topology and arrangement of the different domains are indicated, according to the results of the present study. **b)** Model for triggering peroxisomal translocation upon binding of the receptor-cargo complex to Pex14p/Pex17p at the peroxisomal membrane.

Acknowledgments We are grateful to D. Prumbaum for excellent assistance with electron microscope facilities and S. Tacke for acquisition of the tomographic tilt series. We thank Raphael-Gasper-Schoenenbruecher for his excellent support during SEC-MALS data collection and analysis, as well as fruitful discussions. We also thank Sven Fischer for experiments during the initial phase of this work, Bettina Knapp for technical assistance with LC/MS analyses as well as Carla Schmidt and Jan Commandeur (MS Vision) for discussions on native MS. C.G thanks Stefan Raunser for continuous support and I. Vetter for stimulating discussions. This work was supported by the Max Planck Society, the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) (FOR1905 to C.G., R.E, B.W). Work in the Warscheid lab was also funded by the DFG – Project-ID 278002225 – RTG 2202 and – Project-ID 403222702 – SFB 1381, the Excellence Strategy (CIBSS – EXC-2189 – Project-ID 390939984) and the Excellence Initiative of the German Federal & State Governments (EXC 294, BIOSS). Work included in this study has also been performed in partial fulfillment of the doctoral theses of P.L., T.H. and D.W..

Accession codes

The EM density map has been deposited in the Electron Microscopy Data Bank under accession code EMD-XXX. MS raw data and result files have been deposited to the ProteomeXchange Consortium via the PRIDE repository[46] and are publicly accessible from its website (http://www.ebi.ac.uk/pride) with the dataset identifier PXD016304.

Competing Interests The authors declare no competing interests

Author contributions T.H. and W.G. designed proteins, performed mutational studies and yeast-two hybrid assays. P.L., T.H., D.W., T.J. purified proteins; P.L., T.H. optimized proteins for structural studies; P.L., B.K. screened samples and collected negative stain data; P.L. collected cryo electron microscopy data; P.L. and C.G. processed, refined and analyzed cryo-EM SPA and cryoET data; D.W. optimized proteins for MS studies; D.W., J.B. and F.D. performed crosslinking MS and native MS experiments and analyzed MS data; C.G. wrote the manuscript; R.E., B.W. co-wrote the manuscript. C.G., R.E., B.W. designed the study and managed the project. All authors discussed the results and commented on the manuscript.

References

1. Braverman, N. E. *et al.* Peroxisome biogenesis disorders in the Zellweger spectrum: An overview of current diagnosis, clinical manifestations, and treatment guidelines. *Mol. Genet. Metab.* **117,** 313–321 (2016).
2. Waterham, H. R., Ferdinandusse, S. & Wanders, R. J. A. Human disorders of peroxisome metabolism and biogenesis. *Biochim. Biophys. Acta* **1863,** 922–933 (2016).
3. Lazarow, P. B. & Fujiki, Y. Biogenesis of peroxisomes. *Annu. Rev. Cell Biol.* **1,** 489–530 (1985).
4. Glover, J. R., Andrews, D. W. & Rachubinski, R. A. Saccharomyces cerevisiae peroxisomal thiolase is imported as a dimer. *Proc. Natl. Acad. Sci. U.S.A.* **91,** 10541–10545 (1994).
5. McNew, J. A. & Goodman, J. M. An oligomeric protein is imported into peroxisomes in vivo. *J. Cell Biol.* **127,** 1245–1257 (1994).
6. Hasan, S., Platta, H. W. & Erdmann, R. Import of proteins into the peroxisomal matrix. *Front Physiol* **4,** 261 (2013).
7. Purdue, P. E. & Lazarow, P. B. Peroxisome biogenesis. *Annu. Rev. Cell Dev. Biol.* **17,** 701–752 (2001).
8. Brown, L.-A. & Baker, A. Peroxisome biogenesis and the role of protein import. *J. Cell. Mol. Med.* **7,** 388–400 (2003).

9.   Girzalsky, W., Saffian, D. & Erdmann, R. Peroxisomal protein translocation. *Biochim. Biophys. Acta* **1803,** 724–731 (2010).

10. Albertini, M. *et al.* Pex14p, a peroxisomal membrane protein binding both receptors of the two PTS-dependent import pathways. *Cell* **89,** 83–92 (1997).

11. Elgersma, Y. *et al.* Analysis of the carboxyl-terminal peroxisomal targeting signal 1 in a homologous context in Saccharomyces cerevisiae. *Journal of Biological Chemistry* **271,** 26375–26382 (1996).

12. Erdmann, R. & Blobel, G. Identification of Pex13p a peroxisomal membrane receptor for the PTS1 recognition factor. *J. Cell Biol.* **135,** 111–121 (1996).

13. Gould, S. J. *et al.* Pex13p is an SH3 protein of the peroxisome membrane and a docking factor for the predominantly cytoplasmic PTs1 receptor. *J. Cell Biol.* **135,** 85–95 (1996).

14. Huhse, B. *et al.* Pex17p of Saccharomyces cerevisiae is a novel peroxin and component of the peroxisomal protein translocation machinery. *J. Cell Biol.* **140,** 49–60 (1998).

15. Meinecke, M. *et al.* The peroxisomal importomer constitutes a large and highly dynamic pore. *Nat. Cell Biol.* **12,** 273–277 (2010).

16. Meinecke, M., Bartsch, P. & Wagner, R. Peroxisomal protein import pores. *Biochim. Biophys. Acta* **1863,** 821–827 (2016).

17. Montilla-Martinez, M. *et al.* Distinct Pores for Peroxisomal Import of PTS1 and PTS2 Proteins. *Cell Rep* **13,** 2126–2134 (2015).

18. Miyata, N. & Fujiki, Y. Shuttling mechanism of peroxisome targeting signal type 1 receptor Pex5: ATP-independent import and ATP-dependent export. *Mol. Cell. Biol.* **25,** 10822–10832 (2005).

19. Platta, H. W., Grunau, S., Rosenkranz, K., Girzalsky, W. & Erdmann, R. Functional role of the AAA peroxins in dislocation of the cycling PTS1 receptor back to the cytosol. *Nat. Cell Biol.* **7,** 817–822 (2005).

20. Schliebs, W., Girzalsky, W. & Erdmann, R. Peroxisomal protein import and ERAD: variations on a common theme. *Nat. Rev. Mol. Cell Biol.* **11,** 885–890 (2010).

21. Agne, B. *et al.* Pex8p: an intraperoxisomal organizer of the peroxisomal import machinery. *Mol. Cell* **11,** 635–646 (2003).

22. Chan, A. *et al.* Pex17p-dependent assembly of Pex14p/Dyn2p-subcomplexes of the peroxisomal protein import machinery. *Eur. J. Cell Biol.* **95,** 585–597 (2016).

23. Schell-Steven, A. *et al.* Identification of a novel, intraperoxisomal pex14-binding site in pex13: association of pex13 with the docking complex is essential for peroxisomal matrix protein import. *Mol. Cell. Biol.* **25,** 3007–3018 (2005).

24. Niederhoff, K. *et al.* Yeast Pex14p possesses two functionally distinct Pex5p and one Pex7p binding sites. *Journal of Biological Chemistry* **280,** 35571–35578 (2005).

25. Azevedo, J. E. & Schliebs, W. Pex14p, more than just a docking protein. *Biochim. Biophys. Acta* **1763,** 1574–1584 (2006).

26. Itoh, R. & Fujiki, Y. Functional domains and dynamic assembly of the peroxin Pex14p, the entry site of matrix proteins. *Journal of Biological Chemistry* **281,** 10196–10205 (2006).

27. Barros-Barbosa, A. *et al.* Membrane topologies of PEX13 and PEX14

provide new insights on the mechanism of protein import into peroxisomes. *FEBS J.* **286,** 205–222 (2019).

28. Li, L. *et al.* Pex14/17, a filamentous fungus-specific peroxin, is required for the import of peroxisomal matrix proteins and full virulence of Magnaporthe oryzae. *Mol. Plant Pathol.* **18,** 1238–1252 (2017).
29. Tribet, C., Audebert, R. & Popot, J. L. Amphipols: polymers that keep membrane proteins soluble in aqueous solutions. *Proc. Natl. Acad. Sci. U.S.A.* **93,** 15047–15050 (1996).
30. Miehling, J., Goricanec, D. & Hagn, F. A Split-Intein-Based Method for the Efficient Production of Circularized Nanodiscs for Structural Studies of Membrane Proteins. *Chembiochem* **19,** 1927–1933 (2018).
31. Wagner, T. *et al.* SPHIRE-crYOLO is a fast and accurate fully automated particle picker for cryo-EM. *Communications Biology 2019 2:1* **2,** 218 (2019).
32. Moriya, T. *et al.* High-resolution Single Particle Analysis from Electron Cryo-microscopy Images Using SPHIRE. *J Vis Exp* (2017). doi:10.3791/55448
33. Gao, Y., Cao, E., Julius, D. & Cheng, Y. TRPV1 structures in nanodiscs reveal mechanisms of ligand and lipid action. *Nature* **534,** 347–351 (2016).
34. Gatsogiannis, C. *et al.* Membrane insertion of a Tc toxin in near-atomic detail. *Nature Publishing Group* (2016). doi:10.1038/nsmb.3281
35. Girzalsky, W. *et al.* Pex19p-dependent targeting of Pex17p, a peripheral component of the peroxisomal protein import machinery. *Journal of Biological Chemistry* **281,** 19417–19425 (2006).
36. Brown, L.-A. & Baker, A. Shuttles and cycles: transport of proteins into the peroxisome matrix (review). *Mol. Membr. Biol.* **25,** 363–375 (2008).
37. Brocard, C., Lametschwandtner, G., Koudelka, R. & Hartig, A. Pex14p is a member of the protein linkage map of Pex5p. *EMBO J.* **16,** 5491–5500 (1997).
38. Shimizu, N. *et al.* The peroxin Pex14p. cDNA cloning by functional complementation on a Chinese hamster ovary cell mutant, characterization, and functional analysis. *Journal of Biological Chemistry* **274,** 12593–12604 (1999).
39. Snyder, W. B. *et al.* Pex17p is required for import of both peroxisome membrane and lumenal proteins and interacts with Pex19p and the peroxisome targeting signal-receptor docking complex in Pichia pastoris. *Mol. Biol. Cell* **10,** 4005–4019 (1999).
40. Holroyd, C. & Erdmann, R. Protein translocation machineries of peroxisomes. *FEBS Lett.* **501,** 6–10 (2001).
41. Ma, C. & Subramani, S. Peroxisome matrix and membrane protein biogenesis. *IUBMB Life* **61,** 713–722 (2009).
42. Khoshouei, M., Radjainia, M., Baumeister, W. & Danev, R. Cryo-EM structure of haemoglobin at 3.2 Å determined with the Volta phase plate. *Nat Commun* **8,** 16099 (2017).
43. Erdmann, R. & Schliebs, W. Peroxisomal matrix protein import: the transient pore model. *Nat. Rev. Mol. Cell Biol.* **6,** 738–742 (2005).
44. Salomons, F. A., Kiel, J. A., Faber, K. N., Veenhuis, M. & van der Klei, I. J. Overproduction of Pex5p stimulates import of alcohol oxidase and dihydroxyacetone synthase in a Hansenula polymorpha Pex14 null

mutant. *Journal of Biological Chemistry* **275,** 12603–12611 (2000).

45. Kerssen, D. *et al.* Membrane association of the cycling peroxisome import receptor Pex5p. *Journal of Biological Chemistry* **281,** 27003–27015 (2006).

46. Perez-Riverol, Y. *et al.* The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic Acids Res.* **47,** D442–D450 (2019).

47. Källberg, M. *et al.* Template-based protein structure modeling using the RaptorX web server. *Nature Protocols* **7,** 1511–1522 (2012).

Supplementary Figures

a

b

c

d

e

Supplementary Figure 1. Partial degradation of recombinant Pex14p by cleavage at methionine 51 was prevented by the mutation M51L. (a) Strep$_{II}$-Pex14p/His$_6$-Pex17p-complex analyzed by SDS-PAGE revealed a dominant Pex14p-degradation product (Pex14p*) at an apparent molecular mass of 35 kDa. **(b)** Intensities measured by LC-MS of tryptic peptides recovered from individual gel bands of Pex14p (red line) and Pex14p* (blue line) indicate that the N-terminal part up to methionine 51 (M62 in Strep$_{II}$-Pex14p) was absent in the degradation product. The inset shows the identified peptide representing the putative N-terminus of Pex14p*. Predicted cleavage sites of trypsin are highlighted in red. **(c)** MS/MS spectrum identifying the peptide generated by cleavage N-terminal to methionine 51. Fragment ions of type y (red) and type b (blue) identifying the peptide sequence are marked. **(d)** Recombinant Pex14p, wild-type and M51L were analyzed by SDS-PAGE and Coomassie staining. Proteins in individual gel slices were subjected to in-gel digestion using trypsin and quantitative LC-MS/MS analysis. The band assigned to the degradation product (Pex14p*) was significantly decreased in the M51L compared to wild-type. **(e)** LC-MS intensities of Pex14p peptides recovered from Pex14p* gel slices amounted to 1.6% and 36.1% of the total intensity for both gel slices for M51L and wild-type, respectively.

Supplementary Figure 2. Size-exclusion chromatography and negative stain EM analysis of Pex14p/Pex17p. Gel-filtration chromatogram (left panel) and typical EM micrograph (right panel) of Pex14p/Pex17p in 0.1 % DDM **(a)**, (MSP)1D1dH5 nanodiscs (ratio 1:30) **(b)** and MSP1D1Δ4-6 circular nanodiscs (ratio 1:30) **(c)**. Dashed lines indicate fractions used for EM. Dashed circles in (b) indicate proteo-nanodiscs including multiple copies of Pex14p/Pex17p. Scale bars 100 nm.

Supplementary Figure 3. Pex14p/Pex17p dissociates after exchange of DDM with amphipols A8-35. (a) Size exclusion chromatography profile of Pex14pPex17p in amphipols. Peak I and II correspond to Pex14p and Pex17p, respectively. **(b)** SDS-PAGE of the peak fractions I and II after size exclusion chromatography. **(c)** Representative negative stain electron micrographs of the Pex14p (left panel) and Pex17p (right panel) fractions. Scale bar, 100 nm.

Supplementary Figure 4. EM analysis of Pex14p/Pex17p. (a) Single particle processing workflow for Pex14p/Pex17p structure determination. **(b)** Fourier Shell Correlation (FSC) curves between two independently refined half maps derived after standard refinement of the complete dataset and after signal subtraction of the nanodisc projections from the experimental images and subsequent local refinement (see, a). **(c)** Local resolution map of the Pex14p/Pex17p complex prior signal subtraction. **(d)** Orientation distribution of the particles used in the final refinement round prior signal subtraction.

Supplementary Figure 5. Native MS analysis of recombinant Strep_{II}-Pex14p(M51L)/His_6-Pex17p (a) MS/MS spectra for the Strep_{II}-Pex14p(M51L)/His_6-Pex17p complex, generated by selection of ions of *m/z* 6550 in the quadrupole and subsequent collisional activation by acceleration voltages of 100 V (bottom) or 120 V (top). At a lower acceleration voltage of 100 V, the series corresponding to the mass of Pex17p (blue circles) is predominant, whereas at higher energy (120 V) Pex14p became more abundant (red hexagons). The right part shows the corresponding charge-state deconvoluted spectra. **(b)** Equivalent to (a), but after selection of ions of *m/z* 8050 in the quadrupole. Similarly, at a lower collision energy (120 V), Pex17p was observed as the predominant product, whereas at 150 V Pex14p was more abundant.

Supplementary Figure 6. Chromatogram of light scattering Rayleigh ratio. The molar masses were calculated using the protein conjugate analysis. Superose 6 increase increase 10/300 column was pre equilibrated with buffer 50 mM Tris pH 8.0, 150 mM NaCl and 0.1% DDM.

Supplementary Figure 7. *In vitro* **crosslinking leads to trimers of Pex14p$_{119\text{-}341}$.** Recombinant Pex14p$_{119\text{-}341}$ was cross-linked using BS3 at increasing concentration as indicated and separated by SDS-PAGE followed by Coomassie staining.

Supplementary Figure 8. Intra- and intermolecular cross-links of Pex14p homooligomers. Following chemical crosslinking with BS3 and SDS-PAGE analysis, monomeric, dimeric and trimeric Pex14p$_{119-341}$ forms (see Figure 5a) were subjected to in-gel digestion with trypsin and analyzed by LC-MS/MS. **(a)** MS/MS spectrum identifying a monomer-specific Pex14p intraprotein crosslinked peptide pair (top) and an oligomer-specific interprotein cross-linked peptide pair (bottom). Ions matched to masses of b- and y-ions (see inset) of the longer (green) and shorter

(blue) peptides are marked. **(b)** Representative ion chromatograms showing the elution profiles of the corresponding monomer- (top) and trimer-specific (bottom) cross-linked peptide pairs. **(c)** Quantitative LC-MS analysis of cross-linked amino acid pairs (red dots), regular peptides (gray dots) and mono- (blue triangles down) or loop-linked amino acids (blue triangles up). Normalized $\log_2$ trimer-to-monomer intensity ratios were computed and plotted against their $\log_{10}$ intensities. Identified linkages are annotated with their sequence positions. Blue vertical lines indicate a 10-fold enrichment either in the trimer or monomer sample.

Supplementary Figure 9. Structure prediction for the Pex14p subunit. (a) Sequence-based secondary structure prediction for Pex14p. Red bands and green arrows represent α-helices and β-strands, respectively. The subunit can be roughly divided into four domains (see also Figure 1a): N-terminal α-helical domain (highlighted in red), transmembrane domain (TMD; highlighted in brown), coiled-coil domain (highlighted in blue) and the unstructured C-terminal domain (highlighted in gray) **(b)** Three-dimensional structure of Pex14p as predicted by RaptorX [47] (coloring similar to (a)).

Video 1. Flexibility of Pex14p/Pex17p in 2D. The arrow indicates flexible densities below the nanodisc.

Video 2. CryoEM map of the cytosolic rod. The cryoEM map of the cytosolic domains obtained after signal subtraction of the nanodisc density from the experimental images and subsequent focused 3D local refinement is rotated in order to show the overall structure. α-helixes are fitted within parallel rod-like densities.

Video 3. 3D segmentation of a Pex14p/Pex17p proteoliposome.