Deschloroclozapine: a potent and selective chemogenetic actuator enables rapid neuronal and behavioral modulations in mice and monkeys

Yuji Nagai[1,10], Naohisa Miyakawa[1,10], Hiroyuki Takuwa[1], Yukiko Hori[1], Kei Oyama[1], Bin Ji[1], Manami Takahashi[1], Xi-Ping Huang[2,8], Samuel T. Slocum[2], Jeffrey F. DiBerto[2], Yan Xiong[3], Takuya Urushihata[1], Toshiyuki Hirabayashi[1], Atsushi Fujimoto[1], Koki Mimura[1], Justin G. English[2], Jing Liu[3], Ken-ichi Inoue[4,5], Katsushi Kumata[6], Chie Seki[1], Maiko Ono[1], Masafumi Shimojo[1], Ming-Rong Zhang[6], Yutaka Tomita[7], Tetsuya Suhara[1], Masahiko Takada[4], Makoto Higuchi[1], Jian Jin[3], Bryan L. Roth[2,8,9,*], Takafumi Minamimoto[1,*]

[1]Department of Functional Brain Imaging, National Institute of Radiological Sciences, National Institutes for Quantum and Radiological Science and Technology, Chiba 263-8555 Japan

[2]Department of Pharmacology, University of North Carolina at Chapel Hill School of Medicine, Chapel Hill, NC 27514, U.S.A.

[3]Mount Sinai Center for Therapeutics Discovery, Departments of Pharmacological Sciences and Oncological Sciences, Tisch Cancer Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029 U.S.A.

[4]Systems Neuroscience Section, Primate Research Institute, Kyoto University, Inuyama, Aichi 484-8506, Japan

[5]PRESTO, Japan Science and Technology Agency, Kawaguchi, Saitama, Japan

[6]Department of Radiopharmaceuticals Development, National Institute of Radiological Sciences, National Institutes for Quantum and Radiological Science and Technology, Chiba 263-8555 Japan

[7]Depaprment of Neurology, Keio University School of Medicine, Tokyo 160-8582, Japan

[8]Division of Chemical Biology and Medicinal Chemistry, Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill, NC 27514, USA

[9]National Institute of Mental Health Psychoactive Drug Screening Program (NIMH PDSP), Department of Pharmacology, University of North Carolina at Chapel Hill Medical School, Chapel Hill, NC 27514, USA

[10] These authors contributed equally.

* Correspondence: minamimoto.takafumi@qst.go.jp (T.M.);

bryan_roth@med.unc.edu (B.L.R.)

ABSTRACT

The chemogenetic technology, Designer Receptors Exclusively Activated by Designer Drugs (DREADDs), affords reversible and remote control of cellular signaling, neuronal activity and behavior. Although the combination of muscarinic-based DREADDs with clozapine-N-oxide (CNO) has been widely used, the sluggish kinetics, metabolic liabilities, and potential for off-target effects of CNO represent areas for improvement. Here we provide a new agonist deschloroclozapine (DCZ), which displays high affinity and selectivity for muscarinic-based DREADDs. Positron emission tomography revealed that DCZ selectively bound to and occupied DREADDs in both mice and monkeys. Systemic delivery of low doses of DCZ (1 or 3 µg/kg) enhanced neuronal activity via hM_3Dq within minutes in mice and monkeys. Intramuscular injection of DCZ (100 µg/kg) reversibly induced behavioral deficits in hM_4Di-expressing monkeys. DCZ represents the most potent, selective, metabolically stable and fast-acting DREADD agonist reported with utility in both mice and non-human primates for a variety of applications.

INTRODUCTION

The chemogenetic technology known as Designer Receptors Exclusively

Activated by Designer Drugs (DREADDs) affords a minimally invasive means to

reversibly and remotely control the activity of DREADD-expressing cells by

systemic delivery of the DREADD ligand[1]. Several DREADDs exist, derived from

muscarinic or kappa-opioid receptors[1, 2] and DREADDs have been widely

adopted for neuroscience research[3]. Muscarinic receptor DREADDs are the

most widely used and can be activated by clozapine-N-oxide (CNO). When

expressed in neurons, activation of hM_3Dq induces excitation, while hM_4Di

silences neuronal activity. DREADDs, actuated by CNO, have been successfully

applied in a variety of *in vitro* and *in vivo* contexts, extending to non-human

primate studies to modify the operation of neural networks[4] and behavior[5-7].

Since CNO has modest brain permeability, relatively large systemic doses are

usually used for DREADD activation. Moreover, CNO can be metabolized to

clozapine—a potent, brain permeable DREADD agonist[1, 8]. The CNO

metabolism to clozapine was significant in rodents[8, 9] and in monkeys with

slower delivery (e.g., subcutaneous injection)[10]. In addition to agonist action on

DREADDs, clozapine is a potent ligand for multiple endogenous receptors and

transporters[11]; accordingly, CNO or clozapine administration could produce

off-target side effects that can interfere with controlled chemogenetic studies.

Alternative DREADD agonists such as compound 21 (C21) and perlapine[12, 13],

have been developed in recent years. However, they require relatively large

systemic doses to activate DREADDs *in vivo*, and at high dose may have

off-target actions at histamine and other receptors[13]. Given the broad utility and

popularity of muscarinic-based DREADDs, the development of alternative

agonists suitable for both rodents and primates—a selective, high-affinity,

metabolically stable, and brain-penetrable DREADD agonist—would represent a

substantial advance.

Here we identified a suitable DREADD agonist that we have named DCZ

[deschloroclozapine or

11-(4-methyl-1-piperazinyl)-5H-dibenzo(b,e)(1,4)diazepine] (Fig. 1a, left),

derived from our previous studies on DREADD ligands for muscarinic-based

DREADDs[12]. We were first drawn to DCZ, because while it is structurally similar

to clozapine, it had been previously reported to have substantially lower affinity

at dopaminergic (D_1 and D_2) and serotonergic (5-HT$_{2A}$ and 5-HT$_{2C}$) receptors compared with clozapine[14]. We determined that DCZ has 100-fold improved affinity and stronger agonist potency for hM$_3$Dq and hM$_4$Di relative to CNO or C21, while reducing off-target binding compared with clozapine *in vitro*. Using positron emission tomography (PET), we demonstrated that DCZ rapidly penetrates into the brain and selectively binds to DREADDs, and that DCZ doses for DREADD occupancy are 20- and 60-fold smaller than CNO and C21, respectively. Finally, we demonstrated that DCZ is capable of rapidly (<10 min post-injection) activating hM$_3$Dq and hM$_4$Di in both mice and monkeys without discernible off-target action. Thus, DCZ represents a potent and selective chemogenetic actuator for muscarinic-based DREADDs in mice and primates that is useful for a variety of *in vitro* and *in vivo* contexts with high translational potential.

RESULTS

DCZ selectively binds to DREADDs in vitro and in vivo

We first assessed the binding affinities of DCZ to hM$_3$Dq and hM$_4$Di, and

compared them to those of clozapine, CNO and C21 using radioligand

competition binding assays. DCZ inhibited [^{3}H]quinuclidinyl benzilate (QNB)

binding to hM$_3$Dq and hM$_4$Di with nanomolar affinity (inhibition constants; $^{hM3Dq}K_i$

= 6.3 nM; $^{hM4Di}K_i$ = 4.2 nM), which was comparable to clozapine ($^{hM3Dq}K_i$ = 5.9

nM; $^{hM4Di}Ki$ = 0.89), while CNO and C21 were about 100- and 50-fold weaker,

respectively (CNO, $^{hM3Dq}K_i$ = 680 nM; $^{hM4Di}K_i$ = 360 nM; C21, $^{hM3Dq}K_i$ = 850 nM;

$^{hM4Di}K_i$ = 180 nM)(Table 1). Unlike clozapine (Fig. 1c and Supplementary Table

1), DCZ had negligible affinities for a large number of tested G protein-coupled

receptors (GPCRs), ion channels or transporters (K_i's > 100 nM) and relatively

low affinities for a few endogenous receptors such as muscarinic acetylcholine

($^{hM1}K_i$ = 83 nM; $^{hM5}K_i$ = 55 nM) and serotonin receptors ($^{5-HT2A}K_i$ = 87 nM)(Fig. 1b

and Supplementary Table 1), representing at least an 8-fold selectivity for

muscarinic DREADDs based on antagonist radioligand binding affinities over its

most potent endogenous targets (also see below).

Having discovered that DCZ displays greater selectivity for DREADDs than clozapine *in vitro*, we next examined whether DCZ enters the brain and selectively binds to DREADDs *in vivo*. For these studies we performed PET with radiolabeled DCZ ([^{11}C]DCZ) in two monkeys, which had received an adeno-associated virus type 2 vector (AAV2-CMV-hM$_4$Di) injection, resulting in neuron-specific expression of hM$_4$Di[6] in the right putamen. As a control, an AAV2 vector carrying the kappa-opioid-based DREADD (AAV2-CMV-KORD) was injected into the left putamen (Fig. 1d), because KORD is orthogonal to muscarinic-based DREADDs and does not bind to DCZ (see original description of KORD[2]). Six weeks after injection, when the expression level was expected to have stabilized[6], the monkeys were scanned via PET after intravenous (i.v.) administration of a microdose of [^{11}C]DCZ (Fig. 1e). PET imaging revealed a rapid penetrance of [^{11}C]DCZ into the brain with a high radioactive accumulation in the hM$_4$Di-expressing putamen, while accumulation in the contralateral putamen was insignificant (Fig. 1f), in agreement with the high selectivity of DCZ for DREADDs *in vitro*. Low DCZ uptake was also visible in surrounding cortical areas, potentially reflecting DCZ binding to endogenous receptors with low affinities. Brain permeability and selectivity of DCZ were then directly compared

with those of clozapine and CNO, two DREADD agonists whose radiolabeled

form was available (C21 radiolabeled with [^{11}C] was not available due to its

chemical structure). Consistent with our previous study[6], PET imaging with

microdoses of [^{11}C]clozapine showed moderate uptake at the control putamen

and cortical areas (Fig. 1g, left), likely reflecting its high binding affinity for many

endogenous receptors (Fig. 1c and Supplementary Table 1), while increased

uptake at the hM$_4$Di-expressing putamen region presumably reflected clozapine

binding to hM$_4$Di (Fig. 1g, left). PET imaging with [^{11}C]CNO showed a low

radioactive signal in the brain, indicating modest brain permeability (Fig. 1i).

Indeed, the whole brain concentration of [^{11}C]CNO at 30 min was 0.14% of the

injected dose, which was >40-fold lower than that of the [^{11}C]DCZ or

[^{11}C]clozapine concentration (5.9% or 7.0%, respectively).

The dynamic regional radioactivity reflects the radioligand localization in the

vascular system and brain tissue in the initial phase, while it reflects ligand

binding to tissue in the later phase. In our study, the [^{11}C]DCZ signal at the target

region continued to increase, nearly reaching a plateau at 90 min after the

radioligand injection, whereas the signal on the contralateral side increased

moderately and then decayed rapidly (Fig. 1f, right). These regional differences

in radioactivity were abolished by pretreatment with unlabeled DCZ (1 mg/kg,

i.v.) (Supplementary Fig. 1a), and thus reflect displaceable ligand binding. We

quantified specific [^{11}C]DCZ binding to tissue using the cerebellum as a

reference region, where radioligand kinetics were unchanged by pretreatment

(Supplementary Fig. 1a). On the parametric image representing specific binding,

BP_{ND} (i.e., binding potential relative to non-displaceable uptake), a high

[^{11}C]DCZ binding region was located in the hM$_4$Di-vector injection side of the

putamen (Fig. 2a and Supplementary Fig. 2a) and corresponded to the area

where anti-hM$_4$ immune-labeling was found in post-mortem tissue (Fig. 2b and

Supplementary Fig. 2a), confirming that [^{11}C]DCZ bound to hM$_4$Di *in vivo*. In

addition, high [^{11}C]DCZ binding was also found in the projection target of the

putamen, i.e., the substantia nigra, reflecting hM$_4$Di expression at the axon

terminal that was confirmed by immunohistochemistry (Supplementary Fig. 2b).

To verify *in vivo* DCZ binding to DREADDs in mice, we performed [^{11}C]DCZ-PET

imaging with a transgenic mouse expressing hM$_4$Di under the control of

neuron-specific Thy-1 promoter[15]. [^{11}C]DCZ uptake in the striatum and

hippocampus was upregulated in the transgenic mouse compared with a

wild-type mouse (Supplementary Fig. 1b). On the parametric image, high

[^{11}C]DCZ binding was observed in the frontal and parietal cortices,

hippocampus, and striatum (Supplementary Fig. 2c, left), consistent with high

hM$_4$Di-expression as confirmed by immunofluorescence (Supplementary Fig. 2c,

right). In contrast, no discernible binding was seen in wild-type mouse brains

(Supplementary Fig. 2d) implying low off-target activity at this dose. Taken

together, our PET results suggest that DCZ rapidly enters the brain and

selectively binds to hM$_4$Di expressed in both mice and monkeys.

Systemic low doses of DCZ occupy hM4Di-DREADD in vivo

In previous chemogenetic studies, a range of systemic CNO doses was used to

obtain behavioral effects, suggesting that a required agonist dose may vary in a

manner dependent on the target neuronal population, type of DREADD and

level/localization of expression, as well as several other factors[16]. Similar to most

drugs, prior to saturation a higher agonist dose will afford a stronger

chemogenetic effect, but will also have a higher potential for off-target effects.

Although DCZ has lower levels of off-target binding than clozapine, finding the

effective and safe dose range for systemic DCZ administration is critical for

studies aiming for selective DREADD manipulation. We previously used PET

imaging to measure the relationship between CNO dose and the degree of

hM_4Di receptor occupied (i.e., occupancy), and provided a reasonable upper

limit of CNO dose (≤10 mg/kg); the doses successfully gave rise to behavioral

alterations in monkeys, while higher doses did not increase occupancy[16]. To

measure the relationship between DCZ dose and hM_4Di occupancy, we

performed [^{11}C]DCZ-PET scans 15 min after i.v. bolus injection of

non-radiolabeled DCZ (10, 30, 100 and 1,000 µg/kg) in two monkeys (Fig. 2c).

With increasing doses of non-radiolabeled DCZ, specific binding of [^{11}C]DCZ

decreased at the hM_4Di-vector injection site, while the decrease at the

contralateral control site was limited. We determined the occupancy as a

reduction of BP_{ND} at the target region over the control side relative to baseline.

The relationship between the occupancy of hM_4Di and DCZ dose was

approximated by a Hill equation with a coefficient equal to approximately 1 (Fig.

2d; Online Methods), indicating that DCZ has non-cooperative binding. The data

also suggest that a reasonable dose range of DCZ for monkey studies is ≤100

µg/kg; the occupancy increases linearly based on the logarithmic dose in this

range, while extra doses would contribute to a minimal increase in occupancy as is predicted based on standard receptor-ligand relationships.

Additionally, we performed occupancy studies with CNO and C21 and found that higher doses were required for these agonists to occupy hM$_4$Di (Fig. 2d). The dose-occupancy relationship of C21 indicated that its simple binding property (Hill coefficient, n = 1) was similar to DCZ, while that of CNO (n = 0.44) might reflect complex kinetics of CNO including the metabolism and excretion. We estimated the dose required for 50% occupancy (ED$_{50}$) for DCZ and compared it with those for CNO and C21; the ED$_{50}$ for DCZ was 25 μg/kg, which was 24- and 60-fold smaller than for CNO (630 μg/kg) and C21 (1,500 μg/kg), respectively (Fig. 2d).

We further examined the brain concentration profiles and biostability of DCZ following the systemic DCZ doses. Pharmacokinetic studies demonstrated that DCZ administration (100 μg/kg, i.v.) yielded a maximum concentration of DCZ in CSF (~10 nM) at 30 min post-injection, and this level was maintained for at least 2 h in monkeys (Fig. 3a, right). We note that the DCZ concentration was higher

than its K_i value for hM$_4$Di in antagonist radioligand binding studies (4.2 nM;

Table 1) and far below its K_i value for numerous endogenous channels,

receptors and transporters (> 50 nM; Fig. 1a). We did not detect any major

metabolites of DCZ in CSF (Fig. 3a, right) in monkeys. Taken together, these

results suggest that a low systemic dose of DCZ (i.e. 100 µg/kg) affords a

sufficient concentration of DCZ to be available for hM$_4$Di-DREADD binding *in*

vivo for at least for 2 h without the production of metabolites (Fig. 3d) in

monkeys.

We also found that DCZ was rapidly available in the mouse brain following

intraperitoneal (i.p.) administration (Fig. 3c). The detected DCZ concentration

was about 6-7 nM in CSF and ~100 nM in brain tissue at 30 min after

administration (Fig. 3c). As the free (unbound) fraction of drug is key for

determining the amount of drug that can bind to a target protein, we next

determined brain protein binding. Such analysis revealed that the fraction of

DCZ unbound to mouse brain tissue was about 7.5% (at 1 µM test

concentration), indicating that a major fraction of DCZ bound nonspecifically to

brain proteins as previously described for many CNS-active compounds[17, 18]; the

free DCZ concentration in CSF is comparable in mice (6-7 nM) and monkeys

(~10 nM). Unlike the PK profiles in monkeys, DCZ concentrations in mice rapidly

diminished and were undetectable at 2 h in either brain tissue or CSF. It is not

surprising that the PK properties of DCZ differ in mice and monkeys, as certain

species differences in the PK properties of CNS active-drugs have been well

documented[19]. Although C21, a desmethyl metabolite of DCZ (Fig. 3d), was

abundant in plasma, the amount of this metabolite in CSF was negligible (<0.2

nM, ~ 3% of DCZ at 30 min, Fig. 3c).

DCZ has greater potencies for DREADDs than previous agonists in vitro

We next examined whether DCZ can act as an agonist for muscarinic-based

DREADDs and compared its potency with several previously reported agonists.

For these studies, we first performed *in vitro* Bioluminescence Resonance

Energy Transfer (BRET) assay which directly quantified agonist-induced G

protein activation with minimal potentiation due to so-called 'receptor reserve' by

over-expression or down-stream signal amplification[20]. Using BRET-based

assays, we found that DCZ was a potent agonist for hM_3Dq with an $EC_{50} = 0.13$

(95% confidence interval, 0.09-0.21) nM, which was comparable to clozapine

[^{hM3Dq}EC$_{50}$ = 0.09 (0.06-0.15) nM], but with about 40- and 100-fold greater

potency than C21 [^{hM3Dq}EC$_{50}$ = 5.2 (3.7-7.3) nM] and CNO [^{hM3Dq}EC$_{50}$ = 15

(8.8-25) nM], respectively (Fig. 4a). DCZ was also a potent agonist for hM$_4$Di

[^{hM4Di}EC$_{50}$ = 0.081 (0.042-0.16) nM], which was comparable to clozapine

[^{hM4Di}EC$_{50}$ = 0.051 (0.027-0.097) nM], but with about 30- and 90-fold greater

potency than C21 [^{hM4Di}EC$_{50}$ = 2.6 (1.6-4.4) nM] and CNO [^{hM4Di}EC$_{50}$ = 7.3

(3.6-15) nM] (Fig. 4b), respectively.

Superior *in vitro* potency of DCZ over C21 and CNO was also demonstrated in a

Ca^{2+} mobilization assay for hM$_3$Dq[12](Supplementary Fig. 3a) and for inhibition of

cAMP accumulation for hM$_4$Di (Supplementary Fig. 3b). To further evaluate

potential off-target agonist actions of DCZ, we assessed its agonist activity at

318 endogenous GPCRs using an arrestin recruitment assay platform[21]. While

DCZ was a potent agonist for hM$_3$Dq and hM$_4$Di, it did not display significant

agonistic activity for any of the 318 tested wild-type GPCRs at <10 nM, the

typical concentration that could be reached with the dose used (Supplementary

Fig. 3c-k). Taken together, these results indicate that DCZ's agonism is

apparently selective for hM$_3$Dq and hM$_4$Di.

DCZ selectively and rapidly enhances neuronal activity via

hM3Dq-DREADD in vivo

We next examined the agonistic effect of DCZ on DREADD and its time course

in vivo using two-photon calcium imaging in mice expressing hM$_3$Dq. AAV

vectors carrying the GCaMP6s genes (AAV-DJ-rSyn-GCaMP6s) and hM$_3$Dq

construct (AAV2-CMV-hM$_3$Dq) were co-injected into the barrel cortex (Fig. 4c).

At 28 days post-injection, *in vivo* hM$_3$Dq expression was visualized by

[^{11}C]DCZ-PET as a high radioligand binding region at the injection site (Fig. 4d).

Given its high potency *in vitro*, we used DCZ at a dose of 1 µg/kg. Shortly after

i.p. injections of DCZ, transient and repetitive increases in fluorescence signals

were observed in the soma of hM$_3$Dq-expressing neurons (Figs. 4e and 4f). On

average, relative signal changes from baseline (ΔF/F) rapidly increased (τ = 3

min) and became significant at 5 min post-injection ($p < 0.01$, one-way ANOVA

with post-hoc Dunnett test). ΔF/F achieved their peak at about 10 min, plateaued

for at least 150 min, and then returned to the baseline levels at 4 h post-injection

(Fig. 4g, red; $p = 0.53$). Vehicle injection did not change the activity of the same

neuronal population (Fig. 4g, cyan). The DCZ-induced increases in fluorescent

signals were apparently mediated by hM$_3$Dq because DCZ did not alter the

signal in mice without DREADD expression (Fig. 4g, non-hM$_3$Dq, black).

Although DCZ induced strong activity changes, it did not cause long-term

apparent desensitization to hM$_3$Dq-positive neurons after recovery, since the

neurons demonstrated a reproduced chemogenetic activation by the 2nd DCZ

dose (Supplementary Fig. 4a) and a replicated response to whisker stimulation

(Supplementary Fig. 4b).

To compare the *in vivo* agonist efficacy of DCZ with those of other

ligands, we used a 100-fold higher dose (100 µg/kg, i.p.) of CNO and C21,

because they require higher doses to occupy and activate DREADDs than DCZ

(cf. Figs. 2d and 4a). Although administration of CNO and C21 increased ΔF/F of

hM$_3$Dq-expressing neurons, their peak values were less than 40% of that of DCZ

(Fig. 4g, blue and purple). Moreover, the kinetics of CNO were relatively slow (τ

= 17 min), with the upward trend lasting for 30 min post-injection and then

plateauing (Fig. 4g, blue), as observed in a previous electrophysiological study[22].

While C21 quickly increased ΔF/F (τ = 4 min), the increase became significant

only at 30 min post-injection or later (Fig. 4g, purple). Thus, DCZ rapidly and

reversibly activates hM$_3$Dq-expressing neuronal populations in mice *in vivo* with

higher efficacy than previous DREADD agonists.

Rapid and reversible chemogenetic neuronal control was also observed in
monkeys after systemic DCZ administration. A total of four monkeys were
injected with AAV2-CMV-hM$_3$Dq into the unilateral amygdala (Fig. 5a; one used
for electrophysiology and the other three for activation imaging; see below). At
48-66 days post-injection, *in vivo* hM$_3$Dq expression was visualized by
[^{11}C]DCZ-PET as a high radioligand binding region at the injection site (Fig. 5b),
which was confirmed by immunohistochemistry (Supplementary Fig. 5). We
recorded local field potential (LFP) from the hM$_3$Dq-positive area of one monkey
using a multi-site linear probe, the location of which was confirmed by CT-PET
fusion image (Fig. 5b). To the best of our knowledge, there has been no prior
report on a monkey chemogenetic study using hM$_3$Dq. Therefore, we applied the
same dose of DCZ (1 µg/kg) as in mice to avoid potential risks of excitotoxicity or
seizure due to excessive neuronal activation. In the amygdala, LFP gamma
band activity, which is known to correlate with firing of local populations of
neurons[23], increased significantly from baseline by administration of DCZ ($p <$
0.01, t-test), but not by vehicle ($p = 0.95$; Fig. 5c,d). As a control, when we

placed an electrode outside the hM$_3$Dq-positive sites, significant changes in gamma power were not observed after DCZ administration (Fig. 5d, p = 0.64). LFP-gamma power rapidly increased (τ = 4 min) from baseline and became significant at 5 min after DCZ administration, and this significant increase lasted for at least 45 min (Fig. 5e, p < 0.01, one-way ANOVA with post-hoc Dunnett test).

DCZ selectively induces hM3Dq-mediated metabolic activity

Given that the administration of DCZ affords the selective activation of neuronal populations via hM$_3$Dq, we next performed a PET study with [^{18}F]fluorodeoxyglucose (FDG) to examine whether DCZ induces dose-dependent and DREADD-selective changes in regional brain glucose metabolism, an index of brain neuronal/synaptic activation[24-26]. Three monkeys expressing hM$_3$Dq in the unilateral amygdala underwent [^{18}F]FDG PET imaging following systemic DCZ or vehicle injection (Fig. 6a-c). We found that FDG uptake at the hM$_3$Dq-positive amygdala significantly increased after injections of 1- and 3-μg/kg DCZ doses (p < 0.01, paired t-test), and this was reproducible without a reduction due to repetitive activations (Fig. 6d). The increase of FDG

uptake occurred in a dose-dependent manner at the hM$_3$Dq-positive amygdala

region, while uptake at the contralateral control area was unchanged (one-way

ANOVA, main effect of dose p < 0.001; Fig. 6d). A voxel-wise statistical analysis

further revealed that significant increase in FDG uptake after DCZ administration

(3 µg/kg, i.v.) occurred exclusively at the hM$_3$Dq-positive area (p < 0.001,

uncorrected; Fis. 6b,c). In control monkeys without hM$_3$Dq vector injection

(non-DREADD; n = 2), administration of DCZ (1 or 100 µg/kg) did not change

FDG uptake in the amygdala (one-way ANOVA, main effect of dose, p = 0.09;

Fig. 6e). A voxel-wise statistical analysis demonstrated that no significant

metabolic change was detected throughout the whole brain following a high

dose of DCZ injection (100 µg/kg) in control monkeys, confirming that off-target

effect of DCZ is undetectable *in vivo* by this measure. These results suggest that

DCZ induces a dose-dependent increase of chemogenetic neuronal excitation

as measured by metabolic change with no significantly detectible off-target

effects.

DCZ selectively induces behavioral deficits in hM4Di-expressing monkeys

Finally, we sought to modify cognitive behavior using inhibitory DREADDs and

DCZ. We targeted the prefrontal cortex (PFC), especially sectors around the

principal sulcus corresponding to Brodmann's area 46, which is responsible for

spatial working memory and executive function. To implement chemogenetic

silencing, two monkeys received multiple injections of an AAV-vector carrying

hM_4Di genes (AAV1-hSyn-hM_4Di-IRES-AcGFP) bilaterally into the dorsal and

ventral banks of the principal sulcus (Fig. 7a). [^{11}C]DCZ-PET confirmed that *in*

vivo hM_4Di expression covered the bilateral target regions (Fig. 7b). We used a

spatial delayed response task (Fig. 7c), which has been frequently employed as

a sensitive probe of spatial working memory[7, 27]. Compared with vehicle

administration, intramuscular (i.m.) administration of DCZ (100 µg/kg)

significantly impaired the performance of the delayed response task (two-way

ANOVA with treatment × delay, main effect of treatment, p < 0.0001 for both

monkeys; Fig. 7d). The impairment was more severe in the trials with longer

delays (two-way ANOVA with treatment × delay, interaction, p < 0.001 for both

monkeys), suggesting that it was attributable to loss of working memory function.

Without a screen during the delay period, DCZ did not affect the monkey's

performance, indicating that the impairment was unlikely attributable to deficits in

motor function, visual perception or general motivation (Supplementary Fig.

6a,b). The impairment of the delayed response task appeared shortly after the

administration (<10 min) and lasted for at least 2 h (t-test, $p < 0.05$ for all test

time periods; Fig. 7e), but disappeared at 24 h (two-way ANOVA with treatment

× delay, main effect of treatment, vehicle vs 24 h after DCZ, $p = 0.45$; DCZ vs 24

h after DCZ, $p < 0.0001$; Fig. 7d). The behavioral effect with DCZ was

reproducible without attenuation due to repetitive DREADD activation (Fig. 7f).

We confirmed that DCZ alone did not produce any significant effects on spatial

working memory in non-DREADD monkeys (n = 2, including one of the two

monkeys prior to vector injection, #245; $p > 0.3$; Supplementary Fig. 6c).

Additional behavioral examinations in three other non-DREADD monkeys using

a reward-size task further corroborated that significant side-effect of DCZ

injection (100 μg/kg, i.m.) was undetectable on motor or motivational functions

(Supplementary Fig. 6d,e). These results demonstrated that DCZ enables a

rapidly and reversibly-induced behavioral change through activating

muscarinic-based DREADDs without significant side effects.

DISCUSSION

Here, we report that the new DREADD agonist DCZ represents a metabolically

stable, extremely potent, highly brain-penetrable, and selective actuator for

hM_3Dq- and hM_4Di-DREADDs in both mice and monkeys. The properties of DCZ

described here demonstrate that it can be adopted as a preferred, fast-acting

chemogenetic actuator with minimal off-target actions, thereby enhancing

opportunities for investigating the causal link between neuronal activity and

behavior with high translational potential.

Several features of DCZ indicate its superiority as a chemogenetic actuator over

prior agonists. First, DCZ exhibits DREADD-selective binding both *in vitro* and *in*

vivo. Radioligand binding assays demonstrated that DCZ is selective for hM_3Dq

and hM_4Di. PET imaging has clearly demonstrated that DCZ has substantially

improved the selectivity compared with clozapine; DCZ has decreased off-target

binding, while it retains the same level of affinities to muscarinic DREADDs as

clozapine. Second, DCZ exhibits good brain concentration profiles and

biostability. PET imaging revealed that systemically administered DCZ occupied

hM_4Di-DREADDs with a similar occupancy to CNO or C21 at ~20- or 60-fold

smaller doses, respectively. Pharmacokinetic studies confirmed that DCZ is rapidly accumulated in mouse brains and monkey CSF, while its metabolites are negligible. Third, DCZ has high potency for muscarinic DREADDs. *In vitro* assays revealed that hM_3Dq could be activated by a substantially lower concentration of DCZ, as compared to CNO and C21. DCZ did not display significant agonist potencies for any of the 318 off-target GPCRs tested; binding assays at a number of GPCRs, channels and transporters screened by the National Institute of Mental Health Psychoactive Drug Screening Program showed DCZ has modest affinity (55 – 87 nM) for a few GPCRs, low affinity (100-1000 nM) for several GPCRs and minimal activity (>1,000 nM) at most CNS targets tested. In short, DCZ has a higher selectivity for DREADDs than clozapine and is capable of occupying and activating DREADDs at a considerably smaller dose than CNO and C21, thus offering a greater effective window than the prior DREADD ligands.

CNO has repeatedly been shown to have relatively slow kinetics for neuronal activation via hM_3Dq; following systemic administration, increases in activity began at around 5-10 min and reached a maximum at 45 min or later[22]. Because

bath application of CNO immediately increased the firing rate of

hM$_3$Dq-expressing neurons in slice preparations[22], the relatively slow activation

appears to reflect the slow kinetics of CNO uptake in the brain. Moreover, the

major effect on DREADDs is exerted by its metabolite clozapine, rather than

CNO itself [8], where the metabolism is significant in rodents[8, 9] and in monkeys

with slower delivery (e.g., subcutaneous injection[10]). Although the concentration

of metabolized clozapine is minimal in monkeys with rapid intravenous delivery

of CNO[6, 28], the brain uptake of CNO is considerably low (cf. Fig. 1i). In contrast,

DCZ rapidly penetrates into the brain and induces a rapid onset of excitability of

hM$_3$Dq-positive neuronal populations, as demonstrated by two-photon calcium

imaging in mice and by electrophysiological recording in monkeys (cf. Figs. 4

and 5). Thus, DCZ combined with muscarinic-based DREADDs would provide a

superior platform for examining the effects of rapid and sustainable

chemogenetic neuronal modulation. For example, single-unit or *in vivo*

patch-clamp recording studies, where long-term (30-60 min) maintenance is

technically demanding, would greatly benefit from the rapid action of the

DCZ-DREADD system. Such a rapid chemogenetic action would also be

valuable for future therapeutic applications, for instance, on-demand seizure

attenuation[29].

The present results indicate that DCZ can also be utilized for *in vivo* neuronal

silencing by activating hM$_4$Di, an inhibitory DREADD. Our data suggest that a

reasonable upper limit of DCZ dose is 100 µg/kg (Fig. 2d), which allows a

sufficient concentration of DCZ available for hM$_4$Di-DREADD binding *in vivo* for

at least 2 h in monkeys. Indeed, a 100 µg/kg dose of DCZ rapidly (<10 min) and

reversibly induced spatial working memory deficits in monkeys expressing hM$_4$Di

in the PFC (Fig. 7). Given the high agonist potency of DCZ (Fig. 4b), smaller

DCZ doses would induce similar or detectable behavioral deficits.

One of the technical difficulties in chemogenetic manipulations, especially for

large animals, is the delivery of transgene specifically into a target neuronal

population as well as the accomplishment of stable transgene expression

throughout the experimental period. Previous studies have shown that PET

imaging with [^{11}C]clozapine or [^{11}C]CNO can visualize hM$_4$Di expression in living

mice and monkeys, providing effective tools for non-invasive monitoring of gene

expression exemplified by studies on gene delivery or neuronal differentiation of

progenitors[6, 15]. However, because these ligands display poor DREADD

selectivity or brain permeability, the application of DREADD-PET in monkeys

was limited to visualization of hM$_4$Di expression in the striatum. The present

study has demonstrated that [^{11}C]DCZ is an exceedingly useful PET ligand for

visualizing both hM$_3$Dq and hM$_4$Di expressions in cortical and subcortical areas

of mice and monkeys, with improved specificity compared with [^{11}C]clozapine

(Fig. 1). The monitoring of transgene expression is beneficial for long-term

behavioral studies and is of great advantage for conducting successful

experiments (cf. Fig. 7). In addition to expression monitoring, PET imaging has

allowed the elucidation of a relationship between a pharmacologically effective

dose of DREADD agonists and DREADD occupancy *in vivo* (Fig. 2d). Moreover,

it has also been demonstrated in our study that [^{11}C]DCZ-PET enables us to

visualize DREADDs expression at axon terminal sites (Supplementary Fig. 2),

providing a powerful means for mapping projection areas for pathway-selective

activity manipulation[30], which can be induced by local infusion of DREADD

agonist; DCZ may be suitable because of its high selectivity and efficacy.

Although PET imaging data have implied that DCZ has off-target binding in

cortical areas possibly to endogenous receptors with low affinities (Fig. 1f and

Supplementary Table 1), the agonist potency of DCZ at DREADDs is in the

sub-nM range and, therefore, when used at a sufficient dose, occupancy on

off-targets (~50 nM or higher) can be predicted to be minimal. Indeed, our data

further suggest that systemic administration of DCZ at 100 µg/kg or smaller

doses would not induce discernible off-target effects on neuronal activity or

behavior in monkeys not expressing the muscarinic-based DREADDs (cf. Fig. 6

and Supplementary Fig. 6). Since the variability of CSF concentration across

subjects and the amount of metabolites for DCZ were much smaller than those

for CNO (Fig. 3a)[6], the potential baseline sensitivity to DCZ would be minimal.

Therefore, DCZ provides a clear advantage over prior agonists due to increasing

reliability by minimizing concerns about potential off-target effects. However, we

cannot completely rule out unanticipated off-target effects because it is currently

impossible to measure the activity of DCZ against every potential endogenous

target. Therefore, as with all other neuromodulation technologies, control

experiments—here using non-DREADD animals—are recommended. In mice,

control experiments are essential since the current study did not examine an

off-target action of DCZ at a high dose where C21, a DCZ metabolite, was

detected at very low concentrations in CSF (<0.2 nM; Fig. 3c). Another caveat

for mouse experiments is that the kinetics of DCZ suggest a relatively shorter

duration of chemogenetic action. Future study will be required to determine

appropriate DCZ doses and the effective time window for mouse chemogenetic

experiments.

In conclusion, the characteristics of DCZ described here—its high selectivity,

high brain-penetrability, and biostability—facilitate the rapid and selective

modulations of neuronal activity and behavior with muscarinic-based DREADDs

in living animals. Given the potential drawbacks of prior DREADD agonists, DCZ

will provide clear benefits for many users of muscarinic DREADDs, with

increasing reliability by removing concerns about potential off-target responses.

Online Methods

Subjects

All experimental procedures involving animals were carried out in accordance with the Guide for the Care and Use of Laboratory Animals (National Research Council of the US National Academy of Sciences) and were approved by the Animal Ethics Committee of the National Institute of Radiological Sciences.

A total of 16 macaque monkeys [7 Rhesus (*Macaca mulatta*), and 9 Japanese monkeys (*Macaca fuscata*); 11 males, 5 females; 2.8-8.0 kg; age 3-10 years] were used (a summary of subjects used in the experiments described in Supplementary Table 2). The monkeys were kept in individual primate cages in an air-conditioned room. A standard diet, supplementary fruits/vegetables and a tablet of vitamin C (200 mg) were provided daily.

Adult hM$_4$Di transgenic mice with C57BL/6j background[15](male, age >12 weeks) and age-matched non-Tg littermates were used for PET scans and histochemical analysis. Wild-type C57BL/6j mice (male, age >12 weeks, Japan SLC Inc., Hamamatsu, Japan) were used in two-photon microscopy, PET scan and pharmacokinetics experiments. All mice were maintained in a 12-h light/dark cycle with ad libitum access to standard diet and water.

Viral vector production

AAV2 (AAV2-CMV-hM$_4$Di, AAV2-CMV-KORD, AAV2-CMV-hM$_3$Dq,

AAV2-CMV-AcGFP; Figs. 1d, 4c, 5a) and AAV1

(AAV1-hSyn-hM$_4$Di-IRES-AcGFP; Fig. 7a) vectors were produced by helper-free

triple transfection procedure, which was purified by affinity chromatography (GE

Healthcare, Chicago, USA). Viral titer was determined by quantitative PCR using

Taq-Man technology (Life Technologies, Waltham, USA). Transfer plasmid was

constructed by inserting cDNA fragment and WPRE sequence into an AAV

backbone plasmid (pAAV-CMV, Stratagene, San Diego, USA). For production of

an AAV-DJ vector (AAV-DJ-rSyn-GCaMP6s), a transfer plasmid containing rat

Synapsin promoter and cDNA encoding GCaMP6s (Addgene plasmid #40753)

was assembled and transfected with helper-free DJ plasmids (Cell Biolabs, San

Diego, USA). Viral particles were purified by HiTrap heparin column (GE

Healthcare). Viral titer was determined by AAVpro® Titration kit ver2 (TaKaRa).

Surgical procedures and viral vector injections

In monkeys, surgeries were performed under aseptic conditions in a fully

equipped operating suite. We monitored body temperature, heart rate, SpO_2 and

tidal CO_2 throughout all surgical procedures. Monkeys were immobilized by

intramuscular (i.m.) injection of ketamine (5-10 mg/kg) and xylazine (0.2-0.5

mg/kg) and intubated with an endotracheal tube. Anesthesia was maintained

with isoflurane (1-3%, to effect). Prior to surgery in monkeys, magnetic

resonance (MR) imaging (7 tesla 400mm/SS system, NIRS/KOBELCO/Brucker)

and X-ray computed tomography (CT) scans (Accuitomo170, J. MORITA CO.,

Kyoto, Japan) were performed under anesthesia (continuous infusion of propofol

0.2-0.6 mg/kg/min, i.v.). Overlay MR and CT images were created using

PMOD® image analysis software (PMOD Technologies Ltd, Zurich, Switzerland)

to estimate stereotaxic coordinates of target brain structures.

   For subcortical injections, monkeys underwent a surgical procedure to open

burr holes (~8 mm diameter) for the injection needle. Viruses were

pressure-injected by 10-μL Hamilton syringe (Model 1701 RN, Hamilton

Company, Reno, USA) with a 30-gauge injection needle and a fused silica

capillary (450 μm OD) to create a 'step' about 500 μm away from the needle tip

to minimize backflow. The Hamilton syringe was mounted into a motorized

microinjector (Legato130, KD Scientific or UMP3T-2, WPI, Sarasota, USA) that

was held by manipulator (Model 1460, David Kopf, Ltd., Tujunga, USA) on the

stereotaxic frame. After the dura mater was opened about 3 mm, the injection

needle was inserted into the brain and slowly moved down 2 mm beyond the

target and then kept stationary for 5 min, after which it was pulled up to the target

location. The injection speed was set at 0.25-0.5 µL/min. After each injection, the

needle remained *in situ* for 15 min to minimize backflow along the needle. Two

monkeys (#209 and #212) had co-injections of AAV vectors (total 6 µL; 3 µL × 2

different depth in a track) carrying hM$_4$Di construct (AAV-CMV-hM$_4$Di, #209: 1.8

× 10e^{13} and #212: 2.6 × 10e^{13} particles/mL) and AcGFP genes

(AAV-CMV-AcGFP, 0.7 × 10e^{13} particles/mL) in one side of the putamen and

co-injections of AAV vectors (total 6 µL; 3 µL × 2 different depth in a track)

carrying kappa-opioid based DREADD construct (AAV-CMV-KORD; 1.8 × 10e^{12}

particles/mL) and AcGFP (AAV-CMV-AcGFP, 0.7 × 10e^{13} particles/mL) in the

other side. Four monkeys (#214, #215, #233, #236) had co-injections of AAV

vectors (total 6 µL; 3 µL × 2 tracks, 2 mm apart rostrocaudally) carrying hM$_3$Dq

construct (AAV-CMV-hM$_3$Dq; 1.2 × 10e^{13} particles/mL) and AcGFP genes

(AAV-CMV-AcGFP, 0.7 × 10e^{13} particles/mL) into the left (#214 and #215) or

right (#233, #236) amygdala.

Two monkeys (#229, #245) were injected AAV-hSyn-hM$_4$Di-IRES-AcGFP (3.8

× 10e^{13} particles/mL) into the bilateral prefrontal cortex (Brodmann's area 46).

After retracting the skin and galea, the frontal cortex was exposed by removing a

bone flap and reflecting the dura mater. Handheld injections were made under

visual guidance through an operating microscope (Leica M220, Leica

Microsystems GmbH, Wetzlar, Germany), with care taken to place the beveled

tip of the Hamilton syringe containing the viral vector at an oblique angle to the

brain surface. One person inserted the needle into the intended area of injection

and another person pressed the plunger to expel approximately 1 µL. Nine

tracks were injected in each hemisphere; one was located 1 mm posterior to the

caudal tip of the principal sulcus, and the others were located along the dorsal (4

tracks) and ventral (4 tracks) bank of the principal sulcus posterior to the rostral

tip of the ascending limb of the arcuate sulcus (Fig. 7a). Viral vectors were

injected at 3 to 5 µL per track depending on the depth. Total amounts of viral

aliquots injected into the right and left hemispheres were 35 and 37 µL for #229,

and 44 and 40 µL for #245, respectively.

In mice, the animals were anesthetized with a mixture of air, oxygen, and

isoflurane (3–5% for induction and 2% for surgery) via a facemask, and a cranial

window (3-4 mm in diameter) was placed over the left somatosensory cortex,

centered at 1.8 mm caudal and 2.5 mm lateral from the bregma, according to the

'Seylaz-Tomita method'[31]. On the day of cranial window surgery, AAV vectors

carrying GCaMP6 genes (AAV-DJ-rSyn-GCaMP6, $3.5 \times 10e^{11}$ particles/mL) and

hM_3Dq construct (AAV2-CMV-hM_3Dq, $1.5 \times 10e^{13}$ particles/mL) were co-injected

into the barrel cortex using glass needles. A custom metal plate was affixed to

the skull with a 7-mm-diameter hole centered over the cranial window. The

method for preparing the chronic cranial window was previously reported in

detail[32].

Radioligand competition binding assays

Radioligand binding assays with membrane preparations to determine binding

affinity were carried out by the National Institute of Mental Health's Psychoactive

Drug Screening Program (NIMH PDSP) (https://pdsp.unc.edu/). Detailed assay

protocols are available at the NIMH PDSP website

(http://pdspdb.unc.edu/pdspWeb/?site=assays).   The NIMH PDSP is directed

by Bryan L Roth, MD, Ph.D., the University of North Carolina at Chapel Hill,

North Carolina, and Program Officer Jamie Driscoll at NIMH, Bethesda, USA.

Bioluminescence Resonance Energy (BRET) experiments

HEK293T cells split into 10-cm plates and maintained in Dulbecco's Modified

Eagle Medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and

1% penicillin-streptomycin (pen-strep) were co-transfected with 1 µg each of

hM_3D or hM_4D and Gαq-RLuc8 or Gαi1-RLuc8, respectively, and 1 µg each of

Gβ1 and Gγ2-GFP2 using TransIT-2020 (Mirus) as transfection reagent. For

negative controls, 1 µg of pcDNA was transfected in place of hM_3D or hM_4D.

After at least 12 h, cells were plated in poly-D-lysine-coated white 96-well

microplates (Greiner) in 100 µL of DMEM supplemented with 1% dialyzed FBS

and 1% pen-strep at a density of 50,000 cells per well. After at least additional

12 h, media were aspirated and replaced with 60 µL of assay buffer (1x Hanks'

Balance Salt Solution, 20 mM HEPES, pH 7.40). Next, 10 µL of 50 µM

coelenterazine 400a (NanoLight Technology) were added. After a 5-min

incubation, 30 µL of 3x drug (in assay buffer containing 0.3 mg/mL ascorbic acid

and 0.3% bovine serum album) were added. After an additional 5-min

incubation, plates were read for luminescence with a Mithras LB 940 mutimodal

microplate reader (Berthold Technologies). Data were analyzed by GraphPad

Prism 7.0 using the built-in dose-response function, and were normalized to the

responses produced by CNO.

In vitro cAMP assays

G_i-mediated inhibition of cAMP production assays were performed in transiently

transfected HEK293T cells. Brieftly, HEK293T cells, transfected overnight with

GloSensor plasmid (Promega) and receptor DNAs, were plated (10-15,000

cells/40 µL per well) in poly-L-lysine-coated white 384-well clear-bottom cell

culture plates in DMEM with 1% dialyzed FBS. After 16-20 h, cells were removed

from medium and stimulated with ligands prepared in HBSS (Hank's balanced

salt solution), 20 mM HEPES, 0.1% BSA, pH 7.4 for 15 min, followed by 0.1 µM

isoproterenol (final concentration) in GloSensor reagent. Luminescence was

read on a Wallac TriLux Microbeta counter (PerkinElmer). Results were

normalized to the CNO activity and analyzed using the built-in dose-response

function in GraphPad Prism 7.0.

In vitro Ca2+ mobilization (FLIPR) assays

HEK293 cells stably expressing hM_3Dq receptors were used for Gq-mediated

calcium mobilization assays. Assays were performed according to published

procedures[12]. More detailed assay protocols are available at the NIMH PDSP

website (http://pdspdb.unc.edu/pdspWeb/?site=assays).

GPCRome screening (PRESTO-Tango) assays

Potential agonist activity at human GPCRome was measured using the

PRESTO-Tango assay as published[21].

Brain protein binding assays

The unbound fraction of DCZ in brain tissue was measured by an equilibrium

dialysis method. PBS solutions containing mouse brain homogenates (20%) and

DCZ (final concentrations of 10, 100 or 1,000 nM) were added in 96-well plate

(Equilibrium Dialyzer MW10K; HARVARD APPARATUS), followed by

equilibrium dialysis (22 h at 37°C). A certain amount of methanol solution

containing sulfaphenazole was added as an internal standard to the separated

filtrate, brain homogenate fractions and pre-dialyzed samples, and centrifuged

(1,700g, 10 min). The supernatant fractions were applied to LC-MS/MS to measure DCZ concentration. The average fraction of unbound brain tissue was calculated for each concentration as reported in the previous publication[33].

Drug administration

DCZ (HY-42110, MedChemExpress or systhesized in house, see below) was dissolved in 1-2% of dimethyl sulfoxide (DMSO) in saline to a final volume of 0.1 mL/kg. CNO (Toronto Research, North York, Canada) was dissolved in 2.5% of DMSO in saline to a final volume of 0.1 mL/kg. C21 (Compound 21, TOCRIS, Bristol, UK) was dissolved in 2% of DMSO in distilled water to a final volume of 0.1 mL/kg. For plasma/CSF analysis, a 23-gauge catheter was placed in the saphenous vein or the spinal canal for acute sampling while the monkey was under ketamine and xylazine anesthesia. DCZ (100 µg/kg) was administered at a rate of 0.2 mL/s intravenously via catheter. For PET blocking and occupancy studies, DCZ (10, 30, 100 or 1,000 µg/kg), CNO solution (100, 300, 1,000, 3,000 or 10,000 µg/kg) or C21 (300, 1,000 or 6,000 µg/kg) was administered intravenously via a saphenous vein catheter 1-15 min before PET imaging. For FDG-PET study, DCZ (1, 3 or 100 µg/kg) solution or vehicle was administered

intravenously 1 min before PET imaging. Fresh solutions were prepared on the

day of usage.

Synthesis of DCZ

1-methylpiperazine (0.33 mL, 2.97 mmol) was added to a solution of

11-chloro-5*H*-dibenzo[*b,e*][1,4]diazepine (0.22 g, 0.96 mmol)[12] in toluene (5 mL).

The resulting solution was refluxed for 2 h. After cooling down to room

temperature, the reaction was concentrated. The resulting residue was purified

by silica gel flash column chromatography with 0–10% MeOH in CH_2Cl_2 to give

the desired product (0.20 g, yield 70%). [1]H NMR (800 MHz, CD3OD) δ7.69 –

7.64 (m, 2H), 7.41 (d, J = 7.9 Hz, 1H), 7.33 – 7.29 (m, 2H), 7.27 (t, *J* = 7.6 Hz,

1H), 7.19 – 7.13 (m, 2H), 4.40 – 3.80 (br, 4H), 3.80 – 3.50 (br, 4H), 3.06 (s, 3H).

HRMS calcd. for $C_{18}H_{21}N_4$: 293.1761; found: 293.1720 $[M + H]^+$.

Radiosynthesis

$[^{11}C]$DCZ was produced as follows. An automated multi-purpose synthesizer

developed in house was used for the present radiosynthesis. The

cyclotron-produced $[^{11}C]CO_2$ was converted to $[^{11}C]CH_3I$, which was distilled and

sent through a silver trifluoromethanesulfonate glass tube under N_2 gas flow to yield [^{11}C]methyl trifluoromethanesulfonate ([^{11}C]CH$_3$OTf). [^{11}C]CH$_3$OTf was then introduced to a reaction vial containing desmethyl precursor (C21, 0.2 mg) in dichloromethane (0.3 mL) under room temperature. The reaction mixture was kept at room temperature for 5 min. HPLC purification was completed on an X-Bridge C$_{18}$ column (10 mm i.d. × 250 mm, Waters, Milford, USA) using CH$_3$CN/H$_2$O/Et$_3$N (40/60/0.1%) at 5.0 mL/min. The radioactive fraction corresponding to [^{11}C]DCZ (t_R: 9.5 min) was collected and formulated to obtain an injectable solution. Synthetic time was about 40 min from the end of bombardment with an averaged radiochemical yield (decay-corrected) of 76.2% based on [^{11}C]CO$_2$ (n = 16). Radiochemical purity was assayed by analytical HPLC (column: CAPCELL PAK C18 UG120 S5, 4.6mmID, 150mm length, Shiseido, Tokyo, Japan) UV at 254 nm; mobile phase: CH$_3$CN/H$_2$O = 40/60 (Et$_3$N 0.1%). Radiochemical purity and molar activity of [^{11}C]DCZ were >98% and 130 ± 45 GBq /μmol (n = 16), respectively.

[^{11}C]clozapine was radiosynthesized from desmethylclozapine by ^{11}C-methylation using ^{11}C-methyl triflate based on the previously described protocol[34], and its radiochemical purity and specific radioactivity at the end of

synthesis exceeded 95% and 37 GBq/μmol, respectively. [^{18}F]FDG was

purchased from Nihon Medi-Physics Co., Ltd. (Tokyo, Japan).

PET imaging

PET scans were performed using microPET Focus 220 scanner (Siemens

Medical Solutions USA, Malvern, USA). Anesthesia was performed as follows: a

mouse was anesthetized with 1-3% isoflurane and a monkey was immobilized

by intramuscular injection of ketamine (5-10 mg/kg) and xylazine (0.2-0.5 mg/kg)

and then maintained in anesthetized condition with isoflurane (1-3%) during all

PET procedures. Transmission scans were performed for about 20 min with a

Ge-68 source for monkey scans. Emission scans were acquired in 3D list mode

with an energy window of 350–750 keV after intravenous bolus injection of

[^{11}C]DCZ (31.9-34.7 MBq for mice and 276.9-386.9 MBq for monkeys),

[^{11}C]clozapine (287.4-353.8 MBq), [^{11}C]CNO (354.6 MBq) or [^{18}F]FDG

(157.6-358.5 MBq). Emission data acquisition lasted 90 min for [^{11}C]DCZ,

[^{11}C]clozapine and [^{11}C]CNO scans and 120 min for [^{18}F]FDG scans. All

list-mode data were sorted into three-dimensional sinograms, which were then

Fourier-rebinned into two-dimensional sinograms (frames × minutes: 5 × 1, 5 ×

2, 5 × 3, and 12 × 5 for [11C]DCZ and [11C]clozapine, 24 × 5 for [18F]FDG).

Images were thereafter reconstructed with filtered back-projection using a

Hanning filter cut-off at a Nyquist frequency (0.5 mm^{-1}). Standardized uptake

value (SUV) was calculated using PMOD® image analysis software as the

regional concentration of radioactivity averaged across the specific time window

after injection of the radioligand. Volumes of interest (VOIs) were placed using

PMOD® image analysis software with reference to the MR image of individual

monkeys or the mouse brain template generated as described previously[15]. In

FDG studies with hM$_3$Dq-expressing monkeys, VOI for the hM$_3$Dq-positive

region was defined as the area where the BP$_{ND}$ value of [11C]DCZ was higher

than 1.0, while that of the control region was placed at the corresponding

contralateral side.

Two-photon laser-scanning microscopy

Awake mice were placed on a custom-made apparatus, and real-time imaging

was conducted by two-photon microscopy (TCS-SP5 MP, Leica Microsystems

GmbH) with an excitation wavelength of 900 nm. An emission signal was

separated by a beam splitter (560/10 nm) and simultaneously detected with a

band-pass filter for SR101 (610/75 nm) and GCaMP6 (525/50 nm). A single

image plane consisted of 512 × 512 pixels, and in-plane pixel size was 0.9 μm ×

0.9 μm. The methods for functional imaging using two-photon microscopy have

been reported in detail[35]. Briefly, continuous image capturing was conducted for

neurons on the surface over the barrel cortex at a rate of 0.25 s per frame for 60

s with a 512 × 512 pixel field of view. Percentage change in green fluorescence

of neurons was manually measured offline with LAS AF software (Leica

Microsystems GmbH). Sensory-evoked neuronal excitation was examined by

means of whisker stimulation as reported previously[35]. Air puff stimulations (15

psi, 50 ms pulse width, 10 s duration, 10 Hz) were given to whiskers on the

contralateral side of scanning 4 times in a day.

Electrophysiology

Electrophysiological recordings were performed while the monkey was seated in

a monkey-chair with its head fixed via a headpost, and maintained under an

anesthetized condition with intravenous infusion of propofol (20 mg/kg/h). Local

field potential (LFP) was recorded using a multi-contact linear probe with twelve

recording contacts separated by 150 μm (Axial Array, FHC, Bowdoin, USA). The

probe was inserted into the brain with a hydraulic micromanipulator (MO-97A,

Narishige, Tokyo, Japan) guided with a stainless steel tube and an implanted

recording chamber system (Crist Instruments, Hagerstown, USA). Chamber

grids were either purchased from Crist Instruments or custom-made with a 3D

printer (Object260 Connex3, Stratasys, Eden Prairie, USA). On every recording

session, the monkey was CT-scanned before the recording probe was removed.

Electrode location relative to the hM_3Dq expression was confirmed by aligning

the CT image with the pre-acquired PET image on PMOD®. The probe was

connected to a multichannel acquisition system (System3, Tucker-Davis

Technologies (TDT), Alachua, USA) running on a Windows PC. For LFP, the

signal was band-pass filtered between 1.5 Hz and 500 Hz, and sampled at 3

kHz. LFP signals were put to further analysis only when multi-unit waveforms

were qualitatively observed in high frequency band (i.e., 400 Hz – 5 kHz) at the

corresponding recording sites. The spectral component of the LFP signal was

analyzed with MATLAB (MathWorks, Natick, MA, USA). Spectral power of each

frequency was computed using a short FFT algorithm within a 60-s window

sliding in 30-s steps. Change in the gamma band power (40 Hz) and its change

relative to that in a 3- or 10-min pre-injection period was quantified. The

spectrogram was smoothed with a Hanning window with 4.3-min half-maximum width for display reasons.

Behavioral testing

Three monkeys (#226, #229, #245) were tested with a spatial delayed response task (Fig. 7c). The protocol was based on previous studies using the Wisconsin general testing apparatus[36, 37]. Behavioral testing was conducted in a sound-attenuated room. Monkeys were seated in a monkey chair from which they could reach out one hand and take food to their mouths. A wooden table with two food-wells was placed in front of the monkeys, and a screen was placed between the monkeys and the table. First, a piece of food reward (raisin or half peanut) was placed in one of the two food-wells, and then both wells were covered with wooden plates. Then, the screen was placed for 0.5, 5 or 10 s, which served as delay periods (0.5-s delay was not introduced for non-DREADD monkeys). The position of the baited well (left or right) was determined pseudo-randomly. After the delay period, the screen was removed and the monkeys were allowed to select either food-well to get the food. The monkeys were allowed to take the food if they reached for the correct food-well and

removed the cover plate. The inter-trial interval was set at 10-15 s. A daily

session lasted about one hour, and consisted of 3-4 blocks of 20-30 trials, which

were interleaved by a 5-min rest period. The behavioral testing began

immediately or 1 h after an i.m. administration of either vehicle (2% DMSO in

saline) or DCZ (100 µg/kg), and was also conducted on the next day (24 h later)

of the DCZ administration. One of the monkeys injected with AAV-hM$_4$Di in the

bilateral PFC (#229) was also tested in a non-memory control task, which was

almost the same as the delayed response task except that the screen was not

placed during the delay period and that the 0.5-s delay was not introduced

(Supplementary Fig. 6a).

Three monkeys without AAV injections (non-DREADD; #224, #228, #230)

were tested with a reward-size task (Supplementary Fig. 6d) using the same

protocol as applied in a previous study[38]. The behavioral testing began 10 min

after an i.m. administration of either vehicle (2% DMSO in saline) or DCZ (100

µg/kg).

Histology and immunostaining

The monkeys were deeply anesthetized with an overdose of sodium

pentobarbital (80 mg/kg, i.v.) and transcardially perfused with saline at 4°C,

followed by 4% paraformaldehyde in 0.1 M phosphate buffered saline (PBS), pH

7.4. The brains were removed from the skull, postfixed in the same fresh fixative

overnight, saturated with 30% sucrose in phosphate buffer (PB) at 4°C, and then

cut serially into 50-µm-thick sections on a freezing microtome. For visualization

of immunoreactive signals of hM$_4$Di, a series of every 6th section was immersed

in 1% skim milk for 1 h at room temperature and incubated overnight at 4°C with

rabbit anti-M$_4$ polyclonal antibody (1:500; H-175, Santa Cruz Biotechnology,

Dallas, USA) in PBS containing 0.1% Triton X-100 and 1% normal goat serum

for 2 days at 4°C. The sections were then incubated in the same fresh medium

containing biotinylated goat anti-rabbit IgG antibody (1:1,000; Jackson

ImmunoResearch, West Grove, PA, USA) for 2 h at room temperature, followed

by avidin-biotin-peroxidase complex (ABC Elite, Vector Laboratories,

Burlingame, CA, USA) for 2 h at room temperature. For visualization of the

antigen, the sections were reacted in 0.05 M Tris-HCl buffer (pH 7.6) containing

0.04% diaminobenzidine (DAB), 0.04% NiCl$_2$, and 0.003% H$_2$O$_2$. The sections

were mounted on gelatin-coated glass slides, air-dried, and cover-slipped. A part

of other sections was Nissl-stained with 1% Cresyl violet. The same protocol

was used for visualization of immunoreactive signals of hM_3Dq with rabbit

anti-M_3 polyclonal antibody (1:200; HPA024106, Atlas Antibodies, Stockholm,

Sweden).

The mice were deeply anesthetized with sodium pentobarbital and then

transcardially perfused with PBS. Brain tissues were removed and fixed with 4%

paraformaldehyde in PB overnight, followed by cryoprotection with 30% sucrose

in PB. 10-µm-thick frozen sections were generated in a cryostat (HM560; Carl

Zeiss, Oberkochen, Germany). Visualization of immunoreactive signals of hM_4Di

was performed using an anti-M_4 antibody (H-175) and fluorophore-conjugated

secondary antibodies (Invitrogen, Carlsbad, CA, USA) as described previously[15].

Images of sections were digitally captured using an optical microscope

equipped with a high-grade charge-coupled device (CCD) camera (Biorevo,

Keyence, Osaka, Japan).

Pharmacokinetics analysis

Four monkeys and 15 mice were used to assess the concentration of DCZ and

its metabolites in plasma, cerebrospinal fluid (CSF) or brain. In monkeys, blood

and CSF were collected at 15, 30, 45, 60, 75, 90, 105, and 120 min after DCZ

administration (100 µg/kg, i.v. or i.m.) under ketamine and xylazine anesthesia.

In mice, after intraperitoneal injection of DCZ (100 µg/kg), blood and brain

samples were collected at 5, 30, 60, 120 min or 24 h, and CSF was collected at

30 and 120 min. Mouse blood was collected by heparinized syringes via heart

under isoflurane anesthesia followed by centrifugation at 10,000 g for 5 min to

obtain the plasma samples, and the brains were removed and frozen by liquid

nitrogen immediately after blood collection. All samples were stocked at -80°C

until analysis.

The protocols for sample pretreatment of CSF and plasma were described

previously[6]. Mouse brains were homogenized in 2-brain-weight water. An

acetonitrile solution (0.8 mL) containing granisetron (0.5 ng/mL) as internal

standard was added to the brain homogenate (0.2 mL), followed by

centrifugation at 10,600 g for 2 min at 4°C. The supernatant was collected and

filtered by solid phase extraction column (Phree[TM]; Shimadzu GLC Ltd, Tokyo,

Japan). The filtrate was dried under nitrogen gas at 40°C and re-dissolved in 5%

acetonitrile (0.2 mL) followed by sonication for 30 s and centrifugation (500 g, 2

min). The supernatant was filtered by solid phase extraction column again

before its application to LC/MS/MS.

Quantification of C21, DCZ, and DCZ-N-oxide was performed by multiple

reaction monitoring (MRM) using a Shimadzu UHPLC LC-30AD system

(Shimadzu Corp., Kyoto, Japan) coupled to a tandem MS AB Sciex Qtrap 6500

system (AB Sciex LLC, Framingham, USA). The following MRM transitions

(Q1/Q3) were used to monitor each compound: C21 (279.0/193.0), DCZ

(293.0/236.0), DCZ-N-oxide (309.0/192.9), granisetron (313.2/138.1). Other

protocols were described previously[6].

STATISTICAL ANALYSES

In vitro assays. All data were analyzed using GraphPad Prism 7 (San Diego,

USA). Inhibition binding data were analyzed according to a one-site binding

model, and the equilibrium dissociation constants (Ki) of unlabeled ligands were

calculated by constraining the radioligand Kd to the values estimated from

saturation binding assays. Concentration–response curves were fitted to a

three-parameter logistic equation. All affinity and potency values were estimated

as logarithms.

PET imaging. To estimate the specific binding of [^{11}C]DCZ, regional binding

potential relative to nondisplaceable radioligand (BP$_{ND}$) was calculated by

PMOD® with an original multilinear reference tissue model (MRTMo)[39] as

described in the following equation:

$$\frac{\int_0^T C(t)\,dt}{C(T)} = \frac{V}{V'} \cdot \frac{\int_0^T C'(t)\,dt}{C(T)} + \frac{V}{V'k_2'} \frac{C'(T)}{C(T)} + b \qquad (1),$$

where C(t) and C'(t) are the regional or voxel time-radioactivity concentrations in

the tissue and reference regions, respectively (kBq/mL), V and V' are the

corresponding total distribution volumes (mL/mL), k_2' (min^{-1}) is the clearance

rate constant from the reference region to plasma, and b is the intercept term,

which becomes the constant for T > t*. In this study, t* was determined as 2 min

for mice studies and 15 min for monkey studies. Eq. 1 allows estimation of three

parameters, β_1 = V/V', β_2 = V/(V'k$_2$'), and β_3 = b by multilinear regression

analysis for T > t*. Assuming that the nondisplaceable distribution volumes in

tissue and reference regions are identical, BP$_{ND}$ (BP$_{ND}$ = V/V' − 1) is calculated

from the first regression coefficient as BP$_{ND}$ = (β_1 − 1).

Estimates of fractional occupancy were calculated by the following

equation:

$$Occ = (^{hM4Di}BP_{BL} - ^{hM4Di}BP_{PT}) / (^{hM4Di}BP_{BL} - ^{CON}BP_{BL}) \qquad (2),$$

where $^{hM4Di}BP_{BL}$ and $^{CON}BP_{BL}$ indicate BP_{ND} at hM$_4$Di-expressing putaminal

region and control putamen under baseline conditions, respectively, while

$^{hM4Di}BP_{PT}$ indicates BP_{ND} at hM$_4$Di-expressing putaminal region under

pretreatment condition. The relationship between occupancy (*Occ*) and agonist

dose (*D$_{agonist}$*) was modeled by the Hill equation

$$Occ = D_{agonist}^{n} / (D_{agonist}^{n} + ED_{50}^{n}) \qquad (3),$$

where *ED$_{50}$* and *n* indicate the agonist dose achieving 50% occupancy and the

Hill coefficient, respectively.

For FDG-PET analysis, dynamic SUV images were motion-corrected and

then were averaged between 30-60 min after injection of the radioligand.

Voxel-wise statistical analyses were performed with SPM12 software (Wellcome

Department of Cognitive Neurology, London, UK; www.fil.ion.ucl.ac.uk) and

MATLAB R2016a (MathWorks Inc., Natick, MA, USA). We used a total of 12

scans including vehicle and DCZ 3 µg/kg pretreatment conditions to detect the

metabolic change by hM$_3$Dq activation and a total of 12 scans including vehicle

and DCZ 100 µg/kg pretreatment conditions to examine the effect of off-target

actions in the non-DREADD monkeys. The averaged SUV images were spatially

normalized into standard template brain MR image[40] after coregistration of

individual MR images. The resulting images were smoothed with a 2.6-mm

Gaussian filter and were then extracted by brain mask. The SUV values for each

scan were corrected using grand mean scaling and by analysis of covariance for

global normalization. Subsequently, repeated measures one-way ANOVA was

applied. The statistical threshold was set at uncorrected $p < 0.001$ (T > 4.50) and

an extent of 100 contiguous voxels per cluster. In VOI-based analysis, we used

a total of 17 scans including vehicle and a DCZ dose of 1 µg/kg, 3 µg/kg

pretreatment conditions obtained from hM_3Dq-expressing monkeys and a total of

16 scans including vehicle, and a DCZ dose of 1 µg/kg, 100 µg/kg pretreatment

conditions obtained from non-DREADD monkeys. The SUV values were

obtained by PMOD® and then were normalized by whole-brain value. Paired

t-test was used to detect the metabolic change by hM_3Dq activation in each

condition. To examine the dose-dependency or off-target actions, one-way

ANOVA followed by post-hoc Tukey-Kramer test was performed.

Neuronal activity. Time-dependent changes in fluorescent signal and LFP power were analyzed according to a one-phase exponential association model using GraphPad Prism 7. Repeated measure ANOVA followed by post-hoc Dunnett test was used to examine differences between baseline and drug-evoked mean neural signals. The signals were displayed after normalization to the baseline levels, but the statistical tests were conducted on the original data.

To examine the long-term effect of hM_3Dq activation on normal neuronal responsiveness, fluorescent signal change within $0.2 - 10.2$ s after whisker stimulation was averaged across 4 trials and compared using pair-wise t-test. DCZ-induced fluorescent signal changes at 10 min after the 1st and 2nd DCZ administrations were compared by pair-wise t-test The second DCZ administration was applied 24 h after the first.

Behavioral testing. To examine the effect of DCZ on the performance of the delayed response task, behavioral measurement (correct rates) was subjected to two-way ANOVA (treatment ✕ delay) and the following post-hoc t-test with Bonferroni correction using GraphPad Prism 7. To examine whether the performance was recovered at 24 h after DCZ administration, the same procedures were conducted between DCZ and post-DCZ, and vehicle and

post-DCZ sessions. To examine the effect of DCZ treatment on the behavior in test time, the data of 5- and 10-s delays were pooled and then compared between treatment conditions by t-test.

To examine the effect of DCZ on the performance of the reward-size task, reaction times and error rates were subjected to two-way ANOVA (treatment × reward size). The total correct trial number for each session was subjected to two-way ANOVA (treatment × subject). For each analysis, effect size eta-squared (η^2) was calculated using R.

ACKNOWLEDGMENTS

We thank R. Suma, J. Kamei, R. Yamaguchi, Y. Matsuda, Y. Sugii, A. Maruyama, T. Okauchi, T. Kokufuta, Y. Iwasawa, T. Watanabe, A. Tanizawa, S. Shibata, N. Nitta, Y. Ozawa, M. Fujiwara, M. Nakano, T. Minamihisamatsu, S. Uchida and S. Sasaki for their technical assistance. We also thank Dr. S. Hiura for 3D printing of grids. This study was supported by MEXT/JSPS KAKENHI Grant Numbers JP15H05917, JP15K12772 and JP18H04037 (to TM), JP16H02454 (to MTakada), JP19K08138 (to YN), and JP18H05018 and JP19K07811 (to NM), by AMED Grant Numbers JP18dm0107146 (to TM),

JP18dm0207003 (to MTakada), JP18dm0107094 and JP18dm0207007 (to TS), JP18dm0307021(to KI), JP18dm0307007 (to TH), by JST PRESTO Grant Number JPMJPR1683 (to KI), by QST President's Strategic Grant (Creative Research)(to NM), by the Cooperative Research Program at PRI, Kyoto Univ, and by National Bio-Resource Project "Japanese Monkeys" of MEXT, Japan and by U24DK116195, the NIMH Psychoactive Drug Screening Program and the Michael Hooker Distinguished Professorship to BLR.

AUTHOR CONTRIBUTIONS

Conceptualization, T.M.; Formal Analysis, Y.N., N.M., K.O., K.M., and T.M.; Investigation, Y.N., N.M., H.T., Y.H., K.O., B.J., M.Takahashi, X.-P.H., S.T.S., J.F.D., X.Y., T.U., J.G.E., J.L., K.K., C.S., M.O., and M.S.; Resources, B.J., X.Y., J.L., K.I., M.Takada, and J.J.; Writing – original draft, Y.N., N.M., and T.M.; Visualization, Y.N., N.M., H.T., Y.H., K.O., B.J., K.M. and T.M.; Supervision, M.-R.Z., T.S., M.Takada, M.H., J.J., B.L.R., and T.M.; Project Administration, B.L.R., and T.M.; Funding Acquisition, Y.N., N.M., T.H., K.I., M.Takada, T.S., J.J., B.L.R., and T.M.; Writing – review & editing, all authors.

Competing Interests statement

The authors declare no competing interests.

REFERENCES

1. Armbruster, B.N., Li, X., Pausch, M.H., Herlitze, S. & Roth, B.L. Evolving the lock to fit the key to create a family of G protein-coupled receptors potently activated by an inert ligand. *Proc Natl Acad Sci U S A* 104, 5163-5168 (2007).

2. Vardy, E.*, et al.* A New DREADD Facilitates the Multiplexed Chemogenetic Interrogation of Behavior. *Neuron* 86, 936-946 (2015).

3. Roth, B.L. DREADDs for Neuroscientists. *Neuron* 89, 683-694 (2016).

4. Grayson, D.S.*, et al.* The Rhesus Monkey Connectome Predicts Disrupted Functional Networks Resulting from Pharmacogenetic Inactivation of the Amygdala. *Neuron* 91, 453-466 (2016).

5. Eldridge, M.A.*, et al.* Chemogenetic disconnection of monkey orbitofrontal and rhinal cortex reversibly disrupts reward value. *Nat Neurosci* 19, 37-39 (2016).

6. Nagai, Y.*, et al.* PET imaging-guided chemogenetic silencing reveals a critical role of primate rostromedial caudate in reward evaluation. *Nature communications* 7, 13605 (2016).

7. Upright, N.A.*, et al.* Behavioral Effect of Chemogenetic Inhibition Is Directly Related to Receptor Transduction Levels in Rhesus Monkeys. *J Neurosci* 38, 7969-7975 (2018).

8. Gomez, J.L.*, et al.* Chemogenetics revealed: DREADD occupancy and activation via converted clozapine. *Science* 357, 503-507 (2017).

9. Manvich, D.F.*, et al.* The DREADD agonist clozapine N-oxide (CNO) is reverse-metabolized to clozapine and produces clozapine-like interoceptive

stimulus effects in rats and mice. *Scientific reports* 8, 3840 (2018).

10. Raper, J*., et al.* Metabolism and Distribution of Clozapine-N-oxide: Implications for Nonhuman Primate Chemogenetics. *ACS Chem Neurosci* 8, 1570-1576 (2017).

11. Roth, B.L., Sheffler, D.J. & Kroeze, W.K. Magic shotguns versus magic bullets: selectively non-selective drugs for mood disorders and schizophrenia. *Nature Reviews Drug Discovery* 3, 353-359 (2004).

12. Chen, X*., et al.* The first structure-activity relationship studies for designer receptors exclusively activated by designer drugs. *ACS Chem Neurosci* 6, 476-484 (2015).

13. Thompson, K.J*., et al.* DREADD Agonist 21 Is an Effective Agonist for Muscarinic-Based DREADDs in Vitro and in Vivo. *ACS Pharmacol Transl Sci* 1, 61-72 (2018).

14. Phillips, S.T*., et al.* Binding of 5H-dibenzo[b,e][1,4]diazepine and chiral 5H-dibenzo[a,d]cycloheptene analogues of clozapine to dopamine and serotonin receptors. *J Med Chem* 37, 2686-2696 (1994).

15. Ji, B*., et al.* Multimodal Imaging for DREADD-Expressing Neurons in Living Brain and Their Application to Implantation of iPSC-Derived Neural Progenitors. *J Neurosci* 36, 11544-11558 (2016).

16. Farrell, M.S. & Roth, B.L. Pharmacosynthetics: Reimagining the pharmacogenetic approach. *Brain Res* 1511, 6-20 (2013).

17. Kalvass, J.C., Maurer, T.S. & Pollack, G.M. Use of plasma and brain unbound fractions to assess the extent of brain distribution of 34 drugs: comparison of unbound concentration ratios to in vivo p-glycoprotein efflux ratios. *Drug*

Metab Dispos 35, 660-666 (2007).

18. Maurer, T.S., DeBartolo, D.B., Tess, D.A. & Scott, D.O. Relationship between exposure and nonspecific binding of thirty-three central nervous system drugs in mice. *Drug Metabolism and Disposition* 33, 175-181 (2005).

19. Martinez, M.N. Factors influencing the use and interpretation of animal models in the development of parenteral drug delivery systems. *AAPS J* 13, 632-649 (2011).

20. Schihada, H.*, et al.* A universal bioluminescence resonance energy transfer sensor design enables high-sensitivity screening of GPCR activation dynamics. *Commun Biol* 1, 105 (2018).

21. Kroeze, W.K.*, et al.* PRESTO-Tango as an open-source resource for interrogation of the druggable human GPCRome. *Nature structural & molecular biology* 22, 362-369 (2015).

22. Alexander, G.M.*, et al.* Remote control of neuronal activity in transgenic mice expressing evolved G protein-coupled receptors. *Neuron* 63, 27-39 (2009).

23. Buzsaki, G., Anastassiou, C.A. & Koch, C. The origin of extracellular fields and currents--EEG, ECoG, LFP and spikes. *Nat Rev Neurosci* 13, 407-420 (2012).

24. Michaelides, M.*, et al.* Whole-brain circuit dissection in free-moving animals reveals cell-specific mesocorticolimbic networks. *Journal of Clinical Investigation* 123, 5342-5350 (2013).

25. Phelps, M.E.*, et al.* Tomographic measurement of local cerebral glucose metabolic rate in humans with (F-18)2-fluoro-2-deoxy-D-glucose: validation of method. *Ann Neurol* 6, 371-388 (1979).

26. Poremba, A*., et al.* Species-specific calls evoke asymmetric activity in the monkey's temporal poles. *Nature* 427, 448-451 (2004).

27. Tsutsui, K.I., Oyama, K., Nakamura, S. & Iijima, T. Comparative Overview of Visuospatial Working Memory in Monkeys and Rats. *Front Syst Neurosci* 10, 99 (2016).

28. Allen, D.C.*, et al.* A Comparative Study of the Pharmacokinetics of Clozapine N-Oxide and Clozapine N-Oxide Hydrochloride Salt in Rhesus Macaques. *The Journal of pharmacology and experimental therapeutics* 368, 199-207 (2019).

29. Katzel, D., Nicholson, E., Schorge, S., Walker, M.C. & Kullmann, D.M. Chemical-genetic attenuation of focal neocortical seizures. *Nature communications* 5, 3847 (2014).

30. Stachniak, T.J., Ghosh, A. & Sternson, S.M. Chemogenetic synaptic silencing of neural circuits localizes a hypothalamus-->midbrain pathway for feeding behavior. *Neuron* 82, 797-808 (2014).

31. Tomita, Y*., et al.* Long-term in vivo investigation of mouse cerebral microcirculation by fluorescence confocal microscopy in the area of focal ischemia. *J Cereb Blood Flow Metab* 25, 858-867 (2005).

32. Tajima, Y*., et al.* Changes in cortical microvasculature during misery perfusion measured by two-photon laser scanning microscopy. *J Cereb Blood Flow Metab* 34, 1363-1372 (2014).

33. Friden, M*., et al.* Measurement of unbound drug exposure in brain: modeling of pH partitioning explains diverging results between the brain slice and brain homogenate methods. *Drug Metab Dispos* 39, 353-362 (2011).

34. Bender, D., Holschbach, M. & Stocklin, G. Synthesis of Nca C-11 Labeled Clozapine and Its Major Metabolite Clozapine-N-Oxide and Comparison of Their Biodistribution in Mice. *Nucl Med Biol* 21, 921-925 (1994).

35. Takuwa, H*., et al.* Reproducibility and variance of a stimulation-induced hemodynamic response in barrel cortex of awake behaving mice. *Brain Res* 1369, 103-111 (2011).

36. Croxson, P.L., Kyriazis, D.A. & Baxter, M.G. Cholinergic modulation of a specific memory function of prefrontal cortex. *Nat Neurosci* 14, 1510-1512 (2011).

37. Goldman, P.S., Rosvold, H.E. & Mishkin, M. Evidence for behavioral impairment following prefrontal lobectomy in the infant monkey. *Journal of comparative and physiological psychology* 70, 454-463 (1970).

38. Minamimoto, T., La Camera, G. & Richmond, B.J. Measuring and modeling the interaction among reward size, delay to reward, and satiation level on motivation in monkeys. *J Neurophysiol* 101, 437-447 (2009).

39. Ichise, M*., et al.* Noninvasive quantification of dopamine D2 receptors with iodine-123-IBF SPECT. *Journal of nuclear medicine : official publication, Society of Nuclear Medicine* 37, 513-520 (1996).

40. McLaren, D.G*., et al.* A population-average MRI-based atlas collection of the rhesus macaque. *Neuroimage* 45, 52-59 (2009).

FIGURE LEGENDS

Fig. 1 | DCZ selectively binds to DREADDs. a, Chemical structure of DCZ and clozapine. **b,c,** Binding affinities of DCZ and clozapine to DREADDs and endogenous receptors, channels and transporters. *Ki* values are the average of at least 3 triplicate experiments with standard deviation values that are 3-fold less than the average. The values are also shown in Table 2. **d,** Illustration representing location of viral vector injections. AAV2-CMV-hM$_4$Di and AAV2-CMV-KORD (as control) were injected into the right and left putamen, respectively. **e,** Illustration representing *in vivo* PET imaging with [^{11}C]DCZ in a monkey. **f,** (left) Coronal section of PET image with [^{11}C]DCZ representing standardized radio-ligand uptake value [SUV; regional radioactivity (Bq/cm^3) × body weight (g) / injected radioactivity (Bq)] between 30-90 min from injection. (right) Time course of regional uptake (mean ± sd) of [^{11}C]DCZ at the hM$_4$Di-expression region in the putamen (hM$_4$Di-PUT), control region in the contralateral putamen (Ctrl-PUT), and the cerebellum, respectively. **g,** Same as **f** but with [^{11}C]clozapine. **h,** Chemical structure of C21 and CNO. **i,** Same as **f** but with [^{11}C]CNO.

Fig. 2 | [^{11}C]DCZ-PET visualizes DREADD expression and measures

agonist dose–occupancy relationship. a, Coronal section of parametric

[^{11}C]DCZ-PET image of specific binding (BP$_{ND}$) overlaying on MR image of a

monkey (#209) expressing hM$_4$Di in the putamen. **b,** An anti-hM$_4$ staining

section corresponding to the image in **a**. The inset is a high magnification image

(scale bar: 100 μm). **c,** Illustration of occupancy study. Monkeys underwent

[^{11}C]DCZ-PET scan 15 min after i.v. injection of nonradiolabeled DCZ, CNO or

C21. **d,** Occupancy of hM$_4$Di is plotted as a function of DCZ, CNO or C21 dose.

Dotted curve is the best-fit Hill equation for the data. ED$_{50}$ indicates the agonist

dose inducing 50% occupancy. Hill coefficient (n) and coefficient of

determination (R^2) as are follows: DCZ, n = 1, R^2 = 0.99; CNO, n = 0.44, R^2 =

0.96; C21, n = 1, R^2 > 0.99.

Fig. 3 | Time concentration profiles of DCZ and its metabolites in monkeys

and mice. a, Plasma and CSF concentrations of DCZ, and its major metabolites

(C21 and DCZ-N-oxide) following intravenous injection of DCZ (100 μg/kg). Data

were collected from 4 monkeys and are shown as mean ± sem. **b**, Same as a,

but following intramuscular injection. **c**, Plasma, brain and CSF concentrations of

DCZ, C21 and DCZ-N-oxide following intraperitoneal injection of DCZ (100

μg/kg). Data were collected from 3 or 4 mice and are shown as mean ± sem. **d**,

Major routes of DCZ metabolism leading to C21 and DCZ-N-oxide and their

structures.

Fig. 4 | *In vitro* potency and *in vivo* efficacy of DCZ for DREADDs. a,b,

BRET-based estimation of agonist potency and efficacy at hM$_3$Dq and hM$_4$Di.

Shown are mean of N=3 separate biological replicates (each with duplicate

technical replicates) of dose-response curves for hM$_3$Dq using a Gq-based

BRET sensor and hM$_4$Di using a Gi1-based BRET sensor. **c,** Illustration of

two-photon calcium imaging and location for co-injection of AAV2-CMV-hM$_3$Dq

and AAV-DJ-rSyn-GCaMP6s. **d,** Parametric image of specific binding of

[^{11}C]DCZ overlaying MR image. **e,** Time-averaged two-photon images of

GCaMP6 of pre-injection (left) and 5 min post DCZ injection (1 μg/kg, i.p.)(right).

f, Raw fluorescence signals in 14 representative neurons from imaging in **e**. au:

arbitrary units. **g,** Fluorescent intensity change (mean ± sem from baseline) as a

function of post-injection time. Data were obtained from n = 88 neurons in four

hM$_3$Dq-expressing mice and n = 30 neurons in two non-hM$_3$Dq mice. Curves

represent exponential fits to the data. Asterisks represent significant difference

(p < 0.01, one-way ANOVA with post-hoc Dunnett) from pre-injection.

Fig. 5 | DCZ rapidly drives activation of hM$_3$Dq-expressing neuronal

population in monkeys. a, Illustration representing viral vector

(AAV2-CMV-hM$_3$Dq) injection into one side of the amygdala. **b,** Coronal PET

image overlaying CT image demonstrated that recording contacts on a

multichannel electrode (shown on right) were located in an hM$_3$Dq-expressing

region, which was visualized as a high [^{11}C]DCZ binding site. The high binding

observed in the dorsomedial surface corresponds to a site of biological reaction

within the recording chamber. **c,** Representative LFP activity changes after DCZ

(left) and vehicle administration (right). **d,** Normalized gamma (40 Hz) power

change (mean ± sem from baseline) in 10-20 min after intravenous

administration of reagents. Data were obtained from hM$_3$Dq-positive region

(DCZ and vehicle; n = 27 and 26 channels, respectively) and non-hM$_3$Dq

expressing sites (DCZ-non-hMD$_3$q; n = 11 channels). Gray bars indicate

baseline. Asterisk represents significant difference (p < 0.001, t-test) from

baseline. **e,** Plots represent normalized power change (mean ± sem from

baseline) as a function of time from vehicle or DCZ administration. Curve

represents exponential fit to the data. Asterisks represent significant difference

($p < 0.01$, one-way ANOVA with post-hoc Dunnett) from baseline.

Fig. 6 | DCZ selectively induces metabolic changes in hM$_3$Dq-expressing

region in a dose-dependent manner. a, Illustration representing DREADD

activation imaging with [^{18}F]FDG. Following administration of DCZ at effective

doses, [^{18}F]FDG was injected and PET scan was performed to assess

DREADD-induced brain metabolic change. **b,** Coronal PET image overlaying

MR image representing specific [^{11}C]DCZ binding (BP$_{ND}$) in a monkey (#215).

Filled and open arrowheads indicate the hM$_3$Dq-expressing and control region in

the amygdala, respectively. **c,** Brain areas with significantly increased [^{18}F]FDG

uptake after DCZ injection (3 µg/kg) compared with vehicle overlay an MRI

template ($p < 0.001$, uncorrected, monkeys #215 and #236). Significance level is

given as a t-value represented on a color scale. **d,** Relationship between the

dose of DCZ administration and standardized FDG uptake (SUVR) in monkeys

expressing hM$_3$Dq in amygdala. Open and filled symbols are the values at

hM$_3$Dq-expressing amygdala and contralateral control region, respectively. * $p <$

0.01, paired t-test; # p < 0.05, one-way ANOVA with post-hoc Tukey test. **e,**

Same as **d**, but for amygdala of control monkeys without hM$_3$Dq-vector injection

(non-DREADD monkeys; #210 and #230).

Fig. 7 | DCZ selectively and rapidly induces spatial working-memory deficit

in monkeys expressing hM$_4$Di in PFC. a, Illustration representing location of

viral vector injections. AAV1-hSyn-hM$_4$Di-IRES-AcGFP was injected bilaterally

into the dorsal and ventral banks of the principal sulcus. **b,** Coronal parametric

image of specific binding of [^{11}C]DCZ overlaying MR image in a monkey (#229).

Open and filled arrowheads represent the dorsal and ventral borders of the

target regions, respectively. **c,** Spatial delayed response task. After the monkey

was shown the baited well (Cue), the screen was placed for 0.5, 5 or 10 s

(Delay). The monkey obtained the food if they reached for the correct food-well

and removed the cover plate (Response). **d,** Correct performance rates (mean ±

sem) of spatial delayed response task with 0.5, 5 and 10 s delays after vehicle

administration (n = 5 sessions; cyan), DCZ administration (100 µg/kg, i.m.; n = 5;

red), and 24 h after DCZ administration (n = 5; purple) in two monkeys

expressing hM$_4$Di in PFC. Note that one of the monkeys (#245) was also

examined prior to vector injection (see Supplementary Fig. 6c, right). The

behavioral testing began immediately after an i.m. administration of either

vehicle (2% DMSO in saline) or DCZ, except for 24 h after DCZ. Each dot

represents the data in each daily session. **e,** Time course of DCZ effect. Correct

performance rates of 5- and 10-s delays as a function of test time for vehicle and

DCZ sessions (mean ± sem; n = 5 sessions) in monkey #229. The behavioral

testings for 0-55 min and 60-115 min were conducted separately. Asterisks

represent significant difference (p < 0.05). **f,** Correct performance rates (mean of

5- and 10-s delays) as a function of session number in two monkeys.

Table 1. | Binding affinities of DCZ, clozapine and CNO to DREADDs and native muscarinic receptors.

compound	K_i (nM) / pK_i (± SEM) hM$_3$Dq	hM$_3$	hM$_4$Di	hM$_4$
DCZ	6.3	230	4.2	210
	8.20 (± 0.15)	6.63 (± 0.06)	8.38 (± 0.27)	6.68 (± 0.04)
clozapine	5.9	21	0.89	10
	8.23 (± 0.26)	7.69 (± 0.14)	9.05 (± 0.13)	7.98 (± 0.10)
CNO	680	2600	360	3700
	6.17 (± 0.07)	5.59 (± 0.06)	6.44 (± 0.03)	5.43 (± 0.18)
C21	850	270	178*	3,631*
	6.07 (± 0.06)	6.57 (± 0.15)	6.75 (± 0.26)	5.44 (± 0.11)

Mean K_i values (upper row) and mean ± SEM of pK_i values (lower row) are shown. The average of at least 3 duplicate experiments with standard deviation (SD) values that are 3-fold less than average. * data from Thompson et al., 2018.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7