Reactomics: using mass spectrometry as a reaction detector

Miao Yu[1], Lauren Petrick[1,2*]

[1] Department of Environmental Medicine and Public Health, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, United States
[2] Institute for Exposomic Research, Icahn School of Medicine at Mount Sinai, New York, NY, 10029, United States
*Corresponding author: Email: lauren.petrick@mssm.edu Phone: +1-212-241-7351. Fax: +1-646-537-9654.

Abstract

Untargeted metabolomics analysis captures chemical reactions among small molecules. Common mass spectrometry-based metabolomics workflows first identify the small molecules significantly associated with the outcome of interest, then begin exploring their biochemical relationships to understand biological fate (environmental studies) or biological impact (physiological response). We suggest an alternative by which biochemical relationships can be directly retrieved through untargeted high-resolution paired mass distance (PMD) analysis without a priori knowledge of the identities of participating compounds. Retrieval is done using high resolution mass spectrometry as a chemical reaction detector, where PMDs calculated from the mass spectrometry data are linked to biochemical reactions obtained via data mining of small molecule and reaction databases, i.e. 'Reactomics'. We demonstrate applications of reactomics including PMD network analysis, source appointment of unknown compounds, and biomarker reaction discovery as a complement to compound discovery analyses used in traditional untargeted workflows. An R implementation of reactomics analysis and the reaction/PMD databases is available as the pmd package (https://yufree.github.io/pmd/).

Introduction

Untargeted metabolomics or non-targeted analysis using high resolution mass spectrometry

(HRMS) is one of the most popular analysis methods for unbiased measurement of organic

compounds [1,2]. A typical metabolomics sample analysis workflow will follow a detection,

annotation, MS/MS validation and/or standards validation process, from which interpretation of

the relationships between these annotated or identified compounds can then be linked to

biological pathways or disease development, for example. However, difficulty annotating or

identifying unknown compounds always limits the interpretation of findings [3]. One practical

solution to this is matching experimentally obtained fragment ions to a mass spectral database [4],

but many compounds remain unreported/absent, thereby preventing annotation. Rules or data

mining-based prediction of in silico fragment ions is successful in many applications [2,5], but

these approaches are prone to overfitting the known compounds, leading to false positives.

Ultimately such workflows require final validation with commercially available or synthetically

generated analytical standards, which may not be available, for unequivocal identification.

Potential molecular structures could be discerned using biochemical knowledge, through the

integration of known relationships between biochemical reactions (e.g. pathway analysis) [3]. Such

methods are readily used to annotate compounds by chemical class. For example, the Referenced

Kendrick Mass Defect (RKMD) was able to predict lipid class using specific mass distances for

lipids and heteroatoms [6], and isotope patterns in combination with specific mass distances

characteristic of halogenated compounds such as +Cl/-H, +Br/-H were used to screen

halogenated chemical compounds in environmental samples [7]. For these examples, known

relationships among compounds were used to annotate unknown compounds, as a

complementary approach to obtaining compound identifications.

The most common relationships among compounds are chemical reactions. Substrate-product

pairs in a reaction form by exchanging functional groups or atoms. Almost all organic

compounds originate from biochemical processes, such as carbon fixation [8,9]. Like base pairing

in DNA [10], organic compounds follow biochemical reaction rules, resulting in characteristic mass

differences between the paired substrates and their products. We present a new concept, paired

mass distance (PMD), that reflects such reaction rules by calculating the mass differences

between two compounds or charged ions. PMD can be used to extract biological inference

without identifying unknown compounds.

Exploiting mass differences for compound identification is not new. Mass distances have been

used to reveal isotopologue information when peaks show a PMD of 1 Da [11], adducts from a

single compound [12] such as PMD 22.98 Da between adducts $[M+Na]^+$ and $[M+H]^+$, or adducts

formed via complex in-source reactions [13] from mass spectrometry data. Such between-

compound information has also been used to make annotations of unknown compounds [4,14], to

classify compounds [15] or to perform pathway-independent metabolomic network analysis [16].

However, these calculations of PMD were used to identify compounds or pathways in ultimately

facilitate interpretations of the relationships between important compounds. Here, we propose

that PMD can be used directly, skipping the step for annotation or identification of individual

compounds, to aggregate information at the reaction level, called 'Reactomics'.

HRMS can directly measure PMDs with the mass accuracy needed to provide reaction level

specificity. Therefore, HRMS has the potential to be used as a reaction detector to enable

reaction level study investigations. Here, we use multiple databases and experimental data to

provide a proof-of-concept for using mass spectrometry in 'reactomics'. We also discuss

potential applications such as PMD network analysis, biomarker reaction discovery, and source

appointment of unknown compounds. We envision that these applications will reveal the

measurable reaction level changes without the need to assign molecular structure to unknown

compounds.

Results and Discussion

Definitions

We first define a reaction PMD (PMD_R) using a theoretic framework. Then we demonstrate how

a PMD_R can be calculated using KEGG reaction R00025 as an example (see Equation 1). There

are three KEGG reaction classes (RC00126, RC02541, and RC02759) associated with this

reaction, which is catalyzed by enzyme 1.13.12.16.

Ethylnitronate + Oxygen + Reduced FMN <=> Acetaldehyde + Nitrite + FMN + Water   [1]

In general, we define a chemical reaction (PMD_R) as follows:

$S_1 + S_2 + \ldots + S_n <=> P_1 + P_2 + \ldots + P_m$ (n >= 1, m >= 1)   [2]

Where: S means substrates and P mean products, and n and m the number of substrates and

products, respectively. A PMD matrix [M1] for this reaction is generated:

	S_1	S_2	...	S_n
P_1	$\lvert S_1\text{-}P_1\rvert$	$\lvert S_2\text{-}P_1\rvert$	...	$\lvert S_n\text{-}P_1\rvert$
P_2	$\lvert S_1\text{-}P_2\rvert$	$\lvert S_2\text{-}P_2\rvert$	...	$\lvert S_n\text{-}P_2\rvert$
...	...	...	...	...
P_m	$\lvert S_1\text{-}P_m\rvert$	$\lvert S_2\text{-}P_m\rvert$	...	$\lvert S_n\text{-}P_m\rvert$

[M1]

For each substrate, S_k, and each product, P_i, we calculate a PMD ($\lvert S_n\text{-}P_m\rvert$) .

Assuming that the minimum PMD would have a similar structure or molecular framework

between substrate and products, we select the minimum numeric PMD for each substrate as the

substrate PMD (PMD_{Sk}) of the reaction (Eq. 2).

$PMD_{Sk} = min(\lvert S_k\text{-}P_1\rvert, \lvert S_k\text{-}P_2\rvert, \ldots, \lvert S_k\text{-}P_m\rvert)$ (1<=k<=n)   [2]

Then, the PMD_R , or overall Reaction PMD, is defined as the set of substrates' PMD(s) (Eq. 3):

$PMD_R = \{PMD_{S1}, PMD_{S2}, \ldots, PMD_{Sn}\}$ [3]

For KEGG reaction R00025, S_1 is Ethylnitronate, S_2 is Oxygen, S_3 is Reduced FMN, P_1 is

Acetylaldehyde, P_2 is Nitrite, P_3 is FMN, P_4 is Water, n=4, and m =3. A PMD matrix [M2] for

this reaction can be seen below, where we define $PMD_{Ethylnitronate}$ = 27.023 Da, PMD_{Oxygen} =

12.036 Da and $PMD_{Reduced\ FMN}$ = 2.016 Da.

	Ethylnitronate	Oxygen	Reduced FMN
Acetaldehyde	29.998 Da	12.036 Da	414.094 Da
Nitrite	27.023 Da	15.011Da	411.120 Da
FMN	382.080 Da	424.115 Da	2.016 Da
H_2O	56.014 Da	13.979 Da	440.110 Da

[M2]

In our example, there are three PMD_R calculated from three PMD_S: PMD_R is 27.023 Da, which

is equivalent to the mass difference between two carbon atoms and three hydrogen atoms: PMD_R

is 12.036 Da for the additions of two carbon atoms and four hydrogen atoms and loss of one

oxygen atom: and PMD_R is 2.016 Da for the addition of two hydrogen atoms. However, other

reactions may have multiple PMD_S that generate the same PMD_R value, such as certain

combination reactions or replacement reactions. In this case, only one value will be kept as

reaction PMD. In addition, each PMD_R has two notations. One is shown as an absolute mass

difference of the substrate-product pairs' exact masses or monoisotopic masses with unit Da.

Another notation is using elemental compositions as the differences between two chemical

formulas. Here, we describe it as an elemental composition instead of chemical formula, because

it also describes the gain and loss of elements, and therefore the neat mass change. In our

example reaction, the PMD_R can also be written as +2C3H, +2C4H/-O, and +2H, respectively.

This elemental composition can be linked to known chemical processes retrieved from a reaction

database, ie. KEGG. For example, +2H represents the elemental composition change of a

reaction involving a double bond breaking such as KEGG example RC00126, and +2C3H

indicates reaction with nitronate monooxygenase (EC:1.13.12.16) or reaction class RC02541.

However, some elemental compositions, such as +2C4H/-H in our example, might not have a

clear mechanism (e.g. no suggested KEGG reaction selection). By this definition, PMD_R can be

generated automatically in terms of elemental compositions or mass units in Da.

We used these definitions to establish reference databases of PMDs. We used KEGG as a

'reaction database'representing common reactions in human endogenous pathways, and we used

human metabolome database (HMDB)[17] as the 'compound database' representing common

reactions between chemicals measured in human biofluids (see S.1 in the supplementary

information).

Qualitative and quantitative PMD analysis with mass spectrometry

PMD can be determined in biological or environmental samples from peaks observed in mass

spectrometry. Mathematically, a PMD of uncharged compounds is equivalent to the PMD of

their charged species observed with a mass spectrometer, as long as both compounds share the

same adducts, neutral losses, and charges. In example reaction [1], reduced FMN has a

monoisotopic mass of 458.1203 Da, while FMN has a monoisotopic mass of 456.1046 Da.

Spectra from HMDB[17] showed that common ions for reduced FMN and FMN using liquid

chromatography (LC)-HRMS in negative mode are typically [M-H]$^-$ with m/z 457.1124 and

455.0968, respectively. The mass distance of the monoisotopic masses is 2.016 Da and the mass

distance of the observed adducts is also 2.016 Da. In cases such as this, mass spectrometry can

be used to detect the PMD of paired compounds, but only for HRMS (see S.1 for data mining,

S.2 for issues dealing with redundant peaks and fragments, and S.3 for discussion of resolution

issues).

In addition to qualitative analysis, peaks that share the same PMD can be summed and used as a

quantitative group measure of that specific 'reaction' in the sample, thereby providing a

description of chemical reaction level changes across samples without annotating individual

compounds. We define two types of PMD across samples: static PMD in which intensity ratios

between the pairs are stable across samples, and dynamic PMD in which the intensity ratios

between pairs change across samples. Only static PMDs, those with similar instrument response,

can be used for quantitative analysis (see Table S.4 for theoretical example). Similar to other

non-targeted analysis[18], an RSD between quantitative pair ratios $< 30\%$ and a high correlation

between the paired peaks' intensity (> 0.6) are suggested to be considered a static PMD.

PMD network analysis

Using the proposed PMD network analysis (see S.4 for details), we can identify metabolites

associated with a known biomarker of interest. In fact, PMD network analysis can also be used in

combination with classic identification techniques to enhance associated networks with targeted

biomarkers. As a proof of concept, we re-analyzed data from a published study to detect the

biological metabolites of exposure to Tetrabromobisphenol A (TBBPA) in pumpkin [19,20] using a

local, recursive search strategy (see Figure 1). Using TBBPA as a target of interest, we searched

for PMDs linked with the debromination process, glycosylation, malonylation, methylation, and

hydroxylation, which are phase II reactions (e.g. primary metabolites) found in the original

paper. Using this PMD network analysis, we identified 22 unique m/z ions of potential TBBPA

metabolites, confirmed by the presence of brominated isotopologue mass spectral patterns

(Figure S1.). This total was 15 more than the 7 unique ions that were described in the original

publication. Such a network was built based on the experimental data and our local fast recursive

search algorithms as shown in S.4. We found that most of the potential metabolites of TBBPA

were found as higher-generation TBBPA metabolites, which are too computationally intensive to

be identified using *in silico* prediction and matching protocols [21].

Figure 1. Metabolites of TBBPA in pumpkin seedlings' root samples. Edges between two nodes

were defined as Pearson's correlation coefficients > 0.6 and shared reactions related PMDs

including 77.91 Da for Debromination, 162.05 Da for Glycosylation, 86 Da for Malonylation,

14.02 Da for Methylation, and 15.99 Da for Hydroxylation. Previous reported metabolites are

labeled.

Source appointment of unknown compounds

When an unknown compound is identified as a potential biomarker, determining whether it is

associated with endogenous biochemical pathways or exogenous exposures can provide

important information toward identification. As can be seen in Tables S1 and S3, high frequency

PMDs from HMDB and KEGG are dominated by reactions with carbon, hydrogen, and oxygen

suggesting links to metabolism pathways. Therefore, if an unknown biomarker is mapped using a

PMD network, connection to these high frequency PMDs would suggest an endogenous link.

However, separation from this network is expected for an exogenous biomarker in which the

reactive enzyme is not in the database, the exogenous compound is secreted in the parent form,

or if it undergoes changes in functional groups such as during phase I and phase II xenobiotic

metabolism processes. In this case, endogenous and exogenous compounds should be separated

by their PMD network in samples.

Topological differences in PMD networks for endogenous and exogenous metabolites were

explored using compounds from The Toxin-Toxin-Target Database (T3DB) [22]. As shown in

Figure 2, the PMD network of compounds was generated based on top ten high frequency PMDs

of 255 endogenous compounds with 223 unique masses and 705 exogenous compounds with 394

unique masses and carcinogenic 1, 2A, or 2B classifications. Most endogenous compounds

(Figure 2, red) were connected into a large network, while the exogenous compounds' networks

were much smaller (Figure 2, blue). Interestingly, most carcinogenic compounds were not

connected by high frequency PMDs. Expanding this beyond just carcinogenic compounds, we

randomly sampled 255 exogenous compounds from a total of 2491 exogenous compounds

available in T3DB, and built a PMD network with the top 10 high frequency PMDs of those 510

compounds (255 exogenous compounds and 255 endogenous compounds). This step was

repeated 1000 times, and the average degree of connection with other nodes was calculated as

4.5 (95% CI [4.3, 4.8]) for endogenous compounds and 1.7 (95% CI [1.2, 2.2]) for exogenous

compounds.

Similar findings were observed for known compounds. We selected caffeine, glucose,

bromophenol, and 5-cholestene as well characterized chemicals that are commonly observed

with mass spectrometry, and paired them with other metabolites in the KEGG reaction database

using the top ten high frequency PMDs from Table S1. As shown in Figure 3, different

topological properties (e.g. number of nodes, average distances, degree, communities, etc.) of

compounds' PMD network were observed for each selected target metabolite. Endogenous

compounds such as glucose or 5-cholestene were highly connected (average degree of node is

3.4 and 3.2, respectively) while exogenous compounds such as caffeine and bromophenol have

more simple networks (average degree of node is 2.2 and 2.4, respectively). Further, the average

PMD edge numbers between all nodes (edges end-to-end) in glucose and 5-cholestene networks

are 9.7 and 6.6, respectively, while the average PMD edge numbers for caffeine and

bromophenol are 3.3 and 1.8, respectively. Larger average PMD edge numbers mean a complex

network structure with lots of nodes while smaller average PMD edge numbers mean a simple

network structure with a few nodes. Based on these estimates, we proposed that unknown

metabolites with average network node degree more than 3 would be likely endogenous

compounds. Similarly, if the unknown compound belongs to a network with longer average

PMD edge numbers , such compounds might also be of endogenous origin.

Figure 2. PMD network of 255 endogenous compounds and 705 exogenous carcinogens from
T3DB database with the top 10 high frequency PMDs network. Compounds without linkage to
other compounds have been removed.
247

248
249 Figure 3. PMD networks for selected compounds from the KEGG reaction database. Networks
250 are limited to relationships with the 10 top frequency PMDs.
251

Biomarker reactions

Reactomics can be used to discover biomarker 'reactions' instead of biomarker 'compounds'.

Unlike typical biomarkers that are a specific chemical compound, biomarker reactions contain all

peaks within a fixed PMD relationship and correlation cutoff. Thus, quantitative PMD analysis

(see S.2) can be used to determine if there are differences between groups (e.g. control or

treatment, exposed or not-exposed) on a reaction level. Such differences are described as a

biomarker 'reaction'.

We used publicly available metabolomics data (MetaboLight ID: MTBLS28) collected on urine

from a study on lung cancer in adults[23]. Four peaks out of 1807 features from 1005 blood

samples (469 cases and 536 controls) generated the quantitative responses of PMD 2.02 Da. This

biomarker reaction (e.g. +2H from our annotated database) was significantly decreased in case

samples compared with the control group (t-test, $p < 0.05$): see Figure 4). The original

publication associated with this dataset did not report any molecular biomarker associated with

this reaction [23], or the metabolites linked with this reaction, suggesting that quantitative PMD

analysis offers additional information on biological differences between the groups on the

reaction level that may be lost when focused on analysis at the chemical level. PMD-level

investigations directly reduce the high dimensional analysis typically performed on a peaks or

features level into low dimensional analysis on the chemical reaction level with explainable

elemental compositions. Furthermore, these results suggest that follow-up analysis in this

population should include targeted analysis of proteins or enzymes linked with +2H changes.

Figure 4. Quantitative PMD analysis identifies PMD 2.02 Da as a potential biomarker reaction
for lung cancer (MTBLS28 dataset).

Conclusion

We provide the theoretical basis and empirical evidence that high resolution mass spectrometry
can be used as a reaction detector through calculation of high resolution paired mass distances
and linkage to reaction databases such as KEGG. Reactomics, as a new concept in
bioinformatics, can be used to find biomarker reactions or develop PMD networks. The major
limitation of reactomics analysis is that mass spectrometry software is designed for analysis of
compounds instead of reactions. In this case, the uncertainty in PMD measurements can not be
captured directly from the instrument, and instead are calculated after data acquisition.
Nevertheless, reactomics techniques provide information on biological changes for new

biological inferences that may not be observed through classic chemical biomarker discovery

strategies.

Acknowledgments

This research was financially supported by NIEHS grants P30ES23515, 1U2CES030859,

R21ES030882, and R01ES031117.

References

1. Zhang, A., Sun, H., Wang, P., Han, Y. & Wang, X. Modern analytical techniques in

metabolomics analysis. *The Analyst* **137**, 293–300 (2012).

2. Hooft, J. J. J. van der, Wandy, J., Barrett, M. P., Burgess, K. E. V. & Rogers, S. Topic

modeling for untargeted substructure exploration in metabolomics. *Proc. Natl. Acad. Sci.*

113, 13738–13743 (2016).

3. Domingo-Almenara, X., Montenegro-Burke, J. R., Benton, H. P. & Siuzdak, G. Annotation:

A Computational Solution for Streamlining Metabolomics Analysis. *Anal. Chem.* **90**, 480–

489 (2018).

4. Guijas, C. *et al.* METLIN: A Technology Platform for Identifying Knowns and Unknowns.

Anal. Chem. **90**, 3156–3164 (2018).

5. Wolf, S., Schmidt, S., Müller-Hannemann, M. & Neumann, S. In silico fragmentation for

computer assisted identification of metabolite mass spectra. *BMC Bioinformatics* **11**, 148

(2010).

6. Lerno, L. A., German, J. B. & Lebrilla, C. B. Method for the Identification of Lipid Classes

Based on Referenced Kendrick Mass Analysis. *Anal. Chem.* **82**, 4236–4245 (2010).

7. Jobst, K. J. *et al.* The use of mass defect plots for the identification of (novel) halogenated

contaminants in the environment. *Anal. Bioanal. Chem.* **405**, 3289–3297 (2013).

8. Bar-Even, A., Noor, E., Lewis, N. E. & Milo, R. Design and analysis of synthetic carbon

fixation pathways. *Proc. Natl. Acad. Sci.* **107**, 8889–8894 (2010).

9. Normile, D. Round and Round: A Guide to the Carbon Cycle. *Science* **325**, 1642–1643

(2009).

10. Donohue, J. & Trueblood, K. N. Base pairing in DNA. *J. Mol. Biol.* **2**, 363–371 (1960).

11. Chokkathukalam, A. *et al.* mzMatch–ISO: an R tool for the annotation and relative

quantification of isotope-labelled mass spectrometry data. *Bioinformatics* **29**, 281–283

(2013).

12. Yu, M., Olkowicz, M. & Pawliszyn, J. Structure/reaction directed analysis for LC-MS based

untargeted analysis. *Anal. Chim. Acta* **1050**, 16–24 (2019).

13. Mahieu, N. G. & Patti, G. J. Systems-Level Annotation of a Metabolomics Data Set Reduces

25□000 Features to Fewer than 1000 Unique Metabolites. *Anal. Chem.* **89**, 10397–10406

(2017).

14. Shen, X. *et al.* Metabolic reaction network-based recursive metabolite annotation for

untargeted metabolomics. *Nat. Commun.* **10**, 1–14 (2019).

15. Burgess, K. E. V., Borutzki, Y., Rankin, N., Daly, R. & Jourdan, F. MetaNetter 2: A

Cytoscape plugin for ab initio network analysis and metabolite feature classification. *J.*

Chromatogr. B **1071**, 68–74 (2017).

16. Grapov, D., Wanichthanarak, K. & Fiehn, O. MetaMapR: pathway independent metabolomic

network analysis incorporating unknowns. *Bioinformatics* **31**, 2757–2760 (2015).

17. Wishart, D. S. *et al.* HMDB 4.0: the human metabolome database for 2018. *Nucleic Acids*

Res. **46**, D608–D617 (2018).

18. Dunn, W. B., Wilson, I. D., Nicholls, A. W. & Broadhurst, D. The importance of

experimental design and QC samples in large-scale and MS-driven untargeted metabolomic

studies of humans. *Bioanalysis* **4**, 2249–2264 (2012).

19. Hou, X. *et al.* Glycosylation of Tetrabromobisphenol A in Pumpkin. *Environ. Sci. Technol.*

(2019) doi:10.1021/acs.est.9b02122.

20. Yu, M. *et al.* Evaluation and reduction of the analytical uncertainties in GC-MS analysis

using a boundary regression model. *Talanta* **164**, 141–147 (2017).

21. Djoumbou-Feunang, Y. *et al.* BioTransformer: a comprehensive computational tool for small

molecule metabolism prediction and metabolite identification. *J. Cheminformatics* **11**, 2

(2019).

22. Wishart, D. *et al.* T3DB: the toxic exposome database. *Nucleic Acids Res.* **43**, D928–D934

(2015).

23. Mathé, E. A. *et al.* Noninvasive urinary metabolomic profiling identifies diagnostic and

prognostic markers in lung cancer. *Cancer Res.* **74**, 3259–3270 (2014).

TOC

352