Drosophila **models of pathogenic copy-number variant genes show global and**

non-neuronal defects during development

Short title: Non-neuronal defects of fly homologs of CNV genes

Tanzeen Yusuff[1,4], Matthew Jensen[1,4], Sneha Yennawar[1,4], Lucilla Pizzo[1], Siddharth

Karthikeyan[1], Dagny J. Gould[1], Avik Sarker[1], Yurika Matsui[1,2], Janani Iyer[1], Zhi-Chun Lai[1,2],

and Santhosh Girirajan[1,3]*

1. Department of Biochemistry and Molecular Biology, Pennsylvania State University,

9   University Park, PA 16802

2. Department of Biology, Pennsylvania State University, University Park, PA 16802

3. Department of Anthropology, Pennsylvania State University, University Park, PA 16802

[4] contributed equally to work

*Correspondence:

Santhosh Girirajan, MBBS, PhD

205A Life Sciences Building

Pennsylvania State University

University Park, PA 16802

E-mail: sxg47@psu.edu

Phone: 814-865-0674

ABSTRACT

While rare pathogenic copy-number variants (CNVs) are associated with both neuronal and non-neuronal phenotypes, functional studies evaluating these regions have focused on the molecular basis of neuronal defects. We report a systematic functional analysis of non-neuronal defects for homologs of 59 genes within ten CNVs and 20 neurodevelopmental genes in *Drosophila*. Using wing-specific knockdown of 136 RNA interference lines, we identified qualitative and quantitative phenotypes in 72/79 homologs, including six lines with lethality and 21 lines with severe wing defects. Assessment of 66 lines for tissue-specific effects showed no correlation between the severity of wing and eye-specific defects. We observed disruptions in cell proliferation and apoptosis in larval wing discs for 23/27 homologs, and altered Wnt, Hedgehog and Notch signaling for 9/14 homologs, including *AATF/Aatf*, *PPP4C/Pp4-19C*, and *KIF11/Klp61F*. These findings were further validated with differences in human tissue-specific expression and network connectivity of CNV genes. Our findings suggest that multiple genes within each CNV differentially affect both global and tissue-specific developmental processes within conserved pathways, and that their roles are not restricted to neuronal functions.

INTRODUCTION

Rare copy-number variants (CNV), or deletions and duplications in the genome, are associated with neurodevelopmental disorders such as autism, intellectual disability (ID), and schizophrenia [1,2]. While dosage alteration of CNV regions contribute predominantly to defects in nervous system development, several CNV-associated disorders also lead to early developmental features involving other organ systems [3,4], including cardiac defects [5,6], kidney malformations[7], craniofacial features [3], and skeletal abnormalities [8]. For example, the 1q21.1 deletion causes variable expression of multiple neuronal and non-neuronal phenotypes, including developmental delay, autism, and schizophrenia as well as craniofacial features, cataracts, cardiac defects, and skeletal abnormalities [9–11]. Additionally, while the 7q11.23 deletion associated with Williams-Beuren syndrome (WBS) causes neuropsychiatric and behavioral features, other non-neuronal phenotypes, including supravalvular aortic stenosis, auditory defects, hypertension, diabetes mellitus, and musculoskeletal and connective tissue anomalies, are also observed among the deletion carriers [12].

Despite the notable prevalence of non-neuronal phenotypes in CNV carriers, functional studies of CNV genes have primarily focused on detailed assessments of neuronal and behavioral phenotypes in model systems. For example, mouse models for the 16p11.2 deletion exhibited post-natal lethality, reduced brain size and neural progenitor cell count, motor and habituation defects, synaptic defects, and behavioral defects [13–15]. Similarly, mouse models for the 3q29 deletion showed decreased weight and brain size, increased locomotor activity and startle response, and decreased spatial learning and memory [16,17]. However, fewer studies have focused on detailed evaluation of non-neuronal phenotypes in functional models of CNV disorders. For example, Arbogast and colleagues evaluated obesity and metabolic changes in

16p11.2 deletion mice, which showed reduced weight and impaired adipogenesis [18]. While

Haller and colleagues showed that mice with knockdown of *MAZ,* a gene within the 16p11.2

deletion region, contribute to the genitourinary defects observed in individuals with the deletion

[19], mouse studies on other homologs of 16p11.2 genes, including *TAOK2*, *KCTD13*, and

MAPK3, have only focused on assessing neuronal defects [20–24]. Furthermore, Dickinson and

colleagues reported a high-throughput analysis of essential genes in mice and identified both

neuronal and non-neuronal phenotypes for individual gene knockouts, including more than 400

genes that lead to lethality [25]. While these efforts aided in implicating novel genes with human

disease, our understanding of how genes associated with neurodevelopmental disorders

contribute towards non-neuronal phenotypes is still limited. Therefore, a large-scale analysis of

non-neuronal phenotypes is necessary to identify specific candidate genes within CNV regions

and associated biological mechanisms that contribute towards these phenotypes.

Drosophila melanogaster is an excellent model system to evaluate homologs of

neurodevelopmental genes, as many developmental processes and signaling pathways are

conserved between humans and flies [26]. In fact, over 75% of human disease genes have

homologs in *Drosophila*, including many genes involved in cellular signaling processes [27,28].

We recently examined the contributions of individual *Drosophila* homologs of 28 genes within

the 16p11.2 and 3q29 deletion regions towards specific neurodevelopmental phenotypes,

including rough eye phenotypes and defects in climbing ability, axon targeting, neuromuscular

junction, and dendritic arborization [29,30]. While these findings implicated multiple genes

within each CNV region towards conserved cellular processes in neuronal tissues, the conserved

role of these genes in non-neuronal tissues is not well understood. The *Drosophila* wing is an

effective model system to evaluate such developmental defects, as key components of conserved

signaling pathways, such as Notch, epidermal growth factor receptor (EGFR), Hegdehog, and

Wnt pathways, were identified using fly wing models [31–37]. Although fly wing phenotypes

cannot be directly translated to human phenotypes, defects observed in fly wings can be used to

assess how homologs of human disease genes alter conserved cellular and developmental

mechanisms. For example, Wu and colleagues showed that overexpression of the *Drosophila*

homolog for *UBE3A*, associated with Angelman syndrome, leads to abnormal wing and eye

morphology defects [38]. Furthermore, *Drosophila* mutant screens for developmental

phenotypes, including wing defects, were used to identify conserved genes for several human

genetic diseases, including Charcot-Marie-Tooth disease and syndromic microcephaly [39].

Kochinke and colleagues also recently performed a large-scale screening of ID-associated genes,

and found an enrichment of wing trichome density and missing vein phenotypes in ID genes

compared to control gene sets [40]. Hence, the fly wing provides a model system that is ideal for

evaluating the contributions of individual homologs of CNV genes towards cellular and

developmental defects.

In this study, we tested tissue-specific and cellular phenotypes of 79 fly homologs of

human genes within ten pathogenic CNV regions and genes associated with neurodevelopmental

disorders. Particularly, we used the adult fly wing to evaluate phenotypes in a non-neuronal

tissue, and observed a wide range of robust qualitative and quantitative wing phenotypes among

the 136 RNA interference (RNAi) lines tested in our study, including size defects, ectopic and

missing veins, severe wrinkling, and lethality. Further analysis of cellular phenotypes revealed

disruptions in conserved developmental processes in the larval imaginal wing disc, including

altered levels of cell proliferation and apoptosis as well as altered expression patterns in the Wnt,

Hedgehog, and Notch signaling pathways. However, we found no correlation in the severity of

phenotypes observed with wing and eye-specific knockdown. Our findings were further

supported by differences in expression patterns and network connectivity of human CNV genes

across different tissues. Our analysis emphasizes the importance of multiple genes within each

CNV region towards both global and tissue-specific developmental processes.

RESULTS

Knockdown of fly homologs of CNV genes contribute to a range of wing defects

Using an RNAi based analysis driven by the bx^{MS1096}-*GAL4* wing-specific driver, we tested a

total of 136 RNAi lines for 59 homologs of genes within pathogenic CNV regions (chromosomal

locations 1q21.1, 3q29, 7q11.23, 15q11.2, 15q13.3, 16p11.2, distal 16p11.2, 16p12.1, 16p13.11,

and 17q12) and 20 homologs of genes associated with neurodevelopmental disorders (**Supp.**

Data 1). Fly homologs of these genes were identified using the DIOPT orthology prediction tool

[41]. We list both the human gene name and the fly gene name for each tested gene as *HUMAN*

GENE/Fly gene (i.e. *KCTD13/CG10465*) as well as the human CNV region for context at first

instance. We scored 20-25 adult wings for five distinct wing phenotypes in each non-lethal

RNAi line, including wrinkled wing, discoloration, ectopic veins, missing veins, and bristle

planar polarity phenotypes (**Fig. 1A; Supp. Data 2**). We first categorized each wing phenotype

based on their severity and performed k-means clustering analysis to categorize each RNAi line

by their overall phenotype severity (**Fig. 1B-C**). We observed four clusters of RNAi lines: 75

lines with no observable qualitative phenotypes (55.2%), 24 lines with mild phenotypes (17.7%),

10 lines with moderate phenotypes (7.4%), 21 lines with severe phenotypes (15.4%), and 6 lines

with lethal phenotypes (4.4%), including *ACACA/ACC* within 17q12, *DLG1/dlg1* within 3q29,

and *STX1A/Syx1A* within 7q11.23 (**Fig. 1B-D; Supp. Data 2**). We observed severe wrinkled

wing phenotypes for 13/79 fly homologs, including *PPP4C/Pp4-19C* within 16p11.2,

ATXN2L/Atx2 within distal 16p11.2, *AATF/Aatf* within 17q12, and *MFI2/Tsf2* within 3q29 (**Fig.**

2A-B, Supp. Data 3). Interestingly, seven out of ten CNV regions contained at least one

homolog that showed lethality or severe wing phenotypes, and five CNV regions (3q29, 16p11.2,

distal 16p11.2, 16p12.1, and 17q12) had multiple homologs showing lethality or severe wing

phenotypes (**Fig. 2A, Supp. Data 3**). For example, RNAi lines for both *UQCRC2/UQCR-C2* and

POLR3E/Sin within 16p12.1 showed lethality. Within the 3q29 region, *NCBP2/Cbp20* and

MFI2/Tsf2 showed severe phenotypes while *DLG1/dlg1* showed lethality. In contrast, 12/20

known neurodevelopmental genes showed no observable wing phenotypes, suggesting that these

genes could be responsible for neuronal-specific phenotypes (**Fig. 2B, Supp. Data 3**). We note

that 18/79 fly homologs showed discordant phenotypes between two or more RNAi lines for the

same gene, which could be due to differences in expression of the RNAi construct among these

lines (**Supp. Data 3**).

Certain qualitative phenotypes exhibited higher frequency in males compared to females.

For example, discoloration (87 lines in males compared to 56 lines in females; $p=1.315\times10^{-4}$,

two-tailed Fisher's exact test) and missing vein phenotypes (92 lines in males compared to 29

lines in females; $p=2.848\times10^{-16}$, two-tailed Fisher's exact test) at any degree of severity were

more commonly observed in males than females (**Supp. Data 2**). In particular, 25/92 lines in

males (compared to 1/29 in females) showed a total loss of the anterior crossvein (ACV) (**Supp.**

Data 2). We further identified 17 RNAi lines that were lethal in males with wing-specific

knockdown of fly homologs. While higher frequencies of wing phenotypes in males could be

due to a sex-specific bias of developmental phenotypes, the increased severity we observed in

males is most likely due to a stronger RNAi knockdown caused by an X-linked dosage

compensation, as the bx^{MS1096}-*GAL4* driver is inserted on the fly X chromosome [42,43].

Next, we measured the total adult wing area and the lengths of six veins (longitudinal L2,

L3, L4, L5, ACV, and posterior crossvein or PCV) in the adult wing for each of the tested RNAi

lines that did not show lethality (or severe wrinkled phenotypes for vein length measurements)

(**Fig. 3A**). Overall, we identified significant wing measurement changes for 89 RNAi lines

compared to controls, which included lines that did not have an observable qualitative wing

phenotype (**Fig. 1D**). A summary of L3 vein lengths is presented in **Fig. 3B**, and the

measurements for the remaining five veins are presented in **Supp. Figure 1** and **Supp. Data. 2**.

We found that 33/61 of the homologs (54%) showed significant changes in L3 vein length,

including 20 homologs with longer vein lengths and 13 homologs with shorter vein lengths

(**Supp. Data 3**). Additionally, 41/74 of the fly homologs (55%) showed changes in wing area

(**Supp. Data 3**), including 36 homologs which showed smaller wing areas and five homologs

showed larger wing areas compared to controls (**Supp. Data 3**). For example, both homologs of

genes within 1q21.1 region, *BCL9*/*lgs* and *FMO5/Fmo-2*, showed decreased wing area and vein

length, potentially mirroring the reduced body length phenotype observed in mouse models of

the deletion [44] (**Fig. 3B-C**). In addition, *PAK2/Pak* within 3q29*, TBX1/org-1* within 22q11.2,

autism-associated *CHD8/kis*, and microcephaly-associated *ASPM/asp* also showed smaller wing

areas and vein lengths (**Fig. 3B-C**). In contrast, *TRPM1/Trpm* within 15q13.3 and the cell

proliferation gene *PTEN/Pten* [45] both showed larger wing areas and vein lengths (**Fig. 3B-C**).

Furthermore, we identified eight homologs that showed no qualitative wing phenotypes but had

significant changes in wing areas and vein lengths (**Supp. Data 3**), including *CCDC101/Sgf29* in

distal 16p11.2*, FMO5/Fmo-2, TRPM1/Trpm, DHRS11/CG9150* in 17q12, and *NSUN5/Nsun5* in

7q11.23 (**Fig. 3B-C; Supp. Data 3**). These results indicate that homologs of certain CNV genes

may influence variations in size without causing adverse wing phenotypes, and may be

specifically implicated towards cellular growth mechanisms.

Homologs of CNV genes show global and tissue-specific effects during development

We previously showed that many of the same fly homologs of CNV genes that showed wing

defects in the current study also contributed towards neuronal phenotypes in the fly eye [29,30],

suggesting a role for these genes in global development. We therefore performed ubiquitous and

eye-specific knockdown of fly homologs to assess tissue-specific effects in comparison to the

wing phenotypes. First, we used the *da-GAL4* driver at 25°C to drive ubiquitous knockdown of

RNAi lines for 31 homologs of CNV genes, including 19 that were previously published [29,30],

and observed complete or partial lethality at larval and pupal stages with knockdown of 10/31

homologs (32.3%) (**Fig. 4A**). Lethal phenotypes have also been documented for 43/130

knockout mouse models of individual CNV genes as well as for the entire deletion (**Supp. Data

4**). For example, mouse models heterozygous for the 16p11.2 deletion showed partial neonatal

lethality, while knockout mouse models of four individual genes within the 16p11.2 region,

including *Ppp4C$^{-/-}$* and *Kif22$^{-/-}$*, showed embryonic lethality [13,46,47]. In our study, the

DLG1/dlg1 line that showed lethality with wing-specific knockdown also exhibited larval

lethality with ubiquitous knockdown, indicating its role in global development (**Fig. 4A**). In

addition, six homologs that showed severe wing phenotypes also showed larval or pupal lethality

with ubiquitous knockdown, including *ALDOA/Ald* and *PPP4C/Pp4-19C* within 16p11.2 and

ATXN2L/Atx2 and *TUFM/mEFTu1* within distal 16p11.2 (**Fig. 4A**). The remaining homologs

that showed lethality with ubiquitous knockdown showed at least a mild qualitative or

quantitative wing phenotype.

We next compared the phenotypes observed with wing-specific knockdown of fly

homologs to their corresponding eye-specific knockdowns to evaluate tissue-specific effects. To

quantitatively assess the phenotypic severity of cellular defects with eye-specific knockdown of

fly homologs, we developed a tool called *Flynotyper* [48] that determines the degree of

disorganization among the ommatidia in the adult eye. We analyzed phenotypic scores obtained

from *Flynotyper* for 66 RNAi lines of 45 fly homologs, including from previously-published

datasets [29,30,48]. We found that 37/45 homologs (82.2%) exhibited both eye and wing-

specific defects (**Fig. 4B, Supp. Fig. 2, Supp. Data 5**). Two homologs with significant eye

phenotypes did not show any wing phenotypes, including *SPNS1/spin* within distal 16p11.2 and

microcephaly-associated *SLC25A19/Tpc1* [49], while five homologs only showed wing-specific

phenotypes, including *CDIPT/Pis* and *YPEL3/CG15309* within 16p11.2, *FBXO45/Fsn* and

OSTalpha/CG6836 within 3q29, and *UQCRC2/UQCR-C2* (**Fig. 4B, Supp. Fig. 2**). In particular,

UQCRC2/UQCR-C2 showed lethality with wing-specific knockdown, suggesting potential

tissue-specific effects of this gene in non-neuronal cells (**Fig. 4B**). While most homologs

contributed towards both eye and wing-specific phenotypes, we observed a wide range of

severity in eye phenotypes that did not correlate with the severity of quantitative or qualitative

wing phenotypes (**Fig. 4C**). For example, *TUFM/mEFTu1* showed a severe wing phenotype but

only a mild increase in eye phenotypic score, while *SH2B1/Lnk,* also within the distal 16p11.2

region, showed severe rough eye phenotypes but only a mild increase in wing size (**Fig. 4D**).

Similarly, *BCL9/lgs* also showed opposing tissue-specific effects with mild qualitative wing

phenotype and severe eye phenotype, suggesting that the role of these homologs towards

development differs across tissue types.

CNV genes show variable expression across different tissues in flies and humans

To assess how expression levels of CNV genes vary across different tissues, we first examined

the expression patterns of fly homologs in larval and adult tissues using the FlyAtlas Anatomical

Microarray dataset [50]. We found that 76/77 homologs with available data were expressed in at

least one larval and adult tissue (**Supp. Fig. 3, Supp. Data 6**). In general, we did not observe a

correlation between wing phenotype severity and expression patterns of homologs in larval or

adult tissues (**Fig. 5A**). For example, 58/77 homologs (75.3%) showed ubiquitous larval

expression, including both fly homologs that showed no qualitative wing phenotypes, such as

KCTD13/CG10465 within 16p11.2 and *FBXO45/Fsn*, and those with severe wing phenotypes,

such as *PPP4C/Pp4-19C* and *NCBP2/Cbp20* (**Fig. 5A, Supp. Fig. 3**). Furthermore, 30/39

homologs (76.9%) that showed eye phenotypes also had ubiquitous larval expression, providing

further support to the observation that genes causing neuronal phenotypes may also contribute to

developmental phenotypes in other tissues (**Supp. Data 5**). Of note, 9/77 homologs (11.7%) did

not have any expression in the larval central nervous system, including *FMO5/Fmo-2*,

BDH1/CG8888 within 3q29, and *TBX6/Doc2* within 16p11.2 (**Fig. 5A, Supp. Fig. 3**). However,

we observed wing phenotypes for 8/9 of these homologs, suggesting that they may contribute to

tissue-specific phenotypes outside of the nervous system. Except for the epilepsy-associated

SCN1A/para [51], which was exclusively expressed in both the larval central nervous system

(CNS) and adult brain tissues, other tested neurodevelopmental genes were also expressed in

non-neuronal tissues (**Fig. 5A**).

We further used the GTEx Consortium dataset [52] to examine tissue-specific expression

of 150 human CNV and known neurodevelopmental genes across six tissues including brain,

heart, kidney, lung, liver, and muscle. We found 121 genes that were expressed in at least one

adult tissue, including 49 genes (32.7%) that showed ubiquitous expression across all six tissues

(**Supp. Data 6**). Of the 112 genes expressed in non-neuronal tissues, 34 did not have any

neuronal expression, including *TBX1*, *FMO5* and *GJA5* within 1q21.1, and *ATP2A1* within distal

16p11.2 (**Fig. 5B**, **Supp. Data 6**). *FMO5* and *TBX1* also showed non-neuronal expression in

Drosophila tissues, suggesting that their tissue-specific expression is highly conserved (**Fig. 5A**).

Other genes showing ubiquitous expression also had preferentially high expression for specific

non-neuronal tissues, including *ALDOA* and *UQCRC2* for muscle and heart (**Fig. 5B**). In

contrast, we found nine genes that were expressed only in the adult brain, including *FAM57B*

and *DOC2A* within 16p11.2, as well as *SCN1A*, which showed similar CNS-only expression in

Drosophila tissues (**Fig. 5B**, **Supp. Data 6**).

Knockdown of fly homologs of CNV genes lead to disruption of cellular processes

The disruption of basic cellular processes in neuronal cells, such as cell proliferation and

apoptosis, have been implicated in neurodevelopmental disorders [53–55]. We previously

identified defects in cell proliferation among photoreceptors neurons in larval eye discs with

knockdown of 16p11.2 homologs, as well as increased apoptosis with knockdown of a subset of

3q29 homologs [29,30]. Here, we explored how these basic cellular processes are altered in non-

neuronal cells, specifically in the developing wing disc, with knockdown of homologs of CNV

genes. We targeted 27 fly homologs that showed a range of adult wing phenotypes for changes in

cell proliferation and apoptosis, using anti-phospho-Histone H3 Ser10 (pH3) and anti-

Drosophila caspase-1 (dcp1), respectively, in the third instar larval wing discs. We identified

23/27 homologs that showed significant increases in apoptotic cells compared to controls,

including seven homologs, such as *PPP4C/Pp4-19C*, *ATXN2L/Atx2*, and *AATF/Aatf* , which

showed dcp1 staining across the entire larval wing pouch (**Fig. 6A-B, Supp. Figs. 4-5, Supp.**

Data 7). In addition, 16/27 genes showed decreased levels of proliferation, including eight

homologs which also showed apoptosis defects, such as *CYFIP1/Sra-1* within 15q11.2,

SH2B1/Lnk, and the microcephaly gene *KIF11/Klp61F* (**Fig. 6A and 6C, Supp. Figs. 4-5, Supp.**

Data 7). All six of the tested homologs with severe adult wing phenotypes showed both

increased apoptosis and decreased proliferation (**Supp. Data 7**). Similarly, 3/4 homologs of

genes showing lethality with wing-specific knockdown also showed defects in apoptosis or

proliferation, with the exception of *ACACA/ACC* (**Supp. Figure 4**, **Supp. Data 7**). As $bx^{MS1096-}$

GAL4 is located on the X-chromosome, we expected to see more severe defects in males

compared with females with knockdown of homologs due to the X-linked dosage compensation

[42,43]. However, knockdown of 3/11 tested homologs with sex-specific differences in adult

wing phenotypes, including *BCL9/lgs*, *CYFIP1/Sra-1*, and *DNAJC30/CG11035* within 7q11.23,

showed significantly decreased levels of cell proliferation in females but no change for males

compared to their respective controls, suggesting a sex-specific effect of these genes for cell

proliferation (**Supp. Fig. 5, Supp. Data 7**). Overall, our results suggest that cell proliferation and

apoptosis are disrupted by reduced expression of homologs of CNV genes in both neuronal and

non-neuronal tissues.

Knockdown of homologs of CNV genes disrupt conserved signaling pathways

Several conserved signaling pathways that are active in a spatial and temporal manner in the

larval wing disc, such as Wnt, Hedgehog, BMP, and Notch signaling, regulate the anterior-

posterior (A/P) and dorsal-ventral (D/V) boundaries to determine accurate morphology and vein

patterning in the adult wing [36,37,56–58]. For example, Wnt and Notch signaling pathways

both act along the D/V boundary to determine cell fate [59,60], while Hedgehog signaling is

dependent upon expression of both engrailed in the posterior compartment and patched along the

A/P border [61,62]. Furthermore, O'Roak and colleagues showed that genes identified from *de*

novo mutations in patients with autism are linked to β-catenin/Wnt pathway [63]. In addition,

familial loss-of-function mutations in the human hedgehog signaling pathway gene *PTCH1* are

implicated in basal cell nevus syndrome, leading to basal cell carcinoma [64,65].

Based on adult wing phenotypes and disruptions to cellular processes, we next tested

whether knockdown of 14 fly homologs disrupt conserved signaling pathways in the third instar

larval wing disc (**Supp. Data 7**). In particular, we evaluated the role of Wnt, Hedgehog, and

Notch signaling pathways by testing the expression patterns of four key proteins within these

pathways, including wingless (Wnt), patched (Hedgehog), engrailed (Hedgehog), and delta

(Notch). We found that 9/14 homologs, including 8/10 homologs showing severe wing

phenotypes or lethality, exhibited disruptions in at least one signaling pathway. For example, five

homologs with severe or lethal phenotypes showed disruptions of all four signaling pathways,

including *AATF/Aatf*, *NCBP2/Cbp20*, *POLR3E/Sin*, *PPP4C/Pp4-19C*, and *KIF11/Klp61F* (**Fig.**

7, Supp. Data 7). Our observations are in concordance with previous findings by Swarup and

colleagues, who showed that *PPP4C/Pp4-19C* is a candidate regulator of Wnt and Notch

signaling pathways in *Drosophila* larval wing discs [66]. Furthermore, two genes from the 3q29

region, *DLG1/dlg1* and *MFI2/Tsf2*, showed altered expression patterns for delta and patched but

not for engrailed, indicating that they selectively interact with the Hedgehog as well as Notch

signaling pathway (**Supp. Fig. 6**). In fact, Six and colleagues showed that Dlg1 directly binds to

the PDZ-binding domain of Delta1 [67]. In contrast, *ACACA/ACC* and *UQCRC2/UQCR-C2*

showed no changes in expression patterns for any of the four signaling proteins tested,

suggesting that the observed lethality could be due to other cellular mechanisms (**Supp. Fig. 6**).

We conclude that several homologs disrupt the expression of key proteins in signaling pathways

in the developing larval wing discs, potentially accounting for the developmental defects

observed in the adult wings.

Connectivity patterns of CNV genes vary across human tissue-specific networks

We examined patterns of connectivity for the nine candidate genes, based on the disruptions of

signaling pathways identified in the developing *Drosophila* wing discs, within the context of

human brain, heart, and kidney-specific gene interaction networks [68]. These tissue-specific

networks were constructed using Bayesian classifier-generated probabilities for pairwise genetic

interactions based on co-expression data [68]. We calculated the lengths of the shortest paths

between each candidate gene and 267 Wnt, Notch, and Hedgehog pathway genes in each

network as a proxy for connectivity (**Supp. Data 8**). In all three networks, each of the candidate

genes were connected to a majority of the tested signaling pathway genes (**Fig. 8A, Supp. Fig.**

7). Interestingly, we observed a higher connectivity (i.e. shorter path distances) between

candidate genes and Wnt and Hedgehog pathway genes in the brain-specific network compared

to the heart and kidney-specific networks (**Fig. 8B**). We further identified enrichments for genes

involved in specific biological processes among the connector genes that were located in the

shortest paths within neuronal and non-neuronal tissue-specific networks (**Fig. 8C, Supp. Data**

8). For example, axon-dendrite transport, dopaminergic signaling, and signal transduction

functions were enriched among connector genes only for the brain-specific network, while

organelle organization and protein ubiquitination were enriched among connector genes only for

kidney and heart networks (**Fig. 8C**). However, several core biological processes, such as cell

cycle, protein metabolism, transcriptional regulation, and RNA processing/splicing, were

enriched among connector genes within all three tissue-specific networks (**Fig. 8C**). Our analysis

highlights that human CNV genes potentially interact with developmental signaling pathways in

a ubiquitous manner, but may affect different biological processes in neuronal and non-neuronal

tissues.

DISCUSSION

We used the *Drosophila* wing as a model to assess how homologs of key CNV genes contribute towards non-neuronal phenotypes. We tested fly homologs of 79 genes and identified multiple homologs within each CNV region that exhibited strong phenotypes indicative of developmental disruptions. Several themes have emerged from our study highlighting the importance of fly homologs of CNV genes towards both global and tissue-specific phenotypes.

First, we found that fly homologs of CNV genes contribute towards developmental phenotypes through ubiquitous roles in neuronal and non-neuronal tissues. Although we did not study models for the entire CNV, nearly all individual fly homologs of CNV genes contribute to wing-specific developmental defects. It is likely that these genes may also contribute to additional phenotypes in other tissues that we did not assess. In fact, a subset of these homologs also showed early lethality with ubiquitous knockdown in addition to severe or lethal wing-specific phenotypes. However, we found no correlation between the severity of the eye and wing phenotypes, suggesting tissue-specific effects of these homologs towards developmental defects. In contrast, fly homologs of known neurodevelopmental genes generally showed milder wing phenotypes compared with eye phenotypes, indicating a more neuronal role for these genes. While our study only examined a subset of CNV genes with *Drosophila* homologs, phenotypic data from knockout mouse models also support a global developmental role for individual CNV genes. In fact, 44/130 knockout models of CNV genes within the Mouse Genome Informatics (MGI) database [69] exhibited non-neuronal phenotypes, including 20 homologs of CNV genes that showed both neuronal and non-neuronal phenotypes (**Supp. Data 4**). For example, knockout mouse models of $Dlg1^{-/-}$ show defects in dendritic growth and branching in the developing nervous system, in addition to craniofacial features and multiple kidney and urinary tract defects

[70–73]. Furthermore, Chapman and colleagues showed that knockout of $Tbx6^{-/-}$ caused defects

in mesodermal and neuronal differentiation early in development, leading to abnormal vascular,

tail bud, and neural tube morphology [74]. These observations further support our findings that

most fly homologs of CNV genes have a global role in development that could account for the

observed non-neuronal defects.

Second, based on tissue-specific phenotypes, we identified fly homologs of CNV genes

that are key regulators of conserved cellular processes important for development. For example,

9/10 homologs with severe or lethal adult wing phenotypes also exhibited defects in cell

proliferation and apoptosis during development. In fact, we found concordance between cellular

processes affected by wing and eye-specific knockdown of homologs of genes within 16p11.2

and 3q29 regions, including decreased proliferation for *MAPK3/rl* and increased apoptosis for

NCBP2/Cbp20 and *DLG1/dlg1* [29,30]. While eye-specific knockdown of *BDH1/CG8888*

showed decreased cell proliferation in larval eye discs [30], we found increased cell proliferation

with wing-specific knockdown, suggesting a tissue-specific effect for this gene. Notably, at least

one fly homolog per CNV region showed defects in cell proliferation or apoptosis, suggesting

these cellular processes are important for development in both neuronal and non-neuronal

tissues. For example, *ATXN2L/Atx2*, *SH2B1/Lnk*, and *CCDC101/Sgf29* each showed decreased

proliferation and increased apoptosis, suggesting a potential shared cellular mechanism for

several genes within the distal 16p11.2 deletion. Furthermore, a subset of these genes also

disrupted multiple signaling pathways, indicating a potential role for these homologs as key

regulators of developmental processes. We specifically identified five homologs whose

knockdown caused disruptions of Wnt, Notch, and hedgehog signaling pathways. Each of these

genes have important roles in cell cycle regulation, apoptosis, transcription, or RNA processing,

based on Gene Ontology annotations [75,76]. In fact, we found that the RNA transport protein

NCBP2/Cbp20 [77], which we recently identified as a key modifier gene for the 3q29 deletion

[30], interfaced with all three signaling pathways. Furthermore, *AATF* disrupts apoptosis and

promotes cell cycle progression through displacement of HDAC1 [78–80], while *PPP4C*

promotes spindle organization at the centromeres during mitosis [81]. While we only evaluated

the role of these genes towards development in a single fly tissue, our additional analysis of

human gene interaction networks showed strong connectivity between the CNV genes and

signaling pathways in multiple neuronal and non-neuronal human tissues. In fact, cell cycle

genes were enriched among the connector genes in all three tissue-specific networks, further

emphasizing the role of cell cycle processes towards developmental phenotypes. Notably, we

also observed certain biological processes enriched among connector genes that were specific to

neuronal or non-neuronal tissues, indicating that genes within CNV regions may affect different

biological processes in a tissue-specific manner.

401   *Overall*, we show that fly homologs of most CNV genes contribute towards global

developmental phenotypes, although exactly how they contribute toward such phenotypes varies

between neuronal and non-neuronal tissues. Previous functional studies for CNV disorders have

focused primarily on identifying candidate genes for the observed neuronal phenotypes. In this

study, we identified several homologs of CNV genes that are responsible for non-neuronal

defects, as well as novel associations between these homologs and conserved biological

processes and pathways. We therefore propose that multiple genes within each CNV region

differentially disrupt conserved cellular pathways and biological processes in neuronal versus

non-neuronal tissues during development (**Fig. 9**). These results are in line with a multigenic

model for CNV disorders, as opposed to models where individual causative genes are

responsible for specific phenotypes [29,30,82]. Our study further exemplifies the utility of

evaluating non-neuronal phenotypes in addition to neuronal phenotypes in functional models of

individual genes and CNV regions associated with developmental disorders, including future

studies in mammalian or cellular model systems. Further studies exploring how CNV genes

interact with each other and with other developmental pathways could more fully explain the

conserved mechanisms underlying global developmental defects and identify potential

therapeutic targets for these disorders.

MATERIALS AND METHODS

Fly stocks and genetics

We tested 59 *Drosophila* homologs for 130 human genes that span across 10 pathogenic CNV

regions associated with neurodevelopmental disorders (1q21.1, 3q29, 7q11.23, 15q11.2, 15q13.3,

16p11.2, distal 16p11.2, 16p12.1, 16p13.11, and 17q12) [83] (**Supp. Data 1**). In addition, we

evaluated fly homologs of 20 human genes known to be in involved in neurodevelopmental

disorders [48,84] (**Supp. Data 1**). These include genes involved in beta-catenin signaling

pathway (5 genes), core genes implicated in neurodevelopmental disorders (8 genes), and genes

associated with microcephaly (7 genes) [85]. We used the DRSC Integrative Ortholog Prediction

Tool (DIOPT, v.7.1) to identify the fly homologs for each human gene [41] (**Supp. Data 1**).

To knockdown individual genes in specific tissues, we used RNA interference (RNAi)

and the *UAS-GAL4* system (**Fig. 1A**), a well-established tool that allows for tissue-specific

expression of a gene of interest [86]. RNAi lines were obtained from Vienna *Drosophila*

Resource Center (VDRC) that include both GD and KK lines. We tested a total of 136 lines in

our final data analysis (**Supp. Data 9**), after eliminating KK lines with additional insertion that

drives the overexpression of the Tiptop (*tio*) transcription factor [87,88]. A complete list of stock

numbers and full genotypes for all RNAi lines used in this study is presented in **Supp. Data 9**.

We used the bx^{MS1096}-*GAL4/FM7c;;UAS-Dicer2/TM6B* driver for wing-specific knockdown and

w^{1118};*GMR-GAL4*;*UAS-Dicer2* driver (Claire Thomas, Penn State University) for eye-specific

knockdown of RNAi lines. Ubiquitous knockdown experiments were performed using the *w;da-*

GAL4;+ driver (Scott Selleck, Penn State University). For all experiments, we used appropriate

GD (w^{1118}, VDRC# 60000) or KK (*y,w^{1118}; P{attP,y$^+$,w$^{3`}$}*, VDRC# 60100) lines as controls to

compare against lines with knockdown of individual homologs. All fly lines were reared on

standard yeast *Drosophila* medium at room temperature. All crosses were set and maintained at

25°C, except for the eye knockdown experiments which were maintained at 30°C.

Phenotypic analysis of adult wing images

Adult progeny were isolated from crosses between RNAi lines and bx^{MS1096}-*GAL4* driver shortly

after eclosion, and kept at 25°C until day 2-5 (**Fig. 1A**). At that point, the progeny were frozen at

-80°C, and were then moved to -20°C prior to imaging and storage. Approximately 20-25

progeny, both male and female, were collected for each RNAi line tested. The adult wings were

plucked from frozen flies and mounted on a glass slide. The slides were covered with a coverslip

and sealed using clear nail polish. Adult wing images were captured using a Zeiss Discovery

V20 stereoscope (Zeiss, Thornwood, NY, USA), with a ProgRes Speed XT Core 3 camera and

CapturePro v.2.8.8 software (Jenoptik AG, Jena, Germany) at 40X magnification.

For each non-lethal RNAi line, we scored the adult wing images for five qualitative

phenotypes, including wrinkled wing, discoloration, missing veins, ectopic veins, and bristle

planar polarity defects, on a scale of 1 (no phenotype) to 5 (lethal) (**Fig. 2C**). Lines showing

severely wrinkled wings or lethality were scored as 4 (severe) or 5 (lethal) for all five

phenotypes. We calculated the frequency of each phenotypic score (i.e. mild bristle polarity,

moderate discoloration) across all of the wing images for each line (**Fig. 2A-B**), and then

performed k-means clustering of these values to generate five clusters for overall wing

phenotypes (**Fig. 1C**). For quantitative analysis of wing phenotypes, we used the Fiji ImageJ

software [89] to calculate the wing area using the Measure Area tool, and calculated the lengths

of longitudinal veins L2, L3, L4, and L5 as well as the anterior and posterior crossveins (ACV

and PCV), by tracing individual veins using the Segmented Line tool (**Fig. 3A, Supp. Data 2**).

We determined discordant homologs when RNAi lines for the same homologs showed

inconsistent wing phenotypes. For each homolog with multiple RNAi lines, we checked

discordance among RNAi lines for no phenotype versus any qualitative or quantitative

phenotypes, followed by discordance for small or large wing measurement phenotypes (**Supp.**

Data 3).

Phenotypic analysis of adult eye images

We crossed RNAi lines with *GMR-GAL4* to achieve eye-specific knockdown of homologs of

CNV and known neurodevelopmental genes. Adult 2-3-day old female progenies from the

crosses were collected, immobilized by freezing at -80°C, and then moved to -20°C prior to

imaging and storage. Flies were mounted on Blu-tac (Bostik Inc, Wauwatosa, WI, USA) and

imaged using an Olympus BX53 compound microscope with LMPLan N 20X air objective using

a DP73 c-mount camera at 0.5X magnification (Olympus Corporation, Tokyo, Japan). CellSens

Dimension software (Olympus Corporation, Tokyo, Japan) was used to capture the eye images,

which were then stacked using the Zerene Stacker software (Zerene Systems LLC, Richland,

WA, USA). All eye images presented in this study are maximum projections of 20 consecutive

optical z-sections, at a z-step size of 12.1μm. Finally. we used our computational method called

Flynotyper (https://flynotyper.sourceforge.net) to quantify the degree of rough eye phenotypes

present due to knockdown of homologs of CNV or neurodevelopmental genes [48]. *Flynotyper*

scores for homologs of 16p11.2 and 3q29, as well as select core neurodevelopmental genes, were

derived from our previous studies [29,30,48].

Immunohistochemistry

Wing imaginal discs from third instar larvae were dissected in 1X PBS. The tissues were fixed using 4% paraformaldehyde and blocked using 1% bovine serum albumin (BSA). The wing discs were incubated with primary antibodies using appropriate dilutions overnight at 4°C. We used the following primary antibodies: mouse monoclonal anti-pHistone3 (S10) (1:100 dilutions, Cell Signaling 9706L), rabbit polyclonal anti-cleaved *Drosophila* Dcp1 (Asp216) (1:100 dilutions, Cell Signaling 9578S), mouse monoclonal anti-Wingless (1:200 dilutions, DSHB, 4D4), mouse monoclonal anti-Patched (1:50 dilutions, DSHB, *Drosophila* Ptc/APA1), mouse monoclonal anti-Engrailed (1:50 dilutions, DSHB, 4D9), and mouse monoclonal anti-Delta (1:50 dilutions, DSHB, C594.9B). Following incubation with primary antibodies, the wing discs were washed and incubated with secondary antibodies at 1:200 dilution for two hours at room temperature. We used the following secondary antibodies: Alexa Fluor 647 dye goat anti-mouse (A21235, Molecular Probes by Invitrogen/Life Technologies), Alexa Fluor 568 dye goat anti-rabbit (A11036, Molecular Probes by Invitrogen/Life Technologies), and Alexa Fluor 568 dye goat anti-mouse (A11031, Molecular Probes by Invitrogen/Life Technologies). All washes and antibody dilutions were made using 0.3% PBS with Triton-X.

Third instar larvae wing imaginal discs were mounted in Prolong Gold antifade reagent with DAPI (Thermo Fisher Scientific, P36930) for imaging using an Olympus Fluoview FV1000 laser scanning confocal microscope (Olympus America, Lake Success, NY). Images were acquired using FV10-ASW 2.1 software (Olympus, Waltham, MA, USA). Composite z-stack images were analyzed using the Fiji ImageJ software [89]. To calculate the number of pH3 positive cells within the wing pouch area of the wing discs, we used the AnalyzeParticles function in ImageJ, while manual counting was used to quantify Dcp1 positive cells. We note

that cell proliferation and apoptosis staining for *NCBP2/Cbp20*, *DLG1/dlg1*, *BDH1/CG8888*, and

FBXO45/Fsn were previously published[30].

Statistical analysis

Significance for the wing area and vein length measurements, cell counts for proliferation and

apoptosis, and *Flynotyper* scores were compared to appropriate GD or KK controls using one-

tailed or two-tailed Mann-Whitney tests. P-values for each set of experiments were corrected for

multiple testing using Benjamini-Hochberg correction. All statistical and clustering analysis was

performed using R v.3.6.1 (R Center for Statistical Computing, Vienna, Austria). Details for the

statistical tests performed for each dataset are provided in **Supp. Data 10**.

Expression data analysis

We obtained tissue-specific expression data for fly homologs of CNV genes from the FlyAtlas

Anatomical Microarray dataset [50]. Raw FPKM (fragments per kilobase of transcript per

million reads) expression values for each tissue were categorized as follows: <10, no expression;

10-100, low expression; 100-500, moderate expression; 500-1000, high expression; and >1000,

very high expression (**Supp. Data 6**). The median expression among midgut, hindgut,

Malpighian tube, and (for adult only) crop tissues was used to represent the overall gut

expression. We similarly obtained human tissue-specific expression data for CNV genes from

the GTEx Consortium v.1.2 RNA-Seq datasets [52]. Median TPM (transcripts per million reads)

expression values for each tissue were categorized as follows: <3, no expression; 3-10, low

expression; 10-25, moderate expression; 25-100, high expression; and >100, very high

expression (**Supp. Data 6**). The median expression among all brain and heart sub-tissues was

used to represent brain and heart expression, while the median expression among all colon,

esophagus, small intestine, and stomach sub-tissues was used to represent digestive tract

expression. Preferential gene expression for a particular tissue within the GTEx dataset was

determined if the expression values for that tissue were greater than the third quartile of all tissue

expression values for that gene, plus 1.5 times the interquartile range. Venn diagrams were

generated using the Venny webtool (http://bioinfogp.cnb.csic.es/tools/venny) (**Supp. Fig. 3**).

Network analysis

We obtained human tissue-specific gene interaction networks for brain, heart, and kidney tissues

from the GIANT network database [68] within HumanBase (https://hb.flatironinstitute.org).

These networks were built by training a Bayesian classifier based on tissue-specific gene co-

expression datasets, which then assigned a posterior probability for interactions between each

pair of genes within the genome for a particular tissue. We downloaded the "Top edge" version

of each tissue-specific network, and extracted all gene pairs with posterior probabilities >0.2 to

create sub-networks containing the top ~0.5% tissue-specific interactions. Next, we identified the

shortest paths in each sub-network between human CNV genes whose fly homologs disrupted

signaling pathways in the larval wing disc and human genes within each disrupted pathway,

using the inverse of the posterior probability as weights for each edge in the network. Gene sets

from the human Notch (KEGG:map04330), Wnt (KEGG:map04310) and Hedgehog pathways

(KEGG:map04340) were curated from the Kyoto Encyclopedia of Genes and Genomes (KEGG)

pathway database [90]. Using the NetworkX Python package [91], we calculated the shortest

distance between each CNV gene and pathway gene, and identified connecting genes that were

within each of the shortest paths for the three tissue-specific networks. We further tested for

enrichment of Gene Ontology (GO) terms (PantherDB GO-Slim) among the connector genes

using the PantherDB Gene List Analysis tool [92]. Lists of the shortest paths and connector

genes in each tissue-specific network, as well as enriched GO terms for sets of connector genes,

are provided in **Supp. Data 8**. Gene networks were visualized using Cytoscape v.3.7.2 [93]

using an edge-weighted spring embedded layout.

DATA AVAILABILITY

All data supporting the findings of this study are available within the paper and its

supplementary information files.

CODE AVAILABILITY

All source code for data analysis in this manuscript, including scripts for k-means clustering of

fly phenotypes and network connectivity of CNV and developmental pathway genes, are

available on the Girirajan lab GitHub page at https://github.com/girirajanlab/CNV_wing_project.

ACKNOWLEDGEMENTS

The authors thank Drs. Scott Selleck and Claire Thomas for providing fly lines for the

experiments, and members of the Girirajan Lab for their helpful discussions and comments on

the manuscript. This work was supported by NIH R01-GM121907 and resources from the Huck

Institutes of the Life Sciences to S.G., and NIH T32-GM102057 to M.J.

CONTRIBUTIONS

T.Y., M.J., S.Y., and S.G. designed the study. T.Y., S.Y., L.P., S.K., D.J.G., A.S., Y.M., J.I., and

Z.C.L. performed the functional experiments. T.Y. and M.J. performed the expression and

network experiments. T.Y., M.J., and S.G. analyzed the data and wrote the manuscript with input

from all authors.

COMPETING OF INTERESTS

The authors declare that they have no competing interests.

REFERENCES

1. Girirajan S, Campbell CD, Eichler EE. Human Copy Number Variation and Complex Genetic Disease. Annu Rev Genet. 2011;45: 203–226. doi:10.1146/annurev-genet-102209-163544

2. Malhotra D, Sebat J. CNVs: harbingers of a rare variant revolution in psychiatric genetics. Cell. 2012;148: 1223–1241. doi:10.1016/j.cell.2012.02.039

3. Cooper GM, Coe BP, Girirajan S, Rosenfeld JA, Vu TH, Baker C, et al. A copy number variation morbidity map of developmental delay. Nat Genet. 2011;43: 838–846. doi:10.1038/ng.909

4. Zhang F, Gu W, Hurles ME, Lupski JR. Copy Number Variation in Human Health, Disease, and Evolution. Annu Rev Genomics Hum Genet. 2009;10: 451–481. doi:10.1146/annurev.genom.9.081307.164217

5. Greenway SC, Pereira AC, Lin JC, Depalma SR, Israel SJ, Mesquita SM, et al. De novo copy number variants identify new genes and loci in isolated sporadic tetralogy of Fallot. Nat Genet. 2009;41: 931–935. doi:10.1038/ng.415

6. Glessner JT, Bick AG, Ito K et al. Increased Frequency of De Novo Copy Number Variations in Congenital Heart Disease by Integrative Analysis of SNP Array and Exome Sequence Data. Circ Res. 2014;115: 884–896. doi:10.1161/CIRCRESAHA.115.304458.

7. Sanna-Cherchi S, Kiryluk K, Burgess KE, Bodria M, Sampson MG, Hadley D, et al. Copy-number disorders are a common cause of congenital kidney malformations. Am J Hum Genet. 2012;91: 987–997. doi:10.1016/j.ajhg.2012.10.007

8. Zahnleiter D, Uebe S, Ekici AB, Hoyer J, Wiesener A, Wieczorek D, et al. Rare copy number variants are a common cause of short stature. PLoS Genet. 2013;9: e1003365. doi:10.1371/journal.pgen.1003365

9. Mefford HC, Sharp AJ, Baker C, Itsara A, Jiang Z, Buysse K, et al. Recurrent rearrangements of chromosome 1q21.1 and variable pediatric phenotypes. N Engl J Med. 2008;359: 1685–99. doi:10.1056/NEJMoa0805384

10. Brunetti-Pierri N, Berg JS, Scaglia F, Belmont J, Bacino CA, Sahoo T, et al. Recurrent reciprocal 1q21.1 deletions and duplications associated with microcephaly or macrocephaly and developmental and behavioral abnormalities. Nat Genet. 2008;40: 1466–1471. doi:10.1038/ng.279

11. Christiansen J, Dyck JD, Elyas BG, Lilley M, Bamforth JS, Hicks M, et al. Chromosome 1q21.1 contiguous gene deletion is associated with congenital heart disease. Circ Res. 2004;94: 1429–35. doi:10.1161/01.RES.0000130528.72330.5c

12. Pober BR. Williams-Beuren syndrome. N Engl J Med. 2010;362: 239–52. doi:10.1056/NEJMra0903074

13. Horev G, Ellegood J, Lerch JP, Son Y-EE, Muthuswamy L, Vogel H, et al. Dosage-dependent phenotypes in models of 16p11.2 lesions found in autism. Proc Natl Acad Sci. 2011;108: 17076–17081. doi:10.1073/pnas.1114042108

14. Pucilowska J, Vithayathil J, Tavares EJ, Kelly C, Karlo JC, Landreth GE. The 16p11.2 Deletion Mouse Model of Autism Exhibits Altered Cortical Progenitor Proliferation and Brain Cytoarchitecture Linked to the ERK MAPK Pathway. J Neurosci. 2015;35: 3190–3200. doi:10.1523/JNEUROSCI.4864-13.2015

15. Portmann T, Yang M, Mao R, Panagiotakos G, Ellegood J, Dolen G, et al. Behavioral

abnormalities and circuit defects in the basal ganglia of a mouse model of 16p11.2
deletion syndrome. Cell Rep. 2014;7: 1077–1092. doi:10.1016/j.celrep.2014.03.036

16. Rutkowski TP, Purcell RH, Pollak RM, Grewenow SM, Gafford GM, Malone T, et al.
Behavioral changes and growth deficits in a CRISPR engineered mouse model of the
schizophrenia-associated 3q29 deletion. Mol Psychiatry. 2019. doi:10.1038/s41380-019-
0413-5

17. Baba M, Yokoyama K, Seiriki K, Naka Y, Matsumura K, Kondo M, et al. Psychiatric-
disorder-related behavioral phenotypes and cortical hyperactivity in a mouse model of
3q29 deletion syndrome. Neuropsychopharmacology. 2019;44: 2125–2135.
doi:10.1038/s41386-019-0441-5

18. Arbogast T, Ouagazzal A-M, Chevalier C, Kopanitsa M, Afinowi N, Migliavacca E, et al.
Reciprocal Effects on Neurocognitive and Metabolic Phenotypes in Mouse Models of
16p11.2 Deletion and Duplication Syndromes. PLOS Genet. 2016;12: e1005709.
doi:10.1371/journal.pgen.1005709

19. Haller M, Au J, O'Neill M, Lamb DJ. 16p11.2 transcription factor MAZ is a dosage-
sensitive regulator of genitourinary development. Proc Natl Acad Sci. 2018;115:
201716092. doi:10.1073/pnas.1716092115

20. Yadav S, Oses-Prieto JA, Peters CJ, Zhou J, Pleasure SJ, Burlingame AL, et al. TAOK2
Kinase Mediates PSD95 Stability and Dendritic Spine Maturation through Septin7
Phosphorylation. Neuron. 2017;93: 379–393. doi:10.1016/j.neuron.2016.12.006

21. Richter M, Murtaza N, Scharrenberg R, White SH, Johanns O, Walker S, et al. Altered
TAOK2 activity causes autism-related neurodevelopmental and cognitive abnormalities
through RhoA signaling. Mol Psychiatry. 2019;24: 1329–1350. doi:10.1038/s41380-018-
0025-5

22. Golzio C, Willer J, Talkowski ME, Oh EC, Taniguchi Y, Jacquemont S, et al. KCTD13 is
a major driver of mirrored neuroanatomical phenotypes of the 16p11.2 copy number
variant. Nature. 2012;485: 363–367. doi:10.1038/nature11091

23. Escamilla CO, Filonova I, Walker AK, Xuan ZX, Holehonnur R, Espinosa F, et al.
Kctd13 deletion reduces synaptic transmission via increased RhoA. Nature. 2017;551:
227–231. doi:10.1038/nature24470

24. Ip JPK, Nagakura I, Petravicz J, Li K, Wiemer EAC, Sur M. Major vault protein, a
candidate gene in 16p11.2 microdeletion syndrome, is required for the homeostatic
regulation of visual cortical plasticity. J Neurosci. 2018;38: 2017–2034.
doi:10.1523/JNEUROSCI.2034-17.2018

25. Dickinson ME, Flenniken AM, Ji X, Teboul L, Wong MD, White JK, et al. High-
throughput discovery of novel developmental phenotypes. Nature. 2016;537: 508–514.
doi:10.1038/nature19356

26. Wangler MF, Yamamoto S, Bellen HJ. Fruit flies in biomedical research. Genetics.
2015;199: 639–653. doi:10.1534/genetics.114.171785

27. Chien S. Homophila: human disease gene cognates in Drosophila. Nucleic Acids Res.
2002;30: 149–151. doi:10.1093/nar/30.1.149

28. Reiter LT, Potocki L, Chien S, Gribskov M, Bier E. A systematic analysis of human
disease-associated gene sequences in Drosophila melanogaster. Genome Res. 2001;11:
1114–1125. doi:10.1101/gr.169101

29. Iyer J, Singh MD, Jensen M, Patel P, Pizzo L, Huber E, et al. Pervasive genetic

interactions modulate neurodevelopmental defects of the autism-Associated 16p11.2 deletion in Drosophila melanogaster. Nat Commun. 2018;9: 2548. doi:10.1038/s41467-018-04882-6

30. Singh MD, Jensen M, Lasser M, Huber E, Yusuff T, Pizzo L, et al. NCBP2 modulates neurodevelopmental defects of the 3q29 deletion in Drosophila and X. laevis models. bioRxiv. 2019; 614750. doi:10.1101/614750

31. Molnar C, Resnik-Docampo M, F. M, Martin M, F. C, de Celis JF. Signalling Pathways in Development and Human Disease: A Drosophila Wing Perspective. Human Genetic Diseases. InTech; 2011. doi:10.5772/23858

32. Dworkin I, Gibson G. Epidermal growth factor receptor and transforming growth factor-β signaling contributes to variation for wing shape in Drosophila melanogaster. Genetics. 2006;173: 1417–1431. doi:10.1534/genetics.105.053868

33. Testa ND, Dworkin I. The sex-limited effects of mutations in the EGFR and TGF-β signaling pathways on shape and size sexual dimorphism and allometry in the Drosophila wing. Dev Genes Evol. 2016;226: 159–171. doi:10.1007/s00427-016-0534-7

34. Yan SJ, Gu Y, Li WX, Fleming RJ. Multiple signaling pathways and a selector protein sequentially regulate Drosophila wing development. Development. 2004;131: 285–298. doi:10.1242/dev.00934

35. Strigini M, Cohen SM. A Hedgehog activity gradient contributes to AP axial patterning of the Drosophila wing. Development. 1997;124: 4697–4705.

36. Diaz de la Loza MC, Thompson BJ. Forces shaping the Drosophila wing. Mech. Dev. 2017;144: 23–32. doi:10.1016/j.mod.2016.10.003

37. Bier E. Drosophila, the golden bug, emerges as a tool for human genetics. Nat Rev Genet. 2005;6: 9–23. doi:10.1038/nrg1503

38. Wu Y, Bolduc F V, Bell K, Tully T, Fang Y, Sehgal A, et al. A Drosophila model for Angelman syndrome. Proc Natl Acad Sci U S A. 2008;105: 12399–12404. doi:10.1073/pnas.0805291105

39. Yamamoto S, Jaiswal M, Charng WL, Gambin T, Karaca E, Mirzaa G, et al. A drosophila genetic resource of mutants to study mechanisms underlying human genetic diseases. Cell. 2014;159: 200–214. doi:10.1016/j.cell.2014.09.002

40. Kochinke K, Zweier C, Nijhof B, Fenckova M, Cizek P, Honti F, et al. Systematic Phenomics Analysis Deconvolutes Genes Mutated in Intellectual Disability into Biologically Coherent Modules. Am J Hum Genet. 2016;98: 149–164. doi:10.1016/j.ajhg.2015.11.024

41. Hu Y, Flockhart I, Vinayagam A, Bergwitz C, Berger B, Perrimon N, et al. An integrative approach to ortholog prediction for disease-focused and other functional studies. BMC Bioinformatics. 2011;12: 357. doi:10.1186/1471-2105-12-357

42. Capdevila J, Guerrero I. Targeted expression of the signaling molecule decapentaplegic induces pattern duplications and growth alterations in Drosophila wings. EMBO J. 1994;13: 4459–4468. doi:10.1002/j.1460-2075.1994.tb06768.x

43. Lindström R, Lindholm P, Palgi M, Saarma M, Heino TI. In vivo screening reveals interactions between Drosophila Manf and genes involved in the mitochondria and the ubiquinone synthesis pathway. BMC Genet. 2017;18: 52. doi:10.1186/s12863-017-0509-3

44. Nielsen J, Fejgin K, Sotty F, Nielsen V, Mørk A, Christoffersen CT, et al. A mouse model of the schizophrenia-associated 1q21.1 microdeletion syndrome exhibits altered

mesolimbic dopamine transmission. Transl Psychiatry. 2017;7: 1261. doi:10.1038/s41398-
017-0011-8

45.   Huang H, Potter CJ, Tao W, Li DM, Brogiolo W, Hafen E, et al. PTEN affects cell size,
cell proliferation and apoptosis during Drosophila eye development. Development.
1999;126: 5365–5372. doi:10.5167/uzh-627

46.   Toyo-oka K, Mori D, Yano Y, Shiota M, Iwao H, Goto H, et al. Protein phosphatase 4
catalytic subunit regulates Cdk1 activity and microtubule organization via NDEL1
dephosphorylation. J Cell Biol. 2008;180: 1133–1147. doi:10.1083/jcb.200705148

47.   Ohsugi M, Adachi K, Horai R, Kakuta S, Sudo K, Kotaki H, et al. Kid-Mediated
Chromosome Compaction Ensures Proper Nuclear Envelope Formation. Cell. 2008;132:
771–782. doi:10.1016/j.cell.2008.01.029

48.   Iyer J, Wang Q, Le T, Pizzo L, Grönke S, Ambegaokar SS, et al. Quantitative assessment
of eye phenotypes for functional genetic studies using Drosophila melanogaster. G3
Genes, Genomes, Genet. 2016;6: 1427–1437. doi:10.1534/g3.116.027060

49.   Rosenberg MJ, Agarwala R, Bouffard G, Davis J, Fiermonte G, Hilliard MS, et al. Mutant
deoxynucleotide carrier is associated with congenital microcephaly. Nat Genet. 2002;32:
175–179. doi:10.1038/ng948

50.   Chintapalli VR, Wang J, Dow JAT. Using FlyAtlas to identify better Drosophila
melanogaster models of human disease. Nat. Genet. 2007;39: 715–720.
doi:10.1038/ng2049

51.   Claes L, Del-Favero J, Ceulemans B, Lagae L, Van Broeckhoven C, De Jonghe P. De
novo mutations in the sodium-channel gene SCN1A cause severe myoclonic epilepsy of
infancy. Am J Hum Genet. 2001;68: 1327–1332. doi:10.1086/320609

52.   Ardlie KG, DeLuca DS, Segrè A V., Sullivan TJ, Young TR, Gelfand ET, et al. The
Genotype-Tissue Expression (GTEx) pilot analysis: Multitissue gene regulation in
humans. Science. 2015;348: 648–660. doi:10.1126/science.1262110

53.   Ernst C. Proliferation and Differentiation Deficits are a Major Convergence Point for
Neurodevelopmental Disorders. Trends Neurosci. 2016;39: 290–299.
doi:10.1016/j.tins.2016.03.001

54.   Marchetto MC, Belinson H, Tian Y, Freitas BC, Fu C, Vadodaria KC, et al. Altered
proliferation and networks in neural cells derived from idiopathic autistic individuals. Mol
Psychiatry. 2017;22: 820–835. doi:10.1038/mp.2016.95

55.   Wei H, Alberts I, Li X. The apoptotic perspective of autism. Int J Dev Neurosci. 2014;36:
13–18. doi:10.1016/j.ijdevneu.2014.04.004

56.   Hartl TA, Scott MP. Wing tips: The wing disc as a platform for studying Hedgehog
signaling. Methods. 2014;68: 199–206. doi:10.1016/j.ymeth.2014.02.002

57.   de Celis JF, García-Bellido A. Roles of the Notch gene in Drosophila wing
morphogenesis. Mech Dev. 1994;46: 109–122. doi:10.1016/0925-4773(94)90080-9

58.   Raftery LA, Umulis DM. Regulation of BMP activity and range in Drosophila wing
development. Curr Opin Cell Biol. 2012;24: 158–65. doi:10.1016/j.ceb.2011.11.004

59.   Diaz-Benjumea FJ, Cohen SM. Serrate signals through Notch to establish a Wingless-
dependent organizer at the dorsal/ventral compartment boundary of the Drosophila wing.
Development. 1995;121: 4215–4225.

60.   Becam I, Milán M. A permissive role of Notch in maintaining the DV affinity boundary of
the Drosophila wing. Dev Biol. 2008;322: 190–198. doi:10.1016/j.ydbio.2008.07.028

61.   Zecca M, Basler K, Struhl G. Sequential organizing activities of engrailed, hedgehog and
decapentaplegic in the Drosophila wing. Development. 1995;121: 2265–2278.
doi:10.5167/uzh-1053

62.   Tabata T, Kornberg TB. Hedgehog is a signaling protein with a key role in patterning
Drosophila imaginal discs. Cell. 1994;76: 89–102. doi:10.1016/0092-8674(94)90175-9

63.   O ' Roak BJ, Vives L, Girirajan S, Karakoc E, Krumm N, Coe BP, et al. Sporadic autism
exomes reveal a highly interconnected protein network of de novo mutations. Nature.
2012;485: 246–250. doi:10.1038/nature10989

64.   Hahn H, Wicking C, Zaphiropoulos PG, Gailani MR, Shanley S, Chidambaram A, et al.
Mutations of the human homolog of drosophila patched in the nevoid basal cell carcinoma
syndrome. Cell. 1996;85: 841–851. doi:10.1016/S0092-8674(00)81268-4

65.   Johnson RL, Rothman AL, Xie J, Goodrich L V., Bare JW, Bonifas JM, et al. Human
homolog of patched, a candidate gene for the basal cell nevus syndrome. Science.
1996;272: 1668–1671. doi:10.1126/science.272.5268.1668

66.   Swarup S, Pradhan-Sundd T, Verheyen EM. Genome-wide identification of phospho-
regulators of Wnt signaling in Drosophila. Development. 2015;142: 1502–1515.
doi:10.1242/dev.116715

67.   Six EM, Ndiaye D, Sauer G, Laâbi Y, Athman R, Cumano A, et al. The Notch ligand
Delta1 recruits Dlg1 at cell-cell contacts and regulates cell migration. J Biol Chem.
2004;279: 55818–55826. doi:10.1074/jbc.M408022200

68.   Greene CS, Krishnan A, Wong AK, Ricciotti E, Zelaya RA, Himmelstein DS, et al.
Understanding multicellular function and disease with human tissue-specific networks.
Nat Genet. 2015;47: 569–576. doi:10.1038/ng.3259

69.   Bult CJ, Blake JA, Smith CL, Kadin JA, Richardson JE, Mouse Genome Database Group.
Mouse Genome Database (MGD) 2019. Nucleic Acids Res. 2019;47: D801–D806.
doi:10.1093/nar/gky1056

70.   Zhou W, Zhang L, Guoxiang X, Mojsilovic-Petrovic J, Takamaya K, Sattler R, et al.
GluR1 controls dendrite growth through its binding partner, SAP97. J Neurosci. 2008;28:
10220–10233. doi:10.1523/JNEUROSCI.3434-08.2008

71.   Iizuka-Kogo A, Ishidao T, Akiyama T, Senda T. Abnormal development of urogenital
organs in Dlgh1-deficient mice. Development. 2007;134: 1799–1807.
doi:10.1242/dev.02830

72.   Mahoney ZX, Sammut B, Xavier RJ, Cunningham J, Go G, Brim KL, et al. Discs-large
homolog 1 regulates smooth muscle orientation in the mouse ureter. Proc Natl Acad Sci U
S A. 2006;103: 19872–19877. doi:10.1073/pnas.0609326103

73.   Caruana G, Bernstein A. Craniofacial dysmorphogenesis including cleft palate in mice
with an insertional mutation in the discs large gene. Mol Cell Biol. 2001;21: 1475–1483.
doi:10.1128/MCB.21.5.1475-1483.2001

74.   Chapman DL, Papaioannou VE. Three neural tubes in mouse embryos with mutations in
the T-box gene Tbx6. Nature. 1998;391: 695–697. doi:10.1038/35624

75.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene ontology:
Tool for the unification of biology. Nat. Genet. 2000;25: 25–29. doi:10.1038/75556

76.   The Gene Ontology Consortium. The Gene Ontology Resource: 20 years and still GOing
strong. Nucleic Acids Res. 2019;47: D330–D338. doi:10.1093/nar/gky1055

77.   Pabis M, Neufeld N, Shav-Tal Y, Neugebauer KM. Binding properties and dynamic

localization of an alternative isoform of the cap-binding complex subunit CBP20.
Nucleus. 2010;1: 412–421. doi:10.4161/nucl.1.5.12839

78. Di Padova M, Bruno T, De Nicola F, Iezzi S, D'Angelo C, Gallo R, et al. Che-1 arrests
human colon carcinoma cell proliferation by displacing HDAC1 from the p21WAF1/CIP1
promoter. J Biol Chem. 2003;278: 36496–36504. doi:10.1074/jbc.M306694200

79. Bruno T, De Angelis R, De Nicola F, Barbato C, Di Padova M, Corbi N, et al. Che-1
affects cell growth by interfering with the recruitment of HDAC1 by Rb. Cancer Cell.
2002;2: 387–399. doi:10.1016/S1535-6108(02)00182-4

80. Page G, Lödige I, Kögel D, Scheidtmann KH. AATF, a novel transcription factor that
interacts with Dlk/ZIP kinase and interferes with apoptosis. FEBS Lett. 1999;462: 187–91.
doi:10.1016/s0014-5793(99)01529-x

81. Sumiyoshi E, Sugimoto A, Yamamoto M. Protein phosphatase 4 is required for
centrosome maturation in mitosis and sperm meiosis in C. elegans. J Cell Sci. 2002;115:
1403–1410.

82. Jensen M, Girirajan S. An interaction-based model for neuropsychiatric features of copy-
number variants. PLoS Genet. 2019;15: e1007879. doi:10.1371/journal.pgen.1007879

83. Girirajan S, Rosenfeld JA, Coe BP, Parikh S, Friedman N, Goldstein A, et al. Phenotypic
Heterogeneity of Genomic Disorders and Rare Copy-Number Variants. N Engl J Med.
2012;367: 1321–1331. doi:10.1056/NEJMoa1200395

84. Coe BP, Girirajan S, Eichler EE. A genetic model for neurodevelopmental disease. Curr
Opin Neurobiol. 2012;22: 829–836. doi:10.1016/j.conb.2012.04.007

85. Nicholas AK, Swanson EA, Cox JJ, Karbani G, Malik S, Springell K, et al. The molecular
landscape of ASPM mutations in primary microcephaly. J Med Genet. 2009;46: 249–253.
doi:10.1136/jmg.2008.062380

86. Brand AH, Perrimon N. Targeted gene expression as a means of altering cell fates and
generating dominant phenotypes. Development. 1993;118: 401–415.
doi:10.1101/lm.1331809

87. Green EW, Fedele G, Giorgini F, Kyriacou CP. A Drosophila RNAi collection is subject
to dominant phenotypic effects. Nat Methods. 2014;11: 222–223. doi:10.1038/nmeth.2856

88. Vissers JHA, Manning SA, Kulkarni A, Harvey KF. A Drosophila RNAi library
modulates Hippo pathway-dependent tissue growth. Nat Commun. 2016;7: 10368.
doi:10.1038/ncomms10368

89. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: An
open-source platform for biological-image analysis. Nat. Methods. 2012;9: 676–682.
doi:10.1038/nmeth.2019

90. Kanehisa M, Sato Y, Furumichi M, Morishima K, Tanabe M. New approach for
understanding genome variations in KEGG. Nucleic Acids Res. 2019;47: D590–D595.
doi:10.1093/nar/gky962

91. Hagberg AA, Schult DA, Swart PJ. Exploring network structure, dynamics, and function
using NetworkX. 7th Python in Science Conference (SciPy 2008). 2008. pp. 11–15.
Available: http://conference.scipy.org/proceedings/SciPy2008/paper_2

92. Mi H, Huang X, Muruganujan A, Tang H, Mills C, Kang D, et al. PANTHER version 11:
Expanded annotation data from Gene Ontology and Reactome pathways, and data analysis
tool enhancements. Nucleic Acids Res. 2017;45: D183–D189. doi:10.1093/nar/gkw1138

93. Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, et al. Cytoscape: A

software Environment for integrated models of biomolecular interaction networks.
Genome Res. 2003;13: 2498–2504. doi:10.1101/gr.1239303
858

 FIGURE LEGENDS

Figure 1. Targeted analysis to identify global developmental phenotypes with knockdown

of homologs of CNV genes. (A) Strategy for identifying non-neuronal phenotypes and

underlying cellular mechanisms for homologs of CNV and known neurodevelopmental genes

using the fly wing as a model system. We evaluated 59 *Drosophila* homologs of genes within 10

CNV regions and 20 known neurodevelopmental genes (79 total homologs). Using the *UAS-*

GAL4 system with wing-specific bx^{MS1096} driver, we knocked down 136 individual RNAi lines

for the CNV and neurodevelopmental homologs, and evaluated qualitative and quantitative

phenotypes. We next clustered RNAi lines based on severity of qualitative phenotypes, and

compared adult wing phenotypes to phenotypes observed with ubiquitous and eye-specific

knockdown of homologs. Furthermore, we evaluated underlying cellular mechanisms for the

observed wing-specific phenotypes, and examined the connectivity patterns of candidate

homologs for developmental phenotypes in multiple human tissue-specific networks. **(B)**

Representative brightfield images of adult wing phenotype severity observed with knockdown of

homologs of CNV genes, based on clustering analysis, are shown. **(C)** Heatmap with k-means

clustering of qualitative phenotypes in adult female wings across 136 RNAi lines is shown. The

color of each cell represents the frequency of individual fly wings (n=20-25 adult wings) for

each RNAi line (x-axis) that show a specific severity (no phenotype, mild, moderate, severe,

lethal) for the five qualitative phenotypes assessed (y-axis; wrinkled wings, ectopic veins,

missing veins, discoloration, bristle planar polarity), as detailed in **Supp. Data 2**. Based on these

data, we identified clusters for no phenotype (n=75 lines), mild (n=24 lines), moderate (n=10

lines), severe (n=21 lines), and lethal (n=6 lines). **(D)** Summary table of qualitative and

quantitative adult wing phenotypes for all tested RNAi lines of homologs of CNV and

neurodevelopmental genes. Quantitative phenotype totals do not include lethal RNAi lines for

both area and vein length. In addition, L3 vein length totals do not include severe RNAi lines.

Figure 2. Qualitative adult wing phenotypes of *Drosophila* homologs of CNV and

neurodevelopmental genes. Heatmaps representing the five qualitative adult wing phenotypes

for all 136 RNAi lines, with **(A)** all 59 tested homologs for 10 CNV regions and **(B)** 20

homologs for neurodevelopmental genes (β-catenin, core neurodevelopmental genes, and

microcephaly genes), are shown. The color of each cell represents the frequency of each of the

five qualitative phenotypes by severity (wrinkled wings, WR; ectopic veins, EV; missing veins,

MV; discoloration, DC; bristle planar polarity, BP), ranging from no phenotype to lethal. **(C)**

Representative brightfield images of adult fly wings (scale bar = 500µm) with wing-specific

knockdown of homologs of CNV and neurodevelopmental genes showing the five assessed

qualitative phenotypes, including discoloration, wrinkled wings, bristle polarity, ectopic veins,

and missing veins are shown. The panels in the bx^{MS1096}-GAL4 control and $C6836^{KK112485}$ images

highlight bristle planar polarity phenotypes for the representative images. Black arrowheads

highlight ectopic veins and white arrowheads highlight missing veins. Genotypes for the images

are: w^{1118}/bx^{MS1096}-GAL4;+; UAS-Dicer2/+, w^{1118}/bx^{MS1096}-GAL4;UAS-Rph^{GD7330} RNAi/+;UAS-

Dicer2/+, w^{1118}/bx^{MS1096}-GAL4;UAS-CG15528^{KK107736} RNAi/+; UAS-Dicer2/+, w^{1118}/bx^{MS1096}-

GAL4;UAS-CG6836^{KK112485} RNAi/+; UAS-Dicer2/+, w^{1118}/bx^{MS1096}-GAL4;+;UAS-

CG14182^{GD2738} RNAi/UAS-Dicer2, and w^{1118}/bx^{MS1096}-GAL4;UAS-kis^{KK100890} RNAi/+; UAS-

Dicer2/+.

Figure 3. Quantitative adult wing phenotypes of *Drosophila* homologs of CNV and

neurodevelopmental genes. (A) Representative brightfield images of adult fly wings (scale bar

= 500µm) with wing-specific knockdown of homologs of CNV and neurodevelopmental genes

with size defects are shown. The bx^{MS1096}-*GAL4* control image highlights the six veins, including

longitudinal veins L2, L3, L4, and L5 as well as the anterior and posterior crossveins (ACV and

PCV), that were measured for quantitative analysis. The dotted line in the control image

represents the total wing area calculated for each RNAi line. Genotypes for the images are:

w^{1118}/bx^{MS1096}-GAL4;+; UAS-Dicer2/+, w^{1118}/bx^{MS1096}-GAL4;UAS-Fmo-2^{KK109203} RNAi/+; UAS-

Dicer2/+, and *w^{1118}/bx^{MS1096}-GAL4;+;UAS-Trpm^{GD4541} RNAi/UAS-Dicer2*. **(B)** Boxplot of L3

vein lengths for knockdown of select homologs in adult fly wings (n = 9-91, *p < 0.05, two-

tailed Mann–Whitney test with Benjamini-Hochberg correction). Vein measurements for all

other longitudinal veins and crossveins (ACV and PCV) for these lines are represented in **Supp**

Fig. 2. **(C)** Boxplot of wing areas for knockdown of select homologs in adult fly wings (n = 9-

91, *p < 0.05, two-tailed Mann–Whitney test with Benjamini-Hochberg correction). All boxplots

indicate median (center line), 25th and 75th percentiles (bounds of box), and minimum and

maximum (whiskers), with red dotted lines representing the control median.

Figure 4. Comparison of wing-specific, eye-specific, and ubiquitous knockdown of

homologs of CNV and known neurodevelopmental genes. (A) Heatmap with the penetrance

of phenotypes with ubiquitous knockdown (*da–GAL4*) of select homologs of CNV genes,

compared to their adult wing-specific (*bx^{MS1096}–GAL4*) phenotypic severity is shown. **(B)**

Boxplots of *Flynotyper*-derived phenotypic scores for adult eyes with eye-specific knockdown

(*GMR-GAL4*) of select homologs of CNV and neurodevelopmental genes, normalized as fold-

change (FC) to control values (n = 7–40, *p < 0.05, one-tailed Mann–Whitney test with

Benjamini-Hochberg correction). The boxplots are arranged by severity of adult wing

phenotypes observed for each RNAi line, while the *Flynotyper* phenotypic scores are categorized

into four severity categories: no change (0–1.1 FC), mild (1.1–1.5 FC), moderate (1.5–2.0 FC),

and severe (>2.0 FC). **(C)** Boxplot showing the average eye phenotypic scores for 66 RNAi lines

of select homologs of CNV and neurodevelopmental genes, normalized as fold-change (FC) to

control values, by wing phenotypic category (n=4–30 RNAi lines per group). We did not observe

any significant changes in eye phenotype severity across the five wing phenotypic categories

(Kruskal-Wallis rank sum test, p=0.567, df = 5, χ^2 = 3.881). Examples of average eye phenotypic

scores for RNAi lines with no phenotype (*para$^{GD3392\_1}$*), mild (*rl^{KK115768}*), and lethal (*dlg1^{GD4689}*)

wing phenotype severity are highlighted in the graph. All boxplots indicate median (center line),

25th and 75th percentiles (bounds of box), and minimum and maximum (whiskers), with red

dotted lines representing the control median. **(D)** Representative brightfield adult eye (scale

bar = 100 μm) and adult wing (scale bar = 500μm) images with tissue-specific knockdown of

homologs of CNV genes are shown. Genotypes for the eye images are: *w^{1118};GMR-GAL4/+;*

UAS-Dicer2/+, w^{1118};GMR-GAL4/UAS-Lnk^{KK105731} RNAi; UAS-Dicer2/+, w^{1118};GMR-

GAL4/UAS-mEFTu1^{GD16961} RNAi; UAS-Dicer2/+. Genotypes for the wing images are:

w^{1118}/bx^{MS1096}-GAL4;+; UAS-Dicer2/+, w^{1118}/bx^{MS1096}-GAL4; UAS-Lnk^{KK105731} RNAi/+; UAS-

Dicer2/+, and *w^{1118}/bx^{MS1096}-GAL4; UAS-mEFTu1^{GD16961} RNAi/+; UAS-Dicer2/+*.

Figure 5. Expression patterns of *Drosophila* homologs and human CNV and

neurodevelopmental genes across multiple tissues. (A) Heatmap with expression of fly

homologs of select CNV and neurodevelopmental genes in multiple *Drosophila* larval and adult

tissues, derived from the FlyAtlas Anatomical Microarray dataset, compared with adult wing

phenotype severity, is shown. Expression values are grouped into no expression (<10 fragments

per kilobase of transcript per million reads, or FPKM), low (10–100 FPKM), moderate (100–500

FPKM), high (500–1000 FPKM), and very high (>1000 FPKM) expression categories. **(B)**

Heatmap with expression of select human CNV and neurodevelopmental genes in multiple adult

tissues, derived from the Genotype-Tissue Expression (GTEx) dataset v.1.2, is shown.

Expression values are grouped into no expression (<3 transcripts per million reads, or TPM), low

(3–10 TPM), moderate (10–25 TPM), high (25–100 TPM), and very high (>100 TPM)

expression categories. X symbols denote preferential expression in a particular tissue (see

Methods). Expression data for all CNV and neurodevelopmental genes are provided in **Supp.**

Data 6.

Figure 6. *Drosophila* **homologs of CNV and neurodevelopmental genes show altered levels**

of apoptosis and proliferation. (A) Larval imaginal wing discs (scale bar = 50 µm) stained with

nuclear marker DAPI, apoptosis marker dcp1, and cell proliferation marker pH3 illustrate altered

levels of apoptosis and cell proliferation due to wing-specific knockdown of select fly homologs

of CNV genes. We quantified the number of stained cells within the wing pouch of the wing disc

(white box), which becomes the adult wing. Additional representative images of select homologs

are presented in **Supp Fig. 5**. Genotypes for the wing images are: w^{1118}/bx^{MS1096}-GAL4;+; UAS-

Dicer2/+, w^{1118}/bx^{MS1096}-GAL4;+; UAS-Aatf^{GD7229} RNAi/UAS-Dicer2, w^{1118}/bx^{MS1096}-GAL4;UAS-

Pp4-19C^{GD9561}/+; UAS-Dicer2/+, w^{1118}/bx^{MS1096}-GAL4;+; UAS-Atx2^{GD11562} RNAi/UAS-Dicer2,

and w^{1118}/bx^{MS1096}-GAL4;+; UAS-Sin^{GD7027} RNAi/UAS-Dicer2. **(B)** Box plot of dcp1-positive

cells in larval wing discs with knockdown of select fly homologs of CNV and

neurodevelopmental genes, normalized to controls (n = 7–18, *p < 0.05, two-tailed Mann–

Whitney test with Benjamini-Hochberg correction). We note that several RNAi lines showed

severe dcp1 staining across the entire wing disc and could not be quantified. The number of dcp1

positive cells were calculated manually. **(C)** Box plot of pH3-positive cells in the larval wing

discs with knockdown of select fly homologs of CNV and neurodevelopmental genes,

normalized to controls (n = 6–18, *p < 0.05, two-tailed Mann–Whitney test with Benjamini-

Hochberg correction). The number of pH3 positive cells were calculated using the

AnalyzeParticles function in ImageJ. All boxplots indicate median (center line), 25th and 75th

percentiles (bounds of box), and minimum and maximum (whiskers), with red dotted lines

representing the control median.

Figure 7. Candidate *Drosophila* homologs of genes within CNV regions interact with

conserved signaling pathways. Larval imaginal wing discs (scale bar = 50 µm) stained with **(A)**

wingless, **(B)** patched, **(C)** engrailed, and **(D)** delta illustrate disrupted expression patterns for

proteins located within the Wnt (wingless), Hedgehog (patched and engrailed), and Notch (delta)

signaling pathways due to wing-specific knockdown of select fly homologs of CNV and

neurodevelopmental genes. Dotted yellow boxes represent expected expression patterns for

signaling proteins in bx^{MS1096}-GAL4 control images. White arrowheads and dotted white boxes

highlight disruptions in expression patterns of signaling proteins with knockdown of CNV or

neurodevelopmental genes. Additional representative images of select homologs are presented in

Supp Fig. 7. Genotypes for the wing images are: *w^{1118}/bx^{MS1096}-GAL4;+; UAS-Dicer2/+,*

w^{1118}/bx^{MS1096}-GAL4;UAS-Pp4-19C^{GD9561}/+; UAS-Dicer2/+, w^{1118}/bx^{MS1096}-GAL4;UAS-

Cbp20^{KK109448}/+; UAS-Dicer2/+, w^{1118}/bx^{MS1096}-GAL4;+; UAS-Sin^{GD7027} RNAi/UAS-Dicer2,

w^{1118}/bx^{MS1096}-GAL4;+; UAS-Aatf^{GD7229} RNAi/UAS-Dicer2, and *w^{1118}/bx^{MS1096}-GAL4;UAS-*

Klp61F^{GD14149}/+; UAS-Dicer2/+.

Figure 8. Connectivity of human CNV genes with conserved signaling pathway genes in

human tissue-specific networks. (A) Representative diagrams of eight human CNV and

neurodevelopmental genes whose fly homologs disrupt the Notch signaling pathway and 57

human Notch signaling genes within kidney, heart, and brain-specific gene interaction networks

are shown. Yellow nodes represent CNV and neurodevelopmental genes, pink nodes represent

Notch signaling pathway genes, and green nodes represent connector genes within the shortest

paths between CNV and Notch pathway genes. **(B)** Violin plots showing the average

connectivity (i.e. inverse of shortest path lengths) of CNV genes to genes in Hedgehog, Notch,

and Wnt signaling pathways across the tested tissue-specific networks (n=322–810 pairwise

interactions, *p < 0.05, two-tailed Welch's t-test with Benjamini-Hochberg correction). **(C)**

Table showing enriched clusters of Gene Ontology (GO) Biological Process terms for connector

genes observed for each signaling pathway in the three tested tissue-specific networks,

categorized by enrichments in ubiquitous, neuronal, and non-neuronal tissues (p<0.05, Fisher's

Exact test with Benjamini-Hochberg correction).

1013

Figure 9. A multigenic model for neuronal and non-neuronal phenotypes associated with

pathogenic CNVs. Schematic of a multigenic model for neuronal and non-neuronal phenotypes

associated with pathogenic CNVs. While a subset of genes within CNV regions contribute

towards tissue-specific phenotypes, a majority of genes contribute towards both neuronal and

non-neuronal phenotypes through disruption of developmental signaling pathways and global

biological processes.

 SUPPLEMENTARY LEGENDS

 Supplementary Figure 1. Quantitative vein length phenotypes for select *Drosophila*

 homologs of CNV and neurodevelopmental genes. Boxplots of longitudinal veins **(A)** L2, **(B)**

 L4, **(C)** L5, and **(D)** anterior crossvein (ACV) and **(E)** posterior crossvein (PCV) lengths for

 knockdown of select homologs in adult fly wings (n = 9-91, *p < 0.05, two-tailed Mann–

 Whitney test with Benjamini-Hochberg correction). All boxplots indicate median (center line),

 25th and 75th percentiles (bounds of box), and minimum and maximum (whiskers), with red

 dotted lines representing the control median.

 Supplementary Figure 2. Comparisons of eye-specific and wing-specific knockdowns for

 select *Drosophila* homologs of CNV and neurodevelopmental genes. Boxplots of *Flynotyper-*

 derived phenotypic scores for 66 tested adult eyes with eye-specific knockdown (*GMR-GAL4*) of

 select homologs of CNV and neurodevelopmental genes, normalized as fold-change (FC) to

 control values (n = 1–40, *p < 0.05, one-tailed Mann–Whitney test with Benjamini-Hochberg

 correction). RNAi lines that do not show any observable qualitative adult wing phenotypes,

 including lines that show wing measurement phenotypes, are represented in **(A)**, and RNAi lines

 with observable mild to lethal qualitative wing phenotypes are represented in **(B)**. All boxplots

 indicate median (center line), 25th and 75th percentiles (bounds of box), and minimum and

 maximum (whiskers), with red dotted lines representing the control median.

 Supplementary Figure 3. Expression of *Drosophila* homologs of CNV and

 neurodevelopmental genes in larval and adult tissues. Venn diagrams representing the

 number of 76/77 tested fly homologs for CNV and neurodevelopmental genes expressed (>10

1043   fragments per kilobase of transcript per million reads, or FPKM) in **(A)** larval (central nervous

1044   system or CNS, gut, trachea, and fat body) and **(B)** adult tissues (brain, gut, heart and fat body),

1045   are shown.

1046

Supplementary Figure 4. Additional *Drosophila* homologs of CNV and neurodevelopmental

genes show altered levels of cell proliferation and apoptosis. Larval imaginal wing discs

(scale bar = 50 µm) stained with nuclear marker DAPI, apoptosis marker dcp1, and cell

proliferation marker pH3 illustrate altered levels of cell proliferation and apoptosis due to wing-

specific knockdown of select fly homologs of CNV genes. We examined changes in the number

of stained cells within the wing pouch of the wing disc (white box), which becomes the adult

wing. Genotypes for the wing images are: w^{1118}/bx^{MS1096}-GAL4;+; UAS-Dicer2/+, w^{1118}/bx^{MS1096}-

GAL4;UAS-Cbp20^{KK109448}/+; UAS-Dicer2/+, w^{1118}/bx^{MS1096}-GAL4;+; UAS-dlg1^{GD4689}

RNAi/UAS-Dicer2, w^{1118}/bx^{MS1096}-GAL4;UAS-CG8888^{GD3777}/+; UAS-Dicer2/+, w^{1118}/bx^{MS1096}-

GAL4;+; UAS-UQCR-C2^{GD11238} RNAi/UAS-Dicer2, w^{1118}/bx^{MS1096}-GAL4;+; UAS-ACC^{GD3482}

RNAi/UAS-Dicer2, w^{1118}/bx^{MS1096}-GAL4;UAS-Klp61F^{GD14149}/+; UAS-Dicer2/+, and

w^{1118}/bx^{MS1096}-GAL4;UAS-Rph^{GD7330} RNAi/+;UAS-Dicer2/+.

1059

Supplementary Figure 5. Select female and male *Drosophila* homologs of CNV and

neurodevelopmental genes show altered levels of cell proliferation and apoptosis. (A) Larval

imaginal wing discs (scale bar = 50 µm) stained with nuclear marker DAPI, apoptosis marker

dcp1, and cell proliferation marker pH3 illustrate altered levels of cell proliferation and apoptosis

due to wing-specific knockdown of select fly homologs of CNV genes in females and males. We

examined changes in the number of stained cells within the wing pouch of the wing disc (white

box), which becomes the adult wing. Genotypes for the wing images are: w^{1118}/bx^{MS1096}-GAL4;+;

UAS-Dicer2/+, w^{1118}/bx^{MS1096}-GAL4;+; UAS-lgs^{GD1241}/UAS-Dicer2, w^{1118}/bx^{MS1096}-GAL4;+;

UAS-Sra-1^{GD11477}/UAS-Dicer2, and w^{1118}/bx^{MS1096}-GAL4;UAS-CG11035^{KK101201} RNAi+; UAS-

Dicer2/+. **(B)** Box plot of dcp1-positive cells in larval wing discs with knockdown of select fly

homologs of CNV and neurodevelopmental genes, normalized to controls (n = 9–13, *p < 0.05,

two-tailed Mann–Whitney test with Benjamini-Hochberg correction). The number of dcp1

positive cells were calculated manually. **(C)** Box plot of pH3-positive cells in the larval wing

discs with knockdown of select fly homologs of CNV and neurodevelopmental genes,

normalized to controls (n = 9–13, *p < 0.05, two-tailed Mann–Whitney test with Benjamini-

Hochberg correction). The number of pH3 positive cells were calculated using the

AnalyzeParticles function in ImageJ. All boxplots indicate median (center line), 25th and 75th

percentiles (bounds of box), and minimum and maximum (whiskers), with red dotted lines

representing the control median.

1079

Supplementary Figure 6. Additional *Drosophila* homologs of genes within CNV regions

interact with conserved signaling pathways to induce developmental phenotypes. Larval

imaginal wing discs (scale bar = 50 µm) stained with **(A)** wingless, **(B)** patched, **(C)** engrailed,

and **(D)** delta illustrate disrupted expression patterns for proteins located within the Wnt

(wingless), Hedgehog (patched and engrailed), and Notch (delta) signaling pathways due to

wing-specific knockdown of additional fly homologs of CNV and neurodevelopmental genes.

Dotted yellow boxes represent expected expression patterns for signaling proteins in bx^{MS1096}-

GAL4 control images. White arrowheads and dotted white boxes highlight disruptions in

expression patterns of signaling proteins with knockdown of CNV or neurodevelopmental genes.

Genotypes for the wing images are: *w^{1118}/bx^{MS1096}-GAL4;+; UAS-Dicer2/+*, *w^{1118}/bx^{MS1096}-*

GAL4;+; UAS-dlg1^{GD4689} RNAi/UAS-Dicer2, *w^{1118}/bx^{MS1096}-GAL4;+; UAS-Tsf2^{GD2442}*

RNAi/UAS-Dicer2, *w^{1118}/bx^{MS1096}-GAL4;+; UAS-Atx2^{GD11562} RNAi/UAS-Dicer2*, *w^{1118}/bx^{MS1096}-*

GAL4;+; UAS-UQCR-C2^{GD11238} RNAi/UAS-Dicer2, *w^{1118}/bx^{MS1096}-GAL4;+; UAS-nudE^{GD15226}*

RNAi/UAS-Dicer2, and *w^{1118}/bx^{MS1096}-GAL4;+; UAS-ACC^{GD3482} RNAi/UAS-Dicer2*.

1094

Supplementary Figure 7. Tissue-specific network diagrams showing connectivity of human

CNV genes with conserved signaling pathway genes. Representative network diagrams of nine

human CNV and neurodevelopmental genes whose fly homologs disrupt the **(A)** Wnt and **(B)**

Hedgehog signaling pathway and 162 human Wnt and 46 human Hedgehog signaling genes

within kidney, heart, and brain-specific gene interaction networks are shown. Yellow nodes

represent CNV and neurodevelopmental genes, pink nodes represent Notch signaling pathway

genes, and green nodes represent connector genes within the shortest paths between CNV and

Notch pathway genes.

1103

1104   **Supplementary Data 1 (Excel file).** *Drosophila* homologs of human CNV and

1105   neurodevelopmental genes as determined using DIOPT v.7.1.

1106

1107   **Supplementary Data 2 (Excel file).** Qualitative and quantitative adult wing phenotypic data for

1108   *Drosophila* homologs of human CNV and neurodevelopmental genes. This file shows the raw

1109   frequencies of severity for the five qualitative wing phenotypes and average areas and vein

1110   lengths for all 136 female and male tested RNAi lines. In addition, this file also includes k-means

1111   clustering analysis for the female RNAi lines.

1112

Supplementary Data 3 (Excel file). Summary of adult wing qualitative and quantitative

phenotypes by *Drosophila* homologs. This file summarizes qualitative k-means clustering and

longitudinal L3 vein length and wing area changes for all 136 RNAi lines by fly homologs. We

define discordant homologs when RNAi lines for the same homologs showed inconsistent wing

phenotypes. For each homolog with multiple RNAi lines, we checked discordance among RNAi

lines for no phenotype versus any qualitative or quantitative phenotypes, followed by

discordance for small or large quantitative phenotypes.

1120

Supplementary Data 4 (Excel file). Phenotypes of mouse knockdown models for homologs of

CNV genes. This file lists lethality and neuronal and non-neuronal phenotypes, categorized using

top-level Mammalian Phenotype Ontology terms, for knockdown models of 130 mouse

homologs of CNV genes derived from the Mouse Genome Informatics (MGI) database.

1125

Supplementary Data 5 (Excel file). Summary of eye-specific and wing-specific phenotypes for

fly homologs. This file summarizes eye-specific and wing-specific phenotypes by severity

category for 66 RNAi lines by fly homologs of CNV and neurodevelopmental genes. Eye

phenotype severity is defined by *Flynotyper* phenotypic scores with fold-change (FC)

normalization to control as follows: no change (0–1.1 FC), mild (1.1–1.5 FC), moderate (1.5–2.0

FC), and severe (>2.0 FC). Wing phenotype severity is defined by k-means clustering for

qualitative phenotypes and quantitative size changes as listed in **Supp. Data 3**.

1133

Supplementary Data 6 (Excel file). Tissue-specific expression of *Drosophila* homologs and human CNV and neurodevelopmental genes. This file lists expression values across multiple fly and human tissues for all 79 *Drosophila* homologs and 150 human genes. Fly expression data (fragments per kilobase of transcript per million reads, or FPKM) was derived from the FlyAtlas Anatomical Microarray dataset, and human expression data (transcripts per million reads, or TPM) was derived from the Genotype-Tissue Expression (GTEx) dataset v.1.2.

Supplementary Data 7 (Excel file). Summary of immunostaining of the larval imaginal wing discs. This file summarizes changes in apoptosis (27 homologs), cell proliferation (27 homologs), and Wnt, Hedgehog, and Notch signaling pathway proteins (14 homologs), along with qualitative and quantitative adult wing phenotypes (as listed in **Supp. Data 2.**), for female and male fly homologs.

Supplementary Data 8 (Excel file). Tissue-specific network connectivity for candidate CNV genes and signaling pathway genes. This file lists the shortest path lengths between nine candidate CNV genes and 265 genes within Wnt, Hedgehog, and Notch signaling pathways for heart, kidney, and brain-specific gene interaction networks, along with the connector genes that are within the shortest paths. Enriched Gene Ontology (GO) Biological Process, Cellular Component, and Molecular Function terms for sets of connector genes for each signaling pathway in each tissue-specific networks are also represented.

Supplementary Data 9 (Excel file). List of *Drosophila* stocks used for experiments, including stock numbers and genotypes.

1157

1158 **Supplementary Data 10 (Excel file).** Statistics for all experimental data. This file shows all

1159 statistical information (sample size, mean/median/standard deviation of datasets, test statistics, p-

1160 values, degrees of freedom, confidence intervals, and Benjamini-Hochberg FDR corrections) for

1161 all data. Statistical information for Kruskal-Wallis test includes factors, degrees of freedom, test

1162 statistics, and post-hoc pairwise Wilcoxon tests with Benjamini-Hochberg correction.

Figure 1

A   Experimental strategy

Wing-specific driver
bx^{MS1096}-GAL4   *UAS-RNAi*
                       (79 genes)

Phenotyping of adult wings

Qualitative analysis	Quantitative analysis
• Wrinkled wings	• Longitudinal veins
• Ectopic veins	(L2,L3,L4,L5)
• Missing veins	• Crossveins (ACV, PCV)
• Discoloration	• Wing area
• Bristle planar polarity	

Tissue-specific effects

- Compare to ubiquitous knockdown
- Compare to eye-specific knockdown
- Gene expression across multiple tissues
  (Human and flies)

Cellular processes and developmental pathways

- Apoptosis
- Cell proliferation
- Disruptions in signaling pathways
- Human tissue-specific network analysis
  (Brain, heart, kidney)

B   Phenotypic analysis of adult wings in *Drosophila*

bx^{MS1096}-GAL4 control   No phenotype   Mild   Moderate   Severe

C   K-means clustering of qualitative wing data

D   Summary of qualitative and quantitative adult wing phenotypes

Fly homologs	RNAi lines	Qualitative phenotypes					Quantitative phenotypes	
		No pheno.	Mild	Moderate	Severe	Lethal	Wing area	L3 vein length
CNV genes (15q11.2, 15q13.3, 16p11.2, Distal 16p11.2, 16p12.1, 16p13.11, 17q12, 1q21.1, 3q29, 7q11.23)	95	45	21	7	16	6	65	53
Known neurodevelopmental genes (β-catenin, core genes, microcephaly)	41	30	3	3	5	0	24	21
Total	136	75	24	10	21	6	89	74

Figure 2

A Qualitative wing phenotypes of homologs of CNV genes

B Qualitative wing phenotypes of homologs of neurodevelopmental genes

C Representative qualitative phenotypes in the adult wing

Figure 3

A

Representative quantitative phenotypes in the adult wing

bx^{MS1096}-GAL4

Smaller area/L3 vein length

Fmo-2^{KK109203}

Larger area/L3 vein length

Trpm$^{GD4541\_2}$

B L3 vein lengths for select homologs

C Wing area for select homologs

Figure 4

A Ubiquitous knockdown

B Eye versus wing phenotypes

C Severity of eye versus wing phenotype

D Representative eye and wing images

Figure 5

A   **Expression patterns of fly homologs of CNV genes across multiple fly tissues**

Figure 6

A Cellular processes in larval wing discs

Apoptosis | Cell proliferation

B Quantification of apoptotic cells

C Quantification of proliferating cells

Figure 7

Disruption of signaling pathways in larval wing discs

Figure 8

A   Human CNV genes interact with Notch signaling pathway genes in multiple tissues

Kidney   *Heart*   *Brain*

○ CNV/neurodevelopmental gene   ○ Signaling pathway gene   ○ Connector gene

B   Average connectivity of CNV genes to genes in signaling pathways

C   GO term enrichment for connector genes in tissue-specific networks

Figure 9

CNV region

Gene A Gene B Gene C Gene D Gene E Gene F

Developmental pathways
Wnt, Notch, Hedgehog

Biological processes

Signal transduction   *Cell cycle*   *RNA processing*
Axonal transport   *Apoptosis*   *Protein degradation*

Neuronal phenotypes
Autism
Epilepsy
Intellectual disability
Micro/macrocephaly

Non-neuronal phenotypes
Cardiac defects
Kidney defects
Craniofacial features
Musculoskeletal defects

A L2 vein length in adult wing

B L4 vein length in adult wing

C L5 vein length in adult wing

D ACV length in adult wing

E PCV length in adult wing

Supp. Figure 2

Supp. Figure 3

A Larval *Drosophila* expression

B Adult *Drosophila* expression

Supp. Figure 4

Cellular processes in larval wing discs

A **Cellular processes in female and male larval wing discs**

B **Apoptosis in larval wing discs**

C **Cell proliferation in larval wing discs**

Disruption of signaling pathways in larval wing discs

A   **Human CNV genes interact with Wnt signaling pathway genes in multiple tissues**

Kidney   *Heart*   *Brain*

B   **Human CNV genes interact with Hedgehog signaling pathway genes in multiple tissues**

Kidney   *Heart*   *Brain*

○ CNV/neurodevelopmental gene   ○ Signaling pathway gene   ○ Connector gene