Glioblastoma stem cells induce quiescence in surrounding neural stem cells via Notch signalling.

Katerina Lawlor[1], Maria Angeles Marques-Torrejon[2], Gopuraja Dharmalingham[3], Yasmine El-Azhar[1], Michael D. Schneider[1], Steven M. Pollard[2§] and Tristan A. Rodríguez[1§]

[1]National Heart and Lung Institute, Imperial College London, Hammersmith Hospital Campus, Du Cane Road, London W12 0NN, United Kingdom.

[2] MRC Centre for Regenerative Medicine & Edinburgh Cancer Research UK Centre, University of Edinburgh, Edinburgh, UK.

[3]MRC London Institute of Medical Sciences, Institute of Clinical Sciences, Imperial College London, UK

[§]Authors for correspondence: steven.pollard@ed.ac.uk and tristan.rodriguez@imperial.ac.uk

Running title: Glioblastoma stem cell competition

Keyword: Neural stem cells, quiescence, glioblastoma, Notch, cell competition

Abstract

There is increasing evidence suggesting that adult neural stem cells (NSCs) are a cell of origin of glioblastoma, the most aggressive form of malignant glioma. The earliest stages of hyperplasia are not easy to explore, but likely involve a cross-talk between normal and transformed NSCs. How normal cells respond to this cross-talk and if they expand or are outcompeted is poorly understood. Here we have analysed the interaction of transformed and wild-type NSCs isolated from the adult mouse subventricular zone neural stem cell niche. We find that transformed NSCs are refractory to quiescence-inducing signals. Unexpectedly, however, we also demonstrate that these cells induce a quiescent-like state in surrounding wild-type NSC. We find that this response is cell-cell contact-dependent and that transformed cells activate the Notch pathway in adjacent wild-type NSCs, an event that stimulates their entry into quiescence. Our findings therefore suggest that oncogenic mutations may be propagated in the stem cell niche not just though cell-intrinsic advantages, but also by outcompeting neighbouring stem cells through signalling repression of their proliferation.

Introduction

In the adult mammalian brain, neural stem cells (NSCs) can be primarily found in specific neurogenic regions. In rodents these are the subgranular zone (SGZ) of the dentate gyrus (DG) and the subventricular zone (SVZ), which lines the lateral ventricles (Doetsch et al., 1999; Quinones-Hinojosa et al., 2006; Sanai et al., 2004). In the SVZ there are both quiescent NSCs and activated NSCs that coexist in the niche (Quinones-Hinojosa et al., 2006). The quiescent NSC subpopulation make up the majority of NSCs in the adult SVZ and are capable of completely regenerating this niche following anti-mitotic drug treatment (Doetsch et al., 1999; Giachino and Taylor, 2009; Pastrana et al., 2009). Tight regulation of the switch between these two states is vital to ensure that the pool of NSCs does not accumulate DNA damage or become exhausted over time and therefore understanding how this is achieved is of great interest. One key event in NSC activation is the upregulation of EGFR, which is highly expressed in activated NSCs and progenitor cells but absent in quiescent NSCs. Thus, cell surface expression of EGFR is often used as a marker to differentiate these two populations (Codega et al., 2014; Pastrana et al., 2009). What feedback mechanisms exist between activated EGFR[+ve] NSCs and quiescent NSCs to ensure stem cell homeostasis is poorly understood.

Over the last couple of decades, evidence has accumulated to suggest that adult NSCs are a cell of origin in certain brain tumours. One possibility is that these cells accumulate driver mutations over time, which compromise the normal controls on their proliferation and migration. These transformed NSCs could then escape the niche and acquire further mutations resulting in tumour formation. This has been hypothesised to be the case in glioblastoma (GBM), the most aggressive subtype of malignant glioma (Louis et al., 2007; Rispoli et al., 2014). In particular, the discovery of a subpopulation of cells within GBM tumours with stem cell characteristics, known as glioblastoma stem cells (GSCs), has lent weight to this notion. This GSC population in GBM tumours share many common features

with adult NSCs, including expression of stem and progenitor markers, such as CD133,
SOX2 and Nestin, self-renewal capacity, and the ability to generate multilineage progeny
(Cheng et al., 2013; Denysenko et al., 2010; Singh et al., 2003; Singh et al., 2004; Stieber et
al., 2014; Zhang et al., 2006). Most importantly, GSCs are also able to generate tumours in
mice which recapitulate all the classical features of GBM, even when injected in very small
numbers (Galli et al., 2004; Singh et al., 2004). For this reason, they are sometimes also
referred to as brain tumour initiating cells.

Adult NSCs can be isolated from the SVZ of murine brains and cultured as 3D neurospheres
or adherent cultures under conditions that promote symmetric self-renewal and prevent
differentiation (Conti et al., 2005; Pollard et al., 2006; Pollard et al., 2009). Furthermore, by
using NSCs engineered with oncogenic drivers, these *in vitro* culture systems can also be
useful to model brain tumour development and cross-comparison with wild-type
counterparts. Here we exploit this system to understand how transforming mutations affect
the quiescent or activation status of NSCs. We find that transformed NSCs are refractory to
quiescent inducing signals. Importantly, we also demonstrate that transformed NSCs signal
via Notch and RBPJ to induce a quiescent-like state in surrounding wild-type NSCs. These
results therefore provide new insight into how oncogenic mutations affect not only the
proliferative potential of transformed cells, but also provide them with a competitive
advantage in the niche by suppressing the proliferation of surrounding cells.

Results

Transformed NSCs are refractory to quiescence-inducing signals.
To understand how transforming mutations affect the quiescent or activation status of NSCs
we compared NSCs derived from wild-type mice to those isolated from *Ink4a/Arf$^{/-}$* mice and
transduced with a retrovirus expressing the EGFRvIII mutation (Bruggeman et al., 2007).
This combination of mutations is frequently observed in human GBM (Crespo et al., 2015)
and induces transformation in NSCs, as indicated by their ability to generate tumours *in vivo*
that recapitulate many of the features of human GBM tumours (Bruggeman et al., 2007;
Marques-Torrejon et al., 2018). We first compared the proliferative potential of both these
cell types. We observed that, when grown in conditions that promote the
activated/proliferative NSC state (Pollard et al., 2006), transformed NCSs displayed a small
growth advantage (Figure 1A). Analysis of the prevalence of BrdU incorporation revealed
that this difference was most likely accounted for by a small difference in the rate of cell
division at day four of culture, when transformed cells divided faster (Supplementary Figure
1A-B). However, overall these results indicate that the pattern of proliferation of wild-type
and *Ink4a/Arf$^{/-}$;* EGFRvIII NSCs (IE-NSCs) in growth-promoting conditions was remarkably
similar.

We next investigated if IE-NSCs have a similar response to quiescence-inducing signals
when compared to wild-type NSCs. BMP signalling is known to induce a reversible
quiescent-like state in hippocampal and embryonic stem cell-derived NSCs (Martynoga et al.,

2013; Mira et al., 2010) and the BMP antagonist Noggin enhances NSC expansion both *in*
vitro and *in vivo* (Lim et al., 2000; Mira et al., 2010; Yousef et al., 2015). For this reason, we
treated wild-type and transformed NSCs with 50ng/ml BMP4. We observed that, although
this induced similar levels of SMAD1/5/8 phosphorylation in both cell types (Supplementary
Figure 1C) and efficiently suppressed the proliferation of wild-type cells, it had little effect
on the growth of the transformed cells (Supplementary Figure 1D). This suggests that IE-
NSCs cells are refractory to the induction of quiescence by BMP signalling activation.

Transformed NSCs suppress the proliferation of surrounding NSCs.

The experiments described above allowed us to test the proliferative potential of wild-type
and transformed cells as isolated cell populations. However, we postulated that in the niche
these cell types are likely to co-exist. For this reason, we next studied the proliferation
patterns of both cell types in a 50:50 co-culture and compared these proliferative patterns to
the growth of the cells in homotypic (separate) cultures. IE-NSCs were GFP-labelled,
allowing us to quantify the relative cell proliferation rates in co-culture using flow cytometry.
Unexpectedly, we observed that, in contrast to the exponential growth observed for both cell
types in separate culture, in co-culture wild-type NSCs showed a significant growth arrest
(Figure 1B). Transformed NSCs therefore do not facilitate normal NSC growth as might be
anticipated; rather, they deploy some active mechanism to suppress their expansion and
outcompete them.

Quantification of the fold change in cell number indicated that the wild-type NSCs increased
their cell number only around 15-fold over six days in co-culture, compared to more than 50-
fold when cultured separately (Figure 1C). This dramatic reduction in the proliferation of
wild-type NSCs in the co-culture condition was also confirmed using BrdU incorporation
assays and quantifying expression mitotic marker phospho-Histone3 (Figure 1D-E and
Supplementary Figure 1C-D). These results suggest that IE-NSCs have an inhibitory effect
on the proliferation of wild-type NSCs. To test this possibility further, co-cultures were
established with varying proportions of each cell type, ranging from 10 to 90%, and the total
number of cells present in co-culture was kept constant. Analysis of the fold change in wild-
type NSC number revealed that the reduction in their proliferation was directly proportional
to the percentage of transformed NSCs present (Figure 1F), to the extent that, when 75% of
the cells in co-culture were transformed, the wild-type NSCs barely increased in number
during the 6-day experiment. Together, these results indicate that IE-NSCs suppress the
proliferation of wild-type NSCs.

Transformed cells induce a quiescent-like state in wild-type NSCs

To gain a deeper insight into the mechanism of proliferative arrest of co-cultured wild-type
NSCs, we compared their transcriptional profile to wild-type NSCs grown in separate culture.
Cells were isolated by FACS after 5 days of co-culture, and RNA-seq was performed.
Analysis of the differential gene expression yielded 1441 differentially expressed genes that
were then studied by Ingenuity Pathway Analysis, which uses functional annotations and
interactions of genes to identify enriched pathways and upstream regulators. It was

immediately apparent that the majority of the top canonical pathways that were enriched in
this comparison were related to cell cycle control and DNA repair (Figure 2A). For example,
the top enriched pathway was 'cell cycle control of chromosomal replication', which includes
important cell cycle-associated genes, such as multiple CDKs, MCM proteins and PCNA.
The genes in this pathway were broadly downregulated in wild-type NSCs in co-culture
compared to separately cultured cells (Figure 2B), consistent with a proliferation arrest when
surrounded by transformed cells.

One possible explanation for this cell cycle arrest is differentiation into a post-mitotic cell
types, such as immature neurons. The expression of NSC markers, together with markers for
intermediate progenitors and differentiated cell types (Zhang and Jiao, 2015), was therefore
explored in our RNA-seq data-set. No broad downregulation of classical NSC markers, such
as *Nestin*, *Gfap* and *Pax6* was observed in co-cultured wild-type NSCs compared to their
separate counterparts. Likewise, markers of intermediate or differentiated cell types, such as
Doublecortin, Olig2, S100b or *Tubb3,* were not upregulated (Supplementary Figure 2A),
indicating there was no increase in differentiation, an observation that was consistent with the
morphology of these cells that was not that of oligodentrocyte progenitor cells,
oligodentrocytes or neurons. Interestingly, we did observe an upregulation in the expression
of the glial markers, *Gfap, Glt1* and *Glast*, which are enriched in quiescent radial glia-like
NSCs (Codega et al., 2014; Llorens-Bobadilla et al., 2015). These observations suggest that
wild-type NSCs are not differentiating, but instead may be driven into a quiescent state.

To explore this last possibility we analysed our RNA-seq data-set for the expression of a
broader set of quiescent and activated NSC markers. We found that the quiescence-associated
transcriptional regulators, *Sox9*, *Id2*, *Id3* and *Klf9,* which are upregulated in quiescent SVZ
NSCs (Llorens-Bobadilla et al., 2015; Morizur et al., 2018), were all also upregulated in the
co-cultured wild-type NSCs (Figure 2C). We also observed that the markers of
activated/proliferating NSCs *Ascl1*, *Egr1*, *Fos* and *Sox11* (Andersen et al., 2014; Llorens-
Bobadilla et al., 2015; Morizur et al., 2018) were downregulated in these cells. When we
compared our RNA-seq data-set with a list of transcription factors and co-factors that have
been found to be differentially expressed between activated and quiescent SVZ NSCs
(Morizur et al., 2018), we found that of the 14 genes enriched in quiescent cells in this study,
10 were upregulated in co-cultured wild-type NSCs, and of the 61 genes enriched in activated
NSCs, 50 were downregulated (Supplementary Figure 2B). Together these findings therefore
support the hypothesis that the co-cultured NSCs are adopting a quiescent phenotype rather
than becoming senescent.

A key feature of a quiescent phenotype is its reversibility. To test if the proliferation arrest of
wild-type NSCs in co-culture is reversible, we sorted wild-type and transformed cells after 5
days in co-culture and re-plated them. In parallel, wild-type NSCs that had been cultured
separately were mixed with transformed cells just before sorting and then re-plated.
Importantly, we observed that previously co-cultured wild-type NSCs displayed a similar
proliferation rate to NSCs that had been separately cultured throughout, as judged by their
growth curves and rate of BrdU incorporation (Figure 2E-F). The reversible nature of the

proliferation arrest of co-cultured wild-type NSCs provides further evidence that they are
entering a quiescent state.

Cell contact is required for the inhibitory effect of transformed cells
To understand the mechanisms underlying the interaction between wild-type and transformed
NSCs, we first analysed the possible involvement of the mTOR signalling pathway. mTOR is
a metabolic regulator that senses growth factor and nutrient inputs, is activated in transit
amplifying progenitor cells and its inhibition induces a reversible quiescence-like phenotype
in adult NSCs (Paliouras et al., 2012). Analysis of the expression of ribosomal protein S6
phosphorylation (p-S6), a read-out of mTOR activity indicated that mTOR signalling levels
were reduced in the co-cultured wild-type NSCs, not only relative to these same cells in
separate culture, but also to transformed NSCs in co-culture (Figure 3A). We next tested if
constitutive mTOR activation was sufficient to prevent these wild-type NSCs from entering
quiescence in co-culture. For this we mutated *Tsc2*, an mTOR repressor in wild-type NSCs,
using CRISPR-Cas9 targeting (Supplementary Figure 2C). However, we found that, although
this was sufficient to sustain strong mTOR pathway activation, $Tsc2^{-/-}$ NSCs still entered a
proliferation arrest when co-cultured with transformed NSCs (Supplementary Figure 2D-F).
This indicates that mTOR inhibition is not the primary reason for the quiescent phenotype of
co-cultured wild-type NSCs.

Next, we investigated an alternative possibility that cell contact is required to induce the
growth arrest of wild-type NSCs. We first used a transwell assay, where the two cell types
are cultured in the same well, separated by a permeable membrane that allows the exchange
of factors in the media while physically separating the cells. When wild-type cells were co-
cultured with transformed cells separated by a transwell, they grew similarly to wild-type
cells in homotypic cultures (Figure 3B). This suggests a requirement for cell-cell contact and
that secreted paracrine factors are not sufficient to induce the quiescence. We further tested
this possibility by using fences, which are metal inserts that divide the well into an inner and
outer ring One cell type is plated in the inner ring and the second in the outer ring (Figure 3C)
and once they are growing adherently, the inserts can be removed leaving behind a small gap
between the two populations. Using this system we compared the proliferation rate of wild-
type NSCs that were separated from transformed NSCs to their proliferation when separated
from other wild-type NSCs or when mixed with transformed cells. We found that the rate of
BrdU incorporation of the wild-type NSCs only decreased when these cells were in physical
contact with transformed cells (Figure 3C). This indicates that cell-cell contact is most likely
required for transformed cells to inhibit the proliferation of wild-type NSCs.

Transformed cells signal via Notch and Rbpj to induce quiescence in wild-type NCSs
The Notch pathway is a highly conserved signalling pathway requiring cell-cell contact that
plays a central role in the regulation of embryonic and adult neurogenesis (Ables et al., 2011;
Urban and Guillemot, 2014). To test the potential role for Notch signalling we first explored
the expression of Notch signalling pathway components in our RNA-seq data. Interestingly,
genes related to the Notch signalling pathway were enriched in the comparison between co-

cultured wild-type and transformed NSCs (p-value = $5.94E^{-6}$, 21/38 genes differentially
expressed with adjusted p-value <0.1). Overall, the majority of Notch-related genes were
upregulated in wild-type NSCs compared to transformed NSCs, including Notch targets and
receptors, while there was some evidence that Notch ligands were more highly expressed on
transformed NSCs in co-culture (Figure 3D). Analysis by real-time qPCR of sorted samples
also revealed that the expression of the Notch target genes *Hes1, Hes5, Hes7, Hey1, Hey2*
and *Nrarp* are upregulated in co-cultured wild-type NSCs (Figure 3E). This suggests that
Notch pathway activity is higher in co-cultured wild-type NSCs than in transformed cells and
raises the possibility that transformed cells may be signalling to wild-type NSCs via this
pathway.

To test the functional significance of this difference in Notch activation, two complementary
γ-secretase inhibitors were used to block Notch signalling. We observed that both LY411575
(LY) and crenigacestat not only reduced Notch target gene expression in NSCs, but also
partially rescued the proliferation arrest of wild-type NSCs in co-culture with transformed
cells. Evidence for this is that the growth curves and BrdU incorporation rates of co-cultured
NSCs treated with the γ-secretase inhibitors were only slightly lower than their growth is
separate culture (Figure 3G-H and Supplementary Figure 3A-E). This suggests that Notch
signalling is required for the induction of quiescence in wild-type NSCs by transformed cells.

To further test the requirement of Notch signalling in wild-type NSCs we mutated *Rbpj*, a
key effector of this pathway, as well as *Notch1* and *Notch2* by CRISPR/Cas9 targeting
(Figure 4A, D and G). Importantly, we found that *Rbpj*$^{-/-}$, *Notch1*$^{-/-}$ or *Notch2*$^{-/-}$ NSCs were
not susceptible to proliferation arrest when co-cultured with IE-NSCs. For example, we
observed that all three Notch pathway mutant cell lines displayed similar growth rates in
separate and co-culture (Supplementary Figure 4A-D). Similarly, the fold change in cell
number and BrdU incorporation rates of co-cultured *Rbpj*$^{-/-}$, *Notch1*$^{-/-}$ and *Notch2*$^{-/-}$ NSCs was
similar to separate culture and significantly higher than that of wild-type NSCs co-cultured
with IE-NSCs (Figure 4B-C, E-F and H-I). Interestingly, both the growth curves and final
cell counts of *Rbpj*$^{-/-}$ NSCs indicated that these cell types even out-grew IE-NSCs in co-
culture (Figure 4B and Supplementary Figure 4B), suggesting that disrupting Notch
signalling may be sufficient to provide NSCs with a competitive advantage against
transformed NSCs. Together, these data indicate that Notch signalling is required for the
suppression of proliferation of wild-type NSCs by transformed cells.

Finally, to exclude the possibility that Notch signalling is required in transformed NSCs, we
mutated *Rbpj* in this cell type (Supplementary Figure 4E). We observed that IE-NSCs still
induced the proliferation arrest of wild-type NSCs, as shown by the growth curves, final cell
numbers and rate of BrdU incorporation (Supplementary Figure 4F-H). Therefore Notch
signalling is not required in transformed cells for their growth inhibition of wild-type NSCs.

Discussion

In conclusion, our data indicate that transformed cells activate Notch signalling in adjacent
wild-type NSCs to induce in these a quiescent state. Importantly, the role of Notch signalling

in regulating the activated versus quiescent state of NSCs is borne out by the work of others.
Increased expression of Notch signalling pathway components in quiescent NSCs compared
to activated NSCs has been widely reported in RNA-sequencing studies (Andersen et al.,
2014; Llorens-Bobadilla et al., 2015; Morizur et al., 2018). Additionally, NOTCH receptor
and RBPJ expression have been previously confirmed in both quiescent and activated NSC
populations in the SVZ, and NOTCH2 has been shown to mediate signals promoting NSC
quiescence within the niche (Engler et al., 2018). The Notch ligand Dll1 has also been
reported to be expressed exclusively in activated NSCs and transit-amplifying cells
(Kawaguchi et al., 2013) and its deletion resulted in excessive production of activated NSCs
and a depletion of the quiescent NSC pool. Similar observations have also been made in
zebrafish, where activated radial glial cells were shown to revert to a quiescent state by Notch
pathway activation (Chapouton et al., 2010). Therefore our results imply that transformed
cells highjack an evolutionarily conserved pathway regulating vertebrate NSC quiescence
and use it to gain a competitive advantage over other NSCs in the niche.

Our work also has implications for glioblastoma tumour progression. The SVZ has been
shown to be a permissive region for glioblastoma growth, as tumours in contact with this
region are associated with decreased patient survival and increased recurrence (Khalifa et al.,
2017; Mistry et al., 2017). This niche has also been suggested to serve as a reservoir for
malignant cells away from the tumour mass and may therefore support tumour recurrence
(Piccirillo et al., 2015). In humans it has been shown that glioblastoma can arise from SVZ
cells with low-level driver mutations (Lee et al., 2018), further highlighting the importance of
the SVZ for oncogenic cell propagation. We therefore speculate that the mechanisms we
describe operate during early premalignant disease to support tumorigenesis. Here,
transformed NSCs provide negative feedback signals to surrounding normal NSCs limiting
their self-renewal and enforcing their quiescence. By this mechanism, oncogenic NSCs can
ensure their preferential self-renewal and differentiation, therefore increasing both their cell
number and their likelihood of giving rise to progenitor cells and exiting the niche. This
competitive advantage may be an important early step in the development of glioma and
glioblastoma and therefore understanding it may help develop therapeutic approaches.

Materials and Methods

Neural Stem Cell Cultures
Adult murine NSCs were derived from the SVZ by the protocol described in [5].
Transformed *Ink4a/Arf*$^{-/-}$; EGFRvIII NSCs (IE-NSCs) were generated by the Maarten van
Lohuizen laboratory (Netherlands Cancer Institute). NSCs were isolated from *Ink4a/Arf*$^{-/-}$
mice and transduced with EGFRvIII pMSCV retrovirus (Bruggeman et al., 2007). All cells
were cultured at 37°C in an atmosphere with 5% CO_2. NSCs were maintained in serum-free
adherent growth conditions on poly-L-lysine-coated plates (Sigma). 1µg/ml Laminin (Sigma)
was also added to the growth media to promote adherent growth. Growth medium was
Dulbecco's Modified Eagle Medium: Nutrient Mixture F12 (DMEM/F12) (Gibco)

supplemented with N2 (1:200, Invitrogen), B27 (1:100, Invitrogen), 1x non-essential amino
acids (Invitrogen), 0.1% BSA (Gibco), 100µM β-mercaptoethanol (Invitrogen), 1.5mg/ml
glucose (Sigma), 15mM HEPES (Gibco), 2mM L-glutamine (Gibco), 100U/ml penicillin-
streptomycin (Gibco), 20ng/ml EGF and 10ng/ml bFGF (both Peprotech).

Co-culture assays
For six-day co-culture assays, 40,000 cells per well were seeded in a 24-well plate either
separately or as a 50:50 mix of WT and transformed NSCs. Standard growth media was used
throughout and was changed on days three, four and five. The proportion of each cell type
was assessed by determining the percentage of GFP-positive cells by flow cytometry. Cells
were harvested with accutase and resuspended in PBS containing 3% FCS. All samples were
transferred into strainer-cap flow cytometer tubes for analysis and 100ng/ml propidium
iodide (PI) was added. Flow cytometry was performed on an LSR-II Analyser (BD
Biosciences) with FACS Diva software (BD Biosciences). Gates were applied to exclude
cellular debris, cell doublets and non-viable PI-positive cells. The percentage of GFP-positive
and GFP-negative cells was determined with gates set using separate cell preparations.
Separate cultures were also mixed and analysed by flow cytometry. Viable cell counts at each
time point were performed using a Vi-Cell XR counter (Beckman Coulter) and calculated for
each cell type in separate and co-culture by applying the percentages from the flow cytometry
analysis.

For transwell assays, cell culture inserts with 0.4µm pores (Merck Millipore) were placed in a
6-well plate and coated with PLL. 140,000 NSCs were seeded in NSC growth media either
onto the insert or below in the well. The growth of WT NSCs seeded below either WT or
transformed NSCs or a 50:50 mix was evaluated over six days. For fence assays, 24-well
plates were coated with PLL and fences (Aix-Scientifics) were placed in each well. 100,000
NSCs were seeded in the inner ring and 400,000 in the outer ring. The fences were removed
24 hours later and the media was replaced. The following day, 10µM BrdU was added to the
wells for two hours. The fences were then replaced into the well and the cells in the inner ring
were detached with accutase and harvested. The cells were then fixed in 2% formaldehyde
and analysed for BrdU expression by flow cytometry as described below.

Flow cytometry immunolabelling
Cells were fixed in 2% formaldehyde and permeabilised in ice-cold 90% methanol. Cells
were incubated with the primary antibody (anti-p-rpS6, CST 5364, 1:200, anti-BrdU, CST
5292, 1:200) for one hour in PBS with 1% BSA. AlexaFluor 546 secondary antibodies
(Molecular Probes) were used for detection) diluted 1:2000 in PBS containing 1% BSA.
Secondary incubation was for 30 minutes. Cell debris and doublets were excluded and
controls with the primary antibody omitted were used to gate positive and negative cells.
Flow cytometry was performed on an LSR-II Analyser (BD Biosciences) with FACS Diva
software (BD Biosciences) and data were analysed using FlowJo software.

For BrdU analysis NSCs were incubated for two hours with 10µM BrdU prior to fixation.
Before blocking, NSCs were then incubated for 45 minutes at 37°C in DNase I solution

consisting of 20µl DNase I (Promega RQ1) in 250µl DNase buffer solution. The
immunolabelling protocol was then followed as detailed above.

FACS
180,000 cells per well were seeded in a 6 well plate either separately or as a 50:50 mix of WT
and transformed NSCs and cultured under the standard assay conditions until day five. GFP-
negative WT NSCs were sorted using a FACS Aria cell sorter (BD Biosciences) from both
separate and mix samples. Gates were applied to exclude cell debris and doublets. Cells were
recovered into growth media at room temperature. Cells were then either lysed for
RNA/protein extraction or, for replating experiments, counted and replated into a 24-well
plate at 100,000 cells per well.

Western blot analysis
Cells were lysed directly in Laemmli buffer (50mM Tris-HCl (pH 6.8), 2% SDS, 10%
glycerol in ddH$_2$O) and heated at 95°C for 10 minutes. Protein quantification was carried out
using a BCA Protein Assay kit (Pierce) according to the manufacturer's instructions. 15-20µg
denatured protein from each sample was run in XT sample buffer (BioRad) with 0.25% β-
mercaptoethanol on 10% or 12% SDS-PAGE Bis-Tris gels (pre-cast BioRad). Proteins were
then blotted onto PVDF membranes (Thermo Fisher Scientific) using vertical wet transfer at
100V for one hour at 4°C. Blocking was performed in 5% milk in TBST buffer and primary
antibody incubation was done overnight at 4^{o}C in TBST containing 5% BSA. The following
antibodies were used at the stated concentration: anti-α-tubulin (CST 3873, 1:1000), anti-β-
actin (CST 5970, 1:1000), anti-RBPJ (CST 5313, 1:1000), anti-Notch1 (CST 3608, 1:1000),
anti-Notch2 (CST 5732, 1:1000), anti-p-SMAD1/5/8 (CST 9511, 1:000), anti-TSC2 (CST
4308, 1:1000). Horseradish peroxidase-conjugated secondary antibodies (Abcam) diluted
1:5000 in 5% milk in TBST were used for detection by the addition of an enhanced
chemiluminescence (ECL) substrate (Promega). Western blot quantification was performed
using Fiji software (Schindelin et al., 2012).

RNA Extraction and Quantitative RT-PCR
RNA was extracted with the RNeasy mini kit (Qiagen) and SuperScript III reverse
transcriptase (Thermo Fisher Scientific) was used for cDNA synthesis according to
manufacturer's instructions. RT-qPCR was performed using SYBR-Green Mastermix with
ROX reference dye (Sigma). A list of all primer sequences used can be found in Table 1
below. RT-qPCR was performed on an ABI 7900HT Fast Real Time PCR machine (Applied
Biosystems) and analysed on SDS Biosystem software. Relative gene expression was
calculated using the ΔΔCt method normalised to β-actin expression (Livak and Schmittgen,
2001).

RNA-sequencing analysis
356 Sorting was performed by FACS at day five of the six-day assay as described above for three
357 biological repeats. PI was added to exclude non-viable cells. Cells were recovered into
358 growth media and then resuspended in RLT lysis buffer (Qiagen). RNA extraction was

performed using the Qiagen RNeasy kit according to the manufacturer's instructions. RNA
samples were quantified using a Qubit fluorometer (Thermo Fisher Scientific) and the quality
assessed by TapeStation electrophoresis (Agilent). mRNA was isolated using oligo dT beads.
mRNA was then fragmented, converted to cDNA and ligated to Illumina adapters. Following
sample indexing, the quality of cDNA libraries was also assessed by TapeStation.
Sequencing was performed using the HiSeq 4000 system (Illumina). Sequencing reads were
aligned to the mouse genome (mm9) using TopHat2 (Kim et al., 2013) and differential
expression was analysed using the DESeq2 package (Love et al., 2014). The resulting gene
sets were analysed using Ingenuity Pathway Analysis (IPA) software (Qiagen) (Kramer et al.,
2014). This utilises the Ingenuity Knowledge Base for functional annotations and interactions
of genes. A 'Core Analysis' was performed for each comparison on genes differentially
expressed with a padj cut-off <0.1, corresponding to a false discovery rate (FDR) of 10%.
This identified enriched Canonical Pathways and Upstream Regulators in each comparison.

Generation of *Rbpj*[-/-], *Notch1*[-/-], *Notch2*[-/-] and *Tsc2*[-/-] NSCs

gRNAs for targeting of *Rbpj*, *Notch1* and *Notch2* were cloned into the pX330-U6-
Chimeric_BB-CBh-hSpCas9 (PX330) expression plasmid (gift from Feng Zhang, Addgene
plasmid #42230) using the protocol provided by the Zhang laboratory (Ran et al., 2013).
gRNAs for targeting *Tsc2* were cloned into gRNA expression plasmids (gift from George
Church, Addgene plasmid #41824) (Mali et al., 2013). These were then co-transfected with
an hCas9 plasmid (gift from George Church, Addgene plasmid #41815). gRNA sequences
can be found in Supplementary Table 2. NSCs were transfected with plasmids by
nucleofection (AMAXA 2B, Lonza) using programme A-033. NSCs were harvested with
Accutase and $2-3 \times 10^6$ cells were resuspended in 100µl mouse neural stem cell nucleofector
buffer (Lonza) with 2µg plasmid DNA. NSCs were either transfected with 2ug PX330 vector
containing gRNAs targeted to *Rbpj*, *Notch1* or *Notch2* or 0.6µg each of hCas9 and the two
gRNA expression plasmids containing the *Tsc2*-targeting gRNAs. NSCs were also co-
transfected with 0.125ug of Puro-pPyCAGIP vector or linear hygromycin marker (Clontech
631625) for selection. Empty vector or Cas9 only controls were also performed. Cells were
recovered post-transfection with pre-warmed NSC growth media into a $10cm^2$ dish. Selection
with 0.25ug/ml puromycin or 250ug/ml hygromycin was performed from 48 hours post-
transfection until the emergence of resistant colonies. These were then picked manually with
a p20 pipette and transferred to a 48 well plate for expansion and analysis.

Quantification and Statistical Analysis

Statistical analysis and data representation were performed using GraphPad Prism software.
Statistical methods used are indicated in the relevant figure legends. No randomisation or
blinding was used in experiments. Sample sizes were selected based on the observed effects
and listed in the figure legends. The sample size (n) is described in the figure legends and
refers to the number of independent replicate experiments performed. Adjusted p values are
displayed as * $p<0.05$, ** $p<0.01$ and *** $p<0.001$.

Data and Code availability

The raw data for the RNA-sequencing of wild-type and transformed NSCs are being deposited in the arrayexpress database and an accession code that will be made available prior to acceptance. The authors declare that all data supporting the findings of this study are available within the article and its supplementary information files or from the corresponding author upon reasonable request.

Key Resources Table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
BrdU	Cell Signalling Technologies	Cat#5292, RRID:AB_10548898
NOTCH1	Cell Signalling Technologies	Cat#3608, RRID:AB_2153354
NOTCH2	Cell Signalling Technologies	Cat#5732, RRID:AB_10693319
p-H3	Merck Millipore	Cat#07-145, RRID:AB_11214453
p-rpS6	Cell Signalling Technologies	Cat#5364, RRID:AB_10694233
P-SMAD1/5/9	Cell Signalling Technologies	Cat#9511, RRID:AB_331671
RBPJ	Cell Signalling Technologies	Cat#5313, RRID:AB_2665555
TSC2	Cell Signalling Technologies	Cat#4308, RRID:AB_10547134
β-Actin	Cell Signalling Technologies	Cat#4970, RRID:AB_2223169
Goat anti-rabbit 546	Thermo Fisher Scientific	Cat#A-11010, RRID:AB_2534077
Goat anti-mouse 546	Thermo Fisher Scientific	Cat#A-11003, RRID:AB_2534071
Goat anti-rabbit HRP	Santa Cruz Biotechnology	Cat#sc-2357, RRID:AB_628497
Goat anti-mouse HRP	Santa Cruz Biotechnology	Cat#sc-516102, RRID:AB_2687626
Chemicals, Peptides, and Recombinant Proteins		
LY411575	Selleckchem	Cat#S2714
Crenigacestat	Selleckchem	Cat#S7169
EGF	Peprotech	Cat#315-09
FGF	Peprotech	Cat#100-18b
Laminin	Sigma	Cat#L2020
Poly-L-Lysine	Sigma	Cat#P1274
Critical Commercial Assays		
RNeasy Mini Kit	Qiagen	Cat #74104
BCA Protein Assay Kit	Pierce	Cat #23225
Mouse neural stem cell nucleofector kit	Lonza	Cat #VPG-1004

Deposited Data		
RNA seq data of NSCs in separate or co-culture	This paper	GEO:
Experimental Models: Cell Lines		
Wild-type NSCs	Pollard Lab, 'ANS7'	
Ink4a/Arf⁻/⁻, EGFRvIII NSCs	Lohuizen Lab, 'IENS'	[8]
Oligonucleotides		
qRT-PCR primers	This paper	Supplementary Table 1
gRNA oligonucleotides	This paper	Supplementary Table 2
Recombinant DNA		
pX330-U6-Chimeric_BB-CBh-hSpCas9	Feng Zhang via Addgene	Addgene #42230 [35]
gRNA cloning plasmid	George Church via Addgene	Addgene #41824 [37]
hCas9 expression plasmid	George Church via Addgene	Addgene #41815 [37]
Linear hygromycin marker	Clontech	Cat #631625
Puro-pPyCAGIP	Rodriguez laboratory	pPyCAGIP
Software and Algorithms		
GraphPad Prism	GraphPad	www.graphpad.com/ scientificsoftware/prism
FIJI (ImageJ)	[38]	Imagej.net/fiji
R		https://www.r-project.org/
TopHat2	[32]	https://ccb.jhu.edu/software/tophat/index.shtml
DESeq2	[33]	https://bioconductor.org/packages/release/bioc/html/DESeq2.html
Ingenuity Pathway Analysis	Qiagen, [34]	https://www.qiagenbioinformatics.com/products/ingenuity-pathway-analysis/

Acknowledgments

406 We would like to thank Stephen Rothery for guidance and advice with confocal microscopy.
407 Gratitude also goes to James Elliot for performing cell sorts. Research in Tristan Rodriguez
408 lab was supported by the MRC project grant (MR/N009371/1), by the Rosetrees Trust grant
409 (M693) and by the British Heart Foundation centre for research excellence. Katerina Lawlor
410 was supported by an NHLI PhD studentship.

Conflicts of interests

413 The authors declare no competing financial or non-financial interests.
414

Figure Legends

Figure 1. **WT NSCs show reduced proliferation in the presence of transformed NSCs.**
A Growth curves of WT and transformed IE NSCs cultured separately over six days. **B** Growth curves of wild-type and transformed NSCs in a 50:50 co-culture. **C** Quantification of fold change in cell number relative to seeding density for each culture condition. WT NSCs dramatically reduce their proliferation while transformed cells increase their proliferation relative to separate culture. N>10, ANOVA followed by Sidak's multiple comparisons test. **D** Proportion of BrdU-positive cells in each culture condition at day five following a two-hour BrdU chase. WT NSCs show greatly reduced BrdU incorporation in co-culture. N=10, ANOVA followed by Sidak's multiple comparisons test. **E** Representative western blot for phospho-histone-3 (p-H3) in WT NSCs sorted at day five and quantification of p-H3 density relative to β-actin. p-H3 expression is markedly reduced in co-cultured WT NSCs. N=3, Student's paired T test. **F** Fold change cell number of WT NSCs cultured in co-cultures with varying proportions of transformed NSCs. WT NSCs show smaller increases in cell number in co-cultures with a greater proportion of transformed NSCs. N=3.

Figure 2. Wild-type NSCs adopt a quiescent phenotype in the presence of transformed NSCs
A RNA sequencing was performed on NSCs sorted by FACS following five days' culture in either homogenous or co-culture conditions. Differential gene expression analysis was performed on these samples from three independent experiments. The top canonical pathways enriched in the comparison of WT NSCs in separate or mixed culture. Pathways related to cell cycle and DNA repair are highly enriched. **B** Heat map showing the fold change values for all genes in the 'Cell Cycle Control of Chromosomal Replication' pathway for this comparison with an adjusted p-value <0.1. **C** Heat map showing the change in gene expression of certain transcription factors (TFs) between the WT NSCs cultured separately or in co-culture. These transcription factors have been particularly highlighted by transcriptional profiling studies that sought to compare quiescent and activated NSCs. Quiescence-associated genes were broadly upregulated in the co-cultured NSCs, while activation-associated genes were broadly downregulated. **E** Growth curves showing cell numbers for three days following replating. **F** Proportion of BrdU-positive cells at day three after re-plating. No difference could be found in the growth of WT NSCs which had previously been co-cultured compared to cells grown separately. N=3. **E** and **F** WT NSCs were cultured for 5 days either in co-culture or separately and then sorted by FACS and replated at 100,000 cells/well.

Figure 3. Quiescence of WT NSCs is induced by direct contact with transformed NSCs and is associated with increased Notch activation
A Flow cytometry plot showing p-rpS6 staining in WT and transformed NSCs in separate and mixed culture and quantification of median fluorescence intensity. WT NSCs show reduced expression specifically in the co-culture condition. N=6, ANOVA followed by Sidak's multiple comparisons test. **B** Growth curves of WT NSCs cultured below transwell

inserts with WT, transformed or mixed cultures. No difference was observed between
conditions. N=3. **C** Percentage of BrdU-positive WT NSCs from the inner ring of the three
different cultures shown. When cells were mixed in each well WT NSCs showed a decrease
in the proportion of proliferating cells (Mix/Mix). However, WT NSCs surrounded by a ring
of transformed NSCs (Trans/WT) did not show a reduction in proliferation relative to WT
NSCs surrounded by WT NSCs (WT/WT). N=3, one-way ANOVA followed by Tukey's
multiple comparisons test. Direct cell contact is therefore required for reduced WT NSC
proliferation **D** Heat map showing changes in gene expression in co-cultured transformed and
WT NSCs. Notch target genes and receptors are upregulated in WT NSCs, while Notch
ligands show a more mixed expression with several downregulated relative to transformed
NSCs. **E** RT-qPCR validation of Notch target gene expression in WT NSCs relative to
transformed NSCs sorted from the co-culture. N=3, two-way ANOVA followed by Sidak's
multiple comparisons test. **F** Fold change cell number quantification of NSC growth with and
without addition of LY411575 at 2µM. N=8, two-way ANOVA with Sidak's multiple
comparisons test. **G** Proportion of BrdU positive cells following a 24 hour incubation with
LY411575. N=4, two-way ANOVA with Sidak's multiple comparisons test. WT NSCs in co-
culture show increased proliferation in the presence of the γ-secretase inhibitor, LY411575.

Figure 4. **Disruption of Notch signalling rescues WT NSC proliferation in co-culture.**
A Western blot showing absence of RBPJ protein in both *Rbpj$^{-/-}$* clones. **B** Fold change cell
number quantification of EV and *Rbpj$^{-/-}$* NSCs in separate and co-culture with transformed
NSCs. EV control NSCs show decreased growth in co-culture but this is not the case for the
Rbpj$^{-/-}$ NSCs. N=4, two-way ANOVA followed by Sidak's multiple comparisons test. **C**
Proportion of BrdU-positive cells in separate and co-culture showing decrease in EV control
but not in *Rbpj$^{-/-}$* NSCs. N=5, two-way ANOVA followed by Sidak's multiple comparisons
test. **D** Western blot showing absence of NOTCH1 protein in both *Notch1$^{-/-}$* clones. **E** Fold
change cell number quantification of EV and *Rbpj$^{-/-}$* NSCs in separate and co-culture with
transformed NSCs. EV control NSCs show decreased growth in co-culture but this is not the
case for the *Notch1$^{-/-}$* NSCs. N=4, two-way ANOVA followed by Sidak's multiple
comparisons test. **F** Proportion of BrdU-positive cells in separate and co-culture showing
decrease in EV control but not in *Notch1$^{-/-}$* NSCs. N=5, two-way ANOVA followed by
Sidak's multiple comparisons test. **G** Western blot showing absence of NOTCH2 protein in
both *Notch2$^{-/-}$* clones. **H** Fold change cell number quantification of EV and *Notch2$^{-/-}$* NSCs in
separate and co-culture with transformed NSCs. EV control NSCs show decreased growth in
co-culture but this is not the case for the *Notch2$^{-/-}$* NSCs. N=4, two-way ANOVA followed
by Sidak's multiple comparisons test. **I** Proportion of BrdU-positive cells in separate and co-
culture showing decrease in EV control but not in *Notch2$^{-/-}$* NSCs. N=5, two-way ANOVA
followed by Sidak's multiple comparisons test.

Supplemental Figure 1
A Representative flow cytometry plot of BrdU staining in separately cultured NSCs at day
five. **B** Time-course of BrdU staining throughout the assay in separate culture. N=5, ANOVA

followed by SIdak's multiple comparisons test. **C** Fold change cell number quantification of
NSCs cultured in the presence or absence of 50ng/ml BMP4. BMP4 treatment reduces the
growth of WT, but not transformed, NSCs. N=4, two-way ANOVA followed by Sidak's
multiple comparisons test. **D** Western blot showing increased p-SMAD1/5/8 expression in
NSCs treated with 50ng/ml BMP4 for 24 hours relative to control NSCs (Con). **E**
Representative flow cytometry plot of BrdU staining in co-cultured NSCs at day five. **F**
Time-course of BrdU staining throughout the assay in co-culture showing reduction in BrdU
incorporation in WT NSCs. N=5, ANOVA followed by Sidak's multiple comparisons test.

Supplemental Figure 2
A Heat map showing changes in gene expression in the 'WT Mix vs WT Sep' comparison.
Classical marker genes for different cell populations are shown including those used
commonly to identify adult NSCs. GFAP $\log_2$ fold change is off the scale at 5.4. **B**
Comparison to published transcriptional analysis of quiescent and activated NSCs. Left -
Transcription factors and co-factors differentially expressed in quiescent and activated NSCs
from [18]. Right – Heat map showing the expression of the genes identified by Morizur et al
in the comparison of WT NSCs in separate (WT Sep) or co-culture (WT Mix). Genes
enriched in quiescent NSCs were broadly upregulated in co-cultured WT NSCs, while those
enriched in activated NSCs were broadly downregulated. **C** Western blot showing absence of
TSC2 expression in *Tsc2*$^{-/-}$ clones. The control Cas9 clone was transfected with Cas9 only
without the gRNAs. **D** Growth curves of transformed NSCs and Tsc2$^{-/-}$ or Cas9 control NSCs
in separate or co-culture. *Tsc2*$^{-/-}$ clones grow more slowly than control and still show reduced
growth in co-culture. N=4. **E** Fold change cell number quantification of growth curves. N=4,
two-way ANOVA followed by Sidak's multiple comparisons test. **F** Expression of p-rpS6 in
Cas9 and *Tsc2*$^{-/-}$ clones as determined by flow cytometry. mTOR activation is increased in
Tsc2$^{-/-}$ NSCs and remains high in co-culture despite reduced growth. N=3.

Supplemental Figure 3
A RT-qPCR analysis of the Notch target genes, *Hes1*, *Hes5* and *Nrarp* in WT NSCs treated
with 2µM LY411575. $\log_2$ fold change relative to control NSCs. N=3. **B** Growth curves of
WT and transformed NSCs cultured in the standard assay with or without LY411575 addition
from day three. **C** RT-qPCR analysis of the Notch target genes, *Hes1*, *Hes5* and *Nrarp* in WT
NSCs treated with 1µM crenigacestat. $\log_2$ fold change relative to control NSCs. N=3. **D**
Growth curves of WT and transformed NSCs cultured in the standard assay with or without
crenigacestat addition from day three. N=8, two-way ANOVA followed by Sidak's multiple
comparisons test. **E** Proportion of BrdU positive cells following a 24 hour incubation with
crenigacestat. N=4, two-way ANOVA with Sidak's multiple comparisons test. The γ-
secretase inhibitor crenigacestat also increases WT NSC proliferation in co-culture.

Supplemental Figure 4
A Growth curves of control EV clone cultured separately or in co-culture with transformed
NSCs. **B**   Growth curves of *Rbpj*$^{-/-}$ clones cultured separately or in co-culture with
transformed NSCs. **C** Growth curves of *Notch1*$^{-/-}$ clones cultured separately or in co-culture

with transformed NSCs. **D** Growth curves of *Notch2*$^{-/-}$ clones cultured separately or in co-
culture with transformed NSCs. N>4. **E** Western blot showing absence of RBPJ protein from
Rbpj$^{-/-}$ transformed NSCs. **F** Fold change cell number of WT NSCs in separate or co-culture
with EV or *Rbpj*$^{-/-}$ transformed NSCs. N=5, two-way ANOVA followed by Sidak's multiple
comparisons test. **G** Proportion of BrdU-positive WT NSCs in separate or co-culture with EV
or *Rbpj*$^{-/-}$ transformed cells. N=3, two-way ANOVA followed by Sidak's multiple
comparisons test. **H** Growth curves of transformed EV and *Rbpj*$^{-/-}$ clones cultured separately
or in co-culture with WT NSCs. Both *Rbpj*$^{-/-}$ clones suppress the proliferation of WT NSCs to
a comparable degree to the EV control transformed cells.

Supplementary Table 1. Primers used in qPCR
545

Target gene	Forward primer	Reverse primer
Hes1	GGAAATGACTGTGAAGCACCTCC	GAAGCGGGTCACCTCGTTCATG
Hes5	GCCCGGGGTTCTATGATATT	GAGTTCGGCCTTCACAAAAG
Hes7	CATCAACCGCAGCCTAGAAGAG	CACGGCGAACTCCAGTATCTCC
Hey1	CCAACGACATCGTCCCAGGTTT	CTGCTTCTCAAAGGCACTGGGT
Hey2	TGAAGATGCTCCAGGCTACAGG	CCTTCCACTGAGCTTAGGTACC
Nrarp	CAGACAGCACTACACCAGTCAG	CCGAAAGCGGCGATGTGTAGC
β-actin	GGCACCACACCTTCTACAATG	GGGGTGTTGAAGGTCTCAAAC

547 **Supplementary Table 2. gRNA sequences for deletion**
548

Target gene	gRNA1	gRNA2
Rbpj	TGCAGTGGACGACGACGAGT	GTGGACGACGACGAGTCGGA
Notch1	GGCGTTCAGTGCCTACACAA	
Notch2	GTCCACCTGCATTGACCGCG	AAGTGCATCGATCACCCGAA
Tsc2	GAAGGCCGGCCTACCTCATT	GCATGGCTCTTACAGGTACA

References

Ables, J.L., Breunig, J.J., Eisch, A.J., and Rakic, P. (2011). Not(ch) just development: Notch
signalling in the adult brain. Nat Rev Neurosci *12*, 269-283.

Andersen, J., Urban, N., Achimastou, A., Ito, A., Simic, M., Ullom, K., Martynoga, B.,
Lebel, M., Goritz, C., Frisen, J.*, et al.* (2014). A transcriptional mechanism integrating inputs
from extracellular signals to activate hippocampal stem cells. Neuron *83*, 1085-1097.

Bruggeman, S.W., Hulsman, D., Tanger, E., Buckle, T., Blom, M., Zevenhoven, J., van
Tellingen, O., and van Lohuizen, M. (2007). Bmi1 controls tumor development in an
Ink4a/Arf-independent manner in a mouse model for glioma. Cancer Cell *12*, 328-341.

Chapouton, P., Skupien, P., Hesl, B., Coolen, M., Moore, J.C., Madelaine, R., Kremmer, E.,
Faus-Kessler, T., Blader, P., Lawson, N.D.*, et al.* (2010). Notch activity levels control the
balance between quiescence and recruitment of adult neural stem cells. J Neurosci *30*, 7961-
7974.

Cheng, L., Huang, Z., Zhou, W., Wu, Q., Donnola, S., Liu, J.K., Fang, X., Sloan, A.E., Mao,
Y., Lathia, J.D.*, et al.* (2013). Glioblastoma stem cells generate vascular pericytes to support
vessel function and tumor growth. Cell *153*, 139-152.

Codega, P., Silva-Vargas, V., Paul, A., Maldonado-Soto, A.R., Deleo, A.M., Pastrana, E.,
and Doetsch, F. (2014). Prospective identification and purification of quiescent adult neural
stem cells from their in vivo niche. Neuron *82*, 545-559.

Conti, L., Pollard, S.M., Gorba, T., Reitano, E., Toselli, M., Biella, G., Sun, Y., Sanzone, S.,
Ying, Q.L., Cattaneo, E.*, et al.* (2005). Niche-independent symmetrical self-renewal of a
mammalian tissue stem cell. PLoS Biol *3*, e283.

Crespo, I., Vital, A.L., Gonzalez-Tablas, M., Patino Mdel, C., Otero, A., Lopes, M.C., de
Oliveira, C., Domingues, P., Orfao, A., and Tabernero, M.D. (2015). Molecular and Genomic
Alterations in Glioblastoma Multiforme. Am J Pathol *185*, 1820-1833.

Denysenko, T., Gennero, L., Roos, M.A., Melcarne, A., Juenemann, C., Faccani, G., Morra,
I., Cavallo, G., Reguzzi, S., Pescarmona, G.*, et al.* (2010). Glioblastoma cancer stem cells:
heterogeneity, microenvironment and related therapeutic strategies. Cell Biochem Funct *28*,
343-351.

Doetsch, F., Caille, I., Lim, D.A., Garcia-Verdugo, J.M., and Alvarez-Buylla, A. (1999).
Subventricular zone astrocytes are neural stem cells in the adult mammalian brain. Cell *97*,
703-716.

Engler, A., Rolando, C., Giachino, C., Saotome, I., Erni, A., Brien, C., Zhang, R., Zimber-
Strobl, U., Radtke, F., Artavanis-Tsakonas, S.*, et al.* (2018). Notch2 Signaling Maintains
NSC Quiescence in the Murine Ventricular-Subventricular Zone. Cell Rep *22*, 992-1002.

Galli, R., Binda, E., Orfanelli, U., Cipelletti, B., Gritti, A., De Vitis, S., Fiocco, R., Foroni,
C., Dimeco, F., and Vescovi, A. (2004). Isolation and characterization of tumorigenic, stem-
like neural precursors from human glioblastoma. Cancer Res *64*, 7011-7021.

Giachino, C., and Taylor, V. (2009). Lineage analysis of quiescent regenerative stem cells in
the adult brain by genetic labelling reveals spatially restricted neurogenic niches in the
olfactory bulb. Eur J Neurosci *30*, 9-24.

Kawaguchi, D., Furutachi, S., Kawai, H., Hozumi, K., and Gotoh, Y. (2013). Dll1 maintains
quiescence of adult neural stem cells and segregates asymmetrically during mitosis. Nat
Commun *4*, 1880.

Khalifa, J., Tensaouti, F., Lusque, A., Plas, B., Lotterie, J.A., Benouaich-Amiel, A., Uro-
Coste, E., Lubrano, V., and Cohen-Jonathan Moyal, E. (2017). Subventricular zones: new
key targets for glioblastoma treatment. Radiat Oncol *12*, 67.

Kim, D., Pertea, G., Trapnell, C., Pimentel, H., Kelley, R., and Salzberg, S.L. (2013).
TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and
gene fusions. Genome Biol *14*, R36.
Kramer, A., Green, J., Pollard, J., Jr., and Tugendreich, S. (2014). Causal analysis approaches
in Ingenuity Pathway Analysis. Bioinformatics *30*, 523-530.
Lee, J.H., Lee, J.E., Kahng, J.Y., Kim, S.H., Park, J.S., Yoon, S.J., Um, J.Y., Kim, W.K.,
Lee, J.K., Park, J*., et al.* (2018). Human glioblastoma arises from subventricular zone cells
with low-level driver mutations. Nature *560*, 243-247.
Lim, D.A., Tramontin, A.D., Trevejo, J.M., Herrera, D.G., Garcia-Verdugo, J.M., and
Alvarez-Buylla, A. (2000). Noggin antagonizes BMP signaling to create a niche for adult
neurogenesis. Neuron *28*, 713-726.
Livak, K.J., and Schmittgen, T.D. (2001). Analysis of relative gene expression data using
real-time quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods *25*, 402-408.
Llorens-Bobadilla, E., Zhao, S., Baser, A., Saiz-Castro, G., Zwadlo, K., and Martin-Villalba,
A. (2015). Single-Cell Transcriptomics Reveals a Population of Dormant Neural Stem Cells
that Become Activated upon Brain Injury. Cell Stem Cell *17*, 329-340.
Louis, D.N., Ohgaki, H., Wiestler, O.D., Cavenee, W.K., Burger, P.C., Jouvet, A.,
Scheithauer, B.W., and Kleihues, P. (2007). The 2007 WHO classification of tumours of the
central nervous system. Acta Neuropathol *114*, 97-109.
Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol *15*, 550.
Mali, P., Yang, L., Esvelt, K.M., Aach, J., Guell, M., DiCarlo, J.E., Norville, J.E., and
Church, G.M. (2013). RNA-guided human genome engineering via Cas9. Science *339*, 823-
826.
Marques-Torrejon, M.A., Gangoso, E., and Pollard, S.M. (2018). Modelling glioblastoma
tumour-host cell interactions using adult brain organotypic slice co-culture. Dis Model Mech
11.
Martynoga, B., Mateo, J.L., Zhou, B., Andersen, J., Achimastou, A., Urban, N., van den
Berg, D., Georgopoulou, D., Hadjur, S., Wittbrodt, J*., et al.* (2013). Epigenomic enhancer
annotation reveals a key role for NFIX in neural stem cell quiescence. Genes Dev *27*, 1769-
1786.
Mira, H., Andreu, Z., Suh, H., Lie, D.C., Jessberger, S., Consiglio, A., San Emeterio, J.,
Hortiguela, R., Marques-Torrejon, M.A., Nakashima, K*., et al.* (2010). Signaling through
BMPR-IA regulates quiescence and long-term activity of neural stem cells in the adult
hippocampus. Cell Stem Cell *7*, 78-89.
Mistry, A.M., Hale, A.T., Chambless, L.B., Weaver, K.D., Thompson, R.C., and Ihrie, R.A.
(2017). Influence of glioblastoma contact with the lateral ventricle on survival: a meta-
analysis. J Neurooncol *131*, 125-133.
Morizur, L., Chicheportiche, A., Gauthier, L.R., Daynac, M., Boussin, F.D., and Mouthon,
M.A. (2018). Distinct Molecular Signatures of Quiescent and Activated Adult Neural Stem
Cells Reveal Specific Interactions with Their Microenvironment. Stem Cell Reports *11*, 565-
577.
Paliouras, G.N., Hamilton, L.K., Aumont, A., Joppe, S.E., Barnabe-Heider, F., and
Fernandes, K.J. (2012). Mammalian target of rapamycin signaling is a key regulator of the
transit-amplifying progenitor pool in the adult and aging forebrain. J Neurosci *32*, 15012-
15026.
Pastrana, E., Cheng, L.C., and Doetsch, F. (2009). Simultaneous prospective purification of
adult subventricular zone neural stem cells and their progeny. Proc Natl Acad Sci U S A *106*,
6387-6392.

Piccirillo, S.G., Spiteri, I., Sottoriva, A., Touloumis, A., Ber, S., Price, S.J., Heywood, R.,
Francis, N.J., Howarth, K.D., Collins, V.P.*, et al.* (2015). Contributions to drug resistance in
glioblastoma derived from malignant cells in the sub-ependymal zone. Cancer Res *75*, 194-
202.
Pollard, S.M., Conti, L., Sun, Y., Goffredo, D., and Smith, A. (2006). Adherent neural stem
(NS) cells from fetal and adult forebrain. Cereb Cortex *16 Suppl 1*, i112-120.
Pollard, S.M., Yoshikawa, K., Clarke, I.D., Danovi, D., Stricker, S., Russell, R., Bayani, J.,
Head, R., Lee, M., Bernstein, M.*, et al.* (2009). Glioma stem cell lines expanded in adherent
culture have tumor-specific phenotypes and are suitable for chemical and genetic screens.
Cell Stem Cell *4*, 568-580.
Quinones-Hinojosa, A., Sanai, N., Soriano-Navarro, M., Gonzalez-Perez, O., Mirzadeh, Z.,
Gil-Perotin, S., Romero-Rodriguez, R., Berger, M.S., Garcia-Verdugo, J.M., and Alvarez-
Buylla, A. (2006). Cellular composition and cytoarchitecture of the adult human
subventricular zone: a niche of neural stem cells. J Comp Neurol *494*, 415-434.
Ran, F.A., Hsu, P.D., Wright, J., Agarwala, V., Scott, D.A., and Zhang, F. (2013). Genome
engineering using the CRISPR-Cas9 system. Nat Protoc *8*, 2281-2308.
Rispoli, R., Conti, C., Celli, P., Caroli, E., and Carletti, S. (2014). Neural stem cells and
glioblastoma. Neuroradiol J *27*, 169-174.
Sanai, N., Tramontin, A.D., Quinones-Hinojosa, A., Barbaro, N.M., Gupta, N., Kunwar, S.,
Lawton, M.T., McDermott, M.W., Parsa, A.T., Manuel-Garcia Verdugo, J.*, et al.* (2004).
Unique astrocyte ribbon in adult human brain contains neural stem cells but lacks chain
migration. Nature *427*, 740-744.
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B.*, et al.* (2012). Fiji: an open-source
platform for biological-image analysis. Nat Methods *9*, 676-682.
Singh, S.K., Clarke, I.D., Terasaki, M., Bonn, V.E., Hawkins, C., Squire, J., and Dirks, P.B.
(2003). Identification of a cancer stem cell in human brain tumors. Cancer Res *63*, 5821-
5828.
Singh, S.K., Hawkins, C., Clarke, I.D., Squire, J.A., Bayani, J., Hide, T., Henkelman, R.M.,
Cusimano, M.D., and Dirks, P.B. (2004). Identification of human brain tumour initiating
cells. Nature *432*, 396-401.
Stieber, D., Golebiewska, A., Evers, L., Lenkiewicz, E., Brons, N.H., Nicot, N., Oudin, A.,
Bougnaud, S., Hertel, F., Bjerkvig, R.*, et al.* (2014). Glioblastomas are composed of
genetically divergent clones with distinct tumourigenic potential and variable stem cell-
associated phenotypes. Acta Neuropathol *127*, 203-219.
Urban, N., and Guillemot, F. (2014). Neurogenesis in the embryonic and adult brain: same
regulators, different roles. Front Cell Neurosci *8*, 396.
Yousef, H., Morgenthaler, A., Schlesinger, C., Bugaj, L., Conboy, I.M., and Schaffer, D.V.
(2015). Age-Associated Increase in BMP Signaling Inhibits Hippocampal Neurogenesis.
Stem Cells *33*, 1577-1588.
Zhang, J., and Jiao, J. (2015). Molecular Biomarkers for Embryonic and Adult Neural Stem
Cell and Neurogenesis. Biomed Res Int *2015*, 727542.
Zhang, Q.B., Ji, X.Y., Huang, Q., Dong, J., Zhu, Y.D., and Lan, Q. (2006). Differentiation
profile of brain tumor stem cells: a comparative study with neural stem cells. Cell Res *16*,
909-915.
692

Figure 1

Figure 2

Figure 3

Figure 4

Supplementary Figure 1

Supplementary Figure 2

Supplementary Figure 3

Supplementary Figure 4