Title

Using oxygen and hydrogen stable isotopes to track the migratory movement of Sharp-shinned Hawks (*Accipiter striatus*) along Western Flyways of North America

Short title

Isotopes track migrating Sharp-shinned Hawks (*Accipiter striatus*) in Western North America

Elizabeth A. Wommack[1,2,3,¶,*], Lisa C. Marrack[4,¶,#a], Stefania Mambelli[3,¶], Joshua M. Hull[2,5], Todd E. Dawson[3,4]

[1]University of Wyoming Museum of Vertebrates, Department of Zoology & Physiology, University of Wyoming, Laramie, Wyoming, United States of America

[2]Golden Gate Raptor Observatory, Sausalito, California, United States of America

[3]Department of Integrative Biology, University of California, Berkeley, California, United States of America

[4]Department of Environmental Science, Policy & Management, University of California, Berkeley, California, United States of America

[5]Department of Animal Science, University of California, Davis, California, United States of America

[#a]Current Address: Tropical Conservation Biology and Environmental Science (**TCBES**) Graduate Program, University of Hawaii, Hilo, Hawaii, United States of America

*Corresponding author
E-mail: ewommack@uwyo.edu (EAW)

[¶]These authors contributed equally to this work.

Abstract

The large-scale patterns of movement for the Sharp-shinned Hawk (*Accipiter striatus*), a small forest hawk found throughout western North America, are largely unknown. However, based on field observations we set out to test the hypothesis that juvenile migratory *A. striatus* caught along two distinct migration routes on opposite sides of the Sierra Nevada Mountains of North America (Pacific Coast and Intermountain Migratory Flyways) come from geographically different natal populations. We applied stable isotope analysis of hydrogen (H) and oxygen (O) of feathers, and large scale models of spatial isotopic variation (isoscapes) to formulate spatially explicit predictions of the origin of the migrant birds. Novel relationships were assessed between the measured hydrogen and oxygen isotope values of feathers from *A. striatus* museum specimens of known origin and the isoscape modeled hydrogen and oxygen isotope values of precipitation at those known locations. We used these relationships to predict the origin regions for birds migrating along the two flyways from the measured isotope values of migrant's feathers and the associated hydrogen and oxygen isotopic composition of precipitation where these feathers were formed. The birds from the two migration routes had overlap in their natal/breeding origins and did not differentiate into fully separate migratory populations, with birds from the Pacific Coast Migratory Flyway showing broader natal geographic origins then those from the Intermountain Flyway. The methodology based on oxygen isotopes had, in general, less predictive power than the one based on hydrogen. There was broad agreement between the two isotope approaches in the geographic assignment of the origins of birds migrating along the Pacific Coast Flyway, but not for those migrating along the Intermountain Migratory Flyway. These results are discussed in terms of their implications for conservation

66 efforts of *A. striatus* in western North America, and the use of combined hydrogen and oxygen

67 stable isotope analysis to track the movement of birds of prey on continental scales.

Introduction

Thousands of bird species migrate, traveling from breeding territories to wintering grounds and

back annually, sometimes over vast distances and geographical features [1]. Such seasonal

movements make the factors that stress populations of migratory birds difficult to track, as

individuals may be affected by changes at any point along their migratory route [2–4]. As a

result, understanding the migratory paths and connections between breeding and wintering sites

is critical for strategizing conservation and preservation actions for particular bird species, such

as raptors which are often secretive and breed over a wide area [5].

Migratory routes taken by different birds that follow similar pathways across continents

are identified as migratory "flyways", which are hypothesized to represent the shortest and least

costly course over wide geographic distances [6]. Differences in the choice of migratory flyway

between and within species may represent evolutionary divergences related to specific

adaptations [7–9]. For some species, individuals that stray from or cross between flyways

experience higher costs to migration, therefore evolutionary associations are likely to exist

between populations of birds and specific migratory flyways [10–12]. One common way to

monitor populations of birds of prey is by tracking migratory movement at watch sites and

banding stations along migratory flyways [5, 13, 14]. Long-term data sets from raptor migration

watch-sites can indicate population dynamics for specific species and populations, which can

provide indications of changes in population size [15–19]. However, it can be difficult to connect

migrating birds to other geographic stages in their life cycle (breeding and wintering) through

these data, and without these connections it may be problematic to understand what factors

underlie population size changes [20].

The Sharp-shinned Hawk (*Accipiter striatus*) is a small, forest hawk found throughout

North America [21]. During breeding, *A. striatus* prefers dense coniferous and deciduous forests,

making nesting sites difficult to identify and survey, and breeding birds problematic to monitor

[21–23]. Heavy persecution resulting from the shooting of thousands along migration routes each

year in the late 19[th] and early 20[th] centuries [22, 24], combined with the effects of

dichlorodiphenyltrichloroethane (DDT) used as a pesticide in the early 20[th] century, resulted in

A. striatus numbers decreasing steadily across the species' range [25]. However, over the past

several decades, data from population and migratory monitoring of the species have shown

inconsistent and contrary trends of both significant increases and decreases [26, 13]. Since North

America's smallest hawk is difficult to track on its breeding grounds, most population

monitoring occurs at watch and banding sites along migratory flyways. Connecting migratory

flyways and watch sites with specific breeding ranges will allow for a greater understanding of

wider populations trends for this secretive bird of prey.

Raptors in the western side of the North American continent are believed to travel along

three migratory flyways: the Pacific Coast, the Intermountain, and the Rocky Mountain Flyways

[12]. Band recovery data of *A. striatus* trapped at migration sites along each of the flyways have

shown demographic population differentiation along each route [12, 15, 27]. However, sample

sizes for band returns are often small and may therefore represent biased information [28].

Endogenous markers, such as genetic analysis and stable isotope analysis (SIA), are not able to

provide the same level of precision on locality as extrinsic markers (i.e. bands, satellite

telemetry), but are easier and less expensive to use with large sample sizes, and can provide a

larger return in data for wide geographic areas [29–35]. Integrating information from both levels

of analysis can provide the best account of current population trends, and allow researchers to

connect migratory flyway monitoring data with information on geographical breeding areas.

115   Over the past few decades, a number of studies have used SIA of naturally occurring

hydrogen (H) of feathers in order to estimate migratory patterns and ecological connectivity

among habitats for a broad range of species of birds [36]. In comparison, only recently with

advances in continuous flow pyrolysis techniques have reliable SIA of oxygen (O) of organic

materials been possible. Therefore, much less data exist on the additional information that can be

obtained by measuring both elements to trace animal movements from different tissue types,

such as feathers and hair [37–40].

122   Keratin, the main constituent of feathers, remains chemically inert following its synthesis,

and it incorporates hydrogen and oxygen from consumed water and dietary sources. The stable

hydrogen and oxygen isotopic compositions in the keratin of a feather (δ^2H_F and $\delta^{18}O_F$ values)

therefore become markers of the environmental conditions and will vary geographically due to

the spatial variation in the stable hydrogen and oxygen isotopic composition of meteoric

precipitation (δ^2H_P and $\delta^{18}O_P$ values) [41]. This variation is caused primarily by isotope effects

associated with evaporation and condensation processes [42–44] and correlates inversely with

latitude and elevation across the continents [45]. Additionally, because hydrogen and oxygen are

incorporated or "routed" into organic compounds like keratin along different metabolic

pathways, the δ^2H and $\delta^{18}O$ of feather keratin may not provide the same information even if

these elements had their origin from the same water source.

133   Linking the isotopic composition of animal tissue to geographic locations of origin is

based on important assumptions, such as the presence of known and constant isotope effects

associated with tissue synthesis, and an understanding of the relationship between the isotopic

compositions of the tissue and that of the environmental food web and water source signals [46,

39, 74]. However, work on some species of birds suggests a decoupling between hydrogen and

oxygen isotope compositions in food webs that might affect the usefulness of $\delta^{18}O_F$

measurements for assignment of bird origin [48, 49, 39].

Previous SIA studies have used δ^2H_F values to estimate the timing and pattern of

migration for *A. striatus* from western North America [50, 51]. Birds migrating along the

Intermountain Flyway were found to originate from the Northern Rocky Mountain Range (Idaho

(ID) and Montana (MT)) in the United States of America (USA) north through British Columbia

(BC, Canada) [51]. Specific origins were not determined for birds caught along the Rocky

Mountain Flyway, but it has been shown that *A. striatus* followed a chain migration pattern

where individuals from lower latitudes migrated earlier than those from higher latitudes [50].

However, no work has been done to look at the origins of *A. striatus* from the Pacific Coast

Flyway or has incorporated two isotopes to trace the origin of birds from neighboring flyways.

Migration can be a difficult and dangerous behavior for birds, and specific adaptations for

breeding populations that use different flyways would be anticipated to lead to population

differentiation [52]. If flyways represent unique migratory paths for different populations, then it

can be predicted that *A. striatus* migrating along neighboring flyways should originate from

separate breeding origins, as hinted at by their band results [12, 15]. However, no genetic

difference in mitochondrial control region sequences has been found between *A. striatus* caught

on migration along either flyway [53], suggesting that either birds that utilize the different routes

overlap and intermix in their breeding populations or that mitochondrial markers lack the

variability to resolve population genetic differentiation among flyways. Use of nuclear

microsatellite loci has demonstrated differentiation between the intermountain and Pacific

flyways in another wide-ranging raptor, the Red-tailed Hawk (*Buteo jamaicensis*) [54],

suggesting that similar differences among flyways may exist in *A. striatus*.

In this study, we used the variation in δ^2H_F and $\delta^{18}O_F$ values of *A. striatus* and large scale

models of spatial isotopic variation (isoscapes)for hydrogen and oxygen to: a) examine the origin

of *A. striatus* caught along the Pacific Coast Flyway in comparison to birds caught along the

Intermountain Flyway, and b) investigate the usefulness of oxygen isotopes to determine the

origin of raptorial birds of prey. First, we established relationships, separately for hydrogen and

oxygen, between δ^2H_F and $\delta^{18}O_F$ values of *A. striatus* museum specimens from known natal

locations, to predict the isotope values of precipitation at the origin regions of birds migrating

along the Pacific Coast and Intermountain Flyways from the measured isotope values of

migrant's feathers. Prediction of precipitation hydrogen and oxygen isotope composition and

assignment of migrating bird origin locations were accomplished using an internet-based

environmental water isoscape of Western North America (http://isomap.org) [55]. Finally,

predictions of sites of origin based on δ^2H_P and $\delta^{18}O_P$ values were also compared.

Assuming that the choice of migratory flyway is driven by evolutionary processes [11],

and that it would be costly for individuals to jump between migratory routes, we predicted that *A.*

striatus caught along the Pacific Coast Flyway would originate from further west then those

caught along the Intermountain Flyway, and show little overlap in their determined natal origins.

However, instead the SIA revealed that birds from the Pacific Coast Flyway origins extended

further east then expected, including into the Rocky Mountain Range and the western interior of

North America. We report that both hydrogen and oxygen isotopic analysis with feathers can be

used to determine the origin of migratory birds of prey, but that caution must be taken when

interpreting the outcome.

Materials and methods

Sample collection

Contour feathers of *A. striatus* from the ventral sternal feather track ($n = 23$ juveniles (Table 1,

Fig 1) and $n = 25$ adults (Table S1, S1 Fig) were sampled from museum specimens of known

collection locality. *A. striatus* follow a complex basic strategy of molt, and have a limited to

absent preformative molt which occurs for hatch-year and second-year birds between December

and April of their first year. Adult molt (the definitive prebasic molt) occurs primarily on

breeding grounds, starting during egg laying and incubation [21, 56]. As a result of this molt

strategy both juvenile and adult *A. striatus* contour feathers have a high probability of being

grown either in the location where they were hatched or where they were breeding. Contour

feathers in ventral sternal feather track were selected for the small size of the feather, the ease of

repeatability of selection of feathers from the same track for collection from individual birds, and

as contour feathers have been used successfully to track movement of *A. striatus* in previous

work [51]. Feather samples of *A. striatus* were obtained from the following museums in the

USA: California Academy of Sciences (CAS), San Francisco, CA; Charles R. Conner Museum,

Pullman (CRCM), WA; Museum of Vertebrate Zoology (MVZ), University of California,

Berkeley, CA; San Diego Natural History Museum (SDNHM), San Diego, CA; University of

Wyoming Museum of Vertebrates (UWYMV), Laramie, WY. In order to acquire samples that

were representative of feathers grown on breeding and natal geographic regions of interest,

specimens were constrained by date of collection (15th March – 30th August), and feather wear

[56, 21, 57]. Only feathers with little or no wear were from museum specimens were selected to

guarantee that they were grown in the present year and at the location of collection. Feathers

were also sampled from migrating juvenile *A. striatus* banded at the Marin Headlands, Marin

County, CA (Golden Gate Raptor Observatory (GGRO), Pacific Coast Flyway, $n = 20$), and the

Goshutes Mountains, Elko County, NV (HawkWatch International (Goshutes), Intermountain

Flyway, $n = 10$) between August and October 1999 (see [53] for sampling protocol). The

migratory birds used in this study were collected with samples previously used to analyze the

genetic population structure of *A. striatus*, and so are known to be from the western genetic

population [53]. However, the specific origin of migrating individuals was unknown. No live

birds were handled for the described study. Sampling of birds at banding stations was performed

in 1999, and feathers used in this study were from archived collections.

Fig 1. Map of sample locations for museum specimens of juvenile Sharp-shinned Hawk
(*Accipiter striatus*) feathers.
Museum specimens ($n = 23$) mapped in reference to the species known range in Western North
America (light gray), and suitable forest habitat (dark gray). Juvenile samples ($n = 23$) are shown
as triangles. Collection sites of migratory bird samples are indicated as GGRO (Golden Gate
Raptor Observatory) and GOSH (Goshutes Mountains). The dominant migratory flyways of
western North America are indicated by dashed lines and modified from Hoffman et al. (2002)
for illustrative purposes. State and country boundaries are modified from public domain GIS
files, US Census Bureau (2016) and Natural Earth (2020). Species range acquired with
permission from BirdLife International and NatureServe (2015), and data to create the GIS
biome layer acquired from Brown, Bennan, and Unmack (2007).

225
226 **Table 1. Hydrogen and oxygen stable isotope composition of feathers of museum juvenile**
227 **Sharp-shinned Hawk (*Accipiter striatus*) specimens (δ^2H_f and $\delta^{18}O_f$ values (‰)), and**
228 **IsoMAP isoscape modeled stable hydrogen and oxygen isotope composition of precipitation**
229 **(δ^2H_p and $\delta^{18}O_p$ values (‰)) of known natal origin.**

[a]Museum	[b]Specimen no.	Life stage	[c]State	Latitude	Longitude	$\delta^2 H_f$ (‰)	$\delta^{18}O_f$ (‰)	$\delta^2 H_p$ (‰)	$\delta^{18}O_p$ (‰)
MVZ	169006	Juvenile	AK	61.0561111	-149.79722	-80.3	12.88	-124.9	-13.23
MVZ	99707	Juvenile	BC	54.0167	-132.15	-39.3	12.09	-95.5	-12.07
MVZ	32217	Juvenile	CA	36.537017	-121.9262	-18.7	16.23	-36.9	-4.36
MVZ	144621	Juvenile	ID	48.4797	-116.8483	-46.7	16.99	-102.8	-12.79
MVZ	15622	Juvenile	BC	48.9895	-124.8	-70.5	8.76	-72.0	-8.95
MVZ	81827	Juvenile	BC	49.6833	-124.9333	-44.0	13.03	-71.9	-10.02
MVZ	99705	Juvenile	BC	54.0167	-132.15	-44.9	12.69	-95.5	-12.07
MVZ	99706	Juvenile	BC	54.0167	-132.15	-58.3	13.07	-95.5	-12.07
MVZ	30835	Juvenile	CA	39.6863371	-123.48519	-12.1	22.22	-51.2	-5.65
MVZ	9775	Juvenile	AK	57.03139	-132.8536	-65.1	11.60	-122.4	-14.60
CAS	58055	Juvenile	CA	39.14057	-120.2011	-60.5	15.76	-83.0	-5.98
CAS	87247	Juvenile	CA	41.78846	-124.1668	-34.9	13.12	-49.2	-5.55
CAS	98830	Juvenile	CA	38.49657	-122.9394	-26.0	16.72	-42.4	-4.76
CRCM	57-367	Juvenile	WA	48.66223	-117.9829	-75.4	13.04	-97.6	-12.18
CRCM	57-368	Juvenile	WA	48.66223	-117.9829	-81.1	13.26	-97.6	-12.18
CRCM	89-229	Juvenile	WA	46.79025	-117.2521	-87.0	10.05	-94.2	-9.72
CRCM	89-223	Juvenile	WA	46.73064	-117.1625	-67.4	11.41	-94.2	-10.01
UWYMV	1352	Juvenile	WY	44.27615	-110.4736	-103.7	8.62	-102.5	-11.99
UWYMV	2521	Juvenile	WY	43.83333	-110.7	-75.1	13.41	-100.0	-11.47
UWYMV	2547	Juvenile	CO	40.39162	-106.9051	-62.2	15.97	-82.7	-9.67
SDNHM	52830	Juvenile	WA	47.8674	-122.516	-35.6	10.90	-69.0	-8.40
SDNHM	54234	Juvenile	AZ	35.1957	-111.6326	-26.0	17.19	-52.0	-5.78
SDNHM	18938	Juvenile	CA	35.50708	-118.3434	-85.3	10.08	-64.8	-4.72

[a]Museums: CAS = California Academy of Sciences, San Francisco, CA, USA; CRCM = Charles R. Conner
Museum, Pullman, WA, USA; MVZ = Museum of Vertebrate Zoology, University of California, Berkeley, CA,
USA; SDNHM = San Diego Natural History Museum, San Diego, CA, USA; and UWYMV = University of
Wyoming Museum of Vertebrates: Laramie, WY, USA.
[b]Specimens: Full information on each specimen can be obtained by taking the specimen number and searching for it
in the online databases in http://vertnet.org/index.html and https://arctosdb.org/

 ^cState or Provinces: AK = Alaska, USA; AZ = Arizona, USA; BC = British Colombia, Canada; CA = California,
246 USA; CO = Colorado, USA; ID = Idaho, USA; NM = New Mexico, USA; NV = Nevada, USA; OR = Oregon,
247 USA; UT = Utah, USA; WA = Washington, USA; WY = Wyoming, USA.
248
249

Sample preparation and SIA

Feathers were cleaned by immersion in a chloroform:methanol 2:1 (v/v) solution for 24 hours,

and then again for 1 hour, to remove lipids and mites. After each wash the feathers were air dried

for 24 hours [58–59]. The barbs were removed from the rachis, minced, and 1.5-2.0 mg of

feather material was packed into 3.5 x 5 mm silver capsules. To avoid the possibility of

intrafeather variation found in distal versus proximal samples for some raptor feathers, barbs for

oxygen and hydrogen samples were removed from corresponding sides of each contour feather

[57]. Previous work on other species of raptors has found that breeding condition and age may

affect δ^2H_F values differently in juvenile versus adults [60–62]. Therefore, adult samples were

removed from analysis reflecting these concerns. Only samples from juvenile individuals were

analyzed for both $\delta^{18}O_F$ and δ^2H_F values (museum: $n = 23$, migratory: $n = 21$). However, as there

are no published results of the differences for stable oxygen isotope measurements between adult

and juvenile birds of prey, we present the $\delta^{18}O$ values for adult *A. striatus* samples of know

origin ($n = 25$) as Supporting Information in this paper (S1 Table, S1 Fig).

The stable isotope abundances are presented in δ notation as deviations from the

standard reference (V-SMOW) in parts per mil (‰) according to the following equation: $\delta X =$

($R_{sample}/R_{standard}$) -1) where X represents ^2H or ^{18}O, and R the ratio of the heavy and light isotope

(e.g., $^{18}O/^{16}O$) in the sample and in the standard, respectively.

The $\delta^{18}O$ measurements were performed at the Center for Stable Isotope

Biogeochemistry (University of California, Berkeley, CA, USA) using a PYRO Cube

(Elementar, Hanau, Germany) interfaced to a Thermo Delta V mass spectrometer (Thermo

Fisher Scientific Inc., Waltham, Massachusetts, USA). The δ^2H measurements were carried out

at the Cornell Isotope Laboratory (Cornell University, Ithaca, NY, USA) using a Temperature

Conversion Elemental Analyzer (TC/EA) interfaced to a Thermo Delta V mass spectrometer

(both from Thermo Fisher Scientific Inc., Waltham, Massachusetts, USA). Both measurements

were based on pyrolysis of the sample carried out in reactors kept at 1350 °C, with GC

temperature kept at 90°C for hydrogen analysis, and CO trap temperature kept at 40 °C and then

increased to 130 °C for release of CO for oxygen analysis.

Reference materials of kudu horn and caribou hoof keratins (KHS and CBS) were used

for normalization (δ^{18}O value = +20.3 ‰ and +3.8‰, respectively [63]; δ^2H value = -35.3 ‰

and -157.0 ‰, respectively [64]), and an in-house keratin material was used as quality control

material for both hydrogen and oxygen stable isotope measurements. The precision of the

analysis was ± 2.8‰ and 0.30‰ for hydrogen and oxygen, respectively. To correct the measured

hydrogen isotope ratios for the contribution of exchangeable hydrogen atoms to the total number

of hydrogen atoms in feathers (~15%) [47, 63, 65], samples were allowed to equilibrate with the

laboratory ambient atmosphere along with KHS and CBS standards for a minimum of 72 hours

before isotopic analysis [65].

A linear regression approach was used to examine the relationship between measured

δ^2H$_F$ and δ^{18}O$_F$ values from *A. striatus* feathers for museum samples of known natal/breeding

origin as well as in birds sampled at the two migratory banding sites.

291   H and O precipitation Isoscapes for *Accipiter striatus* museum

292   samples of known natal/breeding origin

IsoMAP, a web resource for isoscapes modeling, (http://www.waterisotopes.org)[66], was used

to predict the hydrogen and oxygen isotopic compositions of precipitation (δ^2H_P and $\delta^{18}O_P$

values) at the locations where museum specimens were collected. Isoscapes are gridded surfaces

representing spatially explicit isotope distributions across a landscape [46, 67]. IsoMAP software

creates water isoscapes by using precipitation isotope data from the Global Network for Isotopes

in Precipitation (GNIP) database administered by the International Atomic Energy Association

and World Meteorological Organization [68]. Within IsoMAP, numerous parameters can be

selected to create an isoscape model that is the best fit for a specific study area and time period

since the geographical variation of δ^2H_P and $\delta^{18}O_P$ values depend on a range of geographic and

meteorological effects including latitude, season, elevation, and regional air mass circulation.

Independently for each isotope, we modeled the geographic distribution of δ^2H_P and $\delta^{18}O_P$ values

across North America using precipitation data collected from 1960–1999 during the months from

March to September to represent the plant growing season [68, 69]. Comparisons of different

precipitation isotope models available in IsoMAP revealed that the most robust isoscapes that

represented δ^2H_P and $\delta^{18}O_P$ values across North America were kriging interpolation models. The

δ^2H_P isoscape produced was based on 117 stations, had resolution of 9x9 km, a correlation

parameter of 0.93, and included the variables elevation (ETOPO, $P < 0.001$), latitude ($P < 0.001$)

and longitude ($P = 0.06$) (available as IsoMAP job key 50333, S2 Fig [70]). The most robust

$\delta^{18}O_P$ isoscape was based on 120 stations, had resolution of 9x9 km, a correlation parameter of

0.92, and included the variables elevation (ETOPO, $P < 0.001$), latitude ($P < 0.001$) and

longitude ($P = 0.05$) (available as IsoMAP job key 63026, S2 Fig [71]).

The hydrogen and oxygen isoscape precipitation models were further modified using the

spatial software ArcGIS (ESRI 2010). In particular, we limited the spatial isotopic predictions to

the known habitat range of *A. striatus* using a geographic information system (GIS) layer of

range delineation provided by BirdLife International and NatureServe [72]. In addition, because

this species is found to nest specifically in forests [21], we applied a GIS biome layer to exclude

non-breeding habitat, such as tundra, open water, desert, and grassland from potential sites of

origin [73].

Predicting the origin of migrating *Accipiter striatus*

Because of isotope discrimination during feather formation and other effects, the isotope values

of feathers may not directly reflect the stable hydrogen and oxygen isotopic composition of the

environmental water at the site where they are formed [74–75]. We estimated the magnitude of

such discrimination factors for *A. striatus* feathers by calculating the linear relationship between

the isotopic compositions of feathers from museum specimens of known origin (δ^2H_F and $\delta^{18}O_F$)

and the corresponding isoscape predicted precipitation values (δ^2H_P and $\delta^{18}O_P$) at the location

where the museum specimens were collected [76–78]. Metadata associated with each specimen

included the uncertainty for the collection location in meters. When multiple isoscape derived

δ^2H_P and $\delta^{18}O_P$ grid values were available within the radius of uncertainty around a collection

location, we used the average isotope values within this radius.

The parameters of the linear regression equations derived from the museum specimens

data were used to predict the isotopic composition of the precipitation at the site of origin of

birds sampled at each of the two migratory banding sites (GGRO and Goshutes). The feather

isotopic compositions of the migrating birds represented the variable *x* while *y* represented an

estimate of the associated δ^2H_P or $\delta^{18}O_P$ values. Resulting δ^2H_P and $\delta^{18}O_P$ values thus

represented the water source isotope compositions expected for the localities where the feathers

of migrating birds were formed. The linear regressions were used as a transfer function to

convert feather isotope values for migratory birds to precipitation isotopes values at the sites of

origin [79]. Many researchers examining the relationship between feather and precipitation

isotopes plot precipitation on the x axis and feather isotopes on the y axis as the dependent

variable, creating a feather isoscape [39, 40, 48, 51]. However, we did not feel justified using this

format of regression equation for the transformation lines in our analysis with the sample size of

museum specimens ($n = 23$) across the entire area of interest. Instead we predicted precipitation

from measured feather values and created a precipitation isoscape to predict probability of origin.

For comparison with previous analyses from the literature we have created the regression

equations in the supporting information (S4 Fig).

To determine if the origin for birds migrating through the GGRO and Goshutes sites

differed, we examined the data in several ways. First, we calculated the frequency distributions

of the predicted precipitation isotope values for the two migratory banding sites. Second, δ^2H_F

and $\delta^{18}O_F$ values for each migrant group (GGRO and Goshutes) were compared statistically

using Welches unequal variance T-test which tests the hypothesis that two populations have the

same means. Additionally, δ^2H_F and $\delta^{18}O_F$ values for each migrant group (GGRO and Goshutes)

were examined with a Kolmogorov-Smirnov distribution test which tests whether the two

populations have the same distribution.

Finally, we utilized the IsoMAP geographic assignment function to produce maps

representing the likelihood of origin for the migrant birds sampled at GGRO and Goshutes. The

assignment function in IsoMAP uses a semi-parametric Bayesian framework to model

probability density surfaces that can be used to determine geographic areas where organic

material, such as bird feathers, were developed [79, 80]. The assignment function requires an

observed sample isotopic composition, a standard deviation associated with the environment to

sample transfer, and an isoscape model in IsoMAP that the sample is compared to. For each of

the birds sampled at the GGRO and Goshutes, δ^2H_P and $\delta^{18}O_P$ values were estimated using the

feather to precipitation linear regressions. The residual standard error (RSE) of each linear

regression was used as the estimate of error (RSE for δ^2H of 18.3‰, and RSE for $\delta^{18}O$ of 3.0).

The precipitation isoscape models described above (IsoMAP job key 50333 and 63026, S2 Fig)

[70, 71] were utilized in the assignment function. Additional uncertainty associated with the

precipitation isoscape is automatically included in the assignment algorithm.

Rather than creating average probability surfaces for each migratory group though the

bulk sample function, probability surfaces were generated for each individual bird [80]. The

individual probability density maps were then averaged for GGRO and Goshutes groups. Using

ArcGIS, individual rasters within each group were summed together and then normalized by the

sum of all cell values in the final density surface. This process resulted in geographic

representations of the likely origin of *A. striatus* migrating along the Pacific Coast and the

Intermountain Flyways. Areas outside the species range and nesting habitat type were not

included as likely origin areas of the migratory birds. To examine the accuracy of the linear

transfer functions and the assignment of origin models, we also generated probability density

surfaces for 10 museum samples of known origin (S5 Fig, Table S2). All statistics were

performed in R (R version 2.14.0) [81].

Results

The stable hydrogen and oxygen isotopic composition of feathers of juvenile *A. striatus*

museum specimens varied from -103.7 to -12.1 ‰ for δ^2H_F and from 8.62 to 22.22 ‰ for $\delta^{18}O_F$

respectively ($n = 23$) (Table 1). A significant and positive relationship was found between the

δ^2H_F and $\delta^{18}O_F$ values from the juvenile feathers ($R^2 = 0.48$, $P < 0.001$, y = 5.34(x) – 128.95)

(Fig 2).

Fig 2. Relationship between the stable isotope composition of museum specimens of Sharp-
shinned Hawks (*Accipiter striatus*). Stable hydrogen (δ^2H_F ‰) and oxygen ($\delta^{18}O_F$ ‰) isotope
values for juvenile museum feather specimens ($n = 23$) of known natal origin.
391

We found positive relationships between the feather isotope values and the isoscape

modeled precipitation isotope values for the *A. striatus* museum specimen of known origin. The

linear regression between δ^2H_F and δ^2H_P values was statistically significant ($n = 23$, $R^2 = 0.46$, P

< 0.001, y = 0.68x – 43.98, RSE = 18.3‰) (Fig 3a), while the linear regressions between $\delta^{18}O_F$

and $\delta^{18}O_P$ values based on feathers of juvenile birds (Fig 3b) were not statistically significant (n

$= 23$, $R^2 = 0.14$, $P = 0.07$, y = 0.39x – 14.67, RSE = 3.0‰).

Fig 3. Relationship between the stable isotopic compositions of museum feathers and the
isoscape model of precipitation. Stable hydrogen (δ^2H_F ‰) and oxygen ($\delta^{18}O_F$ ‰) isotopic
composition of feathers for museum Sharp-shinned Hawk (*Accipiter striatus*) specimens of
known natal origin and the isoscape modeled isotopic compositions of precipitation (δ^2H_P and
$\delta^{18}O_P$ ‰) at the collection locations: (a) δ^2H_F values of birds ($n = 23$) versus δ^2H_P values, (b)
$\delta^{18}O_F$ values of birds ($n = 23$) versus $\delta^{18}O_P$ values.
404

Within the set of migratory birds for which both hydrogen and oxygen isotopic

compositions were measured, there was a positive relationship between δ^2H_F and $\delta^{18}O_F$ values at

both the GGRO and Goshutes banding sites (Fig 4). The linear relationship between δ^2H_F and

$\delta^{18}O_F$ values for the birds captured at the GGRO was very similar to that found for the museum

bird feathers ($n = 14$, $R^2 = 0.42$, $P = 0.01$, y = 6.36x – 146.45, Fig 4a). However, the relationship

between δ^2H_F and $\delta^{18}O_F$ values for the birds migrating through the Goshutes banding site was

not statistically significant ($n = 7$, $R^2 = 0.18$, $P = 0.34$, y = 1.02x – 122.29, Fig 4b).

Fig 4. Relationships between the stable isotope compositions of feathers from birds trapped
along the migratory flyways. Stable oxygen ($\delta^{18}O_F$ ‰) and hydrogen (δ^2H_F ‰) isotope
composition of feathers for juvenile migratory Sharp-shinned Hawk (*Accipiter striatus*)
specimens collected (a) along the Pacific Coast Flyway at the migratory banding site the Golden
Gate Raptor Observatory (GGRO, $n = 14$) and (b) along the Intermountain Flyway at the
migratory banding site Goshute Mountains HawkWatch (Goshutes, $n = 7$).
418
In migratory bird feathers, the range of δ^2H_F values was -105.69 to -4.36 ‰ from the

GGRO ($n = 15$) and -105.40 to -86.60 ‰ at the Goshutes ($n = 7$) (Figure 5a and c, Table 2). The

variation in $\delta^{18}O_F$ values was 8.16 to 21.19 ‰ for the GGRO ($n = 19$) and 14.11 to 22.87 ‰ for

the Goshutes ($n = 10$) (Figure 5b and d, Table 2). The mean δ^2H_F values (-58.6 $\pm$ 32.7 ‰

GGRO, -95.7 $\pm$ 7 ‰ Goshutes) were statistically different between the GGRO and Goshutes

sites (Welch's two-sample t-test, t = 4.13, df = 16.5, $P< 0.001$). There was a significant

difference in the distribution of the δ^2H_F values for the two migratory flyways (Two-sample

Kolmogorov-Smirnov test, D = 0.73, $P = 0.005$). Although a subsample of birds captured at the

GGRO site showed $\delta^{18}O_F$ depleted values compared to the Goshutes birds, the mean $\delta^{18}O_F$ values

(14.78 $\pm$ 3.5 GGRO, 16.44 $\pm$ 2.7 Goshutes) did not differ statistically between sites (Welch's

two-sample t-test, t = -1.4, df = 23.2, $P = 0.16$). Also, we did not detect any significant

difference in the distribution of the $\delta^{18}O_F$ values for the migratory flyways (Two-sample

Kolmogorov-Smirnov test, D = 0.42, $P = 0.16$).

Fig 5. Frequency distribution of the isoscape modeled isotope compositions of feathers of
migratory Sharp-shinned Hawks. Predicted stable hydrogen (δ^2H_F ‰) and oxygen ($\delta^{18}O_F$ ‰)
isotopic compositions of precipitation at the natal origin for migratory Sharp-shinned Hawk
(*Accipiter striatus*) specimens collected (a and b) along the Pacific Coast Flyway at the Golden
Gate Raptor Observatory (GGRO, $n = 17$), and (c and d) along the Intermountain Flyway at the
Goshute Mountains HawkWatch (Goshutes, $n = 10$). Values on the y-axis represent counts of
individual specimens.
439

440 **Table 2. Stable hydrogen and oxygen isotope composition of feathers of migrating Sharp-**
441 **shinned Hawks (*Accipiter striatus*) (δ^2H_f, and $\delta^{18}O_f$ values (‰)) captured at the Golden Gate**
442 **Raptor Observatory (GGRO), and at the Goshute Mountains HawkWatch (Goshutes,), and**

443 **of predicted stable hydrogen and oxygen isotope composition (δ^2H_p and**
444 **$\delta^{18}O_p$ values (‰)) at the migrant's unknown natal origin.**
445

Migratory Banding site	[a]Sample Band number	[b]δ^2H_f (‰)	[b]$\delta^{18}O_f$ (‰)	[b, c]δ^2H_p (‰)	[b, c]$\delta^{18}O_p$ (‰)
GGRO	1363-74740	-70.20	14.99	-91.74	-8.90
	1353-16673	-68.60	16.16	-90.60	-8.45
	1162-13236	-18.50	20.28	-56.58	-6.86
	733-23478	-53.50	12.76	-80.37	-9.76
	1363-74746	-40.70	13.58	-71.67	-9.44
	2003-95304	-77.50	14.65	-96.66	-9.03
	0733-64092	-105.69	12.47	-115.69	-9.87
	1423-50079	-71.40	11.93	-92.54	-10.08
	1433-84282	-97.44	8.66	-110.24	-11.33
	1423-50062	-43.30	17.57	-73.44	-7.90
	2003-95657	-34.73	17.49	-67.60	-7.94
	1433-84273	-9.04	13.42	-50.13	-9.50
	2003-95656	-96.80	8.16	-109.80	-11.53
	2003-95572	-4.36	15.67	-46.94	-8.64
	2003-95363	-87.90		-103.75	
	1152-29272		16.11		-8.47
	2003-95575		21.19		-6.51
	1363-74737		18.86		-7.41
	2003-95653		15.13		-8.85
	1152-29376		11.66		-10.18
Goshutes	1523-88735	-102.50	17.04	-113.70	-8.11
	1162-71742	-95.30	14.70	-108.75	-9.01
	1523-88624	-105.40	15.18	-115.65	-8.83
	1523-88749	-86.6	15.99	-102.87	-8.51
	1162-71901	-87.50	22.87	-103.51	-5.86
	1523-88752	-95.00	17.19	-108.59	-8.05
	1523-88734	-93.90	14.42	-107.84	-9.12
	1162-71551		14.11		-9.24
	1523-88628		18.36		-7.60
	1162-71547		14.58		-9.06

[a] Band numbers represent individual numbers issued by the Bird Banding Laboratory
(http://www.pwrc.usgs.gov/BBL/bblretrv/)
[b] Missing values represent feathers where there was not enough sample from one specimen for both hydrogen and
oxygen analysis.
[c] The δ^2H_p values were predicted using the linear regression equation $\delta^2H_p = 0.68*\delta^2H_f - 43.98$ (RSE =18.3‰).
The $\delta^{18}O_p$ values were predicted using the linear regression equation $\delta^{18}O_p = 0.39*\delta^{18}O_f - 14.67$ (RSE =3.0‰).
452

The predicted δ^2H_P values for migratory birds from the GGRO banding site varied

between -46.94 and -115.69 ‰ (Table 2). The average of these probability density surfaces

shows that the origin of the individuals captured along the Pacific Coast Flyway are most likely

from eastern CA, Oregon (OR), and Washington (WA) as well as some forested areas within

NV, Utah (UT), Colorado (CO), Montana (MT), Idaho (ID), Wyoming (WY), British Columbia

(Canada), and southern Alaska (AK) (Fig 6a). For the birds that migrated through the Goshutes

banding site, the predicted δ^2H_P values were less varied than at the GGRO, ranging between -

102.87 and -115.65‰ (Table 2). The average probability density surface suggests that

Intermountain Flyway captured birds originated from a smaller area focused primarily in eastern

WA, WY, MT, ID, British Columbia (Canada), and in southern AK (Fig 6c). The δ^2H_P

probability surfaces generated for museum samples of known origin showed a high to medium

probability of prediction for the correct localities for almost all the specimens, except for a

specimen from the western slope of the Rocky Mountains in ID (S5 Fig).

Fig 6. Probability density maps of the origin of migrating juvenile Sharp-shinned Hawks
(***Accipiter striatus***). Maps are based on predicted δ^2H_P values (‰) (left panels) and predicted
$\delta^{18}O_P$ values (‰) (right panels) for birds captured (a and b) along the Pacific Coast Flyway at the
Golden Gate Raptor Observatory (GGRO) and (c and d) along the Intermountain Flyway at the
Goshute Mountains HawkWatch (Goshutes). Each map represents the mean of probability
density surfaces created for individual birds sampled at a location and by isotope group. State
and country boundaries are from public domain GIS files US Census Bureau (2016) and Natural
Earth (2020). Species range acquired with permission from BirdLife International and
NatureServe (2015), and data to create the GIS biome layer acquired from Brown, Bennan, and
Unmack (2007).
476
477   Isoscape predicted $\delta^{18}O_P$ values at the natal/breeding sites of the birds captured at the

GGRO site ranged between -6.51 and -11.53‰ (Table 2). The average probability density

surface for birds captured along the Pacific Coast Flyway showed that the natal origins were

most likely from eastern CA, OR, NV, UT, CO, MT, WA, coastal British Columbia (Canada)

and southern AK (Fig 6b). These predicted origins, in general, confirmed the data found from the

δ^2H analysis. Predicted $\delta^{18}O_P$ values at the origin sites of the birds captured at the Goshutes

ranged from -5.86 to -9.24‰ (Table 2). The average probability density surface based on oxygen

isotope composition located the natal origins of birds from the Intermountain Flyway in more

coastal and southern areas then those from the δ^2H analysis, including CA, NV, UT, CO, OR,

WA, British Columbia (Canada), and small portions of AK (Fig 6d). Similar to hydrogen the

$\delta^{18}O_P$ probability density surfaces generated for museum samples of known origin showed high

to medium to medium probability of prediction for most specimens, but showed low probability

of origin for two specimens collected along the western slope of the Rocky Mountains in ID and

WA, and a specimen collected from the southern coast of AK (S5 Fig).

Discussion

Results from the combined SIA of hydrogen and oxygen of *A. striatus* feathers showed

that some raptors migrating along the Pacific Coast Flyway have origins that overlap with those

of raptors migrating along the Intermountain Flyway. Our prediction that juvenile birds that

traveled along each migratory route would come from different and non-overlapping

breeding/natal origins was therefore not supported. Instead it appears that migratory *A. striatus*

juveniles, specifically those that travel along the Pacific Coast Flyway may come from both west

of the Sierra Nevada mountain range and from the northern Rocky Mountain Range and western

interior regions of North America (Fig 6).

This outcome should be taken into consideration for conservation of *A. striatus* in

western North America, as individual populations may not show specific adaptations or fidelity

for movement along a single migratory flyway [7, 10, 11]. Instead, adjustments to their migration

strategy may depend on a multitude of ecological factors, such as minimization of energy cost

and mortality risk [82]. Juvenile raptors have also been found to have a greater degree of

variation in their migratory movement then adults [83], perhaps leading juvenile *A. striatus* to

wander between flyways during their first migration. As a result, movement across multiple

flyways should be heeded when examining data from migratory watch and banding sites, and

when using this data to estimate population size and fluctuations.

In agreement with previous studies, we found that both stable hydrogen and oxygen

isotopic composition of feathers can be used to predict the origin of birds across broad spatial

scales [39, 40]. However, the methodology based on oxygen had, in general, less predictive

power than the one based on hydrogen. Within the overall findings of natal origins for the birds

migrating along western migratory flyways in North America, some differences were found in

the predicted natal sites depending on which isotope and migratory route were examined.

517 Assignment of origin

We found an overlap in natal origin of *A. striatus* for the Flyways based on hydrogen and oxygen

isoscapes (Fig 6). Birds migrating along the Pacific Coast Flyway originated from a larger area

that covered both the coastal and central areas of the species' western range while *A. striatus*

migrating along the Intermountain Flyway had a more limited geographical origin.

Results from both hydrogen and oxygen isotopes point to similar predicted natal areas for

A. striatus migrating along the Pacific Coast Flyway (Fig 6a and b) in forest sites in CA, NV,

OR, WA, ID, and British Columbia (Canada). The origins of the migrants from the

Intermountain Flyway predicted from the oxygen isoscape also included forests in the Sierra

Nevada and Cascade mountains, spanning through CA, OR, and WA up into British Columbia

(Fig 6d). This expands the natal territory of the migratory individuals from the Intermountain

Flyway westward from the hypothesized origins, placing them closer to the west coast of North

America, and creates substantial overlap with the natal origins of those from the Pacific Coast

Flyway. In contrast, the origin sites for the Intermountain Flyway indicated using the hydrogen

isoscape lie along our predicted areas of natal origin, and agree with previously published

hydrogen data from Lott and Smith [51], which identified forests in the states of ID, MT, and

British Columbia as origins for *A. striatus* migrating through this flyway (Fig 6c). As a result, the

hydrogen isoscape revealed overlap in the origins of the two flyways only in the central areas of

the species western range. Modeled high probability of origin for *A. striatus* from δ^2H data was

also found for regions associated with the central migration flyway, including the central

Canadian providences of Alberta and Saskatchewan for both Flyways; but it is unclear if these

values represent a true signal of origin or the lack of differentiation of δ^2H$_P$ values found in the

central northern plains of North America [84]. The origin sites predicted using the oxygen

isoscape data do not show the same high probability for either Flyway. An examination of the

predicted origin of known museum samples did show low probability of origin for some

specimens collected in ID and eastern WA for oxygen, suggesting that there may be bias in the

assignment of origin from δ^{18}O data for that area (S5 Fig). The weak relationship between the

migrant's δ^2H$_F$ and δ^{18}O$_F$ values may account for these discrepancy (Fig 4). However, weak

correlations between δ^2H$_F$ and δ^{18}O$_F$ have not hindered previous studies that looked at the origin

of birds in Europe and Asia [40], and the assignment of origin tests showed high probability for

museum specimens with both hydrogen and oxygen in other areas of the species western range.

The successful use of SIA of hydrogen and oxygen to assign sites of origin for migratory

animals depends on understanding the relationship between isotopic composition in tissues and

the isotopic composition of waters sources within the landscape where these tissues were formed.

Isotopic composition of animal tissues are offset from environmental isotope values due to a

variety of discrimination factors that differ for each element [85]. These discrimination processes

vary among specific isotopes. In this study, the relationship between the stable hydrogen and

oxygen isotope composition of feathers differed depending on sampling locality. This

relationship was stronger for both museum specimens of known origin ($R^2 = 0.48$) and birds

migrating through the Pacific Coast Migratory Flyways ($R^2 = 0.42$) (Fig 2 and Fig 4a), than for

birds traveling along the Intermountain Flyway ($R^2 = 0.18$) (Fig 4b). Research on other species

of vertebrates that compared $\delta^{18}O$ and δ^2H values in feathers, claws, or hair have also found

varied correlation patterns between the two isotopic compositions. Significant correlations

between $\delta^{18}O$ and δ^2H values were found for insectivorous passerines ($R^2 = 0.34$) [39] and

falcons ($R^2 = 0.64$ and $R^2 = 0.48$)[86, 48], as well as for herbivorous mammals ($R^2 = 0.84$, and

0.57) [37]. However, no significant correlations were found between tissue oxygen and hydrogen

isotope compositions for other vertebrate species, including Pumas (*Puma concolor*) [37] and

European Cranes (*Grus grus*) [40]. Our study is the first to report that birds of the same species

sampled along different migrating routes can show different relationships between δ^2H_F and

$\delta^{18}O_F$ values.

The small sample size of birds captured along the Intermountain Flyways may also

account for the poor relationship between the stable hydrogen and oxygen isotope composition in

feathers. These samples were restricted to a single year of migratory data to coincide with

sampling strategy of published genetic data and ensure that all the birds sampled belonged to the

previously characterized *A. striatus* western population [53]. It may be possible that the

individuals analyzed here represented a divergence from the average value for the site that could

have been detected with a larger sample size [48]. However, the predicted origins from our

assignments based on hydrogen isotopes align very closely with previously published work for

A. striatus from the same flyway also based on hydrogen isotopes [51]. This suggests that despite

a small sample size, our results agree with previous findings of origin of birds from the

Intermountain Flyway. In general, increasing the migrant sample sizes from both flyways may

improve the precision in assigning birds to geographic origin. In addition, obtaining similar

results from multiple isotopes over multiple years would provide greater confidence in origin

maps for migratory species and are worth pursuing in future studies.

In summary, birds caught along the Pacific Coast Flyway have origins that overlap with

those birds caught along the Intermountain Flyway, consistent with the absence of population

genetic structure in mitochondrial sequence data among juveniles sampled on these flyways.

However, overlapping regions of origin for migrating juveniles does not preclude the possibility

of fine-scale population structure among regions, as has been seen in other raptors in western

North America [54, 87]. Overall birds that originate from the Rocky Mountain Range of North

America appear to choose to travel through either migration route, but discrepancies in the

predicted origins based on hydrogen and oxygen isotopes encourage caution, and further studies

in how *A. striatus* migrate in western North America.

Feather isotope composition and life history factors

Our understanding is still poor about why the strong correlation between $\delta^{18}O$ and $\delta^{2}H$

values in meteoric (source) water [44] breaks down in the tissues of different groups of

vertebrates. Isotopic variability has been observed in $\delta^{2}H_F$ values for wild American Kestrels

(*Falco sparverius*) at a local scale [60], and in $\delta^{2}H_F$ and $\delta^{18}O_F$ values in laboratory-controlled

groups of House Sparrows (*Passer domesticus*) [77] and Japanese Quail (*Cortunix japonica*)

[88]. This variability is thought to be due to differences in diet and water requirements, seasonal

timing of water use, metabolism, and evaporative cooling effects, among other factors [88, 89].

A variety of factors may have contributed to the different predictive ability of $\delta^{18}O_F$ and

δ^2H_F values in *A. striatus*. Carnivorous animals, including birds, are known to show more

positive δ^2H bone collagen values compared to δ^2H source water values [90], and hydrogen

isotopic composition in organic tissue seems to be influenced more by the diet consumed than

the water used, compared to the oxygen isotopic composition in the same tissue [78, 88, 89, 91].

Oxygen isotopic composition may also be affected by atmospheric and dissolved oxygen in body

water, in addition to diet and environmental water [89]. In addition, consuming prey from

different trophic levels (herbivores vs. insectivores) might affect the hydrogen and oxygen

isotopic compositions differently for a raptor such as *A. striatus*. Elevation may also affect the

δ^2H_P values [92], and changes in elevation performed during seasonal movement may result in

different isotopic compositions then expected [93].Climatic factors may also play a role in the

development of different isotopic compositions for raptorial birds from different habitats on the

continent. Climate and aridity can have profound effects on the degree of variation

(fractionation) of water $\delta^{18}O$ values measured in animal body water [86, 94, 95], as well as the

proportion of drinking versus metabolically produced water in the body-water pool [88].

Moreover, previous work analyzing isotopic information for a variety of raptor species across

North America found δ^2H_F enriched values for birds from the northern ranges of the Rocky

Mountains in ID and MT compared to δ^2H_P values at the same locations [51]. More work is

needed examining the effects of diet and climate on isotopic compositions for birds from

different trophic levels to determine if prey, habitat, or both, may play a role in variation in δ^2H

and $\delta^{18}O$ values.

The age of the animal has also been found to have a significant effect on the δ^2H_F value

for birds, with adults showing more positive isotopic compositions relative to younger animals

[61, 96–97]. Previous studies on breeding populations of known origin suggest a link between

high variability in adult δ^2H_F values, and breeding behavior and physiological effort during the

breeding season [60, 62]. Little work has been done using oxygen variation to determine origin

of birds from different age classes. While we did not assess the origin of adult *A. striatus*

migratory samples, our results for museum samples of known origin are similar between $\delta^{18}O_F$

and predicted $\delta^{18}O_P$ for adult and juvenile *A. striatus* (S3 Fig). Perhaps $\delta^{18}O_F$ values of adult

predatory birds may not be affected by the enrichment seen in tissue δ^2H, possibly because $\delta^{18}O$

values are more strongly influenced by environmental water then diet [89], and are not as

affected by evaporative, metabolic, and respiratory water losses [62].

Research focused on the relationship between the known origin of a bird, its

physiological condition, aspects of its behavior, and its diet and drinking regime will no doubt

improve our understanding of what "sets" the oxygen and hydrogen isotope composition in

feathers. In addition, previous work has identified that the determination of origin for species of

various trophic levels is improved when species specific calibration curves for hydrogen are

included [48, 98]. Ideally, species specific fractionation factors for hydrogen and oxygen

isotopes should be considered along with isotopic variation observed in birds due to differences

in age and trophic level. Further research that examines how these variables affect the

fractionation factors for hydrogen and oxygen isotope analyses will be essential before we can

use hydrogen or oxygen as a reliable tool for examining the origin of all birds. The results from

this study on *A. striatus* are encouraging.

Uncertainties in isoscape modeling methodology

IsoMAP allows users to create and test large scale models of spatial isotopic variation for

a specific area of interest. IsoMAP also allows creation of probability density maps to show

uncertainty due to sample variability. It is known that predictions based on isoscape modeling

methodology might be affected by several factors. The isoscapes created for this study in

IsoMAP.com are based on precipitation isotope ratio values collected through a global network

of stations [68, 70, 71]. Although the global sampling of precipitation isotope ratios used for the

isoscape predictions is spatially and temporally uneven, the coverage in North America is more

thorough then in other areas [66]. Therefore, the trends of δ^2H_P and $\delta^{18}O_P$ values can be

relatively robust on large scales, but they may not capture the variability present at more limited

spatial or temporal scales [67]. This can make attribution challenging or less precise.

Other sources of uncertainty in precipitation isotope values modeled using isoscapes are

due to large grid sizes, the integration of isotope data over multiple years, and the interpolation

model error. Collection location uncertainty, especially for museum specimens that may have a

location description that is not geographically specific, should also be taken into account. Spatial

and temporal resolution could be improved by the addition of quality precipitation isotope data to

the existing network through platforms like IsoMAP [67].

661   **Conclusion**

Our assignment evaluation demonstrates that the hypothesis that juvenile migratory *A.*

striatus birds caught along two distinct migration routes on opposite sides of the Sierra Nevada

Mountains of North America (Pacific Coast and Intermountain Migratory Flyways) come from

different natal populations can be rejected (Fig 6). We found an overlap in the assigned natal

territories of the migrating birds from the two migration routes. Birds captured along the Pacific

Coast Flyway had a range of δ^2H_F and $\delta^{18}O_F$ values that were consistent with precipitation found

not only along the west coast, but also in the western interior regions in the US and Canada. The

birds migrating along the Intermountain Flyway had a more limited geographical origin that also

differed if predicted based on SIA of hydrogen or oxygen. The methodology based on oxygen

appeared to have less predictive power than the one based on hydrogen perhaps because of the

weaker relationships linking feather oxygen isotope ratios to those in precipitation.

We conclude that juvenile migrating *A. striatus* in western North America do not

differentiate into fully separate migratory populations. For this difficult-to-track and secretive

breeding raptor, our data can provide clues to the origins of birds caught along these two

migration routes, and that consideration must be given to both flyways when examining changes

in population size at breeding origins, especially along the Rocky Mountain Range and in the

interior western regions of the species' range. However, further work will need to be done, with

larger sample sizes, to determine what may be driving the lack of correlation found between the

feather hydrogen and oxygen stable isotope compositions of *A. striatus* that migrate through the

Intermountain Flyway.

Results from this study corroborate previous work showing that feather isotopes can be

useful for identifying sites of origin for migrating birds, but also highlight that caution must be

taken when interpreting the outcome, especially if derived by the stable oxygen isotope

composition of feathers. Detailed studies on the sources of isotope variation at stages along the

path of ingestion and assimilation of water into body tissues, including different trophic levels,

life history stages, and geographic complexity, could provide insight for a wider application of

SIA of hydrogen and oxygen to track movement of different organisms, and especially wild

populations. Identification of natal and breeding habitats has important conservation

implications, specifically because the movement of migratory species can often span across large

geographic areas and international borders. Organization of conservation efforts for such species

requires a precise understanding of movement patterns and connections between breeding sites,

migratory pathways, and wintering grounds. The continual development and testing of intrinsic

methods, such as stable isotopes, to track animals using feathers or hair can greatly improve our

insight into how animals, such as raptors like *A. striatus*, travel and move across their ranges.

Acknowledgements

This research is an outgrowth of a project for the class *Stable Isotope Ecology* taught at the

University of California, Berkeley (by T.E. Dawson and S. Mambelli). We thank Hiromi Uno and

Tami Mau for helpful discussion that improved this research, and Paul D. Brooks and Wenbo

Yang for their assistance with IRMS analyses. We thank the following museums and migratory

sites for all of their hard work collecting and providing us with feathers for this study: CAS,

CRCM, MVZ, SDNHM, UWYMV, GGRO, and HawkWatch International Goshutes banding

station. This is contribution #141 for the Golden Gate Raptor Observatory.

705
706
References

1. Rappole JH. The Avian Migrant: The Biology of Bird Migration. New York; Columbia
University Press; 2013.
2. Both C, Visser ME. Adjustment to climate change is constrained by arrival date in a
long-distance migrant bird. Nature. 2001; 411: 296–298.
https://doi.org/10.1038/35077063.
3. Sillett TS, Holmes RT. Variation in survivorship of a migratory songbird throughout its
annual cycle. J Anim Ecol. 2002; 71: 296–308; https://doi.org/10.1046/j.1365-
2656.2002.00599.x.
4. Baker AJ, Gonzalez PM, Piersma T, Niles LJ, de Lima Serrano do Nascimento I,
Atkinson PW, et al. Rapid population decline in red knots: fitness consequences of
decreased refuelling rates and late arrival in Delaware Bay. Proc Biol Sci. 2004; 271:
875–882. https://doi.org/10.1098/rspb.2003.2663.
5. Bildstein KL, Smith JP, Yosef R. Migration Counts and Monitoring. Raptor Research and
Management Techniques. Bird D M, Bildstein KL, editors. Blaine (WA): Hancock House
Publishing; 2007. p.101–116.

6. Newton I. The Migration Ecology of Birds. Boston (MA): Academic Press; 2008. p.19–41.

7. Berthold P, Helbig AJ. The genetics of bird migration - stimulus, timing, and direction. Ibis. 1992; 134: 35–40. https://doi.org/10.1111/j.1474-919X.1992.tb04731.x.

8. Pulido F, Berthold P, Mohr G, Querner U. Heritability of the timing of autumn migration in a natural bird population. Proc R Soc Lond B Biol Sci. 2001; 268: 953–959. DOI: 10.1098/rspb.2001.1602.

9. Bairlein F, Eikenaar C, Schmaljohann H. Routes to genes: unraveling the control of avian migration–an integrated approach using Northern Wheatear *Oenanthe oenanthe* as model organism. J. Ornithol. 2015; 156 (Suppl 1): S3-S14. https://doi.org/10.1007/s10336-015-1224-3.

10. Helbig A. Inheritance of migratory direction in a bird species: a cross-breeding experiment with SE-and SW-migrating blackcaps (*Sylvia atricapilla*). Behav Ecol Sociobiol. 1991; 28: 9–12. https://doi.org/10.1007/BF00172133.

11. Berthold PA. Comprehensive theory for the evolution, control and adaptability of avian migration. Ostrich. 1999; 70: 1–11. https://doi.org/10.1080/00306525.1999.9639744.

12. Hoffman SW, Smith JP, Meehan TD. Breeding grounds, winter ranges, and migratory routes of raptors in the mountain west. J Raptor Res. 2002; 36: 97–110.

13. Farmer CJ, Goodrich LJ, Inzunza ER, Smith JP. Conservation status of North America's birds of prey. In: State of North America's Birds of Prey. Series in Ornithology 3; 2008. p. 303–420.

14. Farmer CJ, Safi K, Barber DR, Newton I, Martell M, Bildstein KL. Efficacy of ,migration counts for monitoring continental populations of raptors: an example using the osprey (*Pandion haliaetus*). Auk. 2010; 127: 863–870. https://doi.org/10.1525/auk.2010.09152.

15. Goodrich LJ, Smith JP. Raptor migration in North America. In: State of North America's Birds of Prey. Series in Ornithology 3; 2008. pp. 37–149.

16. Nagy AC. Population trend indices based on 40 years of autumn counts at Hawk Mountain Sanctuary in North-eastern Pennsylvania. In: R. D. Chancellor, editor. World Conference on Birds of Prey, Vienna Report of Proceedings;. London (UK): International Council for Bird Preservation;1977. pp. 243-252.

17. Kjellén N, Roos G. Population trends in Swedish raptors demonstrated by migration counts at Falsterbo, Sweden 1942–97. Bird Study. 2010; 47: 195–211. https://doi.org/10.1080/00063650009461175.

18. Panuccio M, Martin B, Morganti M, Onrubia A, Ferrer M. Long-term changes in autumn migration dates at the Strait of Gibraltar reflect population trends of soaring birds. Ibis. 2017; 159:, 55–65. https://doi.org/10.1111/ibi.12420.

19. Studds CE, Kendall BE, Murray NJ, Wilson HB, Rogers DI, Clemens RS, et al. Rapid population decline in migratory shorebirds relying on Yellow Sea tidal mudflats as stopover sites. Nat Commun. 2017; 8: 14895. https://doi.org/10.1038/ncomms14895.

20. Faaborg J, Holmes RT, Anders AD, Bildstein KL, Dugger KM, Gauthreaux SAJ, et al. Conserving migratory land birds in the New World: Do we know enough? Ecol Appl. 2010; 20: 398–418. https://doi.org/10.1890/09-0397.1.

21. Bildstein KL, Meyer K. Sharp-shinned Hawk (*Accipiter striatus*). In: Poole A, Gill F, editors. The Birds of North America. Philadelphia (PA): The Birds of North America,

Inc.; 2000. No. 482, https://birdsna.org/Species-Account/bna/species/shshaw/introduction.

22. Bent AC. Life Histories of North American Birds of Prey, Order Falconiformes (Part 1). United States National Museum, Bulletin. 1937; 167: 95-111. https://doi.org/10.5479/si.03629236.167.i.

23. Carlson JE, Piirto DD, Keane JJ, Gill SJ. Estimating site occupancy and detection probabilities for Cooper's and Sharp-Shinned Hawks in the Southern Sierra Nevada. J Raptor Res. 2015; 49: 450–457. https://doi.org/10.3356/rapt-49-04-450-457.1.

24. Bildstein KL. A Brief History of Raptor Conservation in North America. In: State of North America's Birds of Prey. Series in Ornithology 3; 2008. pp. 5-36.

25. Palmer RS. Sharp-shinned Hawk *Accipiter striatus*. In: Handbook of North American Birds; New Haven and London (UK): Yale University Press; 1988. Volume 4, pp. 304-319..

26. Viverette CB, Struve S, Goodrich LJ, Bildstein KL. Decreases in migrating Sharp-shinned Hawks (*Accipiter striatus*) at traditional raptor-migration watch sites in eastern North America. Auk. 1996; 113: 32–40. DOI: 10.2307/4088933 https://www-jstor-org.libproxy.uwyo.edu/stable/4088933.

27. Hull AC. Band Recoveries 2007. In: Pacific Raptor Report. 2008; pp. 29: 14 [Cited 2018 13 August]. Available from: http://www.parksconservancy.org/assets/programs/ggro/pdfs/pacific-raptor-report-2008.pdf.

28. Harn L. Band Recoveries 2008-2009. In: Pacific Raptor Report. 2009; 31: 14-20 [Cited 13 August 2018]. Available from: http://www.parksconservancy.org/assets/programs/ggro/pdfs/pacific-raptor-report-31.pdf.

29. Rubenstein D, Chamberlain C, Holmes R, Ayres M, Waldbauer J, Graves G, et al. Linking breeding and wintering ranges of a migratory songbird using stable isotopes. Science. 2002; 295: 1062–1065. DOI: 10.1126/science.1067124.

30. Clegg S, Kelly J, Kimura M, Smith T. Combining genetic markers and stable isotopes to reveal population connectivity and migration patterns in a Neotropical migrant, Wilson's Warbler (*Wilsonia pusilla*). Mol Ecol. 2003; 12: 819–830. https://doi.org/10.1046/j.1365-294X.2003.01757.x.

31. Rubenstein D, Hobson K. From birds to butterflies: animal movement patterns and stable isotopes. Trends Ecol Evol. 2004; 19: 256–263. https://doi.org/10.1016/j.tree.2004.03.017.

32. Boulet M, Norris DR. Introduction: The past and present of migratory connectivity. Ornithol Monogr. 2006; pp. 1–13. DOI: 10.2307/40166835 https://www.jstor.org/stable/40166835.

33. Inger R, Bearhop S. Applications of stable isotope analyses to avian ecology. Ibis. 2008; 150: 447–461. https://doi.org/10.1111/j.1474-919X.2008.00839.x.

34. Rushing CS, Ryder TB, Saracco JF, Marra PP. Assessing migratory connectivity for a long-distance migratory bird using multiple intrinsic markers. Ecol Appl. 2014; 24: 445–456. https://doi.org/10.1890/13-1091.1.

35. Seifert N, Haase M, Van Wilgenburg SL, Voigt CC, Schmitz Ornés A. Complex migration and breeding strategies in an elusive bird species illuminated by genetic and isotopic markers. J Avian Biol. 2015; 47: 275–287. https://doi.org/10.1111/jav.00751.

36. Hobson KA. Isotope Landscapes for Terrestrial Migration Research. In: Hobson KA,
Wassenaar LI, editors.Tracking Animal Migration with Stable Isotopes, 1st ed. London
(UK): Academic Press; 2008; Volume 2: pp.79-106..
37. Pietsch, S. J., Hobson, K. A., Wassenaar, L. I., and Tütken, T. Tracking Cats: Problems
with Placing Feline Carnivores on δ^{18}O, δD Isoscapes. PLoS ONE 2011, 6, e24601–11,
https://doi.org/10.1371/journal.pone.0024601.
38. Soto DX, Wassenaar LI, Hobson KA. Stable hydrogen and oxygen isotopes in aquatic
food webs are tracers of diet and provenance. Funct Ecol. 2013; 27: 535–543.
https://doi.org/10.1111/1365-2435.12054.
39. Hobson KA, Koehler G. On the use of stable oxygen isotope (δ^{18}O) measurements for
tracking avian movements in North America. Ecol Evol. 2015; 5: 799–806.
https://doi.org/10.1002/ece3.1383.
40. Pekarsky S, Angert A, Haese B, Werner M, Hobson KA, Nathan R. Enriching the
isotopic toolbox for migratory connectivity analysis: a new approach for migratory
species breeding in remote or unexplored areas. Divers Distrib. 2015; 21: 416–427.
https://doi.org/10.1111/ddi.12306.
41. Rozanski K, Araguas-Araguas L, Gonfiantini R. Isotopic patterns in modern global
precipitation. In: Swart PK, Lohmann KC, McKenzie J, Savin S. editors .Climate Change
in Continental Isotopic Records. Washington, DC: Am. Geophys. Union; 1993. Volume
78: pp. 1–36.
42. Epstein S, Mayeda T. Variation of O-18 content of waters from natural sources. Geochim
Cosmochim Acta. 1953; 4: 213-224. https://doi.org/10.1016/0016-7037(53)90051-9.
43. Friedman I. Deuterium content of natural water and other substances. Geochim
Cosmochim Acta. 1953; 4: 89-103. https://doi.org/10.1016/0016-7037(53)90066-0.
44. Craig H. Isotopic variations in meteoric waters. Science. 1961; 133: 1702–1703. DOI:
10.1126/science.133.3465.1702.
45. Dansgaard W. Stable isotopes in precipitation. Tellus. 1964; 16: 436–468.
https://doi.org/10.3402/tellusa.v16i4.8993.
46. Bowen GJ. Isoscapes: spatial pattern in isotopic biogeochemistry. Annu Rev Earth Planet
Sci. 2010; 38: 161-187. https://doi.org/10.1146/annurev-earth-040809-152429.
47. Meier-Augenstein W, Hobson KA,Wassenaar LI. Critique: measuring hydrogen stable
isotope abundance of proteins to infer origins of wildlife, food and people. Bioanalysis.
2013; 5: 751–767. https://doi.org/10.4155/bio.13.36.
48. Hobson K, DeMent S, Van Wilgenburg S, Wassenaar L. Origins of American Kestrels
wintering at two southern US sites: an investigation using stable-isotope (δD, δ18O)
methods. J Raptor Res. 2009; 43: 325–337. https://doi.org/10.3356/JRR-08-74.1.
49. Storm-Suke A, Wassenaar LI, Nol E, Norris DR. The influence of metabolic rate on the
contribution of stable-hydrogen and oxygen isotopes in drinking water to quail blood
plasma and feathers. Funct Ecol. 2012; 26: 1111-1119. doi:10.1111/j.1365-
2435.2012.02014.x.
50. Smith RB, Meehan TD, Wolf BO. Assessing migration patterns of sharp-shinned hawks
Accipiter striatus using stable-isotope and band encounter analysis. J Avian Biol. 2003;
34: 387–392. https://doi.org/10.1111/j.0908-8857.2003.03107.x.
51. Lott CA, Smith JP. A geographic-information-system approach to estimating the origin
of migratory raptors in North America using stable hydrogen isotope ratios in feathers.
Auk. 2006; 123: 822. https://doi.org/10.1642/0004-

8038(2006)123[822:AGATET]2.0.CO;2.

52. Oppel S, Dobrev V, Arkumarev V, Saravia V, Bounas A, Kret E, et al. High juvenile
mortality during migration in a declining population of a long-distance migratory raptor.
Ibis. 2015; 157: 545–557. https://doi.org/10.1111/ibi.12258.

53. Hull J, Girman D. Effects of Holocene climate change on the historical demography of
migrating Sharp-shinned Hawks (*Accipiter striatus velox*) in North America. Mol Ecol.
2005; 14: 159–170. https://doi.org/10.1111/j.1365-294X.2004.02366.x.

54. Hull JM, Hull AC, Sacks BN, Smith JP, Ernest HB. Landscape characteristics influence
morphological and genetic differentiation in a widespread raptor (*Buteo jamiacensis*).
Mol Eco. 2008; 17: 810-824.

55. Bowen GJ, West JB. Isotope landscapes for terrestrial migration research. In: Hobson
KA, Wassenaar LI, editors. Tracking animal migration with stable isotopes. Amsterdam
(Netherlands): Elsevier; 2008. pp. 79–105.

56. Pyle P. Identification guide to North American birds, Part II Anatidae to Alcidae. Point
Reyes Station (CA): Slate Creek Press; 2008. pp. 415-419.

57. Smith A, Donohue K, Dufty A Jr. Intrafeather and intraindividual variation in the stable-
hydrogen isotope (δ D) content of raptor feathers. Condor. 2008; 110: 500–506.
https://doi.org/10.1525/cond.2008.8515.

58. Paritte J, Kelly J. Effect of cleaning regime on stable-isotope ratios of feathers in
Japanese quail (*Coturnix japonica*). Auk. 2009; 126: 165–174. DOI:
10.1525/auk.2009.07187 https://www.jstor.org/stable/10.1525/auk.2009.07187.

59. Bontempo L, Ceppa F, Ziller L, Pedrini P, Hobson KA, Wassenaar LI, et al. Comparison
of methods for stable isotope ratio (δ ^{13}C, δ ^{15}N, δ ^{2}H, δ ^{18}O) measurements of feathers.
Methods Ecol Evol. 2014; 5: 363–371. https://doi.org/10.1111/2041-210X.12165.

60. Greenwood J, Dawson R. Correlates of Deuterium (δ D) enrichment in the feathers of
adult American Kestrels of known origin. Condor. 2011; 113: 555–564. DOI:
10.1525/cond.2011.100213 https://www.jstor.org/stable/10.1525/cond.2011.100213.

61. De Ruyck C, Hobson KA, Koper N, Larson KW, Wassenaar LI. An appraisal of the use
of hydrogen-isotope methods to delineate origins of migratory Saw-whet Owls in North
America. Condor. 2013; 115: 366–374. https://doi.org/10.1525/cond.2013.120019.

62. Briggs CW, Poulson SR, Collopy MW. Correlation between feather isotopes and body
condition for Swainson's Hawks, and implications for migration studies. J Raptor Res.
2017; 51: 107–114. https://doi.org/10.3356/JRR-15-00002.1.

63. Wassenaar LI, Hobson KA. Two new keratin standards (δ2H, δ18O) for daily laboratory
use in wildlife and forensic isotopic studies, In the 7th Inernational Conference on
Applications of Stable Isotope Technqiues to Eoclogical studies. University of Alaska,
Fairbanks, Alaska, USA. 2010.

64. Soto DX, Koehler G, Wassenaar LI, and Hobson KA. Re-evaluation of the hydrogen
stable isotopic composition of keratin calibration standards for wildlife and forensic
science applications. Rapid Commun. Mass Spectrom. 2017; 31: 1193– 1203. https://doi:
10.1002/rcm.7893.

65. Wassenaar L, Hobson K. Comparative equilibration and online technique for
determination of non-exchangeable hydrogen of keratins for use in animal migration
studies. Isotopes Environ Health Stud. 2003; 39: 211–217.
https://doi.org/10.1080/1025601031000096781.

66. Bowen GJ. Gridded maps of the isotopic composition of meteoric waters. 2019 [cited 28

June 2020]. Available from:
http://wateriso.utah.edu/waterisotopes/pages/information/mapping.html

67. Bowen GJ. Statistical and geostatistical mapping of precipitation water isotope ratios. In: West JB, Bowen G, Dawson T, Tu K, editors. Isoscapes: understanding movement, pattern, and process on Earth through isotope mapping;. Verlag (Netherlands): Springer; 2010. pp. 487.

68. IAEA/WMO Global network for isotopes in precipitation, the GNIP database. 2004. Available from: http://www-naweb.iaea.org/napc/ih/GNIP/IHS_GNIP.html.

69. Hobson KA. Using stable isotopes to trace long-distance dispersal in birds and other taxa. Divers Distrib. 2005; 11: 157-164. https://doi.org/10.1111/j.1366-9516.2005.00149.x.

70. Marrack LC. IsoMAP job key 50333, Isoscapes Modeling, Analysis and Prediction (version 1.0). The IsoMAP Project. 2015. http://isomap.org.

71. Marrack LC. IsoMAP job key 63026, Isoscapes Modeling, Analysis and Prediction (version 1.0). The IsoMAP Project. 2017. http://isomap.org.

72. BirdLife International and NatureServe. Bird species distribution maps of the world; BirdLife International. Cambridge (UK) and Arlington: NatureServe. 2015 [cited 15 January 2016].

73. Brown D, Brennan T, Unmack PA. Digitized biotic community map for plotting and comparing North American plant and animal distributions. CANOTIA. 2007; 3: 1. Available from: http://www.canotia.org/volumes/CANOTIA_2007_Vol3_1_Brown_et_al.pdf.

74. Chamberlain C, Blum J, Holmes R, Feng X, Sherry T, Graves G. The use of isotope tracers for identifying populations of migratory birds. Oecologia. 1996; 109: 132-141. https://doi.org/10.1007/s004420050067.

75. Hobson KA, Wassenaar LI. Linking breeding and wintering grounds of neotropical migrant songbirds using stable hydrogen isotopic analysis of feathers. Oecologia. 1997; 109: 142–148. https://doi.org/10.1007/s004420050068.

76. Hobson KA. Using stable isotopes to trace long-distance dispersal in birds and other taxa. Divers Distrib. 2005; 11: 157-164. https://doi.org/10.1111/j.1366-9516.2005.00149.x.

77. Martinez del Rio C, Wolf N, Carleton C, Gannes Z. Isotopic ecology ten years after a call for more laboratory experiments. Biol Rev Camb Philos Soc. 2009; 84: 91-111. https://doi.org/10.1111/j.1469-185X.2008.00064.x.

78. Wolf N, Bowen G, Martinez del Rio C. The influence of drinking water on the δ D and δ 18O values of house sparrow plasma, blood and feathers. J Exp Biol. 2011; 214. 98–103, doi: 10.1242/jeb.050211.

79. Bowen GJ, Liu Z, Vander Zanden HB, Zhao L, Takahashi G. Geographic assignment with stable isotopes in IsoMAP. Methods Ecol Evol. 2014; 5: 201-206. doi:10.1111/2041-210X.12147

80. Wunder MB. Using isoscapes to model probability surfaces for determining geographic origins. In: West J, Bowen G, Dawson T, Tu KP, editors. Isoscapes; Understanding movement, pattern, and process on Earth through isotope mapping. Dordrecht (Netherlands): Springer; 2010. pp. 251–270.

81. R Development Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing. Vienna (Austria). 2011; ISBN 3-900051-07-0, Available from: http://www.R-project.org/.

82. Hedenström A. Adaptations to migration in birds: behavioral strategies, morphology and scaling effects. Philos Trans R Soc Lond B Biol Sci. 2008; 363: 287-299. doi:10.1098/rstb.2007.2140.

83. Sergio F, Tanferna A, De Stephanis R, Jiménez LL, Blas J, Tavecchia G, et al. Individual improvements and selective mortality shape lifelong migratory performance. Nature. 2014; 515: 410-413. doi:10.1038/nature13696.

84. Bowen GJ, Wassenaar LI, Hobson KA. Global application of stable hydrogen and oxygen isotopes to wildlife forensics. Oecologia. 2005; 143: 337-348.

85. Hobson K, Barnett-Johnson R, Cerling T. Using isoscapes to track animal migration. In: West J, Bowen G, Dawson T, Tu KP, editors. Isoscapes; Understanding movement, pattern, and process on Earth through isotope mapping. Dordrecht (Netherlands): Springer; 2010. pp. 273–298. https://doi.org/10.1007/978-90-481-3354-3_13.

86. Morganti M, Franzoi A, Bontempo L, Sara M. An exploration of isotopic variability in feathers and claws of Lesser Kestrel *Falco naumanni* chicks from southern Sicily. Avocetta. 2016; 40: 23–32.

87. Hull JM, Ernest HV, Harley JA, Fish AM, Hull AC. Differential migration between discrete populations of juvenile Red-Tailed Hawks (*Buteo jamaicensis*). Auk. 2009; 126: 389-396.

88. Wolf N, Newsome SD, Fogel ML, Martinez del Rio C. The relationship between drinking water and the hydrogen and oxygen stable istotope values of tissues in Japanese Quail (*Cortnix japonica*). Auk. 2013; 130: 323-330. Doi:10.1525/auk.2013.12075.

89. Vander Zanden HB, Soto DX, Bowen GJ, Hobson KA. Expanding the isotopic toolbox: applications of hydrogen and oxygen stable isotope ratios to food web studies. Front Ecol Evol. 2016; 4: 3741–20. https://doi.org/10.3389/fevo.2016.00020.

90. Birchall J, O'Connell TC, Heaton THE, Hedges REM. Hydrogen isotope ratios in animal body protein reflect trophic level. J Anim Ecol. 2005; 74: 877–881. https://doi.org/10.1111/j.1365-2656.2005.00979.x.

91. Nielson KE, Bowen GJ. Hydrogen and oxygen in brine shrimp chitin reflect environmental water and dietary isotopic composition. Geochim Cosmochim Acta. 2010; 74: 1812–1822. https://doi.org/10.1016/j.gca.2009.12.025.

92. Hobson KA, Bowen GJ, Wassenaar L, Ferrand Y, Lormee H. Using stable hydrogen and oxygen isotope measurements of feathers to infer geographical origins of migrating European birds. Oecologia. 2004; 141: 477-488.

93. Hobson KA, Wassenaar LI, Milá B, Lovette I, Dingle C, Smith TB. Stable isotopes as indicators of altitudinal distributions and movements in an Ecuadorean hummingbird community. Oecologia. 2003; 136: 302-308.

94. Kohn MJ. Predicting animal δ 18O: accounting for diet and physiological adaptation. Geochim Cosmochim Acta. 1996; 60: 4811–4829. https://doi.org/10.1016/S0016-7037(96)00240-2.

95. Crowley BE, Melin AD, Yeakel JD, Dominy NJ. Do oxygen isotope values in collagen reflect the ecology and physiology of neotropical mammals? Front Ecol Evol. 2015; 3: 431–12. https://doi.org/10.3389/fevo.2015.00127.

96. Smith AD, Dufty AM. Variation in the stable-hydrogen isotope composition of Northern Goshawk feathers: relevance to the study of migratory origins. Condor. 2005; 107: 547–558. https://doi.org/10.1650/0010-5422(2005)107[0547:VITSIC]2.0.CO;2.

97. Gow E, Stutchbury B, Done T, Kyser T. An examination of stable hydrogen isotope (δ

996   D) variation in adult and juvenile feathers from a migratory songbird. Can J Zool. 2012;
997   90: 585–594. https://doi.org/10.1139/z2012-024.
998   98. van Dijk JGB, Meissner W, Klaassen M. Improving provenance studies in migratory
999   birds when using feather hydrogen stable isotopes. J Avian Biol. 2014; 45: 103–108. Doi:
1000   10.1111/j.1600-048X.2013.00232.x.

1001
1002

Supporting Information

1004 **Table S1: Oxygen stable isotope composition of feathers of adult museum Sharp-shinned**
1005 **Hawk (*Accipiter striatus*) specimens ($\delta^{18}O_f$ values (‰)) ($n = 25$), and isoscape modeled stable**
1006 **hydrogen and oxygen isotope composition of precipitation ($\delta^{18}O_p$ values (‰)) of known**
1007 **breeding origin.**

1008

Table S2: Stable isotopic compositions of museum feathers and estimates of precipitation
isotopes used to test assignment of origin models. Stable hydrogen (δ^2H_F ‰) and oxygen
($\delta^{18}O_F$ ‰) isotopic composition of feathers for juvenile museum Sharp-shinned Hawk (*Accipiter*
striatus) specimens of known natal origin and estimated isotopic compositions of precipitation
(δ^2H_P and $\delta^{18}O_P$ ‰) from the transfer functions used to determine assignment of origin. The
states where samples were collected and the museum Specimen IDs are included.

1015

S1 Fig: Map of sample locations for museum specimens of juvenile and adult Sharp-
shinned Hawk (*Accipiter striatus*) feathers. Sampling locations are shown in reference to the
species known range in Western North America (light gray), and suitable breeding forest habitat
(dark gray). Juveniles samples ($n = 23$) are shown as triangles, and adult samples ($n = 25$) are
shown as circles. State and country boundaries are modified from public domain GIS files, US
Census Bureau (2016) and Natural Earth (2020). Species range acquired from Birdlife
International and NatureServe (2015), and data to create the GIS biome layer acquired from
Brown, Bennan, and Unmack (2007).

1024

S2 Fig: Stable hydrogen (δ^2H_P ‰) and oxygen ($\delta^{18}O_P$ ‰) isoscapes created within IsoMAP.
These were used to determine transfer functions for specimens of known origin and performing
assignment of origin for migrating specimens. The hydrogen isoscape (A) produced was based
on 117 stations, had resolution of 9x9 km, a correlation parameter of 0.93, and included the
variables elevation (ETOPO, $P < 0.001$), latitude ($P < 0.001$) and longitude ($P = 0.06$) (available
as IsoMAP job key 50333 (Marrack 2015)). The most robust oxygen isoscape (B) was based on
120 stations, had resolution of 9x9 km, a correlation parameter of 0.92, and included the
variables elevation (ETOPO, $P < 0.001$), latitude ($P < 0.001$) and longitude ($P = 0.05$) (available
as IsoMAP job key 63026 (Marrack 2017)).

1034

S3 Fig: **Relationship between the stable isotopic compositions of museum feathers and the**
isoscape model of precipitation for adult and juvenile Sharp-shinned Hawks (*Accipiter*
***striatus*).** Stable oxygen ($\delta^{18}O_F$ ‰) isotopic composition of feathers for museum Sharp-shinned
Hawk (*Accipiter striatus*) specimens of known natal/breeding origin and isoscape modeled
isotopic composition of precipitation ($\delta^{18}O_P$ ‰) at the collection locations of (a) juvenile (black

dots) birds as well as (b) adult birds (white dots) ($n = 48$). The linear regression for both juvenile
and adult birds is $y = 0.29x – 14.77$, $R^2 = 0.1$, $P = 0.03$ (dashed line). Note that this equation is
similar in slope and intercept to the linear regression relationship for juvenile birds ($y = 0.385x –$
14.67 ($R^2 = 0.14$, $P = 0.07$) (solid line)).
1044
S4 Fig: Relationship between the stable isotopic compositions of museum feathers and the
isoscape model of precipitation. Stable hydrogen (δ^2H_F ‰) and oxygen ($\delta^{18}O_F$ ‰) isotopic
composition of feathers for museum juvenile Sharp-shinned Hawk (*Accipiter striatus*) specimens
($n = 23$) of known natal origin and the isoscape modeled isotopic compositions of precipitation
(δ^2H_P and $\delta^{18}O_P$ ‰) at the collection locations: (a) δ^2H_F values of birds versus δ^2H_P values, (b)
$\delta^{18}O_F$ values of birds versus $\delta^{18}O_P$ values.
1051
S5 Fig: Probability density maps predicting the origin of museum specimens of Sharp-
shinned Hawks (*Accipiter striatus*) with known collection locations. Maps are based on
predicted δ^2H_P values (‰) (left panels) and predicted $\delta^{18}O_P$ values (‰) (right panels) for birds
captured at known locations. Each map represents the probability density surface created for an
individual bird with the known sampling location shown by a circle. State and country
boundaries are from public domain GIS files US Census Bureau (2016) and Natural Earth
(2020). Species range acquired from BirdLife International and NatureServe (2015), and data to
create the GIS biome layer acquired with permission from Brown, Bennan, and Unmack (2007).