Rough and smooth variant *Mycobacterium abscessus* infections are differentially controlled by

host immunity during chronic infection of adult zebrafish

Elinor Hortle[1,2]*, Julia Y Kam[1]*, Elizabeth Krogman[1]*, Sherridan E Warner[1,2], Pradeep

Manuneedhi Cholan[1], Kazu Kikuchi[3,4], James A Triccas[2], Warwick J Britton[1,2], Matt D Johansen[5],

Laurent Kremer[5,6], Stefan H Oehlers[1,2#]

1 Tuberculosis Research Program at the Centenary Institute, The University of Sydney,

Camperdown NSW Australia

2 The University of Sydney, Faculty of Medicine and Health & Marie Bashir Institute,

Camperdown NSW Australia

3 Developmental and Stem Cell Biology Division, Victor Chang Cardiac Research Institute,

Darlinghurst, NSW Australia

4 St. Vincent's Clinical School, University of New South Wales, Kensington, NSW 2052, Australia

5 Centre National de la Recherche Scientifique UMR 9004, Institut de Recherche en Infectiologie

de Montpellier (IRIM), Université de Montpellier, 1919 Route de Mende, 34293, Montpellier,

France.

6 INSERM, IRIM, 34293 Montpellier, France.

*These authors contributed equally and are listed in alphabetical order.

[#] Corresponding author: Dr Stefan Oehlers, stefan.oehlers@sydney.edu.au

Author contributions

EH: qPCR analyses; JK: histological analysis; EK: dose finding, survival experiments, histological

analysis; SW: CFU recovery assays; PMC: histological analysis; KK: provided reagents; JAT &

WJB: supervision of study; MDJ: conceived study, supervision of study, wrote manuscript; LK:

conceived study, provided reagents, supervision of study, wrote manuscript; SHO: conceived study,

performed experiments, supervision of study, wrote manuscript.

Keywords: non-tuberculous mycobacterium, rapid-growing mycobacteria, animal model, zebrafish,

glycopeptidolipids

Abstract

Infections caused by *Mycobacterium abscessus* are increasing in prevalence within patient groups

with respiratory comorbidities including Cystic Fibrosis or Chronic Obstructive Pulmonary Disease.

Initial colonisation by the smooth colony *M. abscessus* (S) can be followed by an irreversible

genetic switch into a highly inflammatory rough colony *M. abscessus* (R), often associated with a

decline in pulmonary function. Currently available animal models such as the embryonic zebrafish,

have largely explored the role of innate immunity in the pathogenesis of *M. abscessus*, and

demonstrated that infection with the R variant produces a hyperinflammatory infection due to the

presence of large extracellular cords, whereas the S variant produces a chronic persistent infection.

However, our understanding of the role of adaptive immunity in *M. abscessus* pathogenesis is

largely unknown. Here, we have used intraperitoneal infection of adult zebrafish to model *M.*

abscessus pathogenesis in the context of fully functioning host immunity. We find infection with

the R variant penetrates host organs causing an inflammatory immune response leading to necrotic

granuloma and abscess formation within 2 weeks. The R bacilli are targeted by T cell-mediated

immunity and burden is progressively reduced. Strikingly, the S variant colonises host internal

surfaces at high loads and is met with a robust innate immune response but little T cell-mediated

immunity. Invasive granuloma formation is delayed in S variant infection compared to R variant

infection upon which T cell-mediated immunity is required to control infection. In mixed infections,

the S variant outcompetes the R variant in an adaptive-immunity dependent manner. We also find

the R variant activates innate immunity to detriment of S variant *M. abscessus* in mixed infections.

These findings demonstrate the applicability of the adult zebrafish to model persistent *M. abscessus*

infection and provide insight into the immunopathogenesis of chronic *M. abscessus* infection.

Introduction

Mycobacterium abscessus is an increasingly recognized human pathogen responsible for a wide

array of clinical manifestations including muco-cutaneous infections, disseminated or chronic

pulmonary diseases. The latter is mostly encountered in patients with underlying lung disorders,

such as bronchiectasis or cystic fibrosis (CF). Irrespective of being a rapid-growing mycobacteria

(RGM), *M. abscessus* displays many pathophysiological traits with slow-growing mycobacteria

(SGM), such as *Mycobacterium tuberculosis*. These include the capacity to persist silently within

granulomatous structures and to produce pulmonary caseous lesions (1, 2). In addition, *M.*

abscessus is notorious for being one of the most-drug resistant mycobacterial species, characterized

by a wide panel of acquired and innate drug resistance mechanisms against nearly all anti-tubercular

drugs, as well as many classes of antibiotics (3). Consequently, this explains the complexity and

duration of the treatments and the high level of therapeutic failure (4).

M. abscessus exists either as smooth (S) or a rough (R) colony morphotype variants

associated with distinct clinical outcomes (5). Previous epidemiological studies have highlighted the

association of the R variant, persisting for many years in the infected host, with a rapid decline in

the pulmonary functions (6-8). It is well established that these morphological differences between S

and R variants are dependent on the presence or absence of surface-exposed glycopeptidolipids

(GPL), respectively (5, 9, 10). However, our knowledge of the pathophysiological characteristics

and interactions between R or S variants with the host immune cells remains largely incomplete and

is hampered by the lack of animal models that are permissive to persistent *M. abscessus* infection.

Intravenous injection or aerosol administration of *M. abscessus* in immunocompetent BALB/c

mice fails to establish a persistent infection, typified by a rapid clearance of the bacilli from the

liver, spleen and lungs within 4 weeks (11). Immunosuppression is required to produce a

progressive high level of infection with *M. abscessus* in mice, as shown in nude, SCID (severe

combined immunodeficiency), interferon-gamma (GKO) and granulocyte-macrophage colony-

stimulating factor (GM-CSF) knock-out mice (12).

The contribution of B and T cells in the control of *M. abscessus* infection has been studied in

C57BL/6 mice with $Rag2^{-/-}$, $Cd3e^{-/-}$ and $\mu MT^{-/-}$ knockouts. These studies indicated that infection

control was primarily T cell dependent in the spleen, and both B and T cell dependent in the liver

(13). In addition, IFNg-receptor KO mice ($ifngr1^{-/-}$) were significantly impaired in their control of

M. abscessus both in the spleen and in the liver, with markedly different granulomas and more

pronounced in $TNF^{-/-}$ mice (13). This points to the central role of T-cell immunity, IFNg and TNF

for the control of *M. abscessus* in C57BL/6 mice, similarly to the control of *M. tuberculosis*

infection.

In recent years, alternative non-mammalian models, such as Drosophila (14), Galleria larvae

(15), and zebrafish embryos (16) have been developed to study the chronology and pathology of *M.*

abscessus infection and for *in vivo* therapeutic assessment of drugs active against *M. abscessus*. In

particular, zebrafish embryos have delivered important insights into the pathogenesis of *M.*

abscessus and the participation of innate immunity in controlling infection. The optical

transparency of zebrafish embryos has been used to visualise the formation of large extracellular

cords by the R form *in vivo*, representing a mechanism of immune subversion by preventing

phagocytic destruction and highlighting the importance bacterial virulence factors such as the

dehydratase MAB_4780 and the $MmpL8_{MAB}$ lipid transporter (9, 17, 18). Other studies in zebrafish

embryos have demonstrated the contribution of host TNF signalling and IL8-mediated neutrophil

recruitment for protective granulomatous immunity against *M. abscessus* (19), and the link

between dysfunctional CFTR and vulnerability to *M. abscessus* infection via the macrophage

oxidative response (20).

Adult zebrafish models have been well-described for the study of mycobacterial pathogenesis

by *Mycobacterium marinum*, used as a surrogate for the closely related *M. tuberculosis*, and the

human pathogen *Mycobacterium leprae* (21-23). Encompassing a fully functional immune system,

previous studies in adult zebrafish with pathogenic mycobacteria such as *M. marinum* have

unravelled the interplay between innate and adaptive immunity in mycobacterial granuloma

formation and function.

Herein, we addressed whether adult zebrafish may be a useful host to analyse and compare

the chronology of infection with *M. abscessus* S and R variants and to study the contribution of the

T cell-mediated immunity and granulomatous response in *M. abscessus* infection.

Results

<u>Adult zebrafish can be chronically infected with *M. abscessus.*</u>

To establish the susceptibility of adult zebrafish to infection by *M. abscessus*, we performed a dose

escalation experiment up to 10^6 CFU per animal with the rough (R) and smooth (S) variants of the

reference strain CIP104536^T. While animals underwent a period of sickness behaviour within the

first week of infection, we did not observe mortality for up to 4 weeks post infection (wpi) (data not

shown). To determine if *M. abscessus* produces a persistent infection in adult zebrafish, we

performed CFU recovery on animals infected with a standard dose of 10^5 CFU (Figure 1A). We

observed a progressive clearance of the R variant with a 1-log reduction in burden from 1 day post

infection (dpi) to 4 wpi (P=0.018). Conversely, the S variant was recovered at a consistent burden

across the 4 week duration of the experiment (1 dpi vs 4 wpi, P=0.12).

<u>Adult zebrafish mount a robust inflammatory response to *M. abscessus* infection.</u>

We hypothesised that the better survival of the S variant compared to the R variant could be

attributed to reduced immunogenicity of the CIP104536 S cell surface and differential modes of

growth in macrophages (24). To test this hypothesis, we performed qPCR detection of zebrafish

immune gene expression from whole body homogenates of infected animals (25). We observed

higher innate immunity-associated *il1b* transcription in the *M. abscessus* S-infected group from

early in infection and a late trend to increases in *il1b* transcription in 4 wpi *M. abscessus* R-infected

animals indicating a stronger early innate immune response to the S variant (Figure 1B).

The cytokine *tumour necrosis factor* (*tnfa*) is essential for the granulomatous containment of

M. abscessus in zebrafish embryos (19). There was a non-statistically significant trend to increased

transcription of *tnfa* in whole body homogenates from animals infected with either variant of *M.*

abscessus compared to uninfected animals (Figure 1C). We next took advantage of the

TgBAC(tnfa:GFP)^{pd1028} zebrafish line (26), where GFP expression is driven by the *tnfa* promoter, to

investigate if *tnfa* expression is linked to granuloma formation. Expression of GFP was enhanced

around either variant of *M. abscessus* and was tightly concentrated around organised granulomas

demonstrating a conserved induction of *tnfa* expression during *M. abscessus* granuloma formation

in adult zebrafish (Figure 1D and 1E).

Adult zebrafish contain *M. abscessus* R within granulomas.

We next performed histology on adult zebrafish infected with fluorescent *M. abscessus* R. At 3 dpi,

bacteria were diffusely spread through the peritoneal cavity with occasional foci of infection located

external to peritoneal organs. From 10 dpi to 2 wpi we noted a heterogeneous mix of unorganised

lesions (Figure 2A) and organised lesions with stereotypical concentric rings of nuclei around a

central focus of bacteria and necrotic debris in all animals (Figure 2B). We also observed the

appearance of very large abscess-like granulomas filled with fluorescent bacteria and necrotic

debris measuring over 600 µm at a rate of no more than 1 per infected animal from 2 wpi onwards

(Figure 2C). Oil red O staining revealed the accumulation of foam cells in cellular rim of *M.*

abscessus R granulomas (Figure 2D), consistent with immunopathology seen in

immunocompromised mice infected with *M. abscessus* (12). Although we observed a fairly stable

proportion of organised granulomas in *M. abscessus* R-infected adult zebrafish from 10 to 28 dpi

(Figure 2E), the proportion of fluorescent bacteria associated with organised granulomas

significantly increased from 10 to 28 dpi corresponding to the appearance of abscesses at 14 dpi

(Figure 2F).

Adult zebrafish contain *M. abscessus* S infection prior to delayed granuloma formation.

Histological analysis of adult zebrafish infected with fluorescent *M. abscessus* S revealed

significantly less tissue damage than the equivalent *M. abscessus* R infection up to 2 wpi. *M.*

abscessus S was observed to grow freely in mesenteric spaces and form poorly organised cellular

granulomas at 2 wpi (Figure 3A). We observed the appearance of tissue-invasive organised

granulomas at 4 wpi (Figure 3B). Although these granulomas could reach similar size to the very

large abscess-like granulomas seen in *M. abscessus* R infection, there was little Oil Red O staining

in *M. abscessus* S granulomas indicating a lack of foam cell formation (Figure 3B).

We observed a significant increase in the proportion of organised granulomas in S-infected

adult zebrafish from 2 to 4 wpi (Figure 3C). Similarly, the proportion of fluorescent bacteria

associated with organised granulomas significantly also increased from 2 to 4 wpi corresponding to

the consolidation of *M. abscessus* S into granulomas (Figure 3C).

Adaptive immunity differentially controls infection by *M. abscessus* variants in adult zebrafish.

Given the requirement for T cells to maintain granuloma structure in adult zebrafish *M. marinum*

infection (23), we next asked if there was T cell involvement around *M. abscessus* granulomas

using the recently described *TgBAC(lck:GFP)*[vcc4] zebrafish line (23, 27). We observed T cell

association and penetration throughout unorganised and organised *M. abscessus* R granulomas, but

T cells were largely excluded from the cores of the very large abscess-like lesions (Figure 4A).

Conversely, we did not observe T cell interaction with *M. abscessus* S growing around peritoneal

organs until invasive granuloma formation after 2 wpi, which was indistinguishable from the T cell

response to *M. abscessus* R infection (Figure 4B).

182   To directly test the requirement of T cells in containing *M. abscessus* we next utilised the

lck[-/-sa410] mutant line which is T cell-deficient. We infected wild type (WT) control and *lck*[-/-sa410]

mutant adult zebrafish with both the S and R variants. T cell-deficient adult zebrafish were

significantly more susceptible to *M. abscessus* R infection with reduced survival over 4 weeks of

infection (P = 0.0005, Log-rank test) (Figure 4C). T cell deficiency had a less pronounced effect on

the survival of animals infected with *M. abscessus* S compared to *M. abscessus* R infection (WT S

vs lck-/- S P = 0.03, Log-rank test), although both groups eventually succumbed to infection at the

same rate after 5 wpi (P = 0.78, Log-rank test) (Figure 4C). Consequently, bacterial burden was

significantly increased in 2 wpi *lck*[-/-sa410] mutants infected with the R (Figure 4D), but not the S

variant (Figure 4E). These observations confirm the highly inflammatory nature of *M. abscessus* R

infection is conserved in the adult zebrafish model and mediated by T-cells. Conversely,

colonisation by *M. abscessus* S is contained by sufficient activation of the innate immune response

until invasive infection between 2 and 4 wpi requires T-cell involvement to contain.

M. abscessus S has a survival advantage over *M. abscessus* R.

To examine our hypothesis that *M. abscessus* S has a survival advantage over *M. abscessus* R in the

adult zebrafish infection model, we performed co-infection of adult zebrafish with equal numbers of

each variant expressing either Wasabi or Tdtomato fluorescent proteins to enable simple tracking

(Figure 5A). Analysis of the ratio of R:S colonies recovered revealed a clear and rapid shift in

population proportions from 1:1 at 1 dpi to 0.5 rough:1 smooth ratio that remained stable at 2 wpi

(Figure 5B). Coinfection did not affect the progressive reduction in *M. abscessus* R burden as near

identical *M. abscessus* R CFUs were recovered from single and mixed-infected animals at 1 and 2

wpi (Figure 5C). Coinfection did cause a decrease in the number of recoverable *M. abscessus* S that

was not observed in single variant infections at 1 and 2 wpi demonstrating a negative effect of R

granuloma formation on the survival of *M. abscessus* S (Figure 5C). Whole mount and histological

examination of bacterial distribution in animals infected with *M. abscessus* R expressing Wasabi

and *M. abscessus* S expressing Tdtomato revealed co-mingling of S and R variants within

granulomas (Figure 5D).

We hypothesised that the T cell response induced by the R variant to antigens shared with

the S variant could be responsible for the clearance of S variant in mixed infections. To test this

hypothesis we infected T-cell deficient *lck*^{-/-sa410} mutant animals with a mixed infection of R and S

variants of *M. abscessus*. CFU recovery from these mixed infections in the T-cell deficient animals

revealed a similar decrease in *M. abscessus* S burden as observed in WT animals (Figure 5E).

Therefore the clearance of *M. abscessus* S variant is mediated by innate immune mechanisms.

Interestingly, *lck*^{-/-sa410} zebrafish had a higher proportion of rough to smooth than in comparable WT

animals, confirming the importance of the adaptive immune response for controlling *M. abscessus*

R variant burden (Figure 5F).

Discussion

In this study, we report for the use of adult zebrafish to probe both host and mycobacterial

determinants of pathogenesis during persistent infection with *M. abscessus*. Infection with the R

and S variants was maintained over months of infection in genetically intact animals, a major

improvement on existing mouse models of *M. abscessus* infection.

While the R variant induces a more robust and aggressive infection than the S variant in

zebrafish embryos (9), this appears to not the case in the adult fish. We observed better clearance of

the R variant and establishment of a higher burden of persistent infection with the S variant. One

possible explanation for the better survival of the S compared to the R is that, in the presence of

effective adaptive immunity, R infection is better controlled due to the induction of a potent Th1

cell-mediated response, as evidenced by the increased expression of the CD3 marker and IFN-

gamma response at later time points. This contribution of the T cell response was further

substantiated using T cell-deficient fish, where infection of *lck*^{-/-} fish with the R bacilli resulted in a

higher bacterial burden than in WT fish, which was not observed with S bacilli. These observations

provide insight into the clinical observation that AIDS patients are not at increased risk of *M.*

abscessus infection to the same degree that AIDS is a risk factor for *M. tuberculosis* and other non-

tuberculous mycobacterium infections such as *Mycobacterium avium*.

It is well known that the intracellular lifestyle of S and R *M. abscessus* variants differ

significantly, resulting in entirely distinct infection scenarios (28). The absence of GPL on the outer

mycomembrane causes corded growth of R variants, resulting in multiple bacilli being

simultaneously phagocytosed by macrophages and overloaded phagosomes that rapidly activate

autophagy pathways (28). Comparatively, the S variant is able to survive for an extended period of

time within the phagosome, producing a chronic and persistent infection (29, 30). As such, these

polar infection responses may explain why the R displays increased granuloma formation at 2 wpi,
compared to S which shows a significantly delayed onset of granuloma formation. Moreover, this
observation matches the superior *in vivo* growth performance of S bacilli compared to R (Fig 1A),
suggesting that the R variant is at an overall disadvantage because of its intrinsic hyper-
inflammatory status and the activation of effective adaptive immunity that results in granuloma
formation. Taken together, our data provides additional evidence for the distinct intracellular fates
of both S and R variants *in vivo*, and further implicates the role of adaptive immunity in granuloma
formation and control of *M. abscessus* infection in an adult zebrafish model.

Our qPCR analysis surprisingly demonstrated that the S variant elicits greater production of
il1b; a key inflammatory cytokine, during the first 2 wpi compared to the R variant (Fig 1B). It is
likely that this observation is the nett effect of progressive increase in bacterial burden of S variants
over time, coupled with the presence of extracellular bacilli and unorganised granuloma formation
observed at 2 wpi, compared to the containment and clearance of the R variant in organised
granulomas.

T cells are critical host determinants in the control of mycobacterial infection (31).
Recruitment of T cells into granulomas are thought to be essential in containing persistent infection,
while T cell deficiencies are associated with greater mycobacterial infection severities (21, 31-33).
Recently, an adult zebrafish infection model for *M. leprae* demonstrated that T cells are essential
for containment of infection (30). We examined the recruitment of T cells within granulomas and
identified that S variant granulomas were marked by the absence of T cell infiltration at 2 wpi,
highlighting the fact that T cells may play a less significant role in S variant infections than those
with R variants. Using the $lck^{-/-}$ mutants, we have shown that adult zebrafish are highly susceptible
to *M. abscessus* infection and succumb to intraperitoneal infection within 40 days in the absence of
T cells, irrespective of bacterial morphotype. Importantly, the R variant displayed an improved *in*
vivo growth performance in the absence of T cells when compared to wild-type zebrafish,
highlighting the role of T cells in the control of R variants. However, this observation was not
maintained with the S variant, which showed no increase in bacterial growth *in vivo* irrespective of
the absence of T cells early in infection.

Our co-infection experiments further support the theory that tissue destruction caused by R
variant activates protective trans-acting host innate immunity that impairs bacterial growth, thereby
restricting S growth. This suggests *M. abscessus* must balance the benefits of R variant
pathogenicity allowing individuals to kill and escape macrophage containment, with the need to
avoid activation of host-protective immunity at a population level.

To date, our understanding of the diverse immune responses between S and R variants have
only been thoroughly described with respect to innate immunity, and currently our knowledge

pertaining to adaptive immunity in *M. abscessus* infection has been poorly characterised. Using this
new adult zebrafish *M. abscessus* infection model, we have shown that S and R variants produce
strikingly different disease phenotypes, which was further exemplified in the absence of a
functioning adaptive immune response. Consequently, these results suggest that the host-pathogen
interactions dictating *M. abscessus* pathogenesis are complex and may implicate adaptive immunity
to a greater extent than originally anticipated. Future work should exploit this new animal model in
combination with zebrafish lacking the Cystic fibrosis transmembrane conductance regulator gene,
and for the development and testing of novel antibiotics and vaccine candidates that may be used
for the treatment of *M. abscessus* infection.
287
Methods
<u>Zebrafish strains and handling</u>
Zebrafish strains used in this study are AB strain wildtype, $TgBAC(tnfa:GFP)^{pd1028}$,
$TgBAC(lck:EGFP)^{vcc4}$, $lck-/-^{sa410}$ (26, 27). Animals were held in a 28°C incubator with a 14:10 hour
light:dark cycle. Animals were infected by intraperitoneal injection with approximately 10^5 CFU *M.*
abscessus, unless otherwise stated, using a 31 G insulin needle and syringe as previously described
(34). Infected zebrafish were recovered into system water and held in 1 L beakers with daily
feeding for the duration of the experiment. Infection experiments were carried out with ethical
approval from the Sydney Local Health District Animal Welfare Committee approval 16-037.
297
<u>*M. abscessus* strains and handling</u>
Rough (R) and smooth (S) variants of *M. abscessus* strain CIP104536^T were grown at 30°C in
Middlebrook 7H9 broth supplemented with 10% Oleic acid/Albumin/Dextrose/Catalase (OADC)
enrichment and 0.05% Tween 80 or on Middlebrook 7H10 agar containing 10% OADC (7H10
OADC). Recombinant *M. abscessus* strains expressing Tdtomato or Wasabi were grown in the
presence of 500 µg/ml hygromycin (9, 19). Homogenous bacterial suspensions for intraperitoneal
injection in adult fish were prepared as previously reported (35).
305
306 <u>Bacterial recovery</u>
307 Animals were euthanised by tricaine anaesthetic overdose and rinsed in sterile water. Individual
308 carcasses were homogenised and serially diluted into sterile water. Homogenates were plated onto
309 7H10 OADC supplemented with 300 µg/ml hygromycin. Plates were grown for at least 4 days at 37
310 degrees.
311
<u>Zebrafish and *M. abscessus* gene expression analysis by qPCR</u>

RNA was extracted from whole fish homogenates and was reverse transcribed with the Applied

Biosystems High Capacity cDNA kit and qPCR was carried out on an Agilent Technologies

Stratagene Mx3005P. Zebrafish gene expression primers were previously described by Hammaren

et al. (25).

Histology

Animals subjected to cryosectioning as previously described (34). Briefly, euthanasia was

performed by tricaine anaesthetic overdose and specimens were fixed for 2-4 days in 10% neutral

buffered formalin at 4°. Specimens were then rinsed in PBS, incubated overnight in 30% sucrose,

incubated overnight in 50/50 30% sucrose and OCT, and finally incubated overnight in OCT prior

to freezing at -80°. Cryosectioning was performed to produce 20 µm thick sections. Sections were

post-fixed for 1-2 minutes in 10% neutral buffered formalin and rinsed in PBS prior to further

processing. Slides for fluorescent imaging were mounted with coverslips using Fluoromount G

containing DAPI. Oil Red O staining was performed as previously described (34, 36). T cells were

detected in sections from *TgBAC(lck:GFP)^{vcc4}* zebrafish by anti-GFP staining (primary antibody:

ab13970, Abcam; secondary antibody: ab150173, Abcam), stained slides were then mounted with

coverslips using Fluoromount G containing DAPI. All imaging was carried out on a Leica

DM6000B microscope.

Statistics

All statistical testing was carried out using Graphpad Prism. Each data point indicates a single

animal unless otherwise stated.

Acknowledgements

We thank the Centenary imaging facility core and Sydney Cytometry staff Drs Kristina Jahn,

Angela Kurz, and David Liu, for their assistance.

Funding: Australian National Health and Medical Research Council CJ Martin Early Career

Fellowship APP1053407 and Project Grant APP1099912; The University of Sydney Fellowship

G197581; NSW Ministry of Health under the NSW Health Early-Mid Career Fellowships Scheme

H18/31086; the Kenyon Family Foundation Inflammation Award; Australian-French Association

for Research and Innovation (AFRAN) Initiative; The University of Sydney Marie Bashir Institute

2019 Seed Funding to SHO. Sydney Medical School Summer Scholarship to JYK. Post-doctoral

fellowship granted by Labex EpiGenMed, an "Investissements d'avenir" program ANR-10-LABX-

12-01 to MDJ. The Fondation pour la Recherche Médicale DEQ20150331719 to LK.

References

1. H. Medjahed, J. L. Gaillard, J. M. Reyrat, Mycobacterium abscessus: a new player in the
mycobacterial field. *Trends Microbiol* **18**, 117-123 (2010).

2. J. F. Tomashefski, Jr., R. C. Stern, C. A. Demko, C. F. Doershuk, Nontuberculous
mycobacteria in cystic fibrosis. An autopsy study. *Am J Respir Crit Care Med* **154**, 523-528
(1996).

3. R. Nessar, E. Cambau, J. M. Reyrat, A. Murray, B. Gicquel, Mycobacterium abscessus: a new
antibiotic nightmare. *The Journal of antimicrobial chemotherapy* **67**, 810-818 (2012).

4. B. E. Ferro *et al.*, Failure of the Amikacin, Cefoxitin, and Clarithromycin Combination
Regimen for Treating Pulmonary Mycobacterium abscessus Infection. *Antimicrob Agents*
Chemother **60**, 6374-6376 (2016).

5. S. T. Howard *et al.*, Spontaneous reversion of Mycobacterium abscessus from a smooth to a
rough morphotype is associated with reduced expression of glycopeptidolipid and
reacquisition of an invasive phenotype. *Microbiology* **152**, 1581-1590 (2006).

6. E. Catherinot *et al.*, Acute respiratory failure involving an R variant of Mycobacterium
abscessus. *J Clin Microbiol* **47**, 271-274 (2009).

7. C. R. Esther, Jr., D. A. Esserman, P. Gilligan, A. Kerr, P. G. Noone, Chronic Mycobacterium
abscessus infection and lung function decline in cystic fibrosis. *J Cyst Fibros* **9**, 117-123
(2010).

8. B. E. Jonsson *et al.*, Molecular epidemiology of Mycobacterium abscessus, with focus on
cystic fibrosis. *J Clin Microbiol* **45**, 1497-1504 (2007).

9. A. Bernut *et al.*, Mycobacterium abscessus cording prevents phagocytosis and promotes
abscess formation. *Proc Natl Acad Sci U S A* **111**, E943-952 (2014).

10. H. Medjahed, J. M. Reyrat, Construction of Mycobacterium abscessus defined
glycopeptidolipid mutants: comparison of genetic tools. *Appl Environ Microbiol* **75**, 1331-
1338 (2009).

11. A. Bernut *et al.*, In vivo assessment of drug efficacy against Mycobacterium abscessus using
the embryonic zebrafish test system. *Antimicrob Agents Chemother* **58**, 4054-4063 (2014).

12. A. Obregon-Henao *et al.*, Susceptibility of Mycobacterium abscessus to antimycobacterial
drugs in preclinical models. *Antimicrob Agents Chemother* **59**, 6904-6912 (2015).

13. M. Rottman *et al.*, Importance of T cells, gamma interferon, and tumor necrosis factor in
immune control of the rapid grower Mycobacterium abscessus in C57BL/6 mice. *Infect*
Immun **75**, 5898-5907 (2007).

14. C. T. Oh, C. Moon, M. S. Jeong, S. H. Kwon, J. Jang, Drosophila melanogaster model for
Mycobacterium abscessus infection. *Microbes Infect* **15**, 788-795 (2013).

15. M. Meir, T. Grosfeld, D. Barkan, Establishment and Validation of Galleria mellonella as a
Novel Model Organism To Study Mycobacterium abscessus Infection, Pathogenesis, and
Treatment. *Antimicrob Agents Chemother* **62** (2018).

16. A. Bernut, J. L. Herrmann, D. Ordway, L. Kremer, The Diverse Cellular and Animal Models
to Decipher the Physiopathological Traits of Mycobacterium abscessus Infection. *Front Cell*
Infect Microbiol **7**, 100 (2017).

17. V. Dubois *et al.*, MmpL8MAB controls Mycobacterium abscessus virulence and production
of a previously unknown glycolipid family. *Proc Natl Acad Sci U S A* **115**, E10147-E10156
(2018).

18. I. Halloum *et al.*, Deletion of a dehydratase important for intracellular growth and cording
renders rough Mycobacterium abscessus avirulent. *Proc Natl Acad Sci U S A* **113**, E4228-
4237 (2016).

19. A. Bernut *et al.*, Mycobacterium abscessus-Induced Granuloma Formation Is Strictly
Dependent on TNF Signaling and Neutrophil Trafficking. *PLoS Pathog* **12**, e1005986 (2016).

20. A. Bernut *et al.*, CFTR Protects against Mycobacterium abscessus Infection by Fine-Tuning
Host Oxidative Defenses. *Cell reports* **26**, 1828-1840 e1824 (2019).

21. C. A. Madigan, J. Cameron, L. Ramakrishnan, A Zebrafish Model of Mycobacterium leprae
Granulomatous Infection. *J Infect Dis* **216**, 776-779 (2017).

22. S. H. Oehlers *et al.*, Interception of host angiogenic signalling limits mycobacterial growth.
Nature **517**, 612-615 (2015).

23. L. E. Swaim *et al.*, Mycobacterium marinum infection of adult zebrafish causes caseating
granulomatous tuberculosis and is moderated by adaptive immunity. *Infect Immun* **74**, 6108-
6117 (2006).

24. A. L. Roux *et al.*, The distinct fate of smooth and rough Mycobacterium abscessus variants
inside macrophages. *Open Biol* **6** (2016).

25. M. M. Hammaren *et al.*, Adequate th2-type response associates with restricted bacterial
growth in latent mycobacterial infection of zebrafish. *PLoS Pathog* **10**, e1004190 (2014).

26. L. Marjoram *et al.*, Epigenetic control of intestinal barrier function and inflammation in
zebrafish. *Proc Natl Acad Sci U S A* **112**, 2770-2775 (2015).

27. K. Sugimoto, S. P. Hui, D. Z. Sheng, M. Nakayama, K. Kikuchi, Zebrafish FOXP3 is required
for the maintenance of immune tolerance. *Dev Comp Immunol* **73**, 156-162 (2017).

28. A.-L. Roux *et al.*, The distinct fate of smooth and rough Mycobacterium abscessus variants
inside macrophages. *Open biology* **6** (2016).

29. A. Bernut *et al.*, *Mycobacterium abscessus* cording prevents phagocytosis and promotes
abscess formation. *Proceedings of the National Academy of Sciences* **111**, 943-952 (2014).

30. L. Ramakrishnan, Revisiting the role of the granuloma in tuberculosis. *Nature Reviews*
Immunology **12**, 352-366 (2012).

31. F. M. Collins, Mycobacterial disease, immunosuppression, and acquired immunodeficiency
syndrome. *Clin Microbiol Rev* **2**, 360-377 (1989).

32. T. Mogues, M. E. Goodrich, L. Ryan, R. LaCourse, R. J. North, The relative importance of T
cell subsets in immunity and immunopathology of airborne Mycobacterium tuberculosis
infection in mice. *J Exp Med* **193**, 271-280 (2001).

33. J. D. Yang *et al.*, Mycobacterium tuberculosis-specific CD4+ and CD8+ T cells differ in their
capacity to recognize infected macrophages. *PLoS Pathog* **14**, e1007060 (2018).

34. T. Cheng, J. Y. Kam, M. D. Johansen, S. H. Oehlers, High content analysis of granuloma
histology and neutrophilic inflammation in adult zebrafish infected with Mycobacterium
marinum. *Micron* 10.1016/j.micron.2019.102782 (2019).

35. A. Bernut *et al.*, Deciphering and Imaging Pathogenesis and Cording of Mycobacterium
abscessus in Zebrafish Embryos. *J Vis Exp* 10.3791/53130 (2015).

36. M. D. Johansen *et al.*, Mycobacterium marinum infection drives foam cell differentiation in
zebrafish infection models. *Dev Comp Immunol* **88**, 169-172 (2018).

Figure Legends

Figure 1: *M. abscessus* establishes chronic infection in adult zebrafish.

A. Enumeration of CFUs from adult zebrafish infected with either the R or the S variant of *M.*
abscessus.

B. Gene expression analysis of zebrafish *interleukin 1b*.

C. Gene expression analysis of zebrafish *tumor necrosis factor*.

D. Examples of R variant *M. abscessus*-Tdtomato lesions in 2 wpi *TgBAC(tnfa:GFP)^{pd1028}* adult
zebrafish.

E. Example of S variant *M. abscessus*-Tdtomato lesions in 4 wpi *TgBAC(tnfa:GFP)^{pd1028}* adult
zebrafish.

Filled arrowheads indicate organised necrotic granulomas with strong *tnfa* expression marked by
GFP, empty arrowheads indicate loose *M. abscessus* lesions with elevated *tnfa* promoter induction.
Scale bars indicate 200 μm.

Statistical tests by one-way ANOVA with Tukey's multiple comparisons test. Data is pooled from
10 animals across 2 independent experiments.

Figure 2: *M. abscessus* R infection causes progressive granulomatous pathology.

A. Stereotypical unorganised granuloma in a 2 wpi adult zebrafish infected with the R variant of *M. abscessus* expressing Tdtomato. Scale bar indicates 100 µm.

B. i. Stereotypical organised granuloma in a 2 wpi adult zebrafish infected with R *M. abscessus*-Tdtomato and ii. Higher magnification image of granuloma wall. Arrowheads indicate epithelised macrophage nuclei surrounding the mycobacterial core. Scale bar indicates 100 µm.

C. Example of a very large abscess-like granuloma in a 2 wpi adult zebrafish infected with R *M. abscessus*-Tdtomato measuring approximately 600 µm in diameter. Scale bar indicates 200 µm.

D. Example of Oil Red O-stained very large abscess-like granuloma in a 2 wpi adult zebrafish infected with R *M. abscessus*. Arrowheads indicate Oil Red O-positive foamy macrophages surrounding the mycobacterial core. Scale bar indicates 200 µm.

E. Quantification of granuloma organisation in adult zebrafish infected with R *M. abscessus*. Data is pooled from 2 animals per timepoint.

F. Quantification of bacterial burden stratified by granuloma organisation in adult zebrafish infected with R *M. abscessus*. Data is pooled from 2 animals per timepoint, statistical testing by Chi-squared test.

Figure 3: *M. abscessus* S infection causes delayed progressive granulomatous pathology.

A. Examples of S *M. abscessus*-Tdtomato lesions from 2 wpi adult zebrafish. i. *M. abscessus* growing free, external to peritoneal organs. ii. Stereotypical example of unorganised cellular granulomas around sites of sparse *M. abscessus* infection.

B. i. Stereotypical large granuloma found in 4 wpi adult zebrafish infected with S *M. abscessus*-Tdtomato. Note lack of punctate bacterial fluorescence in the core of the granuloma compared to loose bacteria to the right. ii. Example of Oil Red O-stained large granuloma in 4 wpi adult zebrafish infected with S *M. abscessus*. Note lack of lipid staining in the rim of the granuloma compared to R infection.

C. i. Quantification of granuloma organisation in adult zebrafish infected with S *M. abscessus*. ii. Quantification of bacterial burden stratified by granuloma organisation in adult zebrafish infected with S *M. abscessus*. Data is pooled from at least 2 animals per timepoint, statistical testing by Chi-squared test.

Figure 4: T cells are necessary to control R but not S *M. abscessus* infection.

A. Examples of T cell recruitment to granulomas in 2 wpi *TgBAC*(*lck*:*EGFP*)^{vcc4} adult zebrafish infected with R *M. abscessus*-Tdtomato . i. Example of an unorganised granuloma. ii. Example of an organised granuloma. iii. Example of a very large abscess-like granuloma. Scale bars indicate 100 µm.

B. Example of lack of T cell recruitment to S *M. abscessus*-Tdtomato in 2 wpi

TgBAC(*lck*:*EGFP*)^{vcc4} adult zebrafish. Scale bar indicates 100 µm.

C. Survival analysis of WT and lck-/-^{sa410} adult zebrafish infected with R or S *M. abscessus*.

D. Enumeration of CFUs from 2 wpi WT and lck-/-^{sa410} adult zebrafish infected with R *M.*

abscessus. Statistical testing by T-test.

E. Enumeration of CFUs from 2 wpi WT and lck-/-^{sa410} adult zebrafish infected with S *M.*

abscessus. Statistical testing by T-test.

Figure 5: S *M. abscessus* has an *in vivo* survival advantage over R *M. abscessus* during chronic

infection.

A. Schema outlining mixed infection experiment.

B. Ratio of R:S CFUs recovered from WT adult zebrafish infected with a mixture of differentially

labelled R and S variant *M. abscessus*.

C. Enumeration of CFUs from the three groups of WT adult zebrafish outlined in panel A. Data is

pooled from two replicates, statistical testing by ANOVA.

D. Representative images of granuloma from adult zebrafish infected with a mixture of R

expressing Wasabi and S expressing Tdtomato *M. abscessus*.

E. Enumeration of CFUs from the three groups of lck-/-^{sa410} adult zebrafish outlined in panel A.

Data is pooled from two replicates, statistical testing by ANOVA.

F. Ratio of R:S CFUs recovered from lck-/-^{sa410} adult zebrafish infected with a mixture of

differentially labelled R and S variants.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Graphical Abstract