Trophic cascade driven by behavioural fine-tuning as naïve prey rapidly

adjust to a novel predator

CHRIS J. JOLLY*[1], ADAM S. SMART[1], JOHN MOREEN[2], JONATHAN K. WEBB[3], GRAEME R.

GILLESPIE[1,4] AND BEN L. PHILLIPS[1]

[1]School of BioSciences, University of Melbourne, Parkville Vic 3010 Australia

[2]Kenbi Rangers, Mandorah NT 0822 Australia

[3]School of Life Sciences, University of Technology Sydney, Broadway NSW 2007 Australia

[4]Flora and Fauna Division, Department of Environment and Natural Resources, Northern Territory

Government, Berrimah NT 0828 Australia

*Correspondence:

C. Jolly

Email: cjolly1@student.unimelb.edu.au

November 2019

Keywords: Behavioural fine-tuning, novel predator, predator-prey dynamics, trophic

cascade

The arrival of novel predators can trigger trophic cascades driven by shifts in prey

numbers. Predators also elicit behavioural change in prey populations, via plasticity and

rapid evolution, and such behavioural responses by prey may also contribute to trophic

cascades. Here we document the effects of a novel predator on the behaviour and

demography of a native prey species. We reveal rapid behavioural responses in the prey

species (grassland melomys, a granivorous rodent) following the introduction of a novel

marsupial predator (northern quoll). Within months of quolls appearing, populations of

melomys exhibited reduced survival and population declines relative to control

populations. Quoll-exposed populations were also significantly shyer than nearby,

predator-free populations of conspecifics. This rapid but generalised response to a novel

threat was replaced over the following two years with more threat-specific antipredator

behaviours (i.e. predator-scent aversion). Predator-exposed populations, however,

remained more neophobic than predator-free populations throughout the study. These

behavioural responses manifested rapidly in changed rates of seed predation by melomys

across treatments. Quoll-exposed melomys populations exhibited lower per-capita seed

take rates, and rapidly developed an avoidance of seeds associated with quoll scent, with

discrimination playing out over a spatial scale of tens of metres. Presumably the

significant and novel predation pressure induced by the introduction of quolls drove

melomys to fine-tune their behavioural response through time as the nature of the threat

became clearer. These behavioural shifts could reflect individual plasticity in behaviour or

may be adaptive shifts from natural selection imposed by quoll predation. Our study

provides a rare insight into the rapid ecological and behavioural shifts enacted by prey to

mitigate the impacts of a novel predator and shows that trophic cascades can be strongly

influenced by behavioural as well as numerical responses.

INTRODUCTION

Predation is one of the most pervasive and powerful forces acting on populations. Not only does predation directly impact a population's demography (Schoener & Spiller 1996), it also imposes natural selection (Abrams 2000). Although pervasive, predation is not a constant. The pressure that predators impose on populations will vary through time and space (Lima & Dill 1990; Sih 1992) as a consequence of tightly coupled predator-prey dynamics; prey switching; or stochastic processes. The fact that predation is not constant, and that antipredator defences may be costly, suggests that flexible, plastic responses to predation pressure will often be favoured (Sih *et al.* 2000; Berger *et al.* 2001). There is, in fact, a great deal of empirical evidence that plastic responses to predation are common (e.g. Relyea 2003; Brown *et al.* 2013; Cunningham *et al.* 2019): investment in antipredator traits across morphology, life-history, and behaviour is often flexible, and dependent on the perceived risk of predation.

As well as impacting prey populations, it is increasingly apparent that predators play a powerful role in structuring communities (Estes *et al.* 2011). Some of our best evidence for this comes from the introduction of predators to naïve communities. Invasive predators can cause extinctions (Medina *et al.* 2011; Woinarski *et al.* 2015; Doherty *et al.* 2016), and alter trophic structures and ecosystem function within recipient communities (Courchamp *et al.* 2003; Simberloff *et al.* 2013). We typically think of these cascading outcomes as purely numeric effects: predators depress the size of prey populations, and the altered numbers of prey can cause cascading numerical changes down trophic levels (e.g. Ripple *et al.* 2001). These numerical effects are undeniably important, but the fact that predators can also elicit phenotypic change in prey populations – through plasticity and natural selection – means that subtler ecological effects may also manifest. Prey species living in a landscape of fear

may forage at different times, or in different places compared with the same species in a

predator-free environment (Laundre *et al.* 2010). Such behavioural shifts will alter all

downstream species interactions in potentially complex ways (e.g. Fortin *et al.* 2005; Suraci

et al. 2016).

Because predator invasions are rarely intentional or anticipated, there is a scarcity of

controlled empirical work on the effects of novel predators on recipient communities and

the mechanisms via which these effects play out (but see Lapiedra *et al.* 2018; Pringle *et al.*

2019). Such tests are needed, however, if we are to predict invasive species impacts,

advance conservation management (Sih *et al.* 2010), and improve our understanding of how

communities are structured (Sax *et al.* 2007). Here, we exploit a conservation-motivated

predator introduction, and show that (as well as numerical effects) rapid shifts in prey

behaviour occurred, and these behavioural shifts caused measurable change in downstream

trophic interactions.

Northern quolls (*Dasyurus hallucatus*) were, until recently, a common predator

across northern Australia. They have declined over the last several decades, following the

general decline in northern Australian mammals (Woinarski *et al.* 2015), thought to be

driven by changes in grazing, fire, and predation regimes (Braithwaite & Griffiths 1994).

More recently, the invasion of toxic invasive prey (cane toads, *Rhinella marina*) has resulted

in dramatic, range-wide population declines in northern quolls (Shine 2010; Oakwood *et al.*

2016). Due to local extinction, northern quolls are now absent from large tracts of their

former range and their ecological function as a medium-sized mammalian predator has

been lost (Moore *et al.* 2019). For their conservation, northern quolls have recently been

introduced to a number of offshore islands where they have never previously existed. We

exploit the introduction of northern quolls to a new island to directly test the effects of

quolls as a novel predator on an island ecosystem and observe how native prey adjust to

mitigate the impacts of their arrival.

In 2017, a population of 55 "toad-smart" northern quolls were introduced to a

25km^2 island off the coast of north-western Northern Territory, Australia (Kelly 2019). Prior

to this introduction, Indian Island (Kabal) lacked mammalian predators, and large native

reptilian predators had recently been reduced to near extinction by the invasion of cane

toads. The introduction of northern quolls provided a unique opportunity to test the effects

of the arrival of a novel predator in a system that has been isolated from major mammalian

predators for thousands of years.

METHODS

Introduction of northern quolls

In May 2017, 55 adult northern quolls were introduced to the north-eastern tip of Indian

Island, Bynoe Harbor, Northern Territory, Australia (12°37'24.60"S, 130°30'0.72"E) to field

test the conservation strategy of targeted gene flow (Kelly & Phillips 2016). Quolls are a

voracious, opportunistic generalist predator (< 1.5kg; Oakwood 1997), and their

introduction presented a unique opportunity to monitor the behavioural and demographic

impacts on grassland melomys (*Melomys burtoni*), a native mammalian granivorous prey

species (mean body mass 56 g, 5.6–103.7g). Immediately prior to the introduction of quolls,

we started monitoring populations of melomys in one woodland and two monsoon vine

thicket plots in the vicinity of where quolls were to be released and radio tracked. After

quolls were introduced and tracked it became immediately apparent that quolls were

largely avoiding monsoon vine thicket sites and these sites were dropped from the on-going

monitoring. For this reason, we only present data from before the introduction of quolls

from one invaded site. Most of our data compare quoll-invaded versus quoll-free sites over

time, commencing within a few months of quoll arrival.

Melomys population monitoring

To determine whether the arrival of a novel predator resulted in demographic impacts

(population size and survival) to native prey species, we monitored populations of melomys

on Indian Island, during four trips occurring in May (site 1) and August 2017 (sites 2–7), April

2018 (sites 1–7), and May 2019 (sites 1–7). Melomys were caught from seven independent

1ha (100 m x 100 m) plots (sites 1–7) spread out across Indian Island using a standard mark-

recapture trapping regime designed for a monitoring project (Begg *et al.* 1983; Kemper *et al.*

1987). Four "impact", quoll-invaded sites were established in the north of the island in the

vicinity of where quolls were released and three "control", quoll-free sites were established

in the south of the island. Sites in the north (quoll-invaded) and south (quoll-free) of the

island were between 8.7 and 9.8km apart (Table 1) and were composed of similar habitat

types. The northern and southern sections of Indian Island are divided by mangrove habitat

which is inundated at high tide. We assumed quolls would be confined to the northern half

of the island for at least a few years. Cage and camera trapping as well as track surveys

confirmed that quolls were present at the "impact" sites and absent from the "control" sites

for the duration of the study (Jolly *et al.* unpub. data).

145 **Table 1.** Pairwise distance matrix between sites on Indian Island, Northern Territory, Australia. Quolls were

146 present at sites 1–4 and quolls were absent at sites 5–7 for the duration of the study.

Distance (m)	Site 1	Site 2	Site 3	Site 4	Site 5	Site 6	Site 7
Site 1							
Site 2	270						
Site 3	260	350					
Site 4	400	300	250				
Site 5	8760	9030	8710	9000			
Site 6	8470	8730	8450	8070	300		
Site 7	9670	9920	9590	9820	1260	1500	

Each of the seven monitoring sites consisted of 100 Elliott traps (Elliott Scientific

Equipment, Upwey, Victoria) spaced at 10 m intervals in a 10 x 10 grid. Most trapping grids

were open for four nights, however, the first trapping grid (site 1, May 2017) was open for

six nights. After four trap nights, the majority of the melomys population had been captured

at least once (Jolly *et al.* 2019). Traps were baited with peanut butter and rolled oats. These

baits were replaced daily for the duration of each trapping session. Traps were checked for

captures early each morning and all traps were cleared within two hours of sunrise.

Captured melomys were weighed (g) and sexed. Before release, each melomys was

implanted with a microchip (Trovan Unique ID100). On successive mornings, all melomys

were scanned (Trovan LID575 Handheld Reader), and any new individuals were

microchipped. On the last morning of each trapping session, all melomys caught were

retained for behavioural assays. Throughout the study 435 individual melomys were

captured and given microchips. Of these, 191 (44%) were caught on the final night of

trapping and were retained for behavioural trials. Only large, healthy juveniles (*n* = 12),

adult males (*n* = 83), and adult non-visibly pregnant females (*n* = 96) were retained for

behavioural experiments. Indian Island is remote and uninhabited by humans, so all

behavioural experiments were conducted in the field under near natural conditions (see

Jolly *et al.* 2019 for detailed experimental procedures).

Modified open field tests

We employed modified open field tests (also referred to as emergence tests: see Brown &

Braithwaite 2004; López *et al.* 2005; Carter *et al.* 2013; Jolly *et al.* 2019) to assess boldness

in grassland melomys and whether the arrival of a novel predator resulted in behavioural

shifts in invaded populations. All open field tests were conducted on the night after the last

trap night (night 5) and in opaque-walled experimental arenas (540mm x 340mm x 370mm).

Experimental arenas were modified plastic boxes that had an inverted Elliott trap sized hole

cut in one end and were illuminated by strings of red LED lights (Jolly *et al.* 2019). Each

experimental arena had natural sand as substrate, and a bait ball located both in the centre

and along one wall of the arena (Jolly *et al.* 2019). After dark, Elliott traps containing a

melomys were inserted into the hole in the side of each experimental arena and melomys

were allowed to habituate for 10 min. At the start of each trial, Elliott trap doors were

locked open – the inverted orientation of the trap prevented them from being triggered

closed. Melomys were given 10 min to explore the open field arena. After 10 min,

individuals were rounded back into their retreat (the Elliott trap) and a novel object (plastic

bowl) was placed at the end of the arena opposite the Elliott trap (Jolly *et al.* 2019).

Melomys were then given a further 10 min to explore the arena and interact with the novel

object. Elliott traps remained open during the open field tests and melomys could shelter

and emerge from them under their own volition. All trials were recorded using a GoPro

HERO 3. A previous study in this system determined that melomys showed repeatable

behaviour between trials (Jolly *et al.* 2019), therefore the data presented in this study were

from a single behavioural trial of each animal (n = 191). Once trials were complete, each

melomys was released at its point of capture.

To measure the boldness of individual melomys, we scored three behaviours

typically associated with boldness and neophobia in rodents (Dielenberg & McGregor 2001;

McGregor *et al.* 2002; Réale *et al.* 2007; Cremona *et al.* 2015): whether melomys fully

emerged from their Elliott trap hide and entered the open arena during the 0–10 min period

(scored 0 or 1, respectively); whether they fully emerged and entered the trial arena during

the 10–20 min period (scored 0 or 1); and whether they interacted (made contact) with the

novel object that was placed in the arena during the 10–20 min period (scored 0 or 1).

Videos were scored by a single observer who was blind to each melomys' origin and

identity. Because interacting with the novel object was predicated on a melomys'

willingness to emerge from their hide during the 10–20 min period, for analysis we

combined their emergence during this period and interaction with the novel object into a

single binary score: 0 (neophobic) = did not emerge or emerged but did not interact with

novel object; or 1 (not neophobic): emerged and interacted with novel object.

Seed removal plots

To assess whether the arrival of a novel predator affected the seed harvesting behaviour of

granivorous melomys, we established seed removal plots at each site and sampled them

each trapping session (night 6). After trapping and open field tests were conducted and

melomys had been returned to their capture location, we set up 81 seed plots at each site

by scraping away leaf litter with a shovel to create bare earth plots. These bare earth plots

were created so that they were located in the centre between four Elliott traps within the

10x10 trapping grid. Just before dark on the night of the seed removal experiment, we

placed a single wheat seed in the centre of each bare earth plot. These seeds were either

unscented, control seeds (n = 40) or predator-scented seeds that had been maintained in a

sealed clip-lock bag filled with freshly collected northern quoll fur (n = 41). The placement of

predator-scented and unscented seeds was alternated so that there was a chequered

arrangement of scented and unscented seeds across the site. To ensure that the predator-

scent was strong enough to be detected by melomys, along with the predator-scented

seeds, we also placed a few strands of quoll fur around the predator-scented seeds. Before

light the next morning, we returned back to each plot and counted the number of seeds of

each scent-type that were removed from the plot. Melomys are the only nocturnal

granivorous animal that occurs on Indian Island, and to avoid diurnal granivorous birds from

removing seeds we conducted this experiment during the night only.

Wildfire on northern Indian Island

Immediately following our monitoring and experiments in August 2017, a wildlife broke out

on northern Indian Island in the vicinity of the four quoll-invaded sites and burnt through all

of the sites. Because of this, our experimental design is confounded by the fact that all of

our quoll-invaded sites were burnt, and all of our quoll-free sites were unburnt. Fire is a

regular disturbance in this landscape (Andersen *et al.* 2005), and previous work has shown

little effect of fire on abundance, survival or recruitment of grassland melomys (Griffiths &

Brook 2015; Liedloff *et al.* 2018). Nonetheless, this confound exists and we proceed with

caution when interpreting the effects of quolls on population size and survival of melomys.

Statistical analysis

During trapping sessions we identified individual melomys that were captured at each site

by their unique microchips. Because melomys on Indian Island have very small home ranges

(tending to be caught in the same or adjacent traps throughout the trapping period: Jolly *et*

al. unpub. data) and since we never observed captures of melomys marked at other sites

(Jolly *et al.* unpub. data), we treated each site as independent with regard to demographics

and behaviour (Table 1).

To estimate between-session survival, we analysed the mark-recapture data to

estimate recapture and survival rates using Cormack-Jolly-Seber models in program MARK.

At each site, there were three primary trapping sessions of four nights, for a total of 12 time

intervals in the input file. Because quolls prey on melomys, we hypothesised that survival

rates of melomys would be lower between trapping sessions at sites with quolls than at

sites without quolls. We included two groups, quoll-free (control) and quoll-invaded

(impact), in the input file. We ran a series of models in MARK to test the following *a priori*

hypotheses: (1) survival rates between sessions are lower at quoll-free sites than at quoll-

invaded sites; (2) survival rates are lower between sessions than within sessions, but are

unaffected by quolls; (3) survival is constant through time; and (4) survival varies through

time. All candidate models were ranked according to their AICc values and associated AIC

weights (Burnham & Anderson 1998).

To test whether the presence of quolls impacted melomys population size, we used

a hierarchical model in which population size was made a function of quoll

presence/absence, capture session, and the interaction between these factors. Population

size at each site during each session is estimated in this process, and we fitted this model in

a Bayesian framework. Our observations consisted of a capture history for each observed

individual over the number of nights at each site for each trapping session. We denoted the

number of individuals at site s during session k as N_{ks}. To estimate N_{ks} we used a closed

population mark-recapture analysis in which each individual, I, was either observed, or not,

according to a Bernoulli distribution:

262   O_{iks} ~ Bernoulli(d_s)

263   Our previous MARK analysis found clear evidence for variation in detection probability

264   across sessions, but detection probabilities of melomys on Indian Island had previously been

265   found not to vary measurably between individuals nor to change over time within a trapping

266   session (Jolly *et al.* 2019). Thus, we made detection probability a function of session

267   according to:

268   logit(d_s) = μ_d + t_s

Where d_s denotes the expected detection probability within session s, μ_d is the expected

detection probability in the first session, and t_s denotes the (categorical) effect of session on

detection.

We used the "data augmentation" method (Tanner & Wong 1987; Royle *et al.* 2007;

Kery & Schaub 2011) in combination with this detection probability to estimate N_{ks} for each

site.session. Using this approach, the data were 'padded' by adding an arbitrary number of

zero-only encounter histories of 'potential' unobserved individuals. The augmented dataset

was modelled as a zero-inflated model (Royle *et al.* 2007) which changes the problem from

estimating a count, to estimating a proportion. This was executed by adding a latent binary

indicator variable, R_{iks}, to classify each row in the augmented data matrix as a 'real'

individual or not, where R_{iks}~ Bernoulli(Ω_{ks}). The parameter Ω_{ks} was estimated from the data,

and $N_{ks} = \Sigma_i R_{iks}$.

We then made Ω_{ks} (which scales with population size) a function of quoll

presence/absence, q_c; session, b_k; and the interaction between the two:

$$\mathrm{logit}(\Omega_{ks}) = \mu_p + q_c + b_k + \gamma_{ck}$$

The model was fitted using Bayesian Markov Chain Monte Carlo (MCMC) methods

and minimally informative priors (Table 2) within the package JAGS (Plummer *et al.* 2017)

using R (R Core Team 2019). Parameter estimates were based on 30,000 iterations with a

thinning interval of 5 following a 10,000 sample burn-in. Three MCMC chains were run, and

model convergence assessed by eye, and using the Gelman-Rubin diagnostic (Gelman &

Rubin 1992a, 1992b).

To assess whether the introduction of quolls affected the behaviour of melomys

populations, we divided the responses of melomys in open field tests into two independent

response variables: the proportion that emerged during the 0-10 min period; and the

proportion that emerged and interacted with the novel object during the 10-20 min period.

We used generalised linear mixed-effects models with binomial errors and a logit link to test

the effect of quoll presence and trapping session, with site included as a random effect, on

the behavioural response variables. *P*-values were obtained by likelihood ratio tests of the

full model with the effect in question against the model without the effect. This analysis was

performed using R with the *lme4* software package (R Core Team 2019).

To assess whether the numerical impact of quolls on melomys affected the seed

harvesting rate of invaded melomys populations, we first examined the relationship

between melomys population size (estimated above) and the total number of seeds

harvested in the control (unscented) seeds. Here we used a simple linear model with

number of seeds harvested as a function of population size, quoll presence/absence and the

interaction between these effects. To test whether there was an additional effect of quoll

presence, beyond their effect on population size, we defined a new variable, Δ, as the

difference in seed take between scented and unscented treatments within each site.session.

Here any effect of melomys density is cancelled out (because density is common to both

treatments within each site.session). Thus, we fitted a model in which Δ is a function of

quoll presence/absence, session and the interaction between these effects. These analyses

were performed using R (R Core Team 2019).

RESULTS

Effect of novel predator on survival

When we assessed the impact of quolls on melomys survival between trapping sessions the

best supported model was one in which survival rates between sessions were lower at

quoll-invaded sites than at quoll-free sites, and recapture rates were session-dependent

(Table 2). All other models were more than 4 AIC units from this best model, and so clearly

inferior descriptions of the data. From the best-supported model, estimates of apparent

survival for the intervals between the capture sessions were substantially higher at quoll-

free sites ($S_{2017–2018}$ = 0.368; $S_{2018–2019}$ = 0.225) than at quoll-invaded sites ($S_{2017–2018}$ = 0.207;

$S_{2018–2019}$ = 0.091; Fig. 1). The differing survival probability between sessions is largely

explained by the time difference between intervals (2017–2018 = 9 months vs. 2018–2019

= 13 months; Fig. 2).

Figure 1. Between trapping session survival (± s.e.) of grassland melomys (*Melomys burtoni*) on Indian Island

in quoll-invaded (*n* = 4) and quoll-free (*n* = 3) populations on Indian Island, Northern Territory, Australia.

Table 2. Results of Cormack-Jolly-Seber analyses used to compare survival (*Phi*) and recapture (*p*) probabilities

of grassland melomys (*Melomys burtoni*) on Indian Island, Northern Territory, Australia. The symbols '.' and '*t*'

refer to constant and time, respectively. Table shows AIC values and associated AIC weights, model likelihood,

number of parameters (*N*), and model deviance. The term 'w/b' indicates that within trapping session survival

rates (s_1-s_3, s_5-s_7, s_9-s_{11}) were constant and equivalent, and different to the between trapping session survival

rates (s_4, s_8). The term 'group w/b' is as above, except that between trapping session survival rates differed

between the two groups.

Model	AICc	Delta AICc	AICc Weights	Model Likelihood	N	Deviance
Phi (group w/b) p(t)	1688.629	0	0.92162	1	16	477.5224
Phi (w/b) p(t)	1693.562	4.933	0.07823	0.0849	14	486.6202
Phi (t) p (t)	1706.168	17.5394	0.00014	0.0002	21	484.553
Phi (g*t) p(t)	1712.896	24.2675	0	0	32	467.664
Phi (t) p (g*t)	1718.863	30.2344	0	0	32	473.631
Phi (t) p (.)	1719.862	31.2339	0	0	12	517.0641
Phi (t) p (g)	1720.751	32.1229	0	0	13	515.8843

     Effect of novel predator on population size

Populations of melomys declined dramatically in quoll-invaded sites in the year following

their introduction but not in quoll-free sites (Fig. 2). A strong negative interaction between

the presence of quolls and trapping session in 2018 (mean = -1.194, 95% credible interval [-

1.732, -0.665]) and 2019 (mean = -1.097, 95% confidence interval [-1.652, -0.551]) suggested

that the presence of quolls dramatically impacted these populations of melomys (Fig. 2;

Table 3).

Figure 2. Posterior mean population sizes ($N_{ks} \pm$ 95% CI) for quoll-invaded and quoll-free populations of

grassland melomys (*Melomys burtoni*) on Indian Island, Northern Territory, Australia. The orange dotted

vertical line denotes the timing of the introduction of quolls. The red dotted vertical line denotes the timing of

an unplanned fire that burnt through the quoll-invaded sites. Estimates assume closure of the population

within each session and detection probability that varies across sessions.

Table 3. Model parameters and their priors including prior distributions, standard deviation, estimated

posterior means and their 95% credible intervals. N denotes normal probability distribution with mean and

standard deviation.

Model Parameters				
Name for parameter	**Parameter**	**Prior (mean, SD)**	**Posterior mean**	**95% credible intervals**
Detection:				
Intercept for detection	μ_d	N (0, 2.71)	-0.938	-1.118, -0.760
Effect of session 2 on detection	t_2	N (0, 2.71)	0.592	0.332, 0.852
Effect of session 3 on detection	t_3	N (0, 2.71)	0.458	0.179, 0.730
Population size:				
Intercept for Omega	μ_P	N (0, 2.71)	-0.906	-1.222, -0.583
Quoll Presence	r_2	N (0, 2.71)	0.698	0.314, 1.093
Trapping Session 2	b_2	N (0, 2.71)	0.061	-0.360, 0.478
Trapping Session 3	b_3	N (0, 2.71)	-0.100	-0.527, 0.339
Interaction 1 [Quoll Presence * Trapping Session 2]	$\gamma_{2,2}$	N (0, 2.71)	-1.194	-1.732, -0.665
Interaction 2 [Quoll Presence * Trapping Session 3]	$\gamma_{2,3}$	N (0, 2.71)	-1.097	-1.652, -0.551

Effects of novel predator on prey behaviour

For the proportion of melomys emerging in open field tests during the 0–10 min period,

there was a significant interaction between quoll presence and trapping session (χ^2 (5) =

4.386, *P* = 0.04; Fig. 3). There was no interaction between quoll presence and trapping

session for the proportion of melomys emerging and interacting with the novel object

during 10–20 min period (χ^2 (5) = 2.567, *P* = 0.109; Fig. 3). The model without this

interaction, however, revealed a significant effect of quoll presence on the neophobic

behaviour of melomys in open field tests, with fewer melomys emerging from hiding and

interacting with the novel object during the 10–20 min period of open field tests from sites

where quolls were present than from sites where quolls were absent (χ^2 (5) = -4.696, P <

0.001; Fig. 3).

Figure 3. Mean proportion (± s.e.) of grassland melomys (*Melomys burtoni*) emerging from hiding during open

field tests from quoll-invaded sites in 2017 (n = 16), 2018 (n = 28) and 2019 (n = 29), and quoll-free sites in

2017 (n = 14), 2018 (n = 35) and 2019 (n = 24) on Indian Island, Northern Territory, Australia.

Effects of novel predator on seed harvesting and predator-scent aversion

There was a very clear positive relationship between melomys density and seed take (t_{18} =

5.112, P < 0.001; Fig. 4) and a clear negative relationship between quoll presence and seed

take (t_{18} = -2.344, P = 0.031; Fig. 4). There was, however, no interaction between melomys

density and quoll presence (t_{18} = -0.251, P = 0.805; Fig. 4). When we looked at the difference

in seed take (Δ) between scent treatments within site.session, a striking pattern emerges, in

which there is a clear interaction between the presence of quolls and session ($F_{3,17}$ = 18.61,

P < 0.001; Fig. 5). Soon after the introduction of quolls to Indian Island in 2017, responses of

melomys to predator-scented seeds were indistinguishable between populations with and

without quolls, however, the difference in the responses of melomys diverged after quolls

had been present for one (2018) and two years (2019), respectively (Fig. 5).

Figure 4. Effect of estimated population size on the number of control, unscented seeds removed from seed

plots (*n* = 21) in quoll-invaded and quolls-free sites. Dotted lines denote the effect of quoll presence on seed

removal rate.

Figure 5. Mean (± s.e.) difference (Δ) between the number of control, unscented seeds and predator-scented

seeds removed by melomys from quoll-invaded (*n* = 3; 2017 & *n* = 4; 2018-19) and quoll-free (*n* = 4; 2017 & *n* =

3; 2018-19) sites during each trapping session.

DISCUSSION

The introduction of northern quolls to Indian Island was associated with lowered survival

(Fig. 1) and an immediate drop in population size (Fig. 2) in quoll-invaded melomys

populations. This numerical effect on melomys density had an immediate impact on seed

predation rates (Fig. 4), because seed take is strongly associated with the density of

melomys in this system. This is a classic trophic cascade: predation suppresses herbivore

density, which reduces the pressure that herbivores place on primary producers. Our study,

however, also reveals an additional, subtler, cascade effect; driven by altered prey

behaviour rather than by altered prey density (Fig. 3 & 5).

Within months of quolls appearing on the island, invaded populations of melomys

were significantly shyer than nearby, predator-free populations of conspecifics (Fig. 3). This

rapid but generalised response to a novel threat appears to have had a subtle effect on seed

predation rates: when we examine unscented seeds, per capita seed take is slightly lower in

quoll-invaded populations (Fig. 4). This generalised response appears to have been

supplemented over time with more threat-specific antipredator behaviours (Fig. 5).

Although the willingness of predator-exposed melomys to emerge from shelter (i.e.

boldness) converged through time with that of predator-free melomys, predator-exposed

melomys continued to be more neophobic than their predator-free conspecifics throughout

the study (Fig. 3). Meanwhile, predator-scent aversion, as evidenced by seed plots, steadily

increased over time (Fig. 5). Presumably the significant and novel predation pressure

induced by the introduction of quolls resulted in selection on behaviour and/or learning in

impacted rodent populations, allowing them to fine-tune their behavioural response

(decrease general shyness, but maintain neophobia, and respond to specific cues) as the

nature of the threat became clearer. These changing behavioural responses imply a

generalised reduction in seed take that also becomes fine-tuned over time, with high risk

sites (those that smell of predators) ultimately displaying substantially much lower seed

take than low risk sites. Thus, we see a reduction in seed take that applies to the entire

landscape becoming a fine-grained response, varying on a spatial scale measured in the tens

of metres.

Predation is a pervasive selective force in most natural systems, driving evolutionary

change in prey morphology, physiology, life history and behaviour. Unlike morphology and

physiology, however, the labile nature of behaviour makes it a particularly powerful trait for

rapid response in a changing world (Réale *et al.* 2007; Dall & Griffith 2014). Behavioural

comparisons of wild populations exposed to differing predation regimes provides some

support for the prediction that reduced boldness would be selected for under high

predation scenarios (Åbjörnsson *et al.* 2004; Bell 2005; Brydges *et al.* 2008) and that the

appearance of novel predators can result in bold individuals becoming shyer (Niemelä *et al.*

2012), however, the opposite pattern of response can also occur (Brown *et al.* 2005; Urban

2007) or behavioural phenotypes can be unrelated to predation regime (Laurila 2000;

Carlson & Langkilde 2014). Interestingly, a number of studies have demonstrated that

individuals from high-predation areas were quicker to emerge (Harris *et al.* 2010) and were

bolder and more aggressive (Bell & Sih 2007; Dingemanse *et al.* 2007) than predator-naïve

conspecifics. Although we found the opposite pattern to this immediately following the

arrival of a novel predator, by the second year after predator introduction we found the

emergence latency and boldness of melomys converging with that of predator-free

populations. It is plausible that with more time they may become bolder than their

predator-free conspecifics (by way of compensating for additional costs of acquiring food).

Thus, it is clear that the behavioural state of the population is dynamic, and it seems likely

this dynamism (and perhaps the capacity of the prey species to identify specific threats) may

explain some of the variation between earlier studies.

Although boldness may change over time, neophobia, as a generalised adaptive

response to predation pressure, is now well supported across a number of studies (Crane *et*

al. 2019). Individuals living under high predation risk scenarios have been shown to typically

display generalized neophobia (Brown *et al.* 2015; Elvidge *et al.* 2016), and neophobia can

increase the survival of predator-naïve individuals in initial encounters with predators

(Ferrari *et al.* 2015; Crane *et al.* 2018). Certainly, in our study, predator-exposed melomys

were significantly more neophobic than their predator-free conspecifics; an effect

maintained throughout the study.

Despite reduced survival, significant population declines, and clear behavioural

changes in invaded populations, it is impossible to determine with certainty from our data

whether changes in the behaviour of predator-invaded melomys populations are the result

phenotypic plasticity (learning) or natural selection. The low between trapping session

survival ($S_{2017–2018}$ = 0.207; $S_{2018–2019}$ = 0.091; Fig. 1) of melomys in quoll-invaded

populations means few individuals survive between sessions, so natural selection is a

possibility, and selection on these behavioural traits is potentially very strong. Although

behavioural changes in predator-invaded populations have been documented in a few

systems where predator introductions have been staged and experimentally controlled

(Lapiedra *et al.* 2018; Blumstein *et al.* 2019; Cunningham *et al.* 2019; Pringle *et al.* 2019),

elucidating whether these observed changes arise because of behavioural plasticity or

natural selection can be exceptionally difficult. Rapid behavioural responses of vulnerable

prey to recovered predators has been observed in a single prey generation, presumably

due to behavioural plasticity (Berger *et al.* 2001; Cunningham *et al.* 2019). Similarly,

behavioural adjustments to an introduced predator have been observed as a result of

natural selection on advantageous behavioural traits (Lapiedra *et al.* 2018). In this study,

although we had measures of individual behaviour, our between session recapture rates of

these individuals was sufficiently low that we had no longitudinal data on the behaviour of

individuals to test whether individuals were adjusting their behaviour or whether natural

selection was resulting in population-level change. It thus remains possible (and quite

likely) that both mechanisms were in play.

Although northern quolls represent a novel predator to melomys on Indian Island,

the two species' shared evolutionary history on the northern Australian mainland may

provide some explanation as to why this staged introduction resulted in rapid, finely-tuned

behavioural adjustment in melomys, rather than extinction. Isolation from predators can

rapidly result in the loss of antipredator behaviours from a prey species' behavioural

repertoire (Blumstein & Daniel 2005; Jolly *et al.* 2018a), dramatically increasing an

individual's susceptibility to predation following the introduction of either predator or prey

(Carthey & Banks 2014; Jolly *et al.* 2018b). But such outcomes are not inevitable: length of

isolation, co-evolutionary history, degree of predator novelty, density-dependent effects,

population size, and pre-existing predator-prey associations (Berger *et al.* 2001; Blumstein

2006; Banks & Dickman 2007; Sih *et al.* 2010; Carthey & Banks 2014) are all likely to be

hugely influential in determining whether an invaded population adjusts to the invader or

proceeds towards extinction. Recently, a conservation introduction of Tasmanian devils to

an island previously lacking them found that their possum prey rapidly adjusted their

foraging behaviour to accommodate this newly arrived predator (Cunningham *et al.* 2019).

Despite possums having lived on the island in isolation from devils since the 1950s,

presumably, their long evolutionary history together on mainland Tasmania had them

primed to respond to this predatory archetype (Sih *et al.* 2010; Carthey & Banks 2014;

Cunningham *et al.* 2019). This shared evolutionary history is likely responsible for both

possums' and melomys' ability to rapidly mount appropriate antipredator responses to the

introduction of these predators. The predators are novel within an individual's lifetime, but

the individual's ancestors have encountered them before.

Such considerations are of more than just theoretical interest. While native

Australian mammals clearly possess the ability to rapidly respond to some evolutionary

novel predators under certain circumstances (Carthey & Banks 2014; Blumstein *et al.* 2019),

many native mammals have been unable to mount rapid and/or effective enough defences

to thwart the invasion of evolutionarily novel predators (cats, foxes). This situation has led

to widespread predator-driven extinction on the Australian continent, resulting in Australia

leading the world in modern mammal extinctions (Woinarski *et al.* 2015).

Although our study documented dramatic population declines in predator-invaded

melomys populations, and we are assigning the causation of these declines to the

introduction of quolls, we need to address the unplanned, confounding factor of the fire

that burnt through northern Indian Island after completion of our population monitoring

and behavioural assays in 2017 (Fig. 2). Such fires are commonplace in the Australian wet-

dry tropics (Russell-Smith & Yates 2007); a regular disturbance that is often rapidly offset by

the annual monsoon driven wet season. Since our sites are composed of grass-free

woodland, the fire that burnt through them mostly burnt leaf-litter (though it reached the

mid-storey in other parts of the island). While this likely reduced the short-term availability

of food and cover for melomys, it is unlikely to directly explain the demographic effects we

observed (Fig. 1 & 2). A previous study investigating the effect of fire regimes on native

mammals in savanna woodland in Kakadu National Park, Northern Territory was unable to

detect an effect of fire frequency or intensity on the survival or recruitment of grassland

melomys, despite finding fire impacts in all other co-occurring native mammals studied

(Griffiths & Brook 2015). Interestingly, even in a system where fire is much more infrequent

and significantly more intense (e.g. mesic habitats of eastern Australia), grassland melomys

were found to be relatively unaffected by a wildfire that caused significant impacts to a co-

occurring native rodent, and any demographic impacts felt by melomys were entirely absent

within months of the fire (Liedloff *et al.* 2018). Additionally, the most dramatic behavioural

difference (boldness and neophobia) between quoll-invaded and quoll-free sites was

observed immediately prior to the occurrence of the fire (early August *vs.* mid-August 2017;

Fig 2 & 3). For the behavioural changes we observed that were potentially confounded by

fire, such as predator-scent aversion, we would expect to see these effects decreasing with

time since fire if fire was driving this response, instead we see the opposite trend (Fig. 5).

Finally, if food had become strongly limiting as a consequence of the fire, we would expect

to have observed an increase in seed take in the burned (quoll-invaded) sites, instead we

saw a decrease. For these reasons, we suspect the fire was unlikely to be directly

responsible for the demographic effects to melomys we observed, and fire cannot in any

way explain the response we observed to quoll-scented seeds. We, therefore, believe our

interpretation of these changes as being driven mostly by the addition of a novel predator

to the system is the most parsimonious and globally coherent interpretation of the data.

Fires in northern Australian savannas have, however, been shown to reduce cover for prey

species, facilitating and improving the hunting ability of predators (McGregor *et al.* 2014).

There is, therefore, a strong possibility that this fire may have facilitated the hunting

capabilities of quolls, increased the predation pressure and selection on behaviour imposed

by their arrival in this system.

Although our results suggest that invaded melomys populations are beginning to

adjust to the presence of northern quolls as a novel predator on Indian Island, there has

been no sign of demographic recovery from the addition of this predation pressure on the

island (Fig. 1 & 2). Data from our seed removal experiment clearly demonstrated that the

function of melomys as seed harvesters and dispersers scales with density (Fig. 4). Trophic

cascades resulting from the addition and loss of predators from ecosystems has been

observed in a number of systems globally (Ripple *et al.* 2001; Terborgh *et al.* 2001; Estes *et*

al. 2011), and the results can profoundly shape entire systems. As the only rodent and the

dominant granivore in this system, while melomys populations may not go extinct as a

result of quoll invasion, their reduced abundance and weakened ability to harvest and

disperse seeds may have dramatic, yet to be observed, longer-term consequences for the

vegetation structure and ecosystem function of Indian Island (McConkey & O'Farrill 2016).

Currently, grass is a rare vegetation feature on Indian island (though it is a dominant feature

of savanna woodlands generally), and this is quite possibly a result of the high density of

melomys on this (previously) predator-free island. The presence of quolls may well change

that, as both numerical and behaviour responses of melomys cascade down to the grass

community.

Empirical research on the effects of novel predators on recipient communities under

controlled conditions on a landscape-scale is exceptionally difficult and remains relatively

rare. The introduction of threatened predators to landscapes from which they have been

lost (Cunningham *et al.* 2019) or where they are entirely novel (Lapiedra *et al.* 2018),

however, provides a unique opportunity to observe how naïve prey can respond to novel

predators, and the mechanisms by which predators can structure communities. Our study

provides empirical support that some impacted prey populations can adjust rapidly to the

arrival of a novel predator via a generalised behavioural response (decreased boldness)

followed by development of a species-specific antipredator response (behavioural fine-

tuning). The arrival of the novel predator set off a trophic cascade that was driven, not only

by changed prey density, but also by changed prey behaviour. Thus, rapid adaptive shift may

allow prey populations to persist, but large-scale, system-wide changes may still follow.

Acknowledgments

Thanks to Kenbi Traditional Owners (Raelene and Zoe Singh) for land access permission and

Kenbi Rangers for assistance in the field. Special thanks to Kenbi Rangers Brett Bigfoot, Rex

Edmunds, Jack Gardner, Ian McFarlane, Dale Singh, and Rex Singh for continued field

assistance throughout this project. Thanks to Kenbi Ranger Co-ordinator Steven Brown for

logistical support in the field. Thanks to Alana de Laive for graphic design of figures. Thanks

to Ella Kelly and Naomi Indigo for logistical and moral support on the island.

Funding information

This research was funded by an Australian Research Council Linkage Grant (JKW and BP

LP150100722). In kind support was provided by Kenbi Rangers and the Northern Territory

Government Department of Environment and Natural Resources, Flora and Fauna

Division (via GRG). CJJ was supported by an Australian Postgraduate Award and the

Holsworth Wildlife Research Endowment.

Data availability

The datasets generated and analysed during this study are available via figshare.com.

https://doi.org/10.26188/5dd8fe5651e8b is currently embargoed but access can be granted

on request.

REFERENCES

Åbjörnsson, K., Hansson, L.-A. & Brönmark, C. (2004). Responses of prey from habitats with

different predator regimes: local, adaptation and heritability. *Ecology*, 85, 1859–

1866.

Abrams, P.A. (2000). The evolution of predator-prey interactions: theory and evidence.

Annu. Rev. Ecol. Syst., 31, 79–105.

Andersen, A.N., Cook, G.D., Corbett, L.K., Douglas, M.M., Eager, R.W., Russell-Smith, J., *et al.*

(2005). Fire frequency and biodiversity conservation in Australian tropical savannas:

implications from the Kapalga fire experiment. *Austral Ecol*, 30, 155–167.

Banks, P.B. & Dickman, C.R. (2007). Alien predation and the effects of multiple levels of prey

naiveté. *Trends in Ecology & Evolution*, 22, 229–230.

Begg, R., Walsh, B., Woerle, F. & King, S. (1983). Ecology of *Melomys burtoni*, the grassland

melomys (Rodentia: Muridae) at Cobourg Peninsula, N.T. *Wildlife Research*, 10, 259–

267.

Bell, A.M. (2005). Behavioural differences between individuals and two populations of

stickleback (*Gasterosteus aculeatus*): behavioural syndromes. *Journal of Evolutionary*

Biology, 18, 464–473.

Bell, A.M. & Sih, A. (2007). Exposure to predation generates personality in threespined

sticklebacks (*Gasterosteus aculeatus*). *Ecology Letters*, 10, 828–834.

Berger, J., Swenson, J.E. & Persson, I.-L. (2001). Recolonizing carnivores and naive prey:

conservation lessons from pleistocene extinctions. *Science*, 291, 1036–1039.

Blumstein, D.T. (2006). The multipredator hypothesis and the evolutionary persistence of

antipredator behavior. *Ethology*, 112, 209–217.

Blumstein, D.T. & Daniel, J.C. (2005). The loss of anti-predator behaviour following isolation

on islands. *Proceedings of the Royal Society B: Biological Sciences*, 272, 1663–1668.

Blumstein, D.T., Letnic, M. & Moseby, K.E. (2019). *In situ* predator conditioning of naive prey

prior to reintroduction. *Phil. Trans. R. Soc. B*, 374, 20180058.

Braithwaite, R. & Griffiths, A. (1994). Demographic variation and range contraction in the

northern quoll, *Dasyurus hallucatus* (Marsupialia: Dasyuridae). *Wildlife Research*, 21,

203.

Brown, C. & Braithwaite, V.A. (2004). Size matters: a test of boldness in eight populations of

the poeciliid *Brachyraphis episcopi*. *Animal Behaviour*, 68, 1325–1329.

Brown, C., Jones, F. & Braithwaite, V. (2005). *In situ* examination of boldness–shyness traits

in the tropical poeciliid, *Brachyraphis episcopi*. *Animal Behaviour*, 70, 1003–1009.

Brown, G.E., Elvidge, C.K., Ramnarine, I., Ferrari, M.C.O. & Chivers, D.P. (2015). Background

risk and recent experience influences retention of neophobic responses to

predators. *Behav Ecol Sociobiol*, 69, 737–745.

Brown, G.E., Ferrari, M.C.O., Elvidge, C.K., Ramnarine, I. & Chivers, D.P. (2013).

Phenotypically plastic neophobia: a response to variable predation risk. *Proceedings*

of the Royal Society B: Biological Sciences, 280, 20122712–20122712.

Brydges, N.M., Colegrave, N., Heathcote, R.J.P. & Braithwaite, V.A. (2008). Habitat stability

and predation pressure affect temperament behaviours in populations of three-

spined sticklebacks. *J Anim Ecology*, 77, 229–235.

Burnham, K.P. & Anderson, D.R. (1998). *Model selection and inference: a practical*

information-theoretic approach. Springer, New York.

Carlson, B.E. & Langkilde, T. (2014). No evidence of selection by predators on tadpole

boldness. *Behav*, 151, 23–45.

Carter, A.J., Feeney, W.E., Marshall, H.H., Cowlishaw, G. & Heinsohn, R. (2013). Animal

personality: what are behavioural ecologists measuring? *Biological Reviews*, 88, 465–

475.

Carthey, A.J.R. & Banks, P.B. (2014). Naïveté in novel ecological interactions: lessons from

theory and experimental evidence. *Biological Reviews*, 89, 932–949.

Courchamp, F., Chapuis, J.-L. & Pascal, M. (2003). Mammal invaders on islands: impact,

control and control impact. *Biological Reviews*, 78, 347–383.

Crane, A.L., Brown, G.E., Chivers, D.P. & Ferrari, M.C.O. (2019). An ecological framework of

neophobia: from cells to organisms to populations. *Biol Rev*, brv.12560.

Crane, A.L., Chivers, D.P. & Ferrari, M.C.O. (2018). Embryonic background risk promotes the

survival of tadpoles facing surface predators. *PLoS ONE*, 13, e0193939.

Cremona, T., Mella, V.S.A., Webb, J.K. & Crowther, M.S. (2015). Do individual differences in

behavior influence wild rodents more than predation risk? *Journal of Mammalogy*,

96, 1337–1343.

Cunningham, C.X., Johnson, C.N., Hollings, T., Kreger, K. & Jones, M.E. (2019). Trophic

rewilding establishes a landscape of fear: Tasmanian devil introduction increases

risk-sensitive foraging in a key prey species. *Ecography*, ecog.04635.

Dall, S.R.X. & Griffith, S.C. (2014). An empiricist guide to animal personality variation in

ecology and evolution. *Front. Ecol. Evol.*, 2, 1–7.

Dielenberg, R.A. & McGregor, I.S. (2001). Defensive behavior in rats towards predatory

odors: a review. *Neuroscience & Biobehavioral Reviews*, 25, 597–609.

Dingemanse, N.J., Wright, J., Kazem, A.J.N., Thomas, D.K., Hickling, R. & Dawnay, N. (2007).

Behavioural syndromes differ predictably between 12 populations of three-spined

stickleback. *J Anim Ecology*, 76, 1128–1138.

Doherty, T.S., Glen, A.S., Nimmo, D.G., Ritchie, E.G. & Dickman, C.R. (2016). Invasive

predators and global biodiversity loss. *Proc Natl Acad Sci USA*, 113, 11261–11265.

Elvidge, C.K., Chuard, P.J.C. & Brown, G.E. (2016). Local predation risk shapes spatial and

foraging neophobia patterns in Trinidadian guppies. *Curr Zool*, 62, 457–462.

Estes, J.A., Terborgh, J., Brashares, J.S., Power, M.E., Berger, J., Bond, W.J., *et al.* (2011).

Trophic downgrading of planet Earth. *Science*, 333, 301–306.

Ferrari, M.C.O., McCormick, M.I., Meekan, M.G. & Chivers, D.P. (2015). Background level of

risk and the survival of predator-naive prey: can neophobia compensate for predator

naivety in juvenile coral reef fishes? *Proc. R. Soc. B*, 282, 20142197.

Fortin, D., Beyer, H.L., Boyce, M.S., Smith, D.W., Duchesne, T. & Mao, J.S. (2005). Wolves

influence elk movements: behavior shapes a trophic cascade in Yellowstone National

Park. *Ecology*, 86, 1320–1330.

Gelman, A. & Rubin, D.B. (1992a). Inference from iterative simulation using multiple

sequences. *Statistical Science*, 7, 457–472.

Gelman, A. & Rubin, D.B. (1992b). [Practical Markov Chain Monte Carlo]: Rejoinder:

replication without contrition. *Statistical Science*, 7, 503–511.

Griffiths, A.D. & Brook, B.W. (2015). Fire impacts recruitment more than survival of small-

mammals in a tropical savanna. *Ecosphere*, 6, 1–22.

Harris, S., Ramnarine, I.W., Smith, H.G. & Pettersson, L.B. (2010). Picking personalities apart:

estimating the influence of predation, sex and body size on boldness in the guppy

Poecilia reticulata. *Oikos*, 119, 1711–1718.

Jolly, C.J., Webb, J.K. & Phillips, B.L. (2018a). The perils of paradise: an endangered species

conserved on an island loses antipredator behaviours within 13 generations. *Biol.*

Lett., 14, 20180222.

Jolly, C.J., Kelly, E., Gillespie, G.R., Phillips, B. & Webb, J.K. (2018b). Out of the frying pan:

Reintroduction of toad-smart northern quolls to southern Kakadu National Park.

Austral Ecology, 43, 139–149.

Jolly, C.J., Webb, J.K., Gillespie, G.R., Hughes, N.K. & Phillips, B.L. (2019). Bias averted:

personality may not influence trappability. *Behav Ecol Sociobiol*, 73, 129.

Kelly, E. (2019). *Targeted gene flow for conservation: northern quolls and the invasive cane

toad.* PhD Thesis. University of Melbourne, Australia.

Kelly, E. & Phillips, B.L. (2016). Targeted gene flow for conservation. *Conservation Biology*,

30, 259–267.

Kemper, C.M., Kitchener, D.J., Humphreys, W.F., How, R.A., Bradley, A.J. & Schmitt, L.H.

(1987). The demography and physiology of *Melomys* sp. (Rodentia: Muridae) in the

Mitchell Plateau area, Kimberley, Western Australia. *Journal of Zoology*, 212, 533–

562.

Kery, M. & Schaub, M. (2011). *Bayesian Population Analysis using WinBUGS: A Hierarchical

Perspective*. Elsevier Science & Technology, San Diego, United States.

Lapiedra, O., Schoener, T.W., Leal, M., Losos, J.B. & Kolbe, J.J. (2018). Predator-driven

natural selection on risk-taking behavior in anole lizards. *Science*, 360, 1017–1020.

Laundre, J.W., Hernandez, L. & Ripple, W.J. (2010). The landscape of fear: ecological

implications of being afraid. *TOECOLJ*, 3, 1–7.

Laurila, A. (2000). Behavioural responses to predator chemical cues and local variation in

antipredator performance in *Rana temporaria* tadpoles. *Oikos*, 88, 159–168.

Liedloff, A.C., Wilson, J.C. & Engeman, R.M. (2018). The effect of wildfire on population

dynamics for two native small mammal species in a coastal heathland in Queensland,

Australia. *Acta Oecologica*, 88, 58–64.

Lima, S.L. & Dill, L.M. (1990). Behavioral decisions made under the risk of predation: a

review and prospectus. *Can. J. Zool.*, 68, 619–640.

López, P., Hawlena, D., Polo, V., Amo, L. & Martín, J. (2005). Sources of individual shy–bold

variations in antipredator behaviour of male Iberian rock lizards. *Animal Behaviour*,

69, 1–9.

McConkey, K.R. & O'Farrill, G. (2016). Loss of seed dispersal before the loss of seed

dispersers. *Biological Conservation*, 201, 38–49.

McGregor, H.W., Legge, S., Jones, M.E. & Johnson, C.N. (2014). Landscape management of

fire and grazing regimes alters the fine-scale habitat utilisation by feral cats. *PLoS*

ONE, 9, e109097.

McGregor, I.S., Schrama, L., Ambermoon, P. & Dielenberg, R.A. (2002). Not all 'predator

odours' are equal: cat odour but not 2,4,5 trimethylthiazoline (TMT; fox odour)

elicits specific defensive behaviours in rats. *Behavioural Brain Research*, 129, 1–16.

Medina, F.M., Bonnaud, E., Vidal, E., Tershy, B.R., Zavaleta, E.S., Josh Donlan, C., *et al.*

(2011). A global review of the impacts of invasive cats on island endangered

vertebrates. *Glob. Change Biol.*, 17, 3503–3510.

Moore, H.A., Dunlop, J.A., Valentine, L.E., Woinarski, J.C.Z., Ritchie, E.G., Watson, D.M., *et al.*

(2019). Topographic ruggedness and rainfall mediate geographic range contraction

of a threatened marsupial predator. *Divers Distrib*, ddi.12982.

Niemelä, P.T., DiRienzo, N. & Hedrick, A.V. (2012). Predator-induced changes in the boldness

of naïve field crickets, *Gryllus integer*, depends on behavioural type. *Animal*

Behaviour, 84, 129–135.

Oakwood, M. (1997). The Ecology of the Northern Quoll, *Dasyurus hallucatus*. PhD Thesis.

Australian National University, Australia.

Oakwood, M., Woinarski, J.C.Z. & Burnett, S. (2016). *Dasyurus hallucatus* (northern quoll).

The IUCN Red List of Threatened Species 2016:

e.T6295A21947321. http://dx.doi.org/10.2305/IUCN.UK.2016-

2.RLTS.T6295A21947321.en. Downloaded on 31 October 2019.

Plummer, M., Stukalov, A. & Denwood, M. (2017). rjags: Bayesian graphical models using

MCMC. *URL http://CRAN. R-project. org/package= rjags. R package version, 2, 0-4.*

Pringle, R.M., Kartzinel, T.R., Palmer, T.M., Thurman, T.J., Fox-Dobbs, K., Xu, C.C.Y., *et al.*

(2019). Predator-induced collapse of niche structure and species coexistence.

Nature, 570, 58–64.

R Core Team. (2019). *R: A Language and Environment for Statistical Computing.* R

Foundation for Statistical Computing, Vienna.

Réale, D., Reader, S.M., Sol, D., McDougall, P.T. & Dingemanse, N.J. (2007). Integrating

animal temperament within ecology and evolution. *Biological Reviews*, 82, 291–318.

Relyea, R.A. (2003). Predators come and predators go: the reversiblity of predator-induced

traits. *Ecology*, 84, 1840–1848.

Ripple, W.J., Larsen, E.J., Renkin, R.A. & Smith, D.W. (2001). Trophic cascades among wolves,

elk and aspen on Yellowstone National Park's northern range. *Biological

Conservation*, 102, 227–234.

Royle, J.A., Dorazio, R.M. & Link, W.A. (2007). Analysis of multinomial models with unknown

index using data augmentation. *Journal of Computational and Graphical Statistics*,

16, 67–85.

Russell-Smith, J. & Yates, C.P. (2007). Australian savanna fire regimes: context, scales,

patchiness. *Fire Ecology*, 3, 48–63.

Sax, D., Stachowicz, J., Brown, J., Bruno, J., Dawson, M., Gaines, S., *et al.* (2007). Ecological

and evolutionary insights from species invasions. *Trends in Ecology & Evolution*, 22,

465–471.

Schoener, T.W. & Spiller, D.A. (1996). Devastation of prey diversity by experimentally

introduced predators in the field. *Nature*, 381, 691–694.

Shine, R. (2010). The ecological impact of invasive cane toads (*Bufo marinus*) in Australia.

The Quarterly Review of Biology, 85, 253–291.

Sih, A. (1992). Prey uncertainty and the balancing of antipredator and feeding needs. *The*

American Naturalist, 139, 1052–1069.

Sih, A., Bolnick, D.I., Luttbeg, B., Orrock, J.L., Peacor, S.D., Pintor, L.M., *et al.* (2010).

Predator-prey naïveté, antipredator behavior, and the ecology of predator invasions.

Oikos, 119, 610–621.

Sih, A., Ziemba, R. & Harding, K.C. (2000). New insights on how temporal variation in

predation risk shapes prey behavior. *Trends in Ecology & Evolution*, 15, 3–4.

Simberloff, D., Martin, J.-L., Genovesi, P., Maris, V., Wardle, D.A., Aronson, J., *et al.* (2013).

Impacts of biological invasions: what's what and the way forward. *Trends in Ecology*

& Evolution, 28, 58–66.

Suraci, J.P., Clinchy, M., Dill, L.M., Roberts, D. & Zanette, L.Y. (2016). Fear of large carnivores

causes a trophic cascade. *Nature Communications*, 7, 10698.

Tanner, M.A. & Wong, W.H. (1987). The calculation of posterior distributions by data

augmentation. *Journal of the American Statistical Association*, 82, 528–540.

Terborgh, J., Lopez, L., Nunez, P., Rao, M., Shahabuddin, G., Orihuela, G., *et al.* (2001).

Ecological meltdown in predator-free forest fragments. *Science*, 294, 1923–1926.

Urban, M.C. (2007). Risky prey behavior evolves in risky habitats. *Proceedings of the*

National Academy of Sciences, 104, 14377–14382.

Woinarski, J.C.Z., Burbidge, A.A. & Harrison, P.L. (2015). Ongoing unraveling of a continental

fauna: decline and extinction of Australian mammals since European settlement.

Proceedings of the National Academy of Sciences, 112, 4531–4540.