Trophic cascade driven by behavioural fine-tuning as naïve prey rapidly

adjust to a novel predator

CHRIS J. JOLLY*[1], ADAM S. SMART[1], JOHN MOREEN[2], JONATHAN K. WEBB[3], GRAEME R.

GILLESPIE[1,4] AND BEN L. PHILLIPS[1]

[1]School of BioSciences, University of Melbourne, Parkville Vic 3010 Australia

[2]Kenbi Rangers, Mandorah NT 0822 Australia

[3]School of Life Sciences, University of Technology Sydney, Broadway NSW 2007 Australia

[4]Flora and Fauna Division, Department of Environment and Natural Resources, Northern Territory

Government, Berrimah NT 0828 Australia

*Correspondence:

C. Jolly

Email: cjolly1@student.unimelb.edu.au

21   November 2019

23   **Keywords:** Antipredator behaviour, boldness, invasion, neophobia, novel predator,

24   predator-prey dynamics, prey naivety

The arrival of novel predators can trigger trophic cascades driven by shifts in prey numbers. Predators also elicit behavioural change in prey populations, via phenotypic plasticity and/or rapid evolution, and such changes may also contribute to trophic cascades. Here we document rapid demographic and behavioural changes in populations of a prey species (grassland melomys *Melomys burtoni*, a granivorous rodent) following the introduction of a novel marsupial predator (northern quoll *Dasyurus hallucatus*). Within months of quolls appearing, populations of melomys exhibited reduced survival and population declines relative to control populations. Quoll-invaded populations ($n = 4$) were also significantly shyer than nearby, quoll-free populations ($n = 3$) of conspecifics. This rapid but generalised response to a novel threat was replaced over the following two years with more threat-specific antipredator behaviours (i.e. predator-scent aversion). Predator-exposed populations, however, remained more neophobic than predator-free populations throughout the study. These behavioural responses manifested rapidly in changed rates of seed predation by melomys across treatments. Quoll-invaded melomys populations exhibited lower per-capita seed take rates, and rapidly developed an avoidance of seeds associated with quoll scent, with discrimination playing out over a spatial scale of tens of metres. Presumably the significant and novel predation pressure induced by quolls drove melomys populations to fine-tune behavioural responses to be more predator-specific through time. These behavioural shifts could reflect individual plasticity (phenotypic flexibility) in behaviour or may be adaptive shifts from natural selection imposed by quoll predation. Our study provides a rare insight into the rapid ecological and behavioural shifts enacted by prey to mitigate the impacts of a novel predator and shows that trophic cascades can be strongly influenced by behavioural as well as numerical responses.

INTRODUCTION

Predation is one of the most pervasive and powerful forces acting on populations. Not only does predation directly impact a population's demography (Schoener & Spiller 1996), it also imposes natural selection (Abrams 2000). The pressure that predators impose on populations will vary through time and space for many reasons, including tightly coupled predator-prey dynamics, predator movement, prey switching, or stochastic processes (Lima & Dill 1990; Sih 1992). The fact that predation is not constant, and that antipredator defences may be costly, suggests that flexible responses to predation pressure will often be favoured (Sih *et al.* 2000; Berger *et al.* 2001). There is, in fact, a great deal of empirical evidence that flexible responses to predation are common (e.g. Relyea 2003; Brown *et al.* 2013; Cunningham *et al.* 2019). Investment in antipredator traits across morphology, life-history, and behaviour often varies, and is dependent on the perceived risk of predation.

As well as impacting prey populations, it is increasingly apparent that predators play a powerful role in structuring communities (Estes *et al.* 2011). Some of our best evidence for this comes from the introduction of predators to naïve communities. Invasive predators can cause extinctions (Medina *et al.* 2011; Woinarski *et al.* 2015; Doherty *et al.* 2016), and alter trophic structures and ecosystem function within recipient communities (Courchamp *et al.* 2003; Simberloff *et al.* 2013). Cascading outcomes are often thought of as purely numeric effects: predators depress the size of prey populations, and the altered numbers of prey can cause cascading numerical changes down trophic levels (Ripple *et al.* 2001). These numerical effects are undeniably important, but the fact that predators can also elicit phenotypic change in prey populations—through phenotypic plasticity and natural selection—means that subtler ecological effects may also manifest. Prey species living alongside predators may forage at different times, or in different places compared with the same species in a

predator-free environment (Laundre *et al.* 2010). Such behavioural shifts can alter

downstream species interactions in potentially complex ways (Fortin *et al.* 2005; Suraci *et al.*

2016).

Because predator invasions are rarely intentional or anticipated, there is a scarcity of

controlled empirical work on the effects of novel predators on recipient communities and

the mechanisms via which these effects play out (but see Lapiedra *et al.* 2018; Pringle *et al.*

2019). Such tests are needed, however, if we are to predict invasive species impacts, and

improve conservation management (Sih *et al.* 2010a) and our understanding of how

communities are structured via predator invasion (Sax *et al.* 2007).

Northern quolls (*Dasyurus hallucatus*) were, until recently, a common predator

across northern Australia. They have declined over the last several decades, following the

general decline in northern Australian mammals (Woinarski *et al.* 2015), thought to be

driven by changes in grazing, fire, and predation regimes (Braithwaite & Griffiths 1994).

More recently, the invasion of toxic invasive prey (cane toads, *Rhinella marina*) has resulted

in dramatic, range-wide population declines in northern quolls (Shine 2010; Oakwood *et al.*

2016). Due to local extinction, northern quolls are now absent from large tracts of their

former range and their ecological function as a medium-sized mammalian predator has

been lost (Moore *et al.* 2019). For their conservation, northern quolls have recently been

introduced to a number of offshore islands where they have never previously existed.

In 2017, a population of 54 northern quolls were introduced to a 25km^2 island off the

coast of north-western Northern Territory, Australia (Kelly 2019). Prior to this introduction,

Indian Island (Kabal) lacked mammalian predators, and large native reptilian predators had

recently been reduced to near extinction by the invasion of cane toads. We take advantage

of the introduction of northern quolls to a new island to directly test the effects of quolls as

a novel predator on an island ecosystem and observe how native prey populations adjust to

mitigate the impacts of their arrival. Since quolls are an ecologically novel predator on this

island, we predict that this introduction may result in demographic effects (reduced survival

and abundance) in invaded prey populations. If behavioural adjustments are able to reduce

the demographic effects of a novel predator, we predict rapid behavioural changes in quoll-

exposed melomys populations, such as reduced boldness and foraging rates, and avoidance

of predator-scent, may manifest through time.

METHODS

Introduction of northern quolls

In May 2017, 54 adult northern quolls were introduced to the north-eastern tip of Indian

Island, Bynoe Harbor, Northern Territory, Australia (12°37'24.60"S, 130°30'0.72"E) to field

test the conservation strategy of targeted gene flow (Kelly & Phillips 2016). Quolls are a

voracious, opportunistic generalist predator (< 1.5k g; Oakwood 1997), and their

introduction presented an opportunity to monitor the behavioural and demographic

impacts on grassland melomys (*Melomys burtoni*), a native mammalian granivorous prey

species (mean body mass 56 g, 5.6–103.7 g). Immediately prior to the introduction of quolls,

we started monitoring populations of melomys in one woodland and two monsoon vine

thicket plots in the vicinity of where quolls were to be released and radio tracked. After

quolls were introduced and tracked it became immediately apparent that quolls were

largely avoiding monsoon vine thicket sites and, since these sites would neither be effective

"impact" or "control" sites, these sites were dropped from the on-going monitoring.

Because these sites had to be dropped from our monitoring, we missed the opportunity to

implement a robust Before-After Impact-Control design. For this reason, we only present

data from before the introduction of quolls from one invaded site. Most of our data

compare quoll-invaded (impact) versus quoll-free (control) sites over time, commencing

within a few months of quoll arrival.

Melomys population monitoring

To determine whether the arrival of a novel predator resulted in demographic impacts

(population size and survival) to native prey species, we monitored four "impact", quoll-

invaded sites established in the north of Indian island in the vicinity of where quolls were

released and three "control", quoll-free site established in the south of the island (Fig 1).

Populations of melomys on Indian Island were monitored during four trips occurring

immediately prior to the introduction of quolls in May (site 1) 2017, and after the

introduction of quolls August 2017 (sites 2–7), April 2018 (sites 1–7), and May 2019 (sites 1–

7).

Melomys were monitored at seven independent 1ha (100 m x 100 m) plots (sites 1–

7) spread out across Indian Island using a standard mark-recapture trapping regime

designed for a monitoring project (Begg *et al.* 1983; Kemper *et al.* 1987). Sites in the north

(quoll-invaded) and south (quoll-free) of the island were between 8.7 and 9.8km apart (Fig.

1; Table 1) and were composed of similar habitat types. The northern and southern sections

of Indian Island are divided by mangrove habitat which is inundated at high tide. Cage and

camera trapping as well as track surveys confirmed that quolls were present at the "impact"

sites and absent from the "control" sites for the duration of the study (Jolly *et al.* unpub.

data).

Figure 1. Map showing the arrangement of grassland melomys (*Melomys burtoni*) monitoring sites on Indian

Island, Northern Territory, Australia. Quolls were present at the four monitoring sites in the north of the island

and quolls were absent from the three monitoring sites in the south of the island for the duration of the study.

154 **Table 1.** Pairwise distance matrix between sites on Indian Island, Northern Territory, Australia. Quolls were

155 present at sites 1–4 and quolls were absent at sites 5–7 for the duration of the study.

Distance (m)	Site 1	Site 2	Site 3	Site 4	Site 5	Site 6	Site 7
Site 1							
Site 2	270						
Site 3	260	350					
Site 4	400	300	250				
Site 5	8760	9030	8710	9000			
Site 6	8470	8730	8450	8070	300		
Site 7	9670	9920	9590	9820	1260	1500	

Each of the seven monitoring sites consisted of 100 Elliott traps (Elliott Scientific

Equipment, Upwey, Victoria) spaced at 10 m intervals in a 10 x 10 grid. Most trapping grids

were open for four nights, however, the first trapping grid (site 1, May 2017) was open for

six nights. After four trap nights, the majority of the melomys population had been captured

at least once (Jolly *et al.* 2019). Traps were baited with balls of peanut butter, rolled oats

and honey. These baits were replaced daily for the duration of each trapping session. Traps

were checked for captures early each morning and all traps were cleared within two hours

of sunrise.

Captured melomys were weighed (g) and sexed. Before release, each melomys was

implanted with a microchip (Trovan Unique ID100). On successive mornings, all melomys

were scanned (Trovan LID575 Handheld Reader), and any new individuals were

microchipped. On the last morning of each trapping session, all melomys caught were

retained for behavioural assays. Throughout the study 439 individual melomys were

captured and given microchips (melomys caught per site: site 1 = 83; site 2 = 52; site 3 = 63;

site 4 = 59; site 5 = 69; site 6 = 59; and site 7 = 54). Of these, 146 (33%) were caught on the

final night of trapping and were retained for behavioural trials. Only large, healthy juveniles

(*n* = 11), adult males (*n* = 58), and adult non-visibly pregnant females (*n* = 77) were retained

for behavioural experiments. Melomys were retained in their respective Elliott traps and

taken to the field station for diurnal husbandry. They were provided food and water *ad*

libitum until 2 hours prior to testing. At this point, in an attempt to standardise hunger

levels, access to food and water was removed. Indian Island is remote and uninhabited by

humans, so all behavioural experiments were conducted in the field under near natural

conditions (see Jolly *et al.* 2019 for detailed experimental procedures).

Modified open field tests

We employed modified open field tests (also referred to as emergence tests: see Brown &

Braithwaite 2004; López *et al.* 2005; Carter *et al.* 2013; Jolly *et al.* 2019) to assess boldness

in grassland melomys and whether the arrival of a novel predator resulted in behavioural

shifts in invaded populations. All open field tests were conducted on the night after the last

trap night (night 5) and in opaque-walled experimental arenas (540mm x 340mm x 370mm).

Experimental arenas were modified plastic boxes that had an inverted Elliott trap sized hole

cut in one end and were illuminated by strings of red LED lights (Jolly *et al.* 2019). Each

experimental arena had natural sand as substrate, and a rolled ball of universal bait (peanut

butter, oats and honey) located both in the centre and along one wall of the arena (Jolly *et*

al. 2019). After dark, Elliott traps containing a melomys were inserted into the hole in the

side of each experimental arena and melomys were allowed to habituate for 10 min. At the

start of each trial, Elliott trap doors were locked open—the inverted orientation of the trap

prevented them from being triggered closed. Melomys were given 10 min to explore the

open field arena. After 10 min, individuals were rounded back into their retreat (the Elliott

trap) and a novel object (standard red, plastic disposable bowl) was placed at the end of the

arena opposite the Elliott trap (Jolly *et al.* 2019). Melomys were then given a further 10 min

to explore the arena and interact with the novel object. Elliott traps remained open during

the open field tests and melomys could shelter and emerge from them under their own

volition. All trials were recorded using a GoPro HERO 3. A previous study in this system

determined that melomys showed repeatable behaviour between trials (boldness: R [±

95%CI] = 0.67 [0.47, 0.80], P < 0.001; emergence time: R [± 95%CI] = 0.73 [0.53, 0.83], P <

0.001; novel object: R [± 95%CI] = 0.61 [0.209, 0.974], P < 0.001; Jolly et $al.$ 2019), therefore

the data presented in this study were from a single behavioural trial of each animal (n =

146). Once trials were complete, each melomys was released at its point of capture.

To measure the boldness of individual melomys, we scored three behaviours

typically associated with boldness and neophobia in rodents (Dielenberg & McGregor 2001;

McGregor et $al.$ 2002; Réale et $al.$ 2007; Cremona et $al.$ 2015): whether melomys fully

emerged from their Elliott trap hide and entered the open arena during the 0–10 min period

(scored 0 or 1, respectively); whether they fully emerged and entered the trial arena during

the 10–20 min period (scored 0 or 1); and whether they interacted (touched) with the novel

object that was placed in the arena during the 10–20 min period (scored 0 or 1). Videos

were scored by a single observer who was blind to each melomys' origin and identity.

Because interacting with the novel object was predicated on a melomys' willingness to

emerge from their hide during the 10–20 min period, for analysis we combined their

emergence during this period and interaction with the novel object into a single binary

score: 0 (neophobic) = did not emerge or emerged but did not interact with novel object; or

1 (not neophobic): emerged and interacted with novel object.

Seed removal plots

To assess whether the arrival of a novel predator affected the seed harvesting behaviour of granivorous melomys, we established seed removal plots at each site and sampled them each trapping session (night 6). After trapping and open field tests were conducted and melomys had been returned to their capture location, we set up 81 seed plots at each site by scraping away leaf litter with a shovel to create bare earth plots. These bare earth plots were created so that they were located in the centre between four Elliott traps within the 10x10 trapping grid. All seed plots were located randomly with respect to "distances to cover" but were all located on relatively open patches of ground. Sufficient within site replication (n = 81) significantly reduces the likelihood of distance to cover biasing population-level responses to seeds. Just before dark on the night of the seed removal experiment, we placed a single wheat seed in the centre of each bare earth plot. These seeds were either unscented, control seeds (n = 40) or predator-scented seeds that had been maintained in a sealed clip-lock bag filled with freshly collected northern quoll fur (n = 41). The placement of predator-scented and unscented seeds was alternated so that there was a chequered arrangement of scented and unscented seeds across the site. To ensure that the predator-scent was strong enough to be detected by melomys, along with the predator-scented seeds, we also placed a few strands of quoll fur around the predator-scented seeds. Before light the next morning, we returned back to each plot and counted the number of seeds of each scent-type that were removed from the plot. Melomys are the only nocturnal granivorous animal that occurs on Indian Island, and to avoid diurnal granivorous birds from removing seeds we conducted this experiment during the night only.

Wildfire on northern Indian Island

Immediately following our monitoring and experiments in August 2017, a wildlife broke out

on northern Indian Island in the vicinity of the four quoll-invaded sites and burnt through all

of the sites. Because of this, our experimental design is confounded by the fact that all of

our quoll-invaded sites were burnt, and all of our quoll-free sites were unburnt. Fire is a

regular disturbance in this landscape (Andersen *et al.* 2005), and previous work has shown

little effect of fire on abundance, survival or recruitment of grassland melomys (Griffiths &

Brook 2015; Liedloff *et al.* 2018). Nonetheless, this confound exists and we proceed with

caution when interpreting the effects of quolls on population size and survival of melomys.

Statistical analysis

During trapping sessions we identified individual melomys that were captured at each site

by their unique microchips. Because melomys on Indian Island have very small home ranges

(tending to be caught in the same or adjacent traps throughout the trapping period: Jolly *et*

al. unpub. data) and since we never observed captures of melomys marked at other sites

(Jolly *et al.* unpub. data), we treated each site as independent with regard to demographics

and behaviour (Table 1).

To estimate between-session survival, we analysed the mark-recapture data to

estimate recapture and survival rates using Cormack-Jolly-Seber models in program MARK.

At each site, there were three primary trapping sessions of four nights, for a total of 12 time

intervals in the input file. Because quolls prey on melomys, we hypothesised that survival

rates of melomys would be lower between trapping sessions at sites with quolls than at

sites without quolls. We included two groups, quoll-free (control) and quoll-invaded

(impact), in the input file. We ran a series of models in MARK to test the following *a priori*

hypotheses: (1) survival rates between sessions are lower at quoll-free sites than at quoll-

invaded sites; (2) survival rates are lower between sessions than within sessions, but are

unaffected by quolls; (3) survival is constant through time; and (4) survival varies through

time. All candidate models were ranked according to their AICc values and associated AIC

weights (Burnham & Anderson 1998). Models with AICc values < 2 were considered to be

well supported by the data (Burnham & Anderson 1998). We used Akaike's Weights, which

are proportional to the normalized, relative likelihood of each model, and to determine

which of these models was most plausible (Buckland *et al.* 1997).

To test whether the presence of quolls impacted melomys population size, we used

a hierarchical model in which population size was made a function of quoll

presence/absence, capture session, and the interaction between these factors. Population

size at each site during each session is estimated in this process, and we fitted this model in

a Bayesian framework. Our observations consisted of a capture history for each observed

individual over the number of nights at each site for each trapping session. We denoted the

number of individuals at site *s* during session *k* as N_{ks}. To estimate N_{ks} we used a closed

population mark-recapture analysis in which each individual, *i*, was either observed, or not

(O_{iks}), according to a Bernoulli distribution:

287                                $O_{iks} \sim$ Bernoulli(d_s)

288   Where d_s denotes the expected detection probability within session *s*. Our previous MARK

289   analysis found clear evidence for variation in detection probability across sessions, but

290   detection probabilities of melomys on Indian Island had previously been found not to vary

291   measurably between individuals nor to change over time within a trapping session (Jolly *et*

292   *al.* 2019). Thus, we made detection probability a function of session according to:

293   logit(d_s) = $\mu_d + t_s$

Where μ_d is the expected detection probability in the first session, and t_s denotes the

(categorical) effect of session on detection.

We used the "data augmentation" method (Tanner & Wong 1987; Royle et al. 2007;

Kery & Schaub 2011) in combination with this detection probability to estimate N_{ks} for each

site per session (site.session). Using this approach, the data were 'padded' to a given size by

adding an arbitrary number of zero-only encounter histories of 'potential' unobserved

individuals. The augmented dataset was then modelled as a zero-inflated model (Royle et al.

2007) which changes the problem from estimating a count, to estimating a proportion. This

was executed by adding a latent binary indicator variable, R_{iks}, (taking values of either 0 or 1)

to classify each row in the augmented data matrix as a 'real' individual or not, where R_{iks}~

Bernoulli(Ω_{ks}). The parameter Ω_{ks} is the proportion of the padded population that is real,

and $N_{ks} = \Sigma_i R_{iks}$.

We then made Ω_{ks} (which scales with population size) a function of quoll

presence/absence, q_c; session, b_k; and the interaction between the two:

308                    $$\text{logit}(\Omega_{ks}) = \mu_p + q_c + b_k + \gamma_{ck}$$

The model was fitted using Bayesian Markov Chain Monte Carlo (MCMC) methods

and minimally informative priors (Table 2) within the package JAGS (Plummer et al. 2017)

using R (R Core Team 2019). Parameter estimates were based on 30,000 iterations with a

thinning interval of 5 following a 10,000 sample burn-in. Three MCMC chains were run, and

model convergence assessed by eye, and using the Gelman-Rubin diagnostic (Gelman &

Rubin 1992a, 1992b).

To assess whether the introduction of quolls affected the behaviour of melomys

populations, we divided the responses of melomys in open field tests into two independent

response variables: whether individuals emerged or not during the 0-10 min period

(binomial: 0 or 1); and whether individuals emerged and interacted with the novel object or

not during the 10-20 min period (binomial: 0 or 1). We used generalised linear mixed-effects

models with binomial errors and a logit link to test the effect of quoll presence (two levels:

quolls present and quolls absent) and trapping session (continuous), with site included as a

random effect, on the behavioural response variables. *P*-values were obtained by likelihood

ratio tests of the full model with the effect in question against the model without the effect.

This analysis was performed using R with the *lme4* software package (R Core Team 2019).

To assess whether the numerical impact of quolls on melomys affected the seed

harvesting rate of invaded melomys populations, we first examined the relationship

between melomys population size (estimated above) and the total number of control

(unscented) seeds harvested from each site. Here we used a simple linear model with

number of seeds harvested a linear function of population size, quoll presence/absence and

the interaction between these effects. To test whether there was an additional effect of

quoll presence, beyond their effect on population size, we defined a new variable, Δ_{ks}, as

the difference in seed take between scented and unscented treatments within each

site.session. Here any effect of melomys density is cancelled out (because density is

common to both treatments within each site.session). Thus, we fitted a model in which Δ_{ks}

is a function of quoll presence/absence, session and the interaction between these effects.

These analyses were performed using R (R Core Team 2019).

RESULTS

Effect of novel predator on survival

When we assessed the impact of quolls on melomys survival between trapping sessions the

best supported model was one in which survival rates between sessions were lower at

quoll-invaded sites than at quoll-free sites, and recapture rates were session-dependent

(Table 2). All other models were more than 4 AIC units from this best model, and so clearly

inferior descriptions of the data. From the best-supported model, estimates of apparent

survival (S) for the intervals between the capture sessions were substantially higher at

quoll-free sites ($S_{2017–2018}$ = 0.368; $S_{2018–2019}$ = 0.225) than at quoll-invaded sites ($S_{2017–2018}$ =

0.207; $S_{2018–2019}$ = 0.091; Fig. 2). The differing survival probability between sessions is

largely explained by the time difference between intervals (2017–2018 = 9 months vs.

2018–2019 = 13 months; Fig. 3).

Figure 2. Between trapping session survival ($\pm$ 95% CI) of grassland melomys (*Melomys burtoni*) on Indian

Island in quoll-invaded (*n* = 4) and quoll-free (*n* = 3) populations on Indian Island, Northern Territory, Australia.

Table 2. Results of Cormack-Jolly-Seber analyses used to compare survival (*Phi*) and recapture (*p*) probabilities

of grassland melomys (*Melomys burtoni*) on Indian Island, Northern Territory, Australia. The symbols '.' and '*t*'

refer to constant and time, respectively, while 'g' denotes the two groups (quoll free versus quolls present).

Table shows AIC values and associated AIC weights, model likelihood, number of parameters (*N*), and model

deviance. The term 'w/b' indicates that within trapping session survival rates (s_1-s_3, s_5-s_7, s_9-s_{11}) were constant

and equivalent, and different to the between trapping session survival rates (s_4, s_8). The term 'group w/b' is as

above, except that between trapping session survival rates differed between the two groups.

Model	AICc	Delta AICc	AICc Weights	Model Likelihood	N	Deviance
Phi (group w/b) p(t)	1688.629	0	0.92162	1	16	477.5224
Phi (w/b) p(t)	1693.562	4.933	0.07823	0.0849	14	486.6202
Phi (group w/b) p(g x t)	1701.116	12.4873	0.00179	0.0019	27	466.705
Phi (group w/b) p(g x t)	1703.132	14.503	0.00065	0.0007	7	510.5963
Phi (t) p (g*t)	1718.863	30.2344	0	0	32	473.631
Phi (t) p (.)	1719.862	31.2339	0	0	12	517.0641
Phi (t) p (g)	1720.751	32.1229	0	0	13	515.8843

Effect of novel predator on population size

Populations of melomys declined dramatically in quoll-invaded sites in the year following

their introduction but not in quoll-free sites (Fig. 3). We observe a strong negative

interaction between the presence of quolls and trapping session in 2018 (mean = -1.194,

95% credible interval [-1.732, -0.665]) and 2019 (mean = -1.097, 95% confidence interval [-

1.652, -0.551]; Fig. 3; Table 3).

Figure 3. Posterior mean population sizes (N_{ks} ± 95% CI) for quoll-invaded and quoll-free populations of grassland melomys (*Melomys burtoni*) on Indian Island, Northern Territory, Australia. The orange dotted vertical line denotes the timing of the introduction of quolls. The red dotted vertical line denotes the timing of an unplanned fire that burnt through the quoll-invaded sites. In each predator treatment, different sites are denoted by different shaped points. Estimates assume closure of the population within each session and detection probability that varies across sessions.

386   **Table 3.** Model parameters and their priors including prior distributions, standard deviation, estimated

387   posterior means and their 95% credible intervals. N denotes normal probability distribution with mean and

388   standard deviation.

Model Parameters				
Name for parameter	**Parameter**	**Prior (mean, SD)**	**Posterior mean**	**95% credible intervals**
Detection:				
Intercept for detection	μ_d	N (0, 2.71)	-0.94	-1.12, -0.76
Effect of session 2 on detection	t_2	N (0, 2.71)	0.59	0.33, 0.85
Effect of session 3 on detection	t_3	N (0, 2.71)	0.46	0.18, 0.73
Population size:				
Intercept for Omega	μ_P	N (0, 2.71)	-0.91	-1.22, -0.58
Quoll Presence	r_2	N (0, 2.71)	0.70	0.31, 1.09
Trapping Session 2	b_2	N (0, 2.71)	0.06	-0.36, 0.48
Trapping Session 3	b_3	N (0, 2.71)	-0.10	-0.53, 0.34
Interaction 1 [Quoll Presence * Trapping Session 2]	$\gamma_{2,2}$	N (0, 2.71)	-1.19	-1.73, -0.67
Interaction 2 [Quoll Presence * Trapping Session 3]	$\gamma_{2,3}$	N (0, 2.71)	-1.10	-1.65, -0.55

Effects of novel predator on prey behaviour

For the proportion of melomys emerging in open field tests during the 0–10 min period,

there was a significant interaction between quoll presence and trapping session (χ^2 (5) =

4.386, *P* = 0.04; Fig. 4). There was no interaction between quoll presence and trapping

session for the proportion of melomys emerging and interacting with the novel object

during 10–20 min period (χ^2 (5) = 2.567, *P* = 0.109; Fig. 4). The model without this

interaction, however, revealed a significant effect of quoll presence, with fewer melomys

emerging from hiding and interacting with the novel object during the 10–20 min period of

open field tests from sites where quolls were present than from sites where quolls were

absent (χ^2 (5) = -4.696, $P < 0.001$; Fig. 4).

Figure 4. Mean proportion (± 95% CI) of grassland melomys (*Melomys burtoni*) emerging from hiding during

open field tests from quoll-invaded sites in 2017 ($n = 16$), 2018 ($n = 28$) and 2019 ($n = 29$), and quoll-free sites

in 2017 ($n = 14$), 2018 ($n = 35$) and 2019 ($n = 24$) on Indian Island, Northern Territory, Australia.

Effects of novel predator on seed harvesting and predator-scent aversion

Although there was no interaction between melomys density and quoll presence (t_{18} = -

0.251, $P = 0.805$; Fig. 5), there was a very clear positive relationship between melomys

density and seed take ($t_{18} = 5.112$, $P < 0.001$; Fig. 5) and a clear negative relationship

between quoll presence and seed take ($t_{18} = -2.344$, $P = 0.031$; Fig. 5).

Figure 5. Effect of estimated population size on the number of control, unscented seeds removed from seed

plots (*n* = 21) in quoll-invaded and quolls-free sites. Dotted lines denote the effect of quoll presence on seed

removal rate.

When we looked at the difference in seed take (Δ_{ks}) between scent treatments

within site.session, a striking pattern emerges, in which there is a clear interaction between

the presence of quolls and session ($F_{3,17}$ = 18.61, *P* < 0.001; Fig. 6).

Figure 6. Mean ($\pm$ 95% CI) difference (Δ) between the number of predator-scented seeds and control,

unscented seeds removed by melomys from quoll-invaded (*n* = 3; 2017 & *n* = 4; 2018-19) and quoll-free (*n* = 4;

2017 & *n* = 3; 2018-19) sites during each trapping session.

DISCUSSION

The introduction of northern quolls to Indian Island was associated with lowered survival

and an apparent drop in population size in quoll-invaded melomys populations. This

numerical effect on melomys density had an impact on seed predation rates, because seed

take is strongly associated with the density of melomys in this system. This is a classic

trophic cascade: predation suppresses herbivore density, which reduces the pressure that

herbivores place on primary producers. Our study, however, also reveals an additional,

subtler, cascade effect; driven by altered prey behaviour rather than by altered prey

density.

Within months of quolls appearing on the island, invaded populations of melomys

were significantly shyer than nearby, predator-free populations of conspecifics. This rapid

but generalised response to a novel threat appears to have had a subtle effect on seed

predation rates: when we examine unscented seeds, per capita seed take is slightly lower in

quoll-invaded populations. This generalised response appears to have been supplemented

over time with more threat-specific antipredator behaviours. Although the willingness of

predator-exposed melomys to emerge from shelter (i.e. boldness) converged through time

with that of predator-free melomys, predator-exposed melomys continued to be more

neophobic than their predator-free conspecifics throughout the study. Meanwhile,

predator-scent aversion, as evidenced by seed plots, steadily increased over time.

Presumably the significant and novel predation pressure induced by the introduction of

quolls resulted in selection on behaviour and/or learning in impacted rodent populations,

allowing them to fine-tune their behavioural response (decrease general shyness, but

maintain neophobia, and respond to specific cues) as the nature of the threat became

clearer. These changing behavioural responses imply a generalised reduction in seed take

that also becomes fine-tuned over time, with high risk sites (those that smell of predators)

ultimately displaying substantially lower seed take than low risk sites. Thus, we see a

reduction in seed take that applies to the landscape becoming a fine-grained response,

varying on a spatial scale measured in the tens of metres.

Although our study documented dramatic population declines in predator-invaded

melomys populations, and we are assigning the causation of these declines to the

introduction of quolls, we need to address the confounding factors that may affect how we

interpret our results. Firstly, there is an inherent and unavoidable spatial confound in our

study system driven by the location of our study sites. We cannot exclude the possibility

that some of the population change we observe in our predator-invaded populations could

also be due to the population naturally declining towards sustainable levels unrelated to the

addition of a novel predator. It is possible that, by chance, when we started monitoring

populations of melomys, populations at northern sites were at a population peak and were

naturally cycling towards sustainable levels, while southern populations were stable.

However, although we cannot rule this out, such between population differences would be

expected to be driven by differences in resource availability between the locations (e.g.

Dickman *et al.* 1999; Russell & Ruffino 2012). We believe this is unlikely in our study system,

given the relatively close proximity of our sites (<10 km) and the spatially homogenous

climatic conditions that govern the wet-dry monsoonal tropics of northern Australia. Rodent

population cycles in the Australian wet-dry tropics appear to be primarily driven by annual

differences in rainfall between wet seasons, rather than spatial differences within years

(Madsen & Shine 1999). For this reason, we suspect natural population cycles are unlikely to

explain the population change differences we observe in this study.

Additionally, there is the unplanned, confounding factor of the fire that burnt

through northern Indian Island after completion of our population monitoring in 2017. Such

fires are commonplace in the Australian wet-dry tropics (Russell-Smith & Yates 2007); a

regular disturbance that is often rapidly offset by the annual monsoon driven wet season.

Since our sites are composed of grass-free woodland, the fire that burnt through them

mostly burnt leaf-litter (though it reached the mid-storey in other parts of the island). While

this likely reduced the short-term availability of food and cover for melomys, it is unlikely to

directly explain the demographic effects we observed. A previous study investigating the

effect of fire regimes on native mammals in savanna woodland in Kakadu National Park,

Northern Territory was unable to detect an effect of fire frequency or intensity on the

survival or recruitment of grassland melomys, despite finding fire impacts in all other co-

occurring native mammals studied (Griffiths & Brook 2015). Interestingly, even in a system

where fire is much more infrequent and significantly more intense (e.g. mesic habitats of

eastern Australia), grassland melomys were found to be relatively unaffected by a wildfire

that caused significant impacts to a co-occurring native rodent, and any demographic

impacts felt by melomys were entirely absent within months of the fire (Liedloff *et al.* 2018).

Additionally, the most dramatic behavioural difference (boldness and neophobia) between

quoll-invaded and quoll-free sites was observed immediately prior to the occurrence of the

fire (early August *vs.* mid-August 2017). For the behavioural changes we observed that were

potentially confounded by fire, such as predator-scent aversion, we would expect to see

these effects decreasing with time since fire if fire was driving this response, instead we see

the opposite trend. Finally, if food had become strongly limiting as a consequence of the

fire, we would expect to have observed an increase in seed take in the burned (quoll-

invaded) sites, instead we saw a decrease. For these reasons, we suspect the fire was

unlikely to be directly responsible for the demographic effects to melomys we observed,

and fire cannot in any way explain the response we observed to quoll-scented seeds. We,

therefore, believe our interpretation of these changes as being driven mostly by the

addition of a novel predator to the system is the most parsimonious and globally coherent

interpretation of the data.

Predation is a pervasive selective force in most natural systems, driving evolutionary

change in prey morphology, physiology, life history and behaviour. Unlike morphology and

physiology, however, the labile nature of behaviour makes it a particularly powerful trait for

rapid response in a changing world (Réale *et al.* 2007; Sih *et al.* 2010b; Dall & Griffith 2014).

Behavioural comparisons of wild populations exposed to differing predation regimes

provides some support for the prediction that reduced boldness would be selected for

under high predation scenarios (Åbjörnsson *et al.* 2004; Bell 2005; Brydges *et al.* 2008) and

that the appearance of novel predators can result in bold individuals becoming shyer

(Niemelä *et al.* 2012), however, the opposite pattern of response can also occur (Brown *et*

al. 2005; Urban 2007) or behavioural phenotypes can be unrelated to predation regime

(Laurila 2000; Carlson & Langkilde 2014). Interestingly, a number of studies have

demonstrated that individuals from high-predation areas were quicker to emerge (Harris *et*

al. 2010) and were bolder and more aggressive (Bell & Sih 2007; Dingemanse *et al.* 2007)

than predator-naïve conspecifics. Although we found the opposite pattern to this

immediately following the arrival of a novel predator, by the second year after predator

introduction we found the boldness of melomys converging with that of predator-free

populations. Thus, it is clear that the behavioural composition of these populations are

dynamic, and it seems likely this dynamism (and perhaps the capacity of the prey species to

identify specific threats) may explain some of the variation between earlier studies.

Although boldness may change over time, neophobia, as a generalised adaptive

response to predation pressure, is now well supported across a number of studies (Crane *et*

al. 2019). Individuals living under high predation risk scenarios have been shown to typically

display generalized neophobia (Brown *et al.* 2015; Elvidge *et al.* 2016), and neophobia can

increase the survival of predator-naïve individuals in initial encounters with predators

(Ferrari *et al.* 2015; Crane *et al.* 2018). Certainly, in our study, predator-exposed melomys

were significantly more neophobic than their predator-free conspecifics; an effect

maintained throughout the study.

Despite reduced survival, significant population declines, and clear behavioural

changes in invaded populations, it is impossible to determine with certainty from our data

whether changes in the behaviour of predator-invaded melomys populations are the result

phenotypic plasticity (learning) or natural selection. The low between trapping session

survival of melomys in quoll-invaded populations means few individuals survive between

sessions, so natural selection is a possibility, and selection on these behavioural traits is

potentially very strong. Although behavioural changes in predator-invaded populations

have been documented in a few systems where predator introductions have been staged

and experimentally controlled (Lapiedra *et al.* 2018; Blumstein *et al.* 2019; Cunningham *et*

al. 2019; Pringle *et al.* 2019), elucidating whether these observed changes arise because of

behavioural plasticity or natural selection can be exceptionally difficult. Rapid behavioural

responses of vulnerable prey to recovered predators has been observed in a single prey

generation, presumably due to behavioural plasticity (Berger *et al.* 2001; Cunningham *et al.*

2019). Similarly, behavioural adjustments to an introduced predator have been observed as

a result of natural selection on advantageous behavioural traits (Lapiedra *et al.* 2018). In

this study, although we had measures of individual behaviour, our between session

recapture rates of these individuals was sufficiently low that we had no longitudinal data

on the behaviour of individuals to test whether individuals were altering their behaviour or

whether natural selection was resulting in population-level change. It thus remains possible

(and quite likely) that both mechanisms were in play.

Although northern quolls represent a novel predator to melomys on Indian Island,

the two species' shared evolutionary history on the northern Australian mainland may

provide some explanation as to why this staged introduction resulted in rapid, finely-tuned

behavioural adjustment in melomys, rather than extinction. Isolation from predators can

rapidly result in the loss of antipredator behaviours from a prey species' behavioural

repertoire (Blumstein & Daniel 2005; Jolly *et al.* 2018a), dramatically increasing an

individual's susceptibility to predation following the introduction of either predator or prey

(Carthey & Banks 2014; Jolly *et al.* 2018b). But such outcomes are not inevitable: length of

isolation, co-evolutionary history, degree of predator novelty, density-dependent effects,

population size, and pre-existing predator-prey associations (Berger *et al.* 2001; Blumstein

2006; Banks & Dickman 2007; Sih *et al.* 2010a; Carthey & Banks 2014) are all likely to be

hugely influential in determining whether an invaded population adjusts to the invader or

proceeds towards extinction. Recently, a conservation introduction of Tasmanian devils to

an island previously lacking them found that their possum prey rapidly adjusted their

foraging behaviour to accommodate this newly arrived predator (Cunningham *et al.* 2019).

Despite possums having lived on the island in isolation from devils since the 1950s,

presumably, their long evolutionary history together on mainland Tasmania had them

primed to respond to this predatory archetype (Sih *et al.* 2010a; Carthey & Banks 2014;

Cunningham *et al.* 2019). This shared evolutionary history is likely responsible for both

possums' and melomys' ability to rapidly mount appropriate antipredator responses to the

introduction of these predators. The predators are novel within an individual's lifetime, but

the individual's ancestors have encountered them before.

Although our results suggest that invaded melomys populations are beginning to

adjust to the presence of northern quolls as a novel predator on Indian Island, there has

been no sign of demographic recovery from the addition of this predation pressure on the

island. Data from our seed removal experiment clearly demonstrated that the function of

melomys as seed harvesters and dispersers scales with density. Trophic cascades resulting

from the addition and loss of predators from ecosystems has been observed in a number of

systems globally (Ripple *et al.* 2001; Terborgh *et al.* 2001; Estes *et al.* 2011), and the results

can profoundly shape entire systems. As the only rodent and the dominant granivore in this

system, while melomys populations may or may not go extinct as a result of quoll invasion,

their reduced abundance and weakened ability to harvest and disperse seeds may have yet

to be observed, longer-term consequences for the vegetation structure and ecosystem

function of Indian Island (McConkey & O'Farrill 2016). Currently, grass is a rare vegetation

feature on Indian island (though it is a dominant feature of savanna woodlands generally),

and this is quite possibly a result of the high density of melomys on this (previously)

predator-free island. The presence of quolls may well change that, as both numerical and

behaviour responses of melomys cascade down to the grass community.

Empirical research on the effects of novel predators on recipient communities under

controlled conditions on a landscape-scale is exceptionally difficult and remains relatively

rare. The introduction of threatened predators to landscapes from which they have been

lost (Cunningham *et al.* 2019) or where they are entirely novel (Lapiedra *et al.* 2018),

however, provides a unique opportunity to observe how naïve prey can respond to novel

predators, and the mechanisms by which predators can structure communities. Our study

provides empirical support that some impacted prey populations can adjust rapidly to the

arrival of a novel predator via a generalised behavioural response (decreased boldness)

followed by development of a species-specific antipredator response (behavioural fine-

tuning). The arrival of the novel predator appears to have set off a trophic cascade that was

likely driven, not only by changed prey density, but also by changed prey behaviour. Thus,

rapid adaptive shift may allow prey populations to persist, but large-scale, system-wide

changes may still follow.

Acknowledgments

Thanks to Kenbi Traditional Owners (Raelene and Zoe Singh) for land access permission and

Kenbi Rangers for assistance in the field. Special thanks to Kenbi Rangers Brett Bigfoot, Rex

Edmunds, Jack Gardner, Ian McFarlane, Dale Singh, and Rex Singh for continued field

assistance throughout this project. Thanks to Kenbi Ranger Co-ordinator Steven Brown for

logistical support in the field. Thanks to Alana de Laive for graphic design of figures. Thanks

to Ella Kelly and Naomi Indigo for logistical and moral support on the island. We thank two

anonymous reviewers and Professor Réale for their comments via Peer Community in

Ecology, which greatly improved the manuscript.

Funding information

This research was funded by an Australian Research Council Linkage Grant (JKW and BP

LP150100722). Animal ethics approval for this work was provided by the University of

Melbourne (1814518) and Charles Darwin University (A13026) Animal Ethics Committees. In

kind support was provided by Kenbi Rangers and the Northern Territory Government

Department of Environment and Natural Resources, Flora and Fauna

Division (via GRG). CJJ was supported by an Australian Postgraduate Award and the

Holsworth Wildlife Research Endowment.

Data availability

The datasets generated and analysed during this study are available via figshare.com.

https://doi.org/10.26188/5dd8fe5651e8b is currently embargoed but access can be granted

on request.

REFERENCES

Åbjörnsson, K., Hansson, L.-A. & Brönmark, C. (2004). Responses of prey from habitats with

different predator regimes: local, adaptation and heritability. *Ecology*, 85, 1859–

1866.

Abrams, P.A. (2000). The evolution of predator-prey interactions: theory and evidence.

Annu. Rev. Ecol. Syst., 31, 79–105.

Andersen, A.N., Cook, G.D., Corbett, L.K., Douglas, M.M., Eager, R.W., Russell-Smith, J., *et al.*

(2005). Fire frequency and biodiversity conservation in Australian tropical savannas:

implications from the Kapalga fire experiment. *Austral Ecol.*, 30, 155–167.

Banks, P.B. & Dickman, C.R. (2007). Alien predation and the effects of multiple levels of prey

naiveté. *Trends Ecol. Evol.*, 22, 229–230.

Begg, R., Walsh, B., Woerle, F. & King, S. (1983). Ecology of *Melomys burtoni*, the grassland

melomys (Rodentia: Muridae) at Cobourg Peninsula, N.T. *Wildl. Res.*, 10, 259–267.

Bell, A.M. (2005). Behavioural differences between individuals and two populations of

stickleback (*Gasterosteus aculeatus*): behavioural syndromes. *J. Evol. Biol.*, 18, 464–

473.

Bell, A.M. & Sih, A. (2007). Exposure to predation generates personality in threespined

sticklebacks (*Gasterosteus aculeatus*). *Ecol. Lett.*, 10, 828–834.

Berger, J., Swenson, J.E. & Persson, I.-L. (2001). Recolonizing carnivores and naive prey:

conservation lessons from pleistocene extinctions. *Science*, 291, 1036–1039.

Blumstein, D.T. (2006). The multipredator hypothesis and the evolutionary persistence of

antipredator behavior. *Ethology*, 112, 209–217.

Blumstein, D.T. & Daniel, J.C. (2005). The loss of anti-predator behaviour following isolation

on islands. *Proc. R. Soc. B*, 272, 1663–1668.

Blumstein, D.T., Letnic, M. & Moseby, K.E. (2019). *In situ* predator conditioning of naive prey

prior to reintroduction. *Phil. Trans. R. Soc. B*, 374, 20180058.

Braithwaite, R. & Griffiths, A. (1994). Demographic variation and range contraction in the

northern quoll, *Dasyurus hallucatus* (Marsupialia: Dasyuridae). *Wildl. Res.*, 21, 203–

217.

Brown, C. & Braithwaite, V.A. (2004). Size matters: a test of boldness in eight populations of

the poeciliid *Brachyraphis episcopi. Anim. Behav.*, 68, 1325–1329.

Brown, C., Jones, F. & Braithwaite, V. (2005). *In situ* examination of boldness–shyness traits

in the tropical poeciliid, *Brachyraphis episcopi*. *Anim. Behav.*, 70, 1003–1009.

Brown, G.E., Elvidge, C.K., Ramnarine, I., Ferrari, M.C.O. & Chivers, D.P. (2015). Background

risk and recent experience influences retention of neophobic responses to

predators. *Behav. Ecol. Sociobiol.*, 69, 737–745.

Brown, G.E., Ferrari, M.C.O., Elvidge, C.K., Ramnarine, I. & Chivers, D.P. (2013).

Phenotypically plastic neophobia: a response to variable predation risk. *Proc. R. Soc.

B*, 280, 20122712.

Brydges, N.M., Colegrave, N., Heathcote, R.J.P. & Braithwaite, V.A. (2008). Habitat stability

and predation pressure affect temperament behaviours in populations of three-

spined sticklebacks. *J. Anim. Ecol.*, 77, 229–235.

Buckland, S.T., Burnham, K.P. & Augustin, N.H. (1997). Model selection: an integral part of

inference. *Biometrics*, 53, 603–618.

Burnham, K.P. & Anderson, D.R. (1998). *Model selection and inference: a practical

information-theoretic approach*. Springer, New York.

Carlson, B.E. & Langkilde, T. (2014). No evidence of selection by predators on tadpole

boldness. *Behaviour*, 151, 23–45.

Carter, A.J., Feeney, W.E., Marshall, H.H., Cowlishaw, G. & Heinsohn, R. (2013). Animal

personality: what are behavioural ecologists measuring? *Biol. Rev.*, 88, 465–475.

Carthey, A.J.R. & Banks, P.B. (2014). Naïveté in novel ecological interactions: lessons from

theory and experimental evidence. *Biol. Rev.*, 89, 932–949.

Courchamp, F., Chapuis, J.-L. & Pascal, M. (2003). Mammal invaders on islands: impact,

control and control impact. *Biol. Rev.*, 78, 347–383.

Crane, A.L., Brown, G.E., Chivers, D.P. & Ferrari, M.C.O. (2019). An ecological framework of

neophobia: from cells to organisms to populations. *Biol. Rev.*, 218–231.

Crane, A.L., Chivers, D.P. & Ferrari, M.C.O. (2018). Embryonic background risk promotes the

survival of tadpoles facing surface predators. *PLoS ONE*, 13, e0193939.

Cremona, T., Mella, V.S.A., Webb, J.K. & Crowther, M.S. (2015). Do individual differences in

behavior influence wild rodents more than predation risk? *J. Mammal.*, 96, 1337–

1343.

Cunningham, C.X., Johnson, C.N., Hollings, T., Kreger, K. & Jones, M.E. (2019). Trophic

rewilding establishes a landscape of fear: Tasmanian devil introduction increases

risk-sensitive foraging in a key prey species. *Ecography*, 42, 2053–2059.

Dall, S.R.X. & Griffith, S.C. (2014). An empiricist guide to animal personality variation in

ecology and evolution. *Front. Ecol. Evol.*, 2, 1–7.

Dickman, C.R., Mahon, P.S., Masters, P. & Gibson, D.F. (1999). Long-term dynamics of

rodent populations in arid Australia: the influence of rainfall. *Wildl. Res.*, 26, 389–

403.

Dielenberg, R.A. & McGregor, I.S. (2001). Defensive behavior in rats towards predatory

odors: a review. *Neurosci. Biobehav. Rev.*, 25, 597–609.

Dingemanse, N.J., Wright, J., Kazem, A.J.N., Thomas, D.K., Hickling, R. & Dawnay, N. (2007).

Behavioural syndromes differ predictably between 12 populations of three-spined

stickleback. *J. Anim. Ecol.*, 76, 1128–1138.

Doherty, T.S., Glen, A.S., Nimmo, D.G., Ritchie, E.G. & Dickman, C.R. (2016). Invasive

predators and global biodiversity loss. *Proc. Natl. Acad. Sci. USA*, 113, 11261–11265.

Elvidge, C.K., Chuard, P.J.C. & Brown, G.E. (2016). Local predation risk shapes spatial and

foraging neophobia patterns in Trinidadian guppies. *Curr. Zool.*, 62, 457–462.

Estes, J.A., Terborgh, J., Brashares, J.S., Power, M.E., Berger, J., Bond, W.J., *et al.* (2011).

Trophic downgrading of planet Earth. *Science*, 333, 301–306.

Ferrari, M.C.O., McCormick, M.I., Meekan, M.G. & Chivers, D.P. (2015). Background level of

risk and the survival of predator-naive prey: can neophobia compensate for predator

naivety in juvenile coral reef fishes? *Proc. R. Soc. B*, 282, 20142197.

Fortin, D., Beyer, H.L., Boyce, M.S., Smith, D.W., Duchesne, T. & Mao, J.S. (2005). Wolves

influence elk movements: behavior shapes a trophic cascade in Yellowstone National

Park. *Ecology*, 86, 1320–1330.

Gelman, A. & Rubin, D.B. (1992a). Inference from iterative simulation using multiple

sequences. *Stat. Sci.*, 7, 457–472.

Gelman, A. & Rubin, D.B. (1992b). [Practical Markov Chain Monte Carlo]: Rejoinder:

replication without contrition. *Stat. Sci.*, 7, 503–511.

Griffiths, A.D. & Brook, B.W. (2015). Fire impacts recruitment more than survival of small-

mammals in a tropical savanna. *Ecosphere*, 6, 1–22.

Harris, S., Ramnarine, I.W., Smith, H.G. & Pettersson, L.B. (2010). Picking personalities apart:

estimating the influence of predation, sex and body size on boldness in the guppy

Poecilia reticulata. *Oikos*, 119, 1711–1718.

Jolly, C.J., Kelly, E., Gillespie, G.R., Phillips, B. & Webb, J.K. (2018b). Out of the frying pan:

Reintroduction of toad-smart northern quolls to southern Kakadu National Park.

Austral Ecol., 43, 139–149.

Jolly, C.J., Webb, J.K., Gillespie, G.R., Hughes, N.K. & Phillips, B.L. (2019). Bias averted:

personality may not influence trappability. *Behav. Ecol. Sociobiol.*, 73, 129.

Jolly, C.J., Webb, J.K. & Phillips, B.L. (2018a). The perils of paradise: an endangered species

conserved on an island loses antipredator behaviours within 13 generations. *Biol.*

Lett., 14, 20180222.

Kelly, E. (2019). *Targeted gene flow for conservation: northern quolls and the invasive cane*

toad. PhD Thesis. University of Melbourne, Australia.

Kelly, E. & Phillips, B.L. (2016). Targeted gene flow for conservation. *Conserv. Biol.*, 30, 259–

267.

Kemper, C.M., Kitchener, D.J., Humphreys, W.F., How, R.A., Bradley, A.J. & Schmitt, L.H.

(1987). The demography and physiology of *Melomys* sp. (Rodentia: Muridae) in the

Mitchell Plateau area, Kimberley, Western Australia. *J. Zool.*, 212, 533–562.

Kery, M. & Schaub, M. (2011). *Bayesian Population Analysis using WinBUGS: A Hierarchical*

Perspective. Elsevier Science & Technology, San Diego, United States.

Lapiedra, O., Schoener, T.W., Leal, M., Losos, J.B. & Kolbe, J.J. (2018). Predator-driven

natural selection on risk-taking behavior in anole lizards. *Science*, 360, 1017–1020.

Laundre, J.W., Hernandez, L. & Ripple, W.J. (2010). The landscape of fear: ecological

implications of being afraid. *TOECOLJ*, 3, 1–7.

Laurila, A. (2000). Behavioural responses to predator chemical cues and local variation in

antipredator performance in *Rana temporaria* tadpoles. *Oikos*, 88, 159–168.

Liedloff, A.C., Wilson, J.C. & Engeman, R.M. (2018). The effect of wildfire on population

dynamics for two native small mammal species in a coastal heathland in Queensland,

Australia. *Acta Oecol.*, 88, 58–64.

Lima, S.L. & Dill, L.M. (1990). Behavioral decisions made under the risk of predation: a

review and prospectus. *Can. J. Zool.*, 68, 619–640.

López, P., Hawlena, D., Polo, V., Amo, L. & Martín, J. (2005). Sources of individual shy–bold

variations in antipredator behaviour of male Iberian rock lizards. *Anim. Behav.*, 69,

1–9.

Madsen, T. & Shine, R. (1999). Rainfall and rats: Climatically-driven dynamics of a tropical

rodent population. *Austral Ecol.*, 24, 80–89.

McConkey, K.R. & O'Farrill, G. (2016). Loss of seed dispersal before the loss of seed

dispersers. *Biol. Conserv.*, 201, 38–49.

McGregor, I.S., Schrama, L., Ambermoon, P. & Dielenberg, R.A. (2002). Not all 'predator

odours' are equal: cat odour but not 2,4,5 trimethylthiazoline (TMT; fox odour)

elicits specific defensive behaviours in rats. *Behav. Brain Res.*, 129, 1–16.

Medina, F.M., Bonnaud, E., Vidal, E., Tershy, B.R., Zavaleta, E.S., Josh Donlan, C., *et al.*

(2011). A global review of the impacts of invasive cats on island endangered

vertebrates. *Glob. Change Biol.*, 17, 3503–3510.

Moore, H.A., Dunlop, J.A., Valentine, L.E., Woinarski, J.C.Z., Ritchie, E.G., Watson, D.M., *et al.*

(2019). Topographic ruggedness and rainfall mediate geographic range contraction

of a threatened marsupial predator. *Divers. Distrib.*, 25, 1818–1831.

Niemelä, P.T., DiRienzo, N. & Hedrick, A.V. (2012). Predator-induced changes in the boldness

of naïve field crickets, *Gryllus integer*, depends on behavioural type. *Anim. Behav.*,

84, 129–135.

Oakwood, M. (1997). *The ecology of the northern quoll,* Dasyurus hallucatus*. PhD Thesis.

Australian National University, Australia.

Oakwood, M., Woinarski, J.C.Z. & Burnett, S. (2016). *Dasyurus hallucatus* (northern quoll).

The IUCN Red List of Threatened Species 2016:

e.T6295A21947321. http://dx.doi.org/10.2305/IUCN.UK.2016-

2.RLTS.T6295A21947321.en. Downloaded on 31 October 2019.

Plummer, M., Stukalov, A. & Denwood, M. (2017). rjags: Bayesian graphical models using

MCMC. *URL http://CRAN. R-project. org/package= rjags. R package version, 2, 0–4.*

Pringle, R.M., Kartzinel, T.R., Palmer, T.M., Thurman, T.J., Fox-Dobbs, K., Xu, C.C.Y., *et al.*

(2019). Predator-induced collapse of niche structure and species coexistence.

Nature, 570, 58–64.

R Core Team. (2019). *R: A Language and Environment for Statistical Computing.* R

Foundation for Statistical Computing, Vienna.

Réale, D., Reader, S.M., Sol, D., McDougall, P.T. & Dingemanse, N.J. (2007). Integrating

animal temperament within ecology and evolution. *Biol. Rev.*, 82, 291–318.

Relyea, R.A. (2003). Predators come and predators go: the reversiblity of predator-induced

traits. *Ecology*, 84, 1840–1848.

Ripple, W.J., Larsen, E.J., Renkin, R.A. & Smith, D.W. (2001). Trophic cascades among wolves,

elk and aspen on Yellowstone National Park's northern range. *Biol. Conserv.*, 102,

227–234.

Royle, J.A., Dorazio, R.M. & Link, W.A. (2007). Analysis of multinomial models with unknown

index using data augmentation. *J. Comput. Graph. Stat.*, 16, 67–85.

Russell, J.C. & Ruffino, L. (2012). The influence of spatio-temporal resource fluctuations on

insular rat population dynamics. *Proc. R. Soc. B*, 279, 767–774.

Russell-Smith, J. & Yates, C.P. (2007). Australian savanna fire regimes: context, scales,

patchiness. *Fire Ecol.*, 3, 48–63.

Sax, D., Stachowicz, J., Brown, J., Bruno, J., Dawson, M., Gaines, S., *et al.* (2007). Ecological

and evolutionary insights from species invasions. *Trends Ecol. Evol.*, 22, 465–471.

Schoener, T.W. & Spiller, D.A. (1996). Devastation of prey diversity by experimentally

introduced predators in the field. *Nature*, 381, 691–694.

Shine, R. (2010). The ecological impact of invasive cane toads (*Bufo marinus*) in Australia. *Q.*

Rev. Biol., 85, 253–291.

Sih, A. (1992). Prey uncertainty and the balancing of antipredator and feeding needs. *Am.*

Nat., 139, 1052–1069.

Sih, A., Bolnick, D.I., Luttbeg, B., Orrock, J.L., Peacor, S.D., Pintor, L.M., *et al.* (2010a).

Predator-prey naïveté, antipredator behavior, and the ecology of predator invasions.

Oikos, 119, 610–621.

Sih, A., Stamps, J., Yang, L.H., McElreath, R. & Ramenofsky, M. (2010b). Behavior as a key

component of integrative biology in a human-altered world. *Integr. Comp. Biol.*, 50,

934–944.

Sih, A., Ziemba, R. & Harding, K.C. (2000). New insights on how temporal variation in

predation risk shapes prey behavior. *Trends Ecol. Evol.*, 15, 3–4.

Simberloff, D., Martin, J.-L., Genovesi, P., Maris, V., Wardle, D.A., Aronson, J., *et al.* (2013).

Impacts of biological invasions: what's what and the way forward. *Trends Ecol. Evol.*,

28, 58–66.

Suraci, J.P., Clinchy, M., Dill, L.M., Roberts, D. & Zanette, L.Y. (2016). Fear of large carnivores

causes a trophic cascade. *Nature Comm.*, 7, 10698.

Tanner, M.A. & Wong, W.H. (1987). The calculation of posterior distributions by data

augmentation. *J. Am. Stat. Assoc.*, 82, 528–540.

Terborgh, J., Lopez, L., Nunez, P., Rao, M., Shahabuddin, G., Orihuela, G., *et al.* (2001).

Ecological meltdown in predator-free forest fragments. *Science*, 294, 1923–1926.

Urban, M.C. (2007). Risky prey behavior evolves in risky habitats. *Proc. Natl. Acad. Sci. USA*,

104, 14377–14382.

Woinarski, J.C.Z., Burbidge, A.A. & Harrison, P.L. (2015). Ongoing unraveling of a continental

fauna: decline and extinction of Australian mammals since European settlement.

Proc. Natl. Acad. Sci. USA, 112, 4531–4540.

820