Running head: Top predator changes isolation effect

Title: Top predator introduction changes the effects of spatial isolation on freshwater community structure

Authors:

Rodolfo Mei Pelinson[1,*]

Mathew A. Leibold[2]

Luis Schiesari[1,3]

[1] Programa de Pós-Graduação em Ecologia, Instituto de Biociências, Universidade de São Paulo, São Paulo, São Paulo, Brazil.

[2] Department of Biology, University of Florida, Gainesville, Florida, 32611, U.S.A.

[3] Escola de Artes, Ciências e Humanidades, Universidade de São Paulo, São Paulo, São Paulo, Brazil.

*Corresponding Author: E-mail: rodolfopelinson@gmail.com

Abstract

The importance of local selective pressures on community structure is predicted to increase

with spatial isolation when species favored by local conditions also have higher dispersal rates.

In freshwater habitats, the introduction of predatory fish can produce trophic cascades because

fish tend to prey upon intermediate predatory taxa, such as predatory insects, which indirectly

benefits herbivores and detritivores. Similarly, spatial isolation is known to limit predatory

insect's colonization rates more strongly than of herbivores and detritivores, thus generating

similar effects. Here we tested the hypothesis that the effect of introduced predatory fish on

macroinvertebrate community structure increases across a gradient of spatial isolation by

conducting a field experiment where artificial ponds with and without fish (the Redbreast

Tilapia) were constructed at three different distances from a source wetland. Overall results show

that fish do reduce the abundance of predatory insects but have no effect on the abundance of

herbivores and detritivores. Spatial isolation, however, does strengthen the trophic cascade

caused by dispersal limitation of predatory insects, but only in the absence of fish. More

importantly, macroinvertebrate communities with and without fish tend to diverge more strongly

at higher spatial isolation, however, this pattern was not due to an increase in the magnitude of

the effect of fish, as initially hypothesized, but to a change in the effect of isolation in the

presence of fish. We argue that as spatial isolation increases, suitable prey, such as predatory

insects become scarce and fish increases predation pressure upon herbivores and detritivores,

dampening the positive effect of spatial isolation on them. Our results highlight the importance

of considering interspecific variation in dispersal and multiple trophic levels to better understand

the processes generating metacommunity structures.

Keywords: metacommunity, community ecology, dispersal, isolation, macroinvertebrates,

trophic cascade, fish introduction, Tilapia, aquaculture, freshwater ecology.

Introduction

The presence of top predators has been widely recognized to cause strong changes in

species abundance patterns and community structure (*e.g.* Paine 1966, Holt et al. 1994, Leibold

1996, Frank et al. 2005, Newsome et al. 2017). Less frequently acknowledged, however, is that

such changes, as of any other local selective pressure, can be modulated by dispersal (Leibold

and Chase 2018, Guzman et al. 2019). Classic metacommunity theory recognizes that the

frequency and intensity of dispersal can modulate the relative importance of ecological drift (*i.e.*

demographic stochasticity) and niche selection processes, such as the presence of top predators,

in structuring metacommunities (Leibold et al. 2004). When species dispersal rates are low for

all species involved, or patches are isolated, stochastic events are likely to cause communities to

drift towards multiple different compositions that aren't necessarily related to these local

selective pressures (Leibold and Chase 2018). In contrast, if species dispersal rates are

sufficiently high for all species involved or patches are highly connected, the constant arrival of

migrants should override the effects of drift and niche selection making communities more

similar to each other (*i.e.* the mass-effect; Mouquet and Loreau 2003, Leibold and Chase 2018).

Thus, the presence of a top predator would be expected to cause greater change in prey

community structure at intermediate levels of connectivity. However, these predictions may not

be realized if there is interspecific variation in dispersal rates (Vellend et al. 2014, Guzman et al.

2019). Variation in dispersal can actually intensify the consequences of niche selection if the

traits that confer higher fitness within a set of local environmental conditions (*e.g.* ability to

escape predation) are positively correlated with traits that confer higher dispersal rates (Vellend

et al. 2014).

Freshwater communities are strongly influenced by the presence of predatory fish

(Hrbáček 1962, Brooks and Dodson 1965, Wellborn et al. 1996). In the absence of fish,

predatory invertebrates such as aquatic beetles and dragonfly larvae are often top predators.

Compared to fish, they are usually less efficient, gape-limited sit-and-wait predators that

consume smaller prey (Wellborn et al. 1996). However, when present, fish, which are generally

large visually-oriented predators, tend to preferentially consume large prey, which frequently

happen to be those predatory insects (Wellborn et al. 1996, McCauley 2008). Therefore, by

preferentially consuming predatory insects, fish can produce a trophic cascade that results in an

increase in the abundance of small herbivores and detritivores (Diehl 1992, Goyke and Hershey

1992).

Freshwater insects can also vary substantially in dispersal and colonization rates and such

variation is often correlated with trophic level (Bilton et al. 2001, Shulman and Chase 2007,

Chase and Shulman 2009, Shurin et al. 2009, Guzman et al. 2019). For example, predatory

insects tend to have larger body sizes than their prey, and thus higher locomotory ability

(McCann et al. 2005). However, predatory insects also tend to have smaller population sizes

(Cohen et al. 2003) and longer generation times than their prey, which makes colonization events

in spatially isolated ponds rarer (Chase and Shulman 2009). They can also be indirectly

disfavored by habitat isolation if their prey is dispersal-limited or unable to reach high population

sizes (Hein and Gillooly 2011). Insects that are herbivores and detritivores, by contrast, tend to

have shorter generation times and larger population sizes, which can increase their number of

dispersal events when compared to predatory insects (Poff et al. 2006). Moreover, the smaller

body sizes of herbivores and detritivores may greatly expand their dispersal range by wind

transport (Muehlbauer et al. 2014). Therefore, the relative decrease in colonization rates across a

gradient of spatial isolation may be stronger for predatory insects. An important outcome of this

steeper decline in colonization rates for higher trophic level insects is that spatial isolation can

promote changes in community structure that are essentially analogous to trophic cascades, as

the abundance of predators would decline and that of herbivores and detritivores increase in

more isolated habitats (Shulman and Chase 2007, Chase and Shulman 2009). Therefore, because

traits that confer lower dispersal rates to predatory insects are correlated with traits that make

them susceptible to predation by fish, spatial isolation is likely to strengthen the cascading effect

caused by the presence of fish.

Our study aimed at experimentally assessing whether and how spatial isolation can change

the effects of the introduction of a generalized fish predator on freshwater insect community

structure. We designed our study to test the following hypotheses with special attention to the

role of interspecific variation in dispersal: First, the presence of predatory fish would change

insect communities mostly as a consequence of a cascading effect by preferentially preying upon

larger predatory insects, decreasing their abundance (Fig. 1a) and causing the abundance of

herbivores and detritivores to increase (Fig. 1b). Second, spatial isolation would promote a

change in community structure due to a similar cascading effect, by reducing the abundance of

predatory insects (Fig. 1a), which can successfully reach more isolated habitats less frequently

than their prey, also increasing the abundance of herbivores and detritivores (Fig. 1b). Finally,

and third, increasing isolation should intensify the effect of fish on community structure because

the ecological traits that strengthen the cascading effects in both cases are considered positively

correlated (Fig. 1c).

Methods

We conducted our field experiment at the Estação Ecológica de Santa Bárbara (EESB) in

Águas de Santa Bárbara, São Paulo, Brazil (22º48'59" S, 49º14'12" W). The EESB is a 2,712-ha

protected area predominantly covered with open savanna habitats, with smaller portions of

seasonal semideciduous forests, *Pinus* spp., and *Eucalyptus* spp. plantations (Melo and Durigan

2011). Soils are sandy and climate is Koeppen´s Cwa, i.e., warm temperate with dry winters and

hot summers (Kottek et al. 2006). Mean annual rainfall is ~1250mm with a distinct rainy season

from October to March (January being the wettest month with ~200mm rainfall) and a dry

season from April to September (July being the driest month with ~25mm rainfall; (Hijmans et

al. 2005). In the EESB the experiment was implemented in an area covered by second-growth

cerrado *sensu stricto*, a moderately dense, open-canopy savanna habitat (Melo and Durigan

2011).

Experimental units consisted of ~1,200L artificial ponds dug into the ground and lined

with a 0.5 mm thick, high-density polyethylene geomembrane to retain water. Each pond was 4

m long, 1m wide and 40 cm deep. Walls were vertical along the length of the pond; 1 m-long

ramps terminating at ground level at each short side of the pond provided shallow microhabitats

for freshwater organisms and escape for terrestrial fauna that eventually fell into the water. Two

roof tiles were placed at the waterline in each of the short sides to provide shelter and/or

oviposition habitat. Three 30 cm-long, 10 cm-wide PVC pipes were placed in the water to

provide shelter for fishes (see Appendix S1).

Experimental design

The experiment followed a fully factorial design crossing fish presence (presence/absence)

with spatial isolation (three levels of isolation). The isolation treatment was achieved by

establishing 8 artificial ponds along each of three parallel transects 30m, 120m, and 480m from a

source wetland consisting of a stream (Riacho Passarinho) and its marshy floodplain (see

Appendix S1). Within each transect, the distance between adjacent artificial ponds was 30 m.

The artificial ponds were naturally colonized by organisms dispersing in the landscape including

phytoplankton, zooplankton, and macroinvertebrates. The chosen distances from the source

habitat and their spacing intervals represent a steep spatial isolation gradient for most aquatic

insects (Wilcox 2001, Trekels et al. 2011; see Appendix S2). The well-drained sandy soils

ensured that no other ponds and puddles formed during the rainy season at our study site, making

the near stream and its floodplain the most likely source of aquatic insects colonizing the

artificial ponds. Each fish-by-distance treatment was replicated four times for a total of 24

artificial ponds.

The experiment ran from 18-Jan-2017 to 24-Apr-2017. Between 18 and 25-Jan-2017

mesocosms were filled with well water, which does not contain already established plankton or

macroinvertebrate communities. On 28-Jan-2017 we added to each mesocosm 1000g (wet mass)

of leaf litter composed of equal amounts of grass and tree leaf litter to provide structural

complexity for benthic organisms. On 29-Jan-2017 we added to each mesocosm 15g of dog

chow to provide an initial pulse of nutrients. At the same day we added one Redbreast Tilapia

(*Coptodon rendalli*, standard length 99.2 mm ± 5.9 mm, wet mass 40.2 g ± 8.8 g, mean ± SD,

N=12) per fish treatment pond, collected in a small reservoir outside the EESB. Over the course

of the experiment, we monitored ponds for fish presence, replacing missing fish as soon as

noticed. Fish dying within the first days of the experiment, possibly due to handling stress, were

immediately replaced. In the following weeks, mesocosms water became turbid and it was not

always possible to assess fish presence without netting. Because netting could represent a

considerable disturbance to freshwater communities, we waited until the end of each sampling

survey to seine the ponds and thereby assess fish presence. Two fishes were found to be missing

by the end of the first and second surveys (see Appendix S3). No fish died between the second

and third surveys. Therefore, every pond with fish had at least 4 weeks of fish presence prior to

the final survey.

We selected the Redbreast Tilapia for three reasons. First, tilapias are generalized

omnivorous predators that, even though many times are described as having a preference for

feeding upon macrophytes (Rao et al. 2015) and zooplankton (Lazzaro 1991), can also prey upon

macroinvertebrates when available (Chimbari et al. 1997; also confirmed in a pilot laboratory

experiment we conducted, see Appendix S4). Second, they are hardy species capable of

surviving in a wide range of environmental conditions including low oxygen levels and a broad

range of temperatures (Caulton 1977, Tran-Duy et al. 2008), conditions likely to be found in our

shallow artificial ponds. Third, we understand that the spatial context at which natural and

constructed ponds occur is important when assessing the impacts of invasive Tilapias on

freshwater communities. The Redbreast Tilapia is, along with the Nile Tilapia (*Oreochromis*

niloticus), one of the most widely introduced fishes in the world for aquaculture and recreational fisheries (Britton and Orsi 2012). These African species represented ~11% (6.1 million tons) of the entire freshwater fish production in the world and ~40% (0.6 million tons) in the Americas in 2017 (FAO 2019). In Brazil, Redbreast and Nile Tilapias are found in reservoirs and lakes in most river basins, and their spread to new river basins may be a matter of time considering that their stocking is still encouraged by public policies (Zambrano et al. 2011, Britton and Orsi 2012, Pelicice et al. 2014, Daga et al. 2016). Indeed, a very common land management practice in rural Brazil is the construction of dugout or impounded lakes, where the Tilapia is usually the first choice of fish species for stocking.

Freshwater community sampling surveys

To assess the influence of fish presence, spatial isolation, and their interaction on community assembly we conducted three sampling surveys of the artificial ponds after ~3 weeks (18 to 23-Feb-2017), ~8 weeks (23 to 27-Mar-2017), and ~12 weeks (20 to 24-Apr-2017) of the experiment. Freshwater communities were dominated by insects, which were sampled by sweeping half of the pond twice, including both pelagic and benthic habitats, with a hand net (mesh size 1.5 mm). Samples were cleaned of debris and stored in 70% ethanol. We identified and counted all aquatic macroinvertebrates to the lowest reliable taxonomical level using taxonomic keys for South American freshwater insects (Costa et al. 2004, Pereira et al. 2007, Segura et al. 2011, Hamada et al. 2014).

Data analysis

We tested whether the presence of fish, spatial isolation, sampling survey, and their interaction caused any changes in total abundance of predatory insects, and herbivores and detritivores separately through sequential likelihood ratio tests. We modeled species total abundance using generalized linear mixed models (*i.e.* GLMMs) with a negative binomial

distribution, and pond identifications as a random effect term. We also performed pairwise *post-*

hoc tests with Šidák adjustment of p-values for factors with more than two levels, and for

interactions, when they were important in explaining abundance patterns.

Prior to the analysis of community structure, we removed species that occurred in less than

two ponds considering each sampling survey (*i.e.* singletons and doubletons) both because they

are uninformative to general community patterns and because they complicate model parameter

estimation (Warton et al. 2015a). To test the hypothesis that community structure is influenced

by the presence of fish, spatial isolation, and their interaction, we performed sequential

likelihood ratio tests on multivariate community data (Warton et al. 2015b) assuming that species

abundances come from a negative binomial distribution due to observed overdispersion in the

data. The main advantage of this approach, in opposition to a perMANOVA, is the possibility of

accounting for the mean-variance relationship of abundance data, and the better interpretability

of change on community structure by looking at coefficients of the effect of treatments on

individual species and their respective confidence intervals. Additionally, we can test for the

effect of traits on species responses to treatments (*i.e.* model-based fourth corner solution; Brown

et al. 2014, Warton et al. 2015a). Thus, we also tested whether trophic level (*i.e.* strict predator

VS herbivores and detritivores) is a good predictor of the changes in community structure

through likelihood ratio tests. Note here that this analysis differs from tests of total abundance

because it tests for similar responses to treatments of species in each trophic level, whereas total

abundances tend to show patterns that are driven by the more abundant species. A shortcoming

of this approach lies in the difficulties of modeling correlations in species abundances (Warton et

al. 2015a, 2015b). To account for those correlations when computing p-values for the entire

communities, we performed 10,000 bootstraps resamples, using the PIT-trap bootstrap resample

procedure (Warton et al. 2017), shuffling entire rows of the incidence matrix (ponds), keeping

species abundances in the same ponds always together. To account for lack of independence

between the same ponds sampled across time when analyzing all surveys together, ponds were

considered blocks, so in each permutation step we shuffled ponds freely within blocks (*i.e.* only

across time), then we shuffled the entire bocks freely (*i.e.* across fish and isolation treatments).

Because we found significantly different effects of fish and isolation treatments across different

sampling surveys, we repeated the analysis in each sampling surveys separately.

Due to inevitable practical constraints, one limitation of our experimental design is that the

experimental units were not fully spatially isolated from each other within the isolation

treatments, meaning that they were not fully independent from each other. Therefore, we also

tested for the presence of spatial dependence among experimental units using Moran's

Eigenvector Maps (MEMs) (Borcard and Legendre 2002, Dray et al. 2006). We did not find

important effects of neighbor communities on community structures (see Appendix S5 for

details).

In the community structure analysis, a significant interaction between fish and isolation

means that there is either or both a difference in direction (*i.e.* positive or negative effect) or

magnitude of the effect of fish in different isolation treatments. To specifically test for

differences in the size of the effect of fish, regardless of direction, we performed a model-based

unconstrained ordination via generalized linear latent variable models (GLLVM; Niku et al.

2017) with a negative binomial distribution using two latent variables for each of the sampling

surveys (Hui et al. 2015). The latent variables were estimated via variational approximation (Hui

et al. 2017). Here the latent variables are interpreted as the axis of any ordination analysis, such

as non-metric multidimensional scaling (*i.e.* nMDS) (Hui et al. 2015). After performing the

ordination, we computed the centroids of each treatment group and the distance between the

centroids of fish and fishless treatments in each isolation treatment as a measure of the size of the

effect of fish (black bars in Fig. 3). Then we tested whether this distance is significantly different

across all the isolation treatments. To test for that we designed a permutation procedure to only

permute ponds across isolation treatments, keeping the fish treatment constant. This represented

a null scenario where the effect of fish is the same in all isolation treatments. For this analysis,

we corrected p-values for multiple comparisons using the false discovery rate correction (FDR).

We also used those ordinations to visualize the effect of treatments on community structure.

We had to exclude samples from four ponds in the third survey due to mislabeling,

therefore we analyzed 24 ponds for the first and second surveys and 20 for the third one. In the

last survey treatments with fish in 30 m, 120 m, and 480 m, and without fish in 480 m, had only

three replicates (see Appendix S4). All analyses were implemented in software R version 4.0.2.

(R Core Team 2020). Code and data to reproduce the analyses and figures are available as an R

package on github in the following repository: RodolfoPelinson/PredatorIsolationComm.

Results

Total Abundance

Detailed information about the taxa sampled across the experiment is available in

Appendix S6. The total abundance of predatory insects increased over the course of the

experiment, decreased in the presence of fish, and decreased with spatial isolation (Table 1; Fig.

2a). The total abundance of herbivores and detritivores also increased throughout the experiment,

but fish did not affect it. Spatial isolation, however, did increase the total abundance of

herbivores and detritivores, but only in the last survey. This effect was stronger for fishless

ponds (Table 1; Fig. 2b).

Community Structure

The effect of treatments on community structure became clearer and stronger over time

(Table 2). We, therefore, will focus on the results of the last (third) survey for clarity.

Community structure was generally affected by the interaction of the presence of fish and spatial

isolation. More importantly, this interactive effect of treatments was significantly explained by

trophic level (Table 2; Fig. 3). Predatory insects had similar negative responses to the presence

of fish, but only in 120 and 480 m (Fig. 4a). The general lack of this effect at 30 m was mostly

because *Pantala* dragonflies suffered no effect of fish and *Orthemis* dragonflies were positively

affected by it at this distance (Fig. 3; see Appendix S9). Predatory insects also had coherent

negative responses to isolation from 30 to 120 and 480 m distances, but only in ponds with fish

(Fig. 4b and 4c). Different from observed for total abundance, in fishless ponds, some predatory

insects, such as *Pantala* and *Orthemis* dragonflies, were positively affected (Fig. 3; see Appendix

S9). Herbivores and detritivores only responded as a group to the presence of fish at the

intermediate spatial isolation, and this response, contrary to what was expected, was negative

(Fig. 4a). However, they were positively affected by isolation from 30 m to 120 m and 480 m,

but only in fishless ponds (Fig. 4c).

When we tested for differences in the distance between centroids of fish and fishless

communities for the third survey, we found that it was larger at 480 m than at 30 m (Difference

in distance between centroids: 1.63; adj. p: 0.038; Fig. 3) meaning that, as we expected, fish and

fishless ponds are more different from each other in more isolated ponds. The distance between

centroids at 120 m was not significantly different from 30 m (Difference in distance between

centroids: 1.25; adj. p: 0.098; Fig. 3) or 480 m (Difference in distance between centroids: 0.38;

adj. p: 0.601; Fig. 3).

Discussion

Generally, both the presence of fish and spatial isolation had important effects on

freshwater community structure and responses to treatments were different for different trophic

levels. More importantly, our results, as expected, diverged from classic metacommunity models,

which predicts that the effect of local niche selection on community structure should be stronger

at intermediate levels of isolation (Mouquet and Loreau 2003, Leibold et al. 2004, Leibold and

Chase 2018). However, even though we show that the difference between communities with and

without fish only increases with isolation, it does not happen because of a positive correlation in

the responses of predatory insects and herbivores and detritivores to the presence of fish and

isolation.

The presence of fish indeed shifted species composition through a reduction of predatory

insects, notably dytiscid beetles, notonectids, and dragonflies, all relatively large taxa (see

Appendix S9). Contrary to our expectations and previous work (*e.g.* Goyke and Hershey 1992),

however, it did not translate into an increase in abundance of herbivores and detritivores.

Previous work has shown that even though generalist predatory fish frequently prey upon

multiple trophic levels, its effect on predatory insects tend to be disproportionally larger, as they

are easier to catch and have smaller population sizes (Diehl 1992, Batzer et al. 2000). Thus, the

net effect of fish on the abundance of herbivores and detritivores is a sum of both direct negative

and indirect positive effects (Batzer et al. 2000). In our experiment, the direct negative effect of

fish predation upon herbivores and detritivores might have balanced out the indirect positive

effect of the reduced abundance of predatory insects.

298   We initially hypothesized that the abundance of predatory insects should decrease as

spatial isolation increases, and the abundance of their prey should increase in response. We

partially found support for this hypothesis. The total abundance of predatory insects does

decrease as spatial isolation increases, however, not all predatory insects appear to be dispersal

limited. Predatory insects taxa only consistently negatively responded to isolation when fish was

present. When it was absent, there were two clear exceptions to this pattern: *Pantala* and

Orthemis dragonflies. While other predators, such as diving beetles (*i.e. Rhantus*) and water

striders (*i.e. Microvelia*), exhibited a strong decrease in abundance in both fish and fishless

ponds as spatial isolation increased, both of these dragonflies were positively affected by

isolation in the absence of fish. Dragonflies are known to have a great dispersal ability; thus it is

not surprising that at the scale of our experiment they would suffer small or no negative effects at

all of isolation (McCauley 2006). We believe that the positive effects observed might be a net

consequence of the absence of other predatory insects in more isolated ponds, which could have

released those dragonflies from competition or made these ponds more attractive for adults to lay

their eggs. Such effect did not happen in ponds where fish was present likely because fish

increased predation pressure upon dragonflies as other suitable prey became scarce. This might

also explain why *Orthemis* dragonflies consistently had higher abundances in ponds with fish in

low isolation treatments, compared to fishless ponds. It is possible that the higher availability of

more suitable prey (*i.e.* dytiscid beetles and notonectids) in low isolation decreased predation

rate upon *Orthemis*, allowing it to have a greater abundance in ponds with fish. Indeed, some

dragonfly species are known to exhibit different vulnerability to predation depending on body

size and antipredatory behavior, allowing them to coexist with fish (Johnson 1991, McPeek

1998, De Marco et al. 1999, Johansson 2000, Hopper 2001, McCauley 2008).

The reduction in total abundance of predatory insects across the isolation gradient indeed

had effects that are analogous to a trophic cascade, causing the abundance of herbivores and

detritivores to increase, but this effect was only clear in fishless ponds both considering total

abundance and community structure patterns. We argue that the most plausible explanation for

this pattern lies in the fact that, because the general abundance of predatory insects decreases in

more isolated ponds, predation pressure by fish upon herbivores and detritivores may also

increase with isolation, again, counteracting the indirect positive effect of the reduction in the

abundance of predatory insects.

Our main hypothesis was that the effect of fish would be stronger as spatial isolation

increases, causing fish and fishless communities to have greater differences in structure at more

isolated ponds (Fig. 1c). We did find this pattern (Fig. 3), however, we initially predicted it

would be a consequence of a reinforced effect of fish by spatial isolation. As discussed above,

herbivores and detritivores did not have similar responses to isolation and presence of generalist

fish. Instead, when generalist fish was present, it dampened the indirect positive effects of

isolation. Additionally, not all predatory insects were limited by dispersal, which made the ones

that are good dispersers to thrive in more isolated ponds when fish was absent. Therefore, we

argue that the greater differences between communities with and without fish in more isolated

ponds are because the presence of fish changes the net effects of spatial isolation on community

structure. When fish is absent, spatial isolation generally favors the abundance of aquatic insects

with higher dispersal rates, regardless of their trophic level. When fish is present, spatial

341 isolation has negative effects on the abundance of predatory insects, and a weaker positive effect

342 on the abundance of herbivores and detritivores. These different paths of change in structure

343 were the cause of greater differences among fish and fishless ponds in our higher isolation

344 treatments. Furthermore, those patterns could be generated by either consumptive (*i.e.* direct

345 predation upon available prey) or non-consumptive effects of predators upon preys, like habitat

346 selection (*i.e.* avoidance of ponds with fish or high density of predatory insects; see Binckley and

347 Resetarits 2005, Blaustein et al. 2005, Resetarits 2005), still, it does not change the general

348 conclusions of this work, as both processes represent a negative local selective pressure of

349 predators upon preys.

350  Finally, our results diverge both from simplistic metacommunity models, which consider

351 similar dispersal rates and single trophic levels (Mouquet and Loreau 2003, Leibold et al. 2004,

352 Leibold and Chase 2018), and slightly more complex ones considering variation in dispersal

353 rates (Vellend et al. 2014). We argue that the complexity of our study system, both in terms of

354 interspecific variability in dispersal rates and complex indirect effects involved in trophic

355 interactions, are the main causes of such divergence. Nonetheless, we were able to show that

356 spatial isolation can cause effects that are similar to trophic cascades on freshwater communities,

357 and that the presence of a generalist top predator can dampen those effects. Our results also

358 highlight the importance of considering interspecific variation in dispersal rates and multiple

359 trophic levels to better understand the processes generating community and metacommunity

360 structures in spatially structured landscapes (see Van De Meutter et al. 2007, Vellend et al. 2014,

361 Hill et al. 2017, Guzman et al. 2019).

363 **Acknowledgments**

364  We thank the EESB staff for assistance in pond construction and Luis Vicente Cavalaro,

365 Bianca Valente, Fernanda Simioni, Débora Negrão, Jessika Akane and Suzana Marte for

366 assistance in the community sampling surveys. We thank Tadeu Siqueira and Paulo Inácio Prado

for conceptual and statistical advice, Victor Saito and Erika Shimabukuro for assistance in the

identification of aquatic insects, Renata Pardini and Daniel Lahr for providing lab and office

space, Paulo Roberto Guimarães Junior and Paulo De Marco Júnior for insightful comments on

early versions of the manuscript; The members of the Schiesari lab and Leibold lab for criticism;

and Giselda Durigan for introducing us to the EESB. This study was funded by The São Paulo

Research Foundation (FAPESP, grant #2014/16320-7, Carlos Navas PI, Luis Schiesari Associate

Researcher; FAPESP, grant #2015/18790-3; Luiz Antonio Martinelli PI, Luis Schiesari Co-PI)

and National Council for Scientific and Technological Development (CNPq, grant

#458796/2014-0). RMP was supported by Ph.D. fellowships from FAPESP (grants

#2017/04122-4 and #2018/07714-2) and Coordination of Superior Level Staff Improvement

(CAPES). Experiments were conducted in EESB under the authorization from Instituto Florestal

(COTEC 553/2017) and collection permits from Instituto Chico Mendes de Conservação da

Biodiversidade (ICMBIo 17559-6), following protocols approved by the Research Ethics

Committee of the School of Arts, Sciences and Humanities of the University of São Paulo

(CEUA 003/2016).

References

Batzer, D. P., C. R. Pusateri, and R. Vetter. 2000. Impacts of fish predation on marsh

invertebrates: Direct and indirect effects. Wetlands 20:307–312.

Bilton, D. T., J. R. Freeland, and B. Okamura. 2001. Dispersal in freshwater invertebrates.

Annual Review of Ecology and Systematics 32:159–181.

Binckley, C. A., and W. J. Resetarits. 2005. Habitat selection determines abundance, richness

and species composition of beetles in aquatic communities. Biology Letters 1:370–374.

Blaustein, L., J. Blaustein, and J. Chase. 2005. Chemical detection of the predator Notonecta

irrorata by ovipositing Culex mosquitoes. Journal of Vector Ecology 30:299.

Borcard, D., and P. Legendre. 2002. All-scale spatial analysis of ecological data by means of

principal coordinates of neighbour matrices. Ecological Modelling 153:51–68.

Britton, J. R., and M. L. Orsi. 2012. Non-native fish in aquaculture and sport fishing in Brazil:
economic benefits versus risks to fish diversity in the upper River Paraná Basin. Reviews
in Fish Biology and Fisheries 22:555–565.

Brooks, J. L., and S. I. Dodson. 1965. Predation, body size, and composition of plankton.
Science 150:28–35.

Brown, A. M., D. I. Warton, N. R. Andrew, M. Binns, G. Cassis, and H. Gibb. 2014. The fourth-
corner solution - using predictive models to understand how species traits interact with
the environment. Methods in Ecology and Evolution 5:344–352.

Caulton, M. S. 1977. The effect of temperature on routine metabolism in Tilapia rendalli
boulenger. Journal of Fish Biology 11:549–553.

Chase, J. M., and R. S. Shulman. 2009. Wetland isolation facilitates larval mosquito density
through the reduction of predators. Ecological Entomology 34:741–747.

Chimbari, M. J., H. Madsen, and J. Ndamba. 1997. Simulated Field Trials to Evaluate the Effect
of Sargochromis codringtoni and Tilapia rendalli on Snails in the Presence and Absence
of Aquatic Plants. Journal of Applied Ecology 34:871.

Cohen, J. E., T. Jonsson, and S. R. Carpenter. 2003. Ecological community description using the
food web, species abundance, and body size. Proceedings of the National Academy of
Sciences 100:1781–1786.

Costa, J. M., L. O. I. de Souza, and B. B. Oldrini. 2004. Chave para identificação das famílias e
gêneros das larvas conhecidas de Odonata do Brasil: comentários e registros
bibliográficos (Insecta, Odonata). Publicações avulsas do Museu Nacional:1–44.

Daga, V., T. Debona, V. Abilhoa, É. A. Gubiani, and J. Vitule. 2016. Non-native fish invasions
of a Neotropical ecoregion with high endemism: a review of the Iguaçu River. Aquatic
Invasions 11:209–223.

De Marco, P., A. O. Latini, and A. P. Reis. 1999. Environmental determination of dragonfly
assemblage in aquaculture ponds. Aquaculture Research 30:357–364.

Diehl, S. 1992. Fish predation and benthic community structure: the role of omnivory and habitat

complexity. Ecology 73:1646–1661.

Dray, S., P. Legendre, and P. R. Peres-Neto. 2006. Spatial modelling: a comprehensive

framework for principal coordinate analysis of neighbour matrices (PCNM). Ecological

Modelling 196:483–493.

FAO. 2019. FAO - Fisheries and Aquaculture Information and Statistics Branch.

http://www.fao.org/fishery/statistics/en.

Frank, K. T., B. Petrie, J. S. Choi, and W. C. Leggett. 2005. Trophic cascades in a formerly cod-

dominated ecosystem. Science 308:1621–1623.

Goyke, A. P., and A. E. Hershey. 1992. Effects of fish predation on larval chironomid (Diptera:

Chironomidae) communities in an arctic ecosystem. Hydrobiologia 240:203–211.

Guzman, L. M., R. M. Germain, C. Forbes, S. Straus, M. I. O'Connor, D. Gravel, D. S.

Srivastava, and P. L. Thompson. 2019. Towards a multi-trophic extension of

metacommunity ecology. Ecology Letters 22:19–33.

Hamada, N., J. L. Nessimian, and R. B. Querino. 2014. Insetos aquáticos na Amazônia brasileira:

taxonomia, biologia e ecologia. Editora do INPA, Manaus.

Hein, A. M., and J. F. Gillooly. 2011. Predators, prey, and transient states in the assembly of

spatially structured communities. Ecology 92:549–555.

Hijmans, R. J., S. E. Cameron, J. L. Parra, P. G. Jones, and A. Jarvis. 2005. Very high resolution

interpolated climate surfaces for global land areas. International Journal of Climatology

25:1965–1978.

Hill, M. J., J. Heino, I. Thornhill, D. B. Ryves, and P. J. Wood. 2017. Effects of dispersal mode

on the environmental and spatial correlates of nestedness and species turnover in pond

communities. Oikos 126:1575–1585.

Holt, R. D., J. Grover, and D. Tilman. 1994. Simple rules for interspecific dominance in systems

with exploitative and apparent competition. The American Naturalist 144:741–771.

Hopper, K. R. 2001. Flexible antipredator behavior in a dragonfly species that coexists with
different predator types. Oikos 93:470–476.

Hrbáček, J. 1962. Species composition and the amount of zooplankton in relation to the fish
stock. ČSAV, Prague.

Hui, F. K. C., S. Taskinen, S. Pledger, S. D. Foster, and D. I. Warton. 2015. Model-based
approaches to unconstrained ordination. Methods in Ecology and Evolution 6:399–411.

Hui, F. K. C., D. I. Warton, J. T. Ormerod, V. Haapaniemi, and S. Taskinen. 2017. Variational
approximations for generalized linear latent variable models. Journal of Computational
and Graphical Statistics 26:35–43.

Johansson, F. 2000. The slow-fast life style characteristics in a suite of six species of odonate
larvae. Freshwater Biology 43:149–159.

Johnson, D. M. 1991. Behavioral ecology of larval dragonflies and damselflies. Trends in
Ecology and Evolution 6:8–13.

Kottek, M., J. Grieser, C. Beck, B. Rudolf, and F. Rubel. 2006. World Map of the Köppen-
Geiger climate classification updated. Meteorologische Zeitschrift:259–263.

Lazzaro, X. 1991. Feeding convergence in South American and African zooplanktivorous
cichlids Geophagus brasiliensis and Tilapia rendalli. Environmental Biology of Fishes
31:283–293.

Leibold, M. A. 1996. A Graphical Model of Keystone Predators in Food Webs: Trophic
regulation of abundance, incidence, and diversity patterns in communities. The American
Naturalist 147:784–812.

Leibold, M. A., and J. M. Chase. 2018. Metacommunity Ecology. Princeton University Press,
Princeton, New Jersey.

Leibold, M. A., M. Holyoak, N. Mouquet, P. Amarasekare, J. M. Chase, M. F. Hoopes, R. D.
Holt, J. B. Shurin, R. Law, D. Tilman, M. Loreau, and A. Gonzalez. 2004. The
metacommunity concept: a framework for multi-scale community ecology. Ecology
Letters 7:601–613.

McCann, K. S., J. B. Rasmussen, and J. Umbanhowar. 2005. The dynamics of spatially coupled

474   food webs. Ecology Letters 8:513–523.

475   McCauley, S. J. 2006. The effects of dispersal and recruitment limitation on community structure

476   of odonates in artificial ponds. Ecography 29:585–595.

477   McCauley, S. J. 2008. Slow, fast and in between: Habitat distribution and behaviour of larvae in

478   nine species of libellulid dragonfly. Freshwater Biology 53:253–263.

479   McPeek, M. A. 1998. The consequences of changing the top predator in a food web: a

480   comparative experimental approach. Ecological Monographs 68:1–23.

481   Melo, A. C. G., and G. Durigan. 2011. Estação Ecológica de Santa Bárbara Plano de Manejo.

482   Page Secretaria do Meio Ambiente. Secretaria do Meio Ambiente.

483   Mouquet, N., and M. Loreau. 2003. Community patterns in source-sink metacommunities. The

484   American Naturalist 162:544–557.

485   Muehlbauer, J. D., S. F. Collins, M. W. Doyle, and K. Tockner. 2014. How wide is a stream?

486   Spatial extent of the potential "stream signature" in terrestrial food webs using meta-

487   analysis. Ecology 95:44–55.

488   Newsome, T. M., A. C. Greenville, D. Ćirović, C. R. Dickman, C. N. Johnson, M. Krofel, M.

489   Letnic, W. J. Ripple, E. G. Ritchie, S. Stoyanov, and A. J. Wirsing. 2017. Top predators

490   constrain mesopredator distributions. Nature Communications 8:15469.

491   Niku, J., D. I. Warton, F. K. C. Hui, and S. Taskinen. 2017. Generalized linear latent variable

492   models for multivariate count and biomass data in ecology. Journal of Agricultural,

493   Biological, and Environmental Statistics 22:498–522.

494   Paine, R. T. 1966. Food web complexity and species diversity. The American Naturalist 100:65–

495   75.

496   Pelicice, F. M., J. R. S. Vitule, D. P. L. Junior, M. L. Orsi, and A. A. Agostinho. 2014. A serious

497   new threat to Brazilian freshwater rcosystems: The Naturalization of Nonnative Fish by

498   Decree. Conservation Letters 7:55–60.

Pereira, D. L. V., A. L. De Melo, and N. Hamada. 2007. Chaves de identificação para famílias e

gêneros de gerromorpha e nepomorpha (Insecta: Heteroptera) na Amazônia Central.

Neotropical Entomology 36:210–228.

Poff, N. L., J. D. Olden, N. K. M. Vieira, D. S. Finn, M. P. Simmons, and B. C. Kondratieff.

2006. Functional trait niches of North American lotic insects: traits-based ecological

applications in light of phylogenetic relationships. Journal of the North American

Benthological Society 25:730–755.

R Core Team. 2020. R: A language and environment for statistical computing. Page R

Foundation for Statistical Computing. Vienna, Austria.

Rao, W., J. Ning, P. Zhong, E. Jeppesen, and Z. Liu. 2015. Size-dependent feeding of

omnivorous Nile tilapia in a macrophyte-dominated lake: implications for lake

management. Hydrobiologia 749:125–134.

Resetarits, W. J. 2005. Habitat selection behaviour links local and regional scales in aquatic

systems. Ecology Letters 8:480–486.

Segura, M. O., F. Valente-Neto, and A. A. Fonseca-Gessner. 2011. Chave de famílias de

coleoptera aquáticos (Insecta) do estado de são paulo, Brasil. Biota Neotropica 11:411–

412.

Shulman, R. S., and J. M. Chase. 2007. Increasing isolation reduces predator:prey species

richness ratios in aquatic food webs. Oikos 116:1581–1587.

Shurin, J. B., K. Cottenie, and H. Hillebrand. 2009. Spatial autocorrelation and dispersal

limitation in freshwater organisms. Oecologia 159:151–159.

Tran-Duy, A., J. W. Schrama, A. A. van Dam, and J. A. J. Verreth. 2008. Effects of oxygen

concentration and body weight on maximum feed intake, growth and hematological

parameters of Nile tilapia, Oreochromis niloticus. Aquaculture 275:152–162.

Trekels, H., F. Van de Meutter, and R. Stoks. 2011. Habitat isolation shapes the recovery of

aquatic insect communities from a pesticide pulse. Journal of Applied Ecology 48:1480–

1489.

Van De Meutter, F., L. De Meester, and R. Stoks. 2007. Metacommunity structure of pond

macroinvertebrates: Effects of dispersal mode and generation time. Ecology 88:1687–

1695.

Vellend, M., D. S. Srivastava, K. M. Anderson, C. D. Brown, J. E. Jankowski, E. J. Kleynhans,

N. J. B. Kraft, A. D. Letaw, A. A. M. Macdonald, J. E. Maclean, I. H. Myers-Smith, A.

R. Norris, and X. Xue. 2014. Assessing the relative importance of neutral stochasticity in

ecological communities. Oikos 123:1420–1430.

Warton, D. I., F. G. Blanchet, R. B. O'Hara, O. Ovaskainen, S. Taskinen, S. C. Walker, and F.

K. C. Hui. 2015a. So many variables: Joint modeling in community ecology. Trends in

Ecology and Evolution 30:766–779.

Warton, D. I., S. D. Foster, G. De'ath, J. Stoklosa, and P. K. Dunstan. 2015b. Model-based

thinking for community ecology. Plant Ecology 216:669–682.

Warton, D. I., L. Thibaut, and Y. A. Wang. 2017. The PIT-trap — A "model-free" bootstrap

procedure for inference about regression models with discrete, multivariate responses.

PLoS ONE 12:e0181790.

Wellborn, G. A., D. K. Skelly, and E. E. Werner. 1996. Mechanisms creating community

structure across a freshwater habitat gradient. Annual Review of Ecology and

Systematics 27:337–363.

Wilcox, C. 2001. Habitat size and isolation affect colonization of seasonal wetlands by predatory

aquatic insects. Israel Journal of Zoology 47:459–476.

Zambrano, L., E. Martínez-Meyer, N. Menezes, and A. T. Peterson. 2011. Invasive potential of

common carp (Cyprinus carpio) and Nile tilapia (Oreochromis niloticus) in American

freshwater systems. Canadian Journal of Fisheries and Aquatic Sciences.

Table 1. Summary of likelihood ratio tests for total abundances of predatory insects and herbivores and detritivores. Bold values represent a

significant effect (p < 0.05).

	Residual Degrees of Freedom	Diff. of Degrees of Freedom	Chi-square	p	*Post-hoc* comparisons
Predatory insects					
Survey	**63**	**2**	**62.012**	**<0.001**	First < Second, First < Third, Second = Third
Fish	**62**	**1**	**12.752**	**<0.001**	
Isolation	**60**	**2**	**19.026**	**<0.001**	30 > 120, 30 > 480, 120 = 480
Survey : Fish	58	2	5.028	0.081	
Survey : Isolation	54	4	4.217	0.377	
Fish : Isolation	52	2	5.153	0.076	
Survey : Fish : Isolation	48	4	8.589	0.072	
Herbivores and detritivores					
Survey	**63**	**2**	**70.3627**	**<0.001**	First < Second < Third
Fish	62	1	0.3095	0.578	
Isolation	60	2	2.5255	0.283	
Survey : Fish	58	2	2.8277	0.243	
Survey : Isolation	**54**	**4**	**11.2050**	**0.024**	First: 30 = 120 = 480; Second: 30 = 120 = 480; Third: 30 = 120, 120 = 480, 30 < 480
Fish : Isolation	**52**	**2**	**6.1092**	**0.047**	Fishless: 30 < 120, 30 < 480, 120 = 480; Fish: 30 = 120 = 480
Survey : Fish : Isolation	48	4	3.6439	0.456	

553 **Table 2.** Summary of likelihood ratio tests for community structure considering all sampling

554 surveys together and separately. Bold values represent a significant effect ($p < 0.05$).

	Residual Degrees of Freedom	Diff. of Degrees of Freedom	Deviance	p
All Sampling Surveys				
Survey	**57**	**2**	**392.8**	**<0.001**
Fish	**56**	**1**	**89.2**	**<0.001**
Isolation	**54**	**2**	**109.2**	**0.001**
Fish : Isolation	**52**	**2**	**120.3**	**<0.001**
Survey : Fish	50	2	50.9	0.075
Survey : Isolation	46	4	78.5	0.247
Survey : Fish : Isolation	**42**	**4**	**81.2**	**0.014**
1st Sampling Survey				
Fish	22	1	19.01	0.104
Isolation	20	2	24.38	0.398
Fish : Isolation	**18**	**2**	**42.60**	**0.024**
(Fish : Isolation) :Trophic Level	197	5	19.26	0.106
2nd Sampling Survey				
Fish	**22**	**1**	**62.28**	**0.002**
Isolation	**20**	**2**	**71.81**	**0.025**
Fish : Isolation	**18**	**2**	**72.15**	**0.016**
(Fish : Isolation) :Trophic Level	**404**	**5**	**33.74**	**0.002**
3rd Sampling Survey				
Fish	**18**	**1**	**49.09**	**0.018**
Isolation	16	2	72.96	0.055
Fish : Isolation	**14**	**2**	**91.12**	**0.015**
(Fish : Isolation) :Trophic Level	**332**	**5**	**33.71**	**0.028**

Figure Legends

Figure 1. Graphical representation describing our initial hypotheses. A: Both fish and spatial

isolation are predicted to have negative effects on the abundance of predatory insects. This is

expected because fish are visually oriented predators, thus they should preferentially prey upon

large-bodied predatory insects. Similarly, even though predatory insects often have higher

dispersal ability, their dispersal rates are lower than of herbivores and detritivores because they

have lower population sizes, thus fewer dispersal events. B: In contrast and as a consequence of

the reduction in the abundance of predatory insects, the abundance of herbivores and detritivores

are predicted to increase in the presence of fish and as spatial isolation increases. C: If species

vary in dispersal rates and good dispersers are favored by local conditions, as shown in B, or bad

dispersers are disfavored by local conditions, as shown in A, the importance of niche selection

processes (*e.g.* the effect of fish) may increase with spatial isolation, increasing the differences

between communities with and without fish (black bars). C is a hypothetical ordination plot

where each symbol represents a community and similar symbols are communities from the same

treatment.

Figure 2. Total abundance of predatory insects (A) and herbivores and detritivores (B) for the

last survey. Plots for the first and second surveys are provided in Appendix S7.

Figure 3. Unconstrained ordination showing pond communities (symbols) and species (filled

circles) for the third survey. Yellow filled circles represent predatory insects and green filled

circles represent herbivores and detritivores. The sizes of the circles are proportional to

maximum body size (the volume of the largest individual of each species in a log-scale).

Abbreviations of names of taxa are provided in Appendix S6. Symbols close to each other have

more similar community structures than those that are further apart. Species are positioned closer

to treatments where their abundance is higher. Note that herbivores and detritivores are more

strongly associated with high isolation treatments, especially fishless ponds, whereas predatory

insects are associated with low isolation treatments, especially fishless ponds. Note also that

Pantala and *Orthemis* dragonflies (Pan and Ort) are detached from the other predatory insects,

Pantala being closer to highly isolated fishless treatments and *Orthemis* to ponds with fish in

low isolation. Black bars show the distance between fish and fishless treatment's centroids in

each isolation treatment. Similar ordination plots for the first and second surveys are provided in

Appendix S8.

Figure 4. 95% Confidence intervals for the effect of fish and isolation on the abundance of

predatory insects and herbivores and detritivores when comparing pairs of treatments for the

third survey of the experiment. Confidence intervals not crossing the zero hatched line were

considered significant effects and colored; blue bars represent positive and red bars negative

responses to a given treatment when compared to a reference treatment. A are effects of the

presence of fish in each isolation treatment. B are effects of isolation in fishless ponds and C in

ponds with fish. In each of the B and C panels, we show the effects of increasing isolation from

30 to 120 m, from 30 to 480 m, and from 120 m to 480 m

Figure 1.

Figure 2.

Figure 3.

Figure 4.