Title: Top predator introduction changes the effects of spatial isolation on freshwater community structure

Authors: Rodolfo Mei Pelinson[1,*], Mathew A. Leibold[2], Luis Schiesari[1,3]

[1] Programa de Pós-Graduação em Ecologia, Instituto de Biociências, Universidade de São Paulo, São Paulo, São Paulo, Brazil.

[2] Department of Biology, University of Florida, Gainesville, Florida, 32611, U.S.A.

[3] Escola de Artes, Ciências e Humanidades, Universidade de São Paulo, São Paulo, São Paulo, Brazil.

*Corresponding Author:
e-mail: rodolfopelinson@gmail.com
Tel: +55 17 981270165

Keywords: metacommunity, community ecology, dispersal, isolation, macroinvertebrates, trophic cascade, fish introduction, Tilapia, aquaculture, freshwater ecology

Statement of authorship: RMP and LS conceived the project and designed the research. RMP collected the data, ran the analysis and wrote the first draft of the paper. MAL and LS contributed to revisions and conceptual manuscript development. All authors gave approval for publication.

Number of words in Abstract: 150

Number of words in main text: 4302

Number of references: 61

Number of figures: 6

Number of Tables: 0

Abstract

Introducing top predators in a community can produce cascading effects due to predation upon larger, frequently intermediate predatory taxa, and indirectly benefit consumers. Similarly, spatial isolation can limit predator's dispersal rates more strongly than that of their prey, thus generating similar trophic cascades. We tested the effects of spatial isolation on the outcome and strength of the cascading effects caused by fish, as our top predator, on macroinvertebrate community structure. Overall results show that fish did reduce the abundance of predatory insects but had no effect on herbivores and detritivores. Spatial isolation caused trophic cascades, but only in the absence of fish. A potential explanation to this is because fish also preyed upon consumers when they increased in abundance. In contrast with traditional theoretical expectations for metacommunities, we found that the difference between communities with and without fish increased with isolation, because fish dampened the cascading effects of spatial isolation.

Introduction

Dispersal is a key driver of community and metacommunity structure (Mouquet & Loreau

2003; Leibold *et al.* 2004; Vellend 2010; Leibold & Chase 2018). Classic metacommunity theory

recognizes that the frequency and intensity of dispersal can determine the relative importance of

ecological drift (*i.e.* stochasticity) and niche selection in structuring metacommunities (Leibold

et al. 2004). When species dispersal rates are low for all species involved, or patches are isolated,

stochastic events are likely to cause communities to drift towards multiple different compositions

that aren't necessarily related to local environmental conditions (Leibold & Chase 2018). In

contrast, if species dispersal rates are sufficiently high for all species involved or patches are

highly connected, the constant arrival of migrants should override the effects of drift or niche

selection making communities more similar to each other irrespective of local environmental

variation (Mouquet & Loreau 2003; Leibold & Chase 2018). Thus, niche selection processes are

more likely to shape community structure at intermediate levels of dispersal rates or connectivity

(Fig. 1A). However, these predictions may not be fully realized if there is also interspecific

variation in dispersal rates among species (Vellend *et al.* 2014; Guzman *et al.* 2019). Variation in

dispersal rates can reduce the effect of stochasticity in community structure because species with

higher dispersal rates will predictably colonize a patch, increasing the predictability of the

subsequent community assembly process (Vellend *et al.* 2014). Additionally, interspecific

variation in dispersal rates can actually intensify the consequences of niche selection if traits that

confer higher fitness within a set of local environmental conditions are positively correlated with

traits that confer higher dispersal frequency or ability (Vellend *et al.* 2014; Fig. 1B).

Freshwater pond species can vary substantially in dispersal and colonization rates (Bilton

et al. 2001; Shulman & Chase 2007; Chase & Shulman 2009; Shurin *et al.* 2009; Guzman *et al.*

2019) and such variation is often correlated with trophic level. For example, predatory insects

tend to have larger body sizes their prey, and thus higher locomotory ability (McCann *et al.*

2005). However, predatory insects also tend to have smaller population sizes (Cohen *et al.* 2003)

and longer generation times than their prey, possibly making colonization events in spatially

isolated ponds rarer (Chase & Shulman 2009). Predators can also be indirectly disfavored by

habitat isolation if their prey is dispersal-limited or unable to reach high population sizes (Hein

& Gillooly 2011). Non-predatory insects, by contrast, have shorter generation times and larger

population sizes, which can increase the number of dispersal events (Poff *et al.* 2006). Moreover,

the smaller body sizes of non-predatory insects may greatly expand their dispersal range by wind

transport (Muehlbauer *et al.* 2014). An important outcome of this negative correlation between

dispersal rate and trophic level is that spatial isolation can lead to trophic cascades, causing

herbivores and detritivores, hereafter referred to as consumers, to be more abundant in more

isolated habitats (Shulman & Chase 2007; Chase & Shulman 2009).

Freshwater community structure is also strongly influenced by the presence of predatory

fish (Hrbáček 1962; Brooks & Dodson 1965; Wellborn *et al.* 1996). In the absence of fish,

predatory invertebrates such as aquatic beetles and dragonfly larvae are often the top predators.

Compared to fish they are usually less efficient, gape-limited sit-and-wait predators that consume

smaller prey (Wellborn *et al.* 1996). However, when present, fish, which are generally large

visually oriented predators, tend to preferentially consume large prey, which frequently happens

to be predatory insects (Wellborn *et al.* 1996; McCauley 2008). This can lead to trophic cascades

like those caused by spatial isolation, leading to an increase in the abundance of small herbivores

and detritivores (Diehl 1992; Goyke & Hershey 1992). Therefore, the consequences of the

presence of fish for aquatic macroinvertebrate communities could be even stronger in more

isolated habitats.

Our study aimed at experimentally assessing whether and how spatial isolation can change

the effects of the introduction of a generalized fish predator on freshwater community structure.

We designed our study to test the following hypotheses with special attention to the role of

interspecific variation in dispersal: First, the presence of predatory fish would promote a

cascading effect by preferentially preying upon larger predatory insects, causing the abundance

of consumers (*i.e.* herbivores and detritivores) to increase. Second, spatial isolation would

promote a similar cascading effect by reducing the abundance of predatory insects, which

frequently have lower dispersal rates, also increasing the abundance of consumers. Finally and

third, increasing isolation should intensify the effect of fish on community structure because the

ecological traits that promote trophic cascades in both cases are positively correlated (Fig. 1B).

We show that the effect of fish not only inceases as ponds become more isolated, it actually

change. We show that because generalist fish also pray upon consumers, it can inhibit the trophic

cascade promoted by spatial isolation, causing communities with and without fish to diverge in

structure as spatial isolation increase.

Methods

We conducted our field experiment at the Estação Ecológica de Santa Bárbara (EESB) in

Águas de Santa Bárbara, São Paulo, Brazil (22º48'59" S, 49º14'12" W). The EESB is a 2,712-ha

protected area predominantly covered with open savanna habitats, with smaller portions of

seasonal semideciduous forests, *Pinus* sp., and *Eucalyptus* sp plantations (Melo & Durigan

2011). Soils are sandy and climate is Koeppen´s Cwa, i.e., warm temperate with dry winters and

hot summers (Kottek *et al.* 2006). Mean annual rainfall is ~1250mm with a distinct rainy season

from October to March (January being the wettest month with ~200mm rainfall) and a dry

season from April to September (July being the driest month with ~25mm rainfall; (Hijmans *et*

al. 2005). In the EESB the experiment was implemented in an area covered by second-growth

cerrado *sensu stricto*, a moderately dense, open-canopy savanna habitat (Melo & Durigan 2011).

Experimental units consisted of ~1,200L artificial ponds dug into the ground and lined

with a 0.5 mm thick, high-density polyethylene geomembrane to retain water. Each pond was 4

m long, 1m wide and 40 cm deep. Walls were vertical along the length of the pond; 1 m-long

ramps terminating at ground level at each short side of the pond provided shallow microhabitats

for freshwater organisms and escape for terrestrial fauna that eventually fell into the water. Two

roof tiles were placed at the waterline in each of the short sides to provide shelter and/or

oviposition habitat. Three 30 cm-long, 10 cm-wide PVC pipes were placed in the water to

provide shelter for fishes.

Experimental design

The experiment followed a fully factorial design crossing fish presence (presence/absence)

with spatial isolation (three levels of isolation). The isolation treatment was achieved by

establishing 8 artificial ponds along each of three parallel transects 30m, 120m, and 480m from a

source wetland consisting of a stream (Riacho Passarinho) and its floodplain (Fig. 2). Within

each transect, the distance between adjacent artificial ponds was 30 m (see Appendix S1 in

Supporting Information) for possible implications of our spatial design). The artificial ponds

were naturally colonized by organisms dispersing in the landscape including phytoplankton,

zooplankton, and macroinvertebrates. The chosen distances from the source habitat and their

spacing intervals represent a steep spatial isolation gradient for most aquatic insects (Wilcox

2001; Trekels *et al.* 2011); see Appendix S2). The well-drained sandy soils ensured that no other

ponds and puddles formed during the rainy season at our study site, making the near stream and

its floodplain the main source of aquatic insects colonizing the artificial ponds. Each fish-by-

distance treatment was replicated four times for a total of 24 artificial ponds.

The experiment ran from 18-Jan-2017 to 24-Apr-2017. Between 18 and 25-Jan-2017

mesocosms were filled with well water, which is unlikely to contain already established plankton

or macroinvertebrate communities. On 28-Jan-2017 we added to each mesocosm 1000g (wet

mass) of leaf litter composed of equal amounts of grass and tree leaf litter to provide structural

complexity for benthic organisms. On 29-Jan-2017 we added to each mesocosm 15g of dog

chow to provide an initial pulse of nutrients. The same day we added one Redbreast Tilapia

(*Coptodon rendalli*, standard length 99.2 mm ± 5.9 mm, wet mass 40.2 g ± 8.8 g, mean ± SD,

N=12) per predator treatment pond, collected in a small reservoir outside the EESB. Over the

course of the experiment, we monitored ponds for fish survival; dead fish were replaced as soon

as noticed (see Appendix S3).

We selected Redbreast Tilapia for three reasons. First, tilapias are generalized omnivorous

predators that, even though many times described as having a preference for feeding upon

macrophytes (Rao *et al.* 2015) and zooplankton (Lazzaro 1991), can also prey upon

macroinvertebrates when they are available (Chimbari *et al.* 1997); also confirmed in a pilot

laboratory experiment we conducted, see Appendix S4). Second, they are hardy species capable

of surviving in a wide range of environmental conditions including low oxygen levels and a

broad range of temperature (Caulton 1977; Tran-Duy *et al.* 2008), conditions likely to be found

in our shallow artificial ponds. Third, the Redbreast Tilapia is, along with the Nile Tilapia

(*Oreochromis niloticus*), one of the most widely introduced fishes in the world for aquaculture

and recreational fisheries (Britton & Orsi 2012). These African species represented ~11% (6.1

million tons) of the entire freshwater fish production in the world and ~40% (0.6 million tons) in

the Americas in 2017 (FAO 2019). In Brazil, Redbreast and Nile Tilapias are found in reservoirs

and lakes in most river basins, and their spread to new river basins may be a matter of time

considering that their stocking is still encouraged by public policies (Zambrano *et al.* 2006;

Britton & Orsi 2012; Pelicice *et al.* 2014; Daga *et al.* 2016). Indeed, a very common land

management practice in rural Brazil is the construction of dugout or impounded lakes, where the

Tilapia is usually the first choice of fish species for stocking.

Freshwater community sampling surveys

To assess the influence of fish presence, spatial isolation, and their interaction on

community assembly we conducted three sampling surveys of the artifitial ponds after ~3 weeks

(18 to 23-Feb-2017), ~8 weeks (23 to 27-Mar-2017) and ~12 weeks (20 to 24-Abr-2017) of the

experiment. Freshwater communities were dominated by insects, which were sampled by

sweeping half of the pond twice, including both pelagic and benthic habitats, with a hand net

(mesh size 1.5 mm). Samples were cleaned of debris and stored in 70% ethanol. We identified

and counted all aquatic macroinvertebrates to the lowest reliable taxonomical level using

taxonomic keys for South American freshwater insects (Costa *et al.* 2004; Pereira *et al.* 2007;

Segura *et al.* 2011; Hamada *et al.* 2014).

Data analysis

To test the hypothesis that community structure is influenced by fish presence, distance to

the source, and their interaction, we used a model-based approach for multivariate data where the

matrix of site-by-raw species abundance data represents community structures (Warton *et al.*

2015b). The main advantages of model-based approaches are the possibility of accounting for the

mean-variance relationship of abundance data, and the better interpretability of data.

Specifically, we can assess which and how species in a community are being influenced by

treatments and test for the effect of traits in their responses (Warton *et al.* 2015a). Prior to this

analysis, we removed species that occurred in less than three ponds considering each sampling

survey (*i.e.* singletons and doubletons) both because they are uninformative to general

community patterns and because they complicate model parameter estimation (Warton *et al.*

2015a).

To test for differences in the effect of treatments on species abundances across different

sampling surveys, we performed likelihood ratio tests to test if the progressive addition of terms

to the model provided a statistically better fit based on values of deviance. We modeled

abundances using a negative binomial distribution because of overdispersion (Appendix S5).

First, we tested if we had any effect of time in species abundances, then we tested if

progressively adding the effect of fish, isolation, and their interaction provided a significantly

better fit to the data. Those tests were always assessed by comparing the best model so far

against the next more complex model. After that, we tested if the effect of treatments were

different across different sampling surveys by adding an interaction between the effect of time

and the interaction between fish and isolation treatments. Because we found significantly

different effects of fish and isolation treatments across different sampling surveys, we repeated

the analysis in each sampling surveys separately. To account for correlations in species

abundances when computing p-values we shuffled entire rows of the incidence matrix (ponds),

keeping species abundances in the same ponds always together. To account for lack of

independence between the same ponds sampled across time, ponds were considered blocks, so in

each permutation step we shuffled ponds freely within blocks (*i.e.* only across time), then we

shuffled the entire bocks freely (*i.e.* across fish and isolation treatments). P-values were

computed using the PIT-trap bootstrap resample procedure, which operates on probability

integral transform residuals (Warton *et al.* 2017). P-values were obtained from 10,000 bootstraps

resamples.

To assess how individual taxa respond to the different treatments we looked at 95%

confidence intervals of estimated parameters for each taxon in each treatment. To see if the

trophic level (*i.e.* strict predator VS consumer) is a good predictor of the changes in community

structure, we further tested if the inclusion of the categorical trait predator/consumer increases

the fit of the models also performing likelihood ratio tests. We considered strict predators only

the predators that were prone to prey upon other sampled macroinvertebrates. Insects that are not

predators at all, that prey mostly upon zooplankton, or that have only a small portion of their diet

based on predation were considered consumers (see Appendix S6). This trait-based approach is

called the model-based fourth corner solution (Brown *et al.* 2014).

A significant interaction between fish and isolation means that there is either or both a

difference in direction (*i.e.* positive or negative effect) or magnitude of the effect of fish in

different isolation treatments. To specifically test for differences in the size of the effect of fish,

regardless of direction, we performed a model-based unconstrained ordination via generalized

linear latent variable models (GLLVM; Niku *et al.* 2017) with a negative binomial distribution

using two latent variables for each of the sampling surveys (Hui *et al.* 2015). The latent variables

were estimated via variational approximation (Hui *et al.* 2017). After performing the ordination,

we computed the centroids of each treatment group, and the distance between the centroids of

fish and fishless treatments in each isolation treatment as a measure of the size of the effect of

fish. Then we tested whether this distance is significantly different across all the isolation

treatments. To test for that we designed a permutation procedure to only permute ponds across

isolation treatments, keeping the fish treatment constant. This represented a null scenario where

the effect of fish is the same in all isolation treatments. We corrected p-values for multiple

comparisons using the false discovery rate (FDR). We also used those ordinations to visualize

the effect of treatments on community structure.

All analyses were implemented in software R version 3.6.1 (R Core Team 2019), using

packages 'mvabund' version 4.0.1 (Wang *et al.* 2012) and 'gllvm' version 1.1.7 (Niku *et al.*

2019). Code and data to reproduce the analysis and figures are available on github in the

following repository: RodolfoPelinson/Pelinson_et_al_Top_predator_isolation.

Results

Mesocosms were colonized by aquatic and semiaquatic insects comprising five orders and

17 families (Odonata: Libellulidae, Coenagrionidae; Coleoptera: Dytiscidae, Hydrophilidae,

Noteridae; Diptera: Ceratopogonidae, Chaoboridae, Chironomidae, Culicidae; Ephemeroptera:

Baetidae, Caenidae, Polymitarcyidae; Hemiptera: Corixidae, Gerridae, Naucoridae,

Notonectidae, Veliidae). The most abundant families, with more than 200 individuals sampled in

the entire experiment, were chironomids (total abundance of 11,558 individuals), veliids (1,097

individuals), libellulid dragonflies (901 individuals) and culicid mosquitoes (707 individuals).

The rarest families, with less than 10 individuals sampled in the entire experiment, were

Gerridae, Naucoridae, Corixidae, Coenagrionidae, Ceratopogonidae, and Hydrophilidae. More

detailed information is available in Appendix S6.

Effects of treatments on community structure became clearer and stronger over time. We

therefore will only show the results of the last (third) survey for clarity. Results for all surveys

are available in Appendix S7. The inclusion of the fish treatment significantly increased the fit of

the explanatory statistical model to community data (Difference of degrees of freedom: 1;

Deviance: 49.09; p: 0.018). The subsequent inclusion of the distance treatment only marginally

improved its fit (Difference of degrees of freedom: 2; Deviance: 72.96; p: 0.056), however, the

inclusion the interactive effect of distance treatments and the fish treatments highly improved the

fit of the model with only the fish treatment (Difference of degrees of freedom: 4; Deviance:

164.1; p: 0.009). More importantly, this interactive effect of treatments was significantly

explained by trophic level (Difference of degrees of freedom: 5; Deviance: 33.71; p: 0.028; Fig.

3).

Presence of fish had a negative impact on the abundance of predators only at 120 and 480

m (Fig. 4A). The general lack of this negative effect on predators at 30 m was mostly because

Pantala dragonflies suffered no effect of fish and *Orthemis* dragonflies were positively affected

by it at this distance (Fig. 5A). Different from what we expected, consumers were not generally

positively affected by fish at the end of the experiment (Fig. 4A).

Spatial isolation negatively affected predators from 30 to 120 and 480 m distances, but

only in ponds with fish (Fig. 4B and C). In fishless ponds, some predators were negatively

affected by isolation, while others, such as *Pantala* and *Orthemis* dragonflies, were positively

affected (Fig. 5A). Consumers, on the other hand, were, as expected, positively affected by

isolation from 30 m to 120 m and 480 m of isolation, but only in fishless ponds (Fig. 4C).

When we tested for differences in the size of the effect of fish on community structure, we

found that the effect of fish at 480 m was larger than at 30 m (Difference in distance between

centroids: 1.63; adj. p: 0.038; Fig. 3) meaning that, as we expected, fish and fishless ponds are

more different from each other in more isolated ponds. The size of the effect of fish at 120 m was

not significantly different from 30 m (Difference in distance between centroids: 1.25; adj. p:

0.098; Fig. 3) or 480 m (Difference in distance between centroids: 0.38; adj. p: 0.601; Fig. 3).

Discussion

Generally, both the presence of fish and spatial isolation had important effects on

freshwater community structure and responses to treatments were different for different trophic

levels. More importantly, our results, as expected, diverged from classic metacommunity models,

which predicts that the effect of environmental filtering and local interactions on community

structure should be stronger at intermediate levels of isolation (Mouquet & Loreau 2003; Leibold

et al. 2004; Leibold & Chase 2018). However, even though we show that the difference between

ponds with and without fish only increases with isolation (Fig. 3), it does not happen because of

a positive correlation in the responses of predatory insects and consumers to the presence of fish

and isolation.

We found that the presence of fish indeed shifts species composition through a reduction of

predatory insects. The most negatively affected taxa were dytiscids beetles, notonectids, and

dragonfly larvae, all of them relatively large taxa (see Appendixes S6). However, contrary to our

expectations and previous work (*e.g.* Goyke & Hershey 1992), consumers were not positively

affected by the presence of fish. We believe that this lack of positive effect of fish on consumers

comes from the fact that tilapias are generalist omnivores that might prey not only upon most

conspicuous prey (*i.e.* predatory insects), but also any available prey, including herbivores and

detritivores. In a pilot experiment (Appendix S4), tilapias readily ate not only large conspicuous

prey such as *Scinax* tadpoles, guppies, belostomatids, nepids and dragonflies, but also much

smaller ones, such as damselflies and mosquito larvae, the latter having about 5 orders of

magnitude less wet mass than the manipulated tilapias. Similar results were observed by (Diehl

1992) when the presence of juvenile perch, a generalist benthic predator, caused biomass of

invertebrate predators to decrease, but did not affect the biomass of consumers, because it was

also preying upon them. Therefore, the same process might have occurred in our experiment

where consumer abundances might suffer the net effects of direct negative effects of predation

by fish, and indirect positive effects caused by the reduction of predatory insects also by fish (*i.e.*

trophic cascade; Fig. 6B).

Orthemis dragonflies were the only exception for the expected negative effects of fish on

predatory insects. They consistently had higher abundance in ponds with fish, but only in low

isolation treatments. Some dragonfly species are known to exhibit different vulnerability to

predation depending on body size and flexible antipredatory behavior, allowing them to coexist

with fish (Johnson 1991; McPeek 1998; Johansson 2000; Hopper 2001; McCauley 2008).

Indeed, *Orthemis* larvae were one of the smallest benthic predators we had in our experiment

(around 20 times smaller than *Pantala*; see Appendix S6) and it has been reported to occur in

high abundance in aquaculture ponds with no vegetation, usually buried into the sediment (de

Marco *et al.* 1999). It is thus possible that higher availability of more suitable prey (*i.e.* other

predatory insects) in low isolation decreased predation rate on *Orthemis*, allowing it to have a

greater abundance in ponds with fish.

We initially hypothesized that the abundance of dispersal-limited predatory insects should

decrease as spatial isolation increases, and the abundance of consumers should increase in

response, as a cascading effect. We partially found support for this hypothesis. Most of the

substantial changes in community structure that we observed because of spatial isolation were

from low (30 m) to higher levels of isolation (120 m and 480 m). Spatial isolation only generally

reduced predatory insect abundance when fish was present, and the positive effect of isolation on

consumer abundance was only observed when it was absent. The lack of a negative effect of

isolation on predatory insects in fishless ponds might be explained by two clear exceptions to the

expected patterns for predatory insects: *Pantala* and *Orthemis* dragonflies. While other

predators, such as diving beetles (*i.e. Rhantus*) and water striders (*i.e. Microvelia*), exhibited a

strong decrease in abundance in both fish and fishless ponds with spatial isolation, both of these

dragonflies were positively affected by isolation in fishless ponds. Dragonflies are known to be

good dispersers, thus it is not surprising that at the scale of our experiment they would suffer

small or no negative effects at all of the isolation distances (McCauley 2006). We believe that

the positive effects observed might be a consequence of the absence of other predatory insects in

more isolated ponds, which could have released those dragonflies from competition or made

these ponds more attractive for adults to lay their eggs (Fig. 6A). Such effect did not happen in

ponds where fish was present, likely because fish increased predation pressure upon dragonflies

as other suitable prey became scarce. Similarly to these dragonflies, if fish indeed prey upon both

consumers and predatory insects, predation pressure upon consumers would be expected to

increase with increasing isolation, explaining why we do not observe the cascading effect upon

consumers when fish is present. Indeed, if we consider predatory insect biomass as an estimate

of food availability for fishes, we observe that average insect biomass per fishless pond

decreases about 17% from low to intermediate and high isolation (19.2 g in low isolation to 15.9

g in intermediate and high isolation; see Appendix S8).

Finally, we found that more isolated communities have a greater difference in community

structure between fish and fishless ponds then less isolated ones. However, we initially predicted

this pattern would be due to correlated responses of predators and consumers to both presence of

fish and spatial isolation (Vellend *et al.* 2014; Guzman *et al.* 2019), which was not what we

observed. We did not find fish to have a stronger similar effect on more spatially isolated

communities, instead, we argue that the presence of fish changed the effects of spatial isolation,

making communities with and without fish to diverge with increasing isolation. We observed

that because fish preys upon both predatory insects and consumers, it can dampen the cascading

effect of spatial isolation on insect consumers by increasing predation pressure upon them in the

absence of more suitable prey (Fig. 6). Also, the presence of fish can dampen any further indirect

positive effects that spatial isolation might have on predatory insects that are less dispersal

limited (Fig. 6). Our results could be due to either consumptive (*i.e.* direct predation upon

available prey) or non-consumptive effects of predators upon preys (*i.e.* avoidance of ponds with

fish or high density of predatory insects through habitat selection; see Binckley & Resetarits

2005; Blaustein *et al.* 2005; Resetarits 2005), which does not change the general conclusions of

this work, as both processes represent a negative selective pressure of predators upon preys.

Conclusion

Our results diverged from those expected for more simplistic metacommunity models (Mouquet & Loreau 2003; Leibold *et al.* 2004; Leibold & Chase 2018; Fig. 1). We argue that the higher complexity of our study system, both in terms of interspecific variability in dispersal rates and responses to the local selective pressure are the main causes of such divergence. Nonetheless, we were able to show that the presence of a generalist top predator can dampen the trophic cascades caused by spatial isolation when consumers have higher dispersal rates than predators (Fig. 6). Our results also highlight the importance of considering interspecific variation in dispersal rates and multiple trophic levels to better understand the processes generating community and metacommunity structures in spatially structured landscapes (see van de Meutter *et al.* 2007; Vellend *et al.* 2014; Hill *et al.* 2017; Guzman *et al.* 2019).

Acknowledgments

We thank the EESB staff for assistance in pond construction and Luis Vicente P. Cavalaro, Bianca S. Valente, Fernanda Simioni, Débora Negrão, Jessika Akane and Suzana Marte for assistance in the community sampling surveys. We thank Tadeu Siqueira and Paulo Inácio Prado for conceptual and statistical advice, Victor Saito and Erika Shimabukuro for assistance in the identification of aquatic insects, Renata Pardini and Daniel Lahr for providing lab and office space, Paulo Roberto Guimarães Junior, Paulo De Marco Júnior for insightful comments on early versions of the manuscript; The members of the Schiesari lab and Leibold lab for criticism; and Giselda Durigan for introducing us to the EESB. This study was funded by Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP, grant #2014/16320-7, Carlos Navas PI, Luis Schiesari Associate Researcher; FAPESP, grant #2015/18790-3; Luiz Antonio Martinelli PI,

Luis Schiesari Co-PI) and Conselho Nacional de Desenvolvimento Científico e Tecnológico

(CNPq, grant #458796/2014-0). RMP was supported by Ph.D. fellowships from FAPESP (grants

#2017/04122-4 and #2018/07714-2) and Coordenação de Aperfeiçoamento de Pessoal de Nível

Superior (CAPES). Experiments were conducted in EESB under the authorization from Instituto

Florestal (COTEC 553/2017) and collection permits from Instituto Chico Mendes de

Conservação da Biodiversidade (ICMBIo 17559-6), following protocols approved by the

Research Ethics Committee of the School of Arts, Sciences and Humanities of the University of

São Paulo (CEUA 003/2016).

References

Bilton, D.T., Freeland, J.R. & Okamura, B. (2001). Dispersal in freshwater invertebrates. *Annual*
Review of Ecology and Systematics, 32, 159–181.

Binckley, C.A. & Resetarits, W.J. (2005). Habitat selection determines abundance, richness and
species composition of beetles in aquatic communities. *Biology Letters*, 1, 370–374.

Blaustein, L., Blaustein, J. & Chase, J. (2005). Chemical detection of the predator Notonecta
irrorata by ovipositing Culex mosquitoes. *Journal of Vector Ecology*, 30, 299.

Britton, J.R. & Orsi, M.L. (2012). Non-native fish in aquaculture and sport fishing in Brazil:
Economic benefits versus risks to fish diversity in the upper River Paraná Basin. *Reviews in*
Fish Biology and Fisheries, 22, 555–565.

Brooks, J.L. & Dodson, S.I. (1965). Predation, body size, and composition of plankton. *Science*,
150, 28–35.

Brown, A.M., Warton, D.I., Andrew, N.R., Binns, M., Cassis, G. & Gibb, H. (2014). The fourth-

corner solution - using predictive models to understand how species traits interact with the

environment. *Methods in Ecology and Evolution*, 5, 344–352.

Caulton, M.S. (1977). The effect of temperature on routine metabolism in Tilapia rendalli

boulenger. *Journal of Fish Biology*, 11, 549–553.

Chase, J.M. & Shulman, R.S. (2009). Wetland isolation facilitates larval mosquito density

through the reduction of predators. *Ecological Entomology*, 34, 741–747.

Chimbari, M.J., Madsen, H. & Ndamba, J. (1997). Simulated Field Trials to Evaluate the Effect

of Sargochromis codringtoni and Tilapia rendalli on Snails in the Presence and Absence of

Aquatic Plants. *The Journal of Applied Ecology*, 34, 871.

Cohen, J.E., Jonsson, T. & Carpenter, S.R. (2003). Ecological community description using the

food web, species abundance, and body size. *Proceedings of the National Academy of*

Sciences of the United States of America, 100, 1781–1786.

Costa, J.M., Souza, L.O.I. de & Oldrini, B.B. (2004). Chave para identificação das famílias e

gêneros das larvas conhecidas de Odonata do Brasil: comentários e registros bibliográficos

(Insecta, Odonata). *Publicações Avulsas do Museu Nacional*, 99, 1–44.

Daga, V.S., Debona, T., Abilhoa, V., Gubiani, É.A. & Vitule, J.R.S. (2016). Non-native fish

invasions of a Neotropical ecoregion with high endemism: A review of the Iguaçu River.

Aquatic Invasions, 11, 209–223.

Diehl, S. (1992). Fish predation and benthic community structure: the role of omnivory and

habitat complexity. *Ecology*, 73, 1646–1661.

FAO. (2019). *FAO - Fisheries and Aquaculture Information and Statistics Branch*. Available at:

http://www.fao.org/fishery/statistics/en. Last accessed 30 September 2019.

Goyke, A.P. & Hershey, A.E. (1992). Effects of fish predation on larval chironomid (Diptera:

Chironomidae) communities in an arctic ecosystem. *Hydrobiologia*, 240, 203–211.

Guzman, L.M., Germain, R.M., Forbes, C., Straus, S., O'Connor, M.I., Gravel, D., *et al.* (2019).

Towards a multi-trophic extension of metacommunity ecology. *Ecology Letters*, 22, 19–33.

Hamada, N., Nessimian, J.L. & Querino, R.B. (2014). *Insetos aquáticos na Amazônia brasileira:*

taxonomia, biologia e ecologia. Editora do INPA, Manaus.

Hein, A.M. & Gillooly, J.F. (2011). Predators, prey, and transient states in the assembly of

spatially structured communities. *Ecology*, 92, 549–555.

Hijmans, R.J., Cameron, S.E., Parra, J.L., Jones, P.G. & Jarvis, A. (2005). Very high resolution

interpolated climate surfaces for global land areas. *International Journal of Climatology*,

25, 1965–1978.

Hill, M.J., Heino, J., Thornhill, I., Ryves, D.B. & Wood, P.J. (2017). Effects of dispersal mode

on the environmental and spatial correlates of nestedness and species turnover in pond

communities. *Oikos*, 126, 1575–1585.

Hopper, K.R. (2001). Flexible antipredator behavior in a dragonfly species that coexists with

different predator types. *Oikos*, 93, 470–476.

Hrbáček, J. (1962). *Species composition and the amount of zooplankton in relation to the fish*

stock. ČSAV, Prague.

Hui, F.K.C., Taskinen, S., Pledger, S., Foster, S.D. & Warton, D.I. (2015). Model-based

approaches to unconstrained ordination. *Methods in Ecology and Evolution*, 6, 399–411.

Hui, F.K.C., Warton, D.I., Ormerod, J.T., Haapaniemi, V. & Taskinen, S. (2017). Variational

Approximations for Generalized Linear Latent Variable Models. *Journal of Computational*

and Graphical Statistics, 26, 35–43.

Johansson, F. (2000). The slow-fast life style characteristics in a suite of six species of odonate

larvae. *Freshwater Biology*, 43, 149–159.

Johnson, D.M. (1991). Behavioral ecology of larval dragonflies and damselflies. *Trends in*

Ecology and Evolution, 6, 8–13.

Kottek, M., Grieser, J., Beck, C., Rudolf, B. & Rubel, F. (2006). World Map of the Köppen-

Geiger climate classification updated. *Meteorologische Zeitschrift*, 15, 259–263.

Lazzaro, X. (1991). Feeding convergence in South American and African zooplanktivorous

cichlids Geophagus brasiliensis and Tilapia rendalli. *Environmental Biology of Fishes*, 31,

283–293.

Leibold, M.A. & Chase, J.M. (2018). *Metacommunity Ecology*. Princeton University Press,

Princeton, New Jersey.

Leibold, M.A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J.M., Hoopes, M.F., *et al.*

(2004). The metacommunity concept: a framework for multi-scale community ecology.

Ecology Letters, 7, 601–613.

de Marco, P., Latini, A.O. & Reis, A.P. (1999). Environmental determination of dragonfly

assemblage in aquaculture ponds. *Aquaculture Research*, 30, 357–364.

McCann, K.S., Rasmussen, J.B. & Umbanhowar, J. (2005). The dynamics of spatially coupled

food webs. *Ecology Letters*, 8, 513–523.

McCauley, S.J. (2006). The effects of dispersal and recruitment limitation on community

structure of odonates in artificial ponds. *Ecography*, 29, 585–595.

McCauley, S.J. (2008). Slow, fast and in between: Habitat distribution and behaviour of larvae in

nine species of libellulid dragonfly. *Freshwater Biology*, 53, 253–263.

McPeek, M.A. (1998). The consequences of changing the top predator in a food web: a

comparative experimental approach. *Ecological Monographs*, 68, 1–23.

Melo, A.C.G. de & Durigan, G. (2011). *Estação Ecológica de Santa Bárbara Plano de Manejo*.

Secretaria do Meio Ambiente. Secretaria do Meio Ambiente.

van de Meutter, F., de Meester, L. & Stoks, R. (2007). Metacommunity structure of pond

macroinvertebrates: Effects of dispersal mode and generation time. *Ecology*, 88, 1687–

1695.

Mouquet, N. & Loreau, M. (2003). Community Patterns in Source-Sink Metacommunities.

American Naturalist, 162, 544–557.

Muehlbauer, J.D., Collins, S.F., Doyle, M.W. & Tockner, K. (2014). How wide is a stream?

Spatial extent of the potential "stream signature" in terrestrial food webs using meta-

analysis. *Ecology*, 95, 44–55.

Niku, J., Hui, F.K.C., Taskinen, S. & Warton, D.I. (2019). gllvm: Fast analysis of multivariate

abundance data with generalized linear latent variable models in r. *Methods in Ecology and*

Evolution, 10, 2173–2182.

Niku, J., Warton, D.I., Hui, F.K.C. & Taskinen, S. (2017). Generalized Linear Latent Variable

Models for Multivariate Count and Biomass Data in Ecology. *Journal of Agricultural,*

Biological, and Environmental Statistics, 22, 498–522.

Pelicice, F.M., Vitule, J.R.S., Lima Junior, D.P., Orsi, M.L. & Agostinho, A.A. (2014). A

Serious New Threat to Brazilian Freshwater Ecosystems: The Naturalization of Nonnative

Fish by Decree. *Conservation Letters*, 7, 55–60.

Pereira, D.L.V., de Melo, A.L. & Hamada, N. (2007). Chaves de identificação para famílias e

gêneros de gerromorpha e nepomorpha (Insecta: Heteroptera) na Amazônia Central.

Neotropical Entomology, 36, 210–228.

Poff, N.L., Olden, J.D., Vieira, N.K.M., Finn, D.S., Simmons, M.P. & Kondratieff, B.C. (2006).

Functional trait niches of North American lotic insects: traits-based ecological applications

in light of phylogenetic relationships. *Journal of the North American Benthological Society*,

25, 730–755.

R Core Team. (2019). R: A language and environment for statistical computing. *R Foundation*

for Statistical Computing.

Rao, W., Ning, J., Zhong, P., Jeppesen, E. & Liu, Z. (2015). Size-dependent feeding of

omnivorous Nile tilapia in a macrophyte-dominated lake: implications for lake

management. *Hydrobiologia*, 749, 125–134.

Resetarits, W.J. (2005). Habitat selection behaviour links local and regional scales in aquatic

systems. *Ecology Letters*, 8, 480–486.

Segura, M.O., Valente-Neto, F. & Fonseca-Gessner, A.A. (2011). Chave de famílias de

coleoptera aquáticos (Insecta) do estado de são paulo, Brasil. *Biota Neotropica*, 11, 411–

412.

Shulman, R.S. & Chase, J.M. (2007). Increasing isolation reduces predator:prey species richness

ratios in aquatic food webs. *Oikos*, 116, 1581–1587.

Shurin, J.B., Cottenie, K. & Hillebrand, H. (2009). Spatial autocorrelation and dispersal

limitation in freshwater organisms, 159, 151–159.

Tran-Duy, A., Schrama, J.W., van Dam, A.A. & Verreth, J.A.J. (2008). Effects of oxygen

concentration and body weight on maximum feed intake, growth and hematological

parameters of Nile tilapia, Oreochromis niloticus. *Aquaculture*, 275, 152–162.

Trekels, H., van de Meutter, F. & Stoks, R. (2011). Habitat isolation shapes the recovery of

aquatic insect communities from a pesticide pulse. *Journal of Applied Ecology*, 48, 1480–

1489.

Vellend, M. (2010). Conceptual synthesis in community ecology. *Quarterly Review of Biology*,

85, 183–206.

Vellend, M., Srivastava, D.S., Anderson, K.M., Brown, C.D., Jankowski, J.E., Kleynhans, E.J.,

et al. (2014). Assessing the relative importance of neutral stochasticity in ecological

communities. *Oikos*, 123, 1420–1430.

Wang, Y., Naumann, U., Wright, S.T. & Warton, D.I. (2012). Mvabund- an R package for

model-based analysis of multivariate abundance data. *Methods in Ecology and Evolution*, 3,

471–474.

Warton, D.I., Blanchet, F.G., O'Hara, R.B., Ovaskainen, O., Taskinen, S., Walker, S.C., *et al.*

(2015a). So Many Variables: Joint Modeling in Community Ecology. *Trends in Ecology*

and Evolution, 30, 766–779.

Warton, D.I., Foster, S.D., De'ath •, G., Stoklosa, J. & Dunstan, P.K. (2015b). Model-based

thinking for community ecology. *Plant Ecology*, 216, 669–682.

Warton, D.I., Thibaut, L. & Wang, Y.A. (2017). The PIT-trap — A "model-free" bootstrap

procedure for inference about regression models with discrete, multivariate responses. *PLoS*

ONE, 12, e0181790.

Wellborn, G.A., Skelly, D.K. & Werner, E.E. (1996). Mechanisms creating community structure

across a freshwater habitat gradient. *Annual Review of Ecology and Systematics*, 27, 337–

63.

Wilcox, C. (2001). Habitat size and isolation affect colonization of seasonal wetlands by

predatory aquatic insects. *Israel Journal of Zoology*, 47, 459–476.

Zambrano, L., Martínez-Meyer, E., Menezes, N. & Peterson, A.T. (2006). Invasive potential of

common carp (Cyprinus carpio) and Nile tilapia (Oreochromis niloticus) in American

freshwater systems. *Canadian Journal of Fisheries and Aquatic Sciences*, 63, 1903–1910.

    Figure Legends

Figure 1. Prediction of experimental effects: Right-hand panels show the importance of niche

selection and ecological drift on community structure along a gradient of dispersal limitation

with (A) and without (B) interspecific variation in dispersal. If species have no variation in

dispersal rates, the importance of niche selection processes on community structure is predicted

to be the highest in intermediate dispersal limitation. However, if species vary in dispersal rates

and good dispersers are also favored by local conditions, they will be disproportionally favored

in more isolated habitats, making the importance of niche selection to increase with dispersal

limitation. Left hand panels show ordination plots with the predicted difference among

macroinvertebrate communities with and without fish along a gradient of spatial isolation. Each

symbol represents a hypothetical community and similar symbols communities from the same

hypothetical treatment. In this case, the greater importance of niche selection causes a larger

difference between fish and fishless communities (black bars) in a given level of isolation.

Figure 2. A. Diagram of the experimental layout. B. Representative photo of one of our

experimental units.

Figure 3. Model-based unconstrained ordination showing pond communities (symbols) and

species (filled circles) for the third sampling survey. Yellow filled circles represent predatory

insects and green filled circles represent consumers. The sizes of the circles are proportional to

maximum body size (the volume of the largest individual of each species in a log-scale).

Abbreviations of names of taxa are provided in Appendix S6. Symbols close to each other have

more similar community structures than those that are further apart. Species are positioned closer

to symbols (*i.e.* communities) where their abundance are higher. Note that consumers are more

strongly associated with high distance treatments, especially fishless ponds, whereas predatory

insects are associated with low distance treatments, especially fishless ponds. Note also that

Pantala and *Orthemis* dragonflies (Pan and Ort) are detached from the other predatory insects.

Figure 4. 95% Confidence intervals for the maximum likelihood estimates (MLE) of the effect

of fish and isolation on the abundance of predators and consumers when comparing pairs of

treatments for the last survey. Confidence intervals not crossing the zero hatched line were

considered significant effects and colored; blue bars represent an increase and red bars a decrease

in abundance from the reference treatment. Note that both predators and consumers were

negatively affected by fish (A). Consumers were positively affected by isolation only in fishless

ponds (B) and predators were negatively affected by isolation only in ponds with fish (C).

Figure 5. Maximum likelihood estimates (MLE) of abundance and their 95% confidence interval

of abundance for the six most abundant predatory taxa (A), and six most abundant consumer taxa

(B). Gray symbols indicate the absence of a taxon in a treatment. Blue triangles are MLEs for

fish treatments and orange balls are MLEs for fishless treatments. The actual estimated

differences are provided in Appendix S9. Note in A that *Pantala* and *Orthemis* dragonflies are

the only exception to the tendency of decreasing abundance of predatory insects along the

isolation gradient in fishless ponds.

Figure 6. Hypothesized processes generating the observed patterns. Black arrows represent the

flow of energy. Red solid arrows represent direct negative effects of predators on lower trophic

levels and blue dashed arrows represent indirect positive effects on lower trophic levels. The

thickness of the arrows represents the proportional amount of energy flowing or the intensity of

the effect. In fishless habitats (A) the increase in spatial isolation causes an increase in the

abundance of predatory insects with higher dispersal ability, but a reduction in the abundance of

those with lower dispersal ability, which results in a cascading effect increasing the abundance of

consumers. The presence of fish (B) however, generally reduces the abundance of predatory

insects, but it also keeps the abundance of consumers unchanged since fish also preys upon them.

As spatial isolation increases, fish prevents the increase in abundance of both consumers and

predatory insects with higher dispersal ability by increasing predation pressure upon them.

Figure 1

A) No interspecific variation in dispersal

B) Interspecific variation in dispersal
with better dispersers favored by local conditions

Figure 2

A)

30 m
120 m
480 m
Source
30m

🔴 Fishless ponds   🟢 Fish ponds   ⬜ Savanna

🟩 Floodplain   🟦 Stream

B)

Figure 3

Figure 4

Figure 5

Figure 6

A) Absence of Fish

Decrease in abundance of predatory insects with lower dispersal rates

Increase in abundance of predatory insects with higher dispersal rates

Increase in abundance of consumers

B) Presence of Fish

Decrease in abundance of predatory insects

Fish inhibits the increase in abundance of consumers

Spatial Isolation