Methylenetetrahydrofolate reductase deficiency alters cellular response after ischemic stroke in male mice

Jamie E. Abato BSc[1], Mahira Moftah[2], Greg O. Cron, PhD [3,4,5], Patrice D. Smith, PhD[2], Nafisa M. Jadavji PhD[1,2]

[1] Department of Biomedical Sciences, Midwestern University, Glendale, US
[2] Department of Neuroscience, Carleton University, Ottawa, Canada
[3] Department of Radiology, University of Ottawa
[4] The Ottawa Hospital Research Institute
[5] Department of Medical Imaging, The Ottawa Hospital

*Corresponding Author:
Nafisa M. Jadavji, PhD
Department of Biomedical Sciences
19555 N 59th Avenue
Glendale, AZ 85308 USA
Phone: 623-572-3665
E-mail: njadav@midwestern.edu

Abstract

Elevated homocysteine concentrations are a risk factor for cardiovascular disease, such as stroke. A common genetic polymorphism in methylenetetrahydrofolate reductase (*MTHFR* 677 C$\rightarrow$T) results in elevated levels of homocysteine. MTHFR plays a critical role in the synthesis of *S*-adenosylmethionine (SAM), a global methyl donor. Our previous work has demonstrated that *Mthfr*$^{+/-}$ mice, which model the *MTHFR* polymorphism in humans, are more vulnerable to ischemic damage. The aim of this study was to investigate the cellular mechanisms by which the MTHFR-deficiency changes the brain in the context of ischemic stroke injury. In the present study, three-month-old male *Mthfr*$^{+/-}$ and wild-type littermate mice were subjected to photothrombosis (PT) damage. Four weeks after PT damage, animals were tested on behavioral tasks, in *vivo* imaging was performed using T2-weighted MRI, and brain tissue was collected. *Mthfr*$^{+/-}$ animals used their non-impaired forepaw more during to explore the cylinder and had a larger damage volume compared to wild-type littermates. In brain tissue of *Mthfr*$^{+/-}$ mice methionine adenosyltransferase II alpha (MAT2A) protein levels were decreased within the damage hemisphere and increased levels in hypoxia induced factor 1 alpha (HIF-1α) in non-damage hemisphere. There was an increased anti-oxidant response in the damage site as indicated by higher levels of nuclear factor erythroid 2-related factor 2 (Nrf2) and superoxide dismutase 2 (SOD2). Our results suggest that *Mthfr*$^{+/-}$ mice are more vulnerable to PT-induced stroke damage through regulation of the cellular response. The increased anti-oxidant response we observed may be compensatory to more damage.

Keywords

Introduction

Stroke is a leading cause of death worldwide (Durga et al., 2007) and predominantly targets older adults (Katan and Luft, 2018). As the population worldwide ages, the prevalence of stroke is expected to increase (Mozaffarian et al., 2015; Katan and Luft, 2018). Older adults are at a higher risk of stroke, and also have a reduced ability to absorb all the nutrients from their diet (Bottiglieri et al., 2000; Shlisky et al., 2017).

Reduced levels of B-vitamins, a major component of one-carbon metabolism, result in elevated homocysteine concentrations which are associated with increased risk for cardiovascular disease, such as stroke (Castro et al., 2006; Lehotský et al., 2016). B-vitamins include folic acid and vitamin B12 play an important role in reducing levels of homocysteine, through one-carbon metabolism. Metheylenetetrahydrofolate reductase (MTHFR) is an enzyme that catalyzes the irreversible conversion of 5,10-methylenetetrahydrofolate to 5-methyltetrahydrofolate (5-methylTHF). The methyl group from 5-methylTHF is a substrate in the vitamin-B12-dependent methylation of homocysteine to form methionine by methionine synthase. A polymorphism in *MTHFR* (677 C$\rightarrow$T) (Frosst et al., 1995), which is present in 5-20% of North American and European populations (Schneider et al., 1998), results in elevated levels of homocysteine. The polymorphism has been associated with increased risk for vascular disease (Frosst et al., 1995) including ischemic stroke (Casas et al., 2005; Cronin et al., 2005; Song et al., 2016; Azad et al., 2017; Li et al., 2017; Mao and Han, 2018). However, the mechanism through which this increased vulnerability exists is not well defined.

The *Mthfr*$^{+/-}$ mouse models traits of the human polymorphism, including reduced levels of enzyme activity and elevated levels of homocysteine (Chen et al., 2001; Jadavji et al., 2012). Using the MTHFR-deficient mice and the photothrombosis induced ischemic damage, our previous

research has demonstrated that aged *Mthfr*$^{+/-}$ mice are more vulnerable to damage after ischemic

stroke through reduced neuronal and astrocyte viability (Jadavji et al., 2017, 2018, 2019). The

mechanisms through which the *Mthfr*$^{+/-}$ mice are more impaired after ischemic damage remains

uncharacterized. The aim of this study was to investigate the cellular response through which

MTHFR deficiency regulates the brain's vulnerability to ischemic stroke.

Experimental Design

Animals and experimental design

All experiments were conducted in accordance with the guidelines of the Canadian Council

on Animal Care (CCAC). Two cohorts of male C57Bl/6 *Mthfr* heterozygote (*Mthfr*$^{+/-}$, n = 15) and

wild-type littermate controls (*Mthfr*$^{+/+}$, n = 15) mice were housed in standard caging conditions

with *ad libitum* standard mouse chow (Envigo) and water for the duration of the experiment.

Animals were randomly assigned to each of the experimental groups.

Photothrombosis

At 3-months of age, animals were subjected photothrombosis (PT) induced ischemic

damage. Animals were anaesthetized with isoflurane (1.5%) in a 70:30 nitrous oxide: oxygen

mixture. Core body temperature was maintained at 37.0 ± 0.2ºC using an automated heat blanket

with temperature feedback (Harvard Apparatus). Focal ischemic damage was targeted to the

sensorimotor cortex (mediolateral ± 0.24 mm) as previously described (Lee et al., 2004; Jadavji et

al., 2017, 2018). An intraperitoneal injection of 10 mg/kg photoactive Rose Bengal (Sigma) was

administered 5 min prior to a 532 nm green laser (Beta Electronics) exposure on the skull.

Behavioral Testing

Bederson exam

Neurological testing was performed 4 weeks after PT damage using the Bederson exam

(perfect score = 0). The exam evaluates symmetry, forepaw outstretching, climbing, and reflex

(Bederson et al., 1986). Investigators were blinded to the treatment performed all neurological

testing on mice.

Forepaw Asymmetry

A cylinder (19 cm high, 14 cm in diameter) made of thick glass was used to assess forepaw

asymmetry. The mouse was placed in the glass cylinder for 10 min and video was recorded from

the side. Forepaw contacts with the wall of the cylinder were counted. The first 10 movements

were scored during the 10-min test. The final score was calculated as follows: final score =

(number of non-impaired forelimb movement – number of impaired forelimb movement)/ (sum of

all movements). A positive score thus indicated favored use of the non-impaired forelimb, a

negative indicated favored use of the impaired forelimb and a score of zero indicated equal use of

both non-impaired and impaired forelimbs upon rearing and exploration of the cylinder walls

(Theoret et al., 2015).

MRI lesion volume analysis

At four weeks after PT damage, lesion volume was quantified. A 7 Tesla GE/Agilent MR

901 was used to collect the data. Mice were anaesthetized prior to the MRI procedure using

isoflurane. A 2D fast spin echo sequence (FSE) pulse sequence was used for the imaging, with

the following parameters: slice thickness = 0.5 mm, spacing = 0 mm, field of view = 2.5 cm,

matrix = 256 x 256, echo time = 41 ms, repetition time = 4000 ms, echo train length = 8, bandwidth

= 16 kHz, 2 averages, fat saturation, imaging time = 5 minutes.

Western blot

At the completion of experiments, approximately 4 weeks after PT damage, animals were

decapitated to obtain fresh brain tissue collection for Western blot analysis, PT and non-PT regions

(~50 mg) were micro-dissected. RIPA buffer containing phosphatase and proteinase inhibitors

(Cell Signalling) was used to extract proteins. Proteins were separated by SDS-PAGE and

transferred to nitrocellulose membranes. Primary antibodies were; anti-rabbit methionine adenosyl

transferase (MAT2A) (1:2000; AbCam), anti-rabbit hypoxia induced factor 1 alpha (HIF-1α)

(1:500; AbCam), or anti-rabbit glyceraldehyde 3-phosphate dehydrogenase (GAPDH) (1:10 000;

Cell Signalling) overnight. Primary antibodies were visualized with infrared dye goat anti-rabbit

IgG (1:10 000, 800 Channel; LiCor Biosciences). Imaging was performed using Near Infrared

(NIR) detection (Li-Cor Biosciences). Quantification of bands was determined using densitometry

analysis via ImageJ (National Institutes of Health) and bands were normalized to GAPDH.

Immunofluorescence

After the completion of behavioral testing, animals were euthanized using the perfusion

method. Brain tissue was fixed in 4% paraformaldehyde (PFA) for 24 hrs. Using a cryostat, brain

tissue was sectioned at 30 μm thickness then mounted on slides in serial order.

To explore mechanisms of neurodegeneration, levels of oxidative stress using nuclear

factor erythroid 2-related factor 2 (NRF2) and superoxide dismutase 2 (SOD2) were investigated.

Primary antibodies used were rabbit anti-NRF2 (1:100, AbCam), rabbit anti-SOD2 (1:100, Life

Technologies), and mouse anti-neuronal nuclei (NeuN) (1:200, AbCam) diluted in 0.5% Triton X.

Brain tissue was incubated overnight at 4°C with primary antibodies. The following day, secondary

antibodies (1:200, Cell Signaling Technologies) brain tissue was incubated at room temperature

for 2 h and then stained with 4', 6-diamidino-2-phenylindole (DAPI) (1:10, 000). For analysis in

within the ischemic core of brain tissue, co-localization of NRF2 or SOD2 with NeuN labelled

neurons were counted and averaged per animal. Data from five animals per group were analyzed.

Statistics analysis

All data analysis was conducted by two individuals blinded to experimental groups. Data

from the study were analyzed using GraphPad Prism 6.0. Unpaired t-tests were performed on

behavioral testing, MRI, immunofluorescence, and Western Blot experiments between genotype

groups with PT damage. In all analysis, a p-value less than or equal to 0.05 was considered

significant. All data are presented as mean $\pm$ standard error of the mean (SEM).

Results

Forepaw usage impairs in Mthfr$^{+/-}$ mice

There was no difference in the neurological testing score between *Mthfr$^{+/+}$* and *Mthfr$^{+/-}$* mice

(Figure 1A, t (12) = 1.62, p = 0.1). However, *Mthfr$^{+/-}$* mice showed a preference to use their non-

impaired forelimb, while completing the forepaw asymmetry task compared to *Mthfr$^{+/+}$* animals

(Figure 1B, t (12) = 3.46, p < 0.01).

Increased damage volume in Mthfr$^{+/-}$ mice

Quantification of PT lesion volume using T2 weighted MRI revealed that the $Mthfr^{+/-}$ mice

showed a larger lesion volume in the sensorimotor cortex compared to wildtype animals (Figure

1C, t (8) = 3.1, p<0.05).

Altered cellular response within damage area of Mthfr$^{+/-}$ mice

We examined two markers of cellular response after PT damage, the first an enzyme,

MAT2A, which is linked to MTHFR through one-carbon metabolism and is important in the

generation of S-adenosylmethionine (SAM), a global methyl donor. Four weeks after PT damage

we measured levels of MAT2A. In $Mthfr^{+/-}$ mice there was a decrease in levels of MAT2A within

the PT damage hemisphere compared to $Mthfr^{+/+}$ animals (Figure 3A, t (14) = 2.7, p < 0.05). There

were no changes in MAT2A within the non-PT hemisphere between groups (data not shown, t (14)

=11.4, p = 0.58).

HIF-1α is an regulator of the cellular response after hypoxia (Katsimpardi et al., 2014).

Previously we have shown increased levels of HIF-1α $Mthfr^{+/-}$ and $Mthfr^{-/-}$ primary astrocytes after

exposure to hypoxia (Jadavji et al., 2018), as well as in brain tissue of $Mthfr^{+/-}$ mice with ischemic

stroke (Jadavji et al., 2019). In the present study, using an *in vivo* model of MTHFR deficiency

and ischemic damage, we observed increased levels of HIF-1α within the non-damage PT

hemisphere of $Mthfr^{+/-}$ mice (Figure 3B, t (15) = 3.4, p < 0.01). Interestingly, we did not observe

any expression of HIF-1α within the lesion hemisphere of $Mthfr^{+/+}$ and $Mthfr^{+/-}$ animals (data not

shown).

Increased oxidative stress within damage area of Mthfr$^{+/-}$ mice

We measured oxidative stress 4 weeks after damage in sectioned brain tissue of animals,

by assessing levels of nuclear factor (erythroid-derived 2)-like 2 (Nrf2) and superoxide dismutase

2 (SOD2). Representative images of Nrf2 are shown in Figure 4A and B. There were increased

levels of Nrf2 within the damaged area of $Mthfr^{+/-}$ mice (Figure 4C, t (8) = 4.02, p < 0.001).

Representative SOD2 images are shown in figures 4D and 4E. $Mthfr^{+/-}$ mice had increased levels

of SOD2 in brain tissue (Figure 4F, t (8) = 2.57, p < 0.05).

Discussion

Elevated levels of homocysteine are a risk factor for stroke (Castro et al., 2006). Previously

we have reported that aged $Mthfr^{+/-}$ mice are more vulnerable to ischemic damage (Jadavji et al.,

2018). We have also shown this increased vulnerability *in vitro* using primary neurons (Jadavji et

al., 2017) and astrocytes (Jadavji et al., 2018). In the present study, we have confirmed our previous

findings that $Mthfr^{+/-}$ mice have a more severe motor impairment after ischemic stroke using the

forepaw asymmetry task. The heterozygote mice also had increased damage volume within the

sensorimotor cortex. To understand the cellular response after ischemic damage, we report reduced

levels of MAT2A $Mthfr^{+/-}$ mice within the damage area. MAT2A is an enzyme linked to MTHFR

through one-carbon metabolism and involved in generating SAM within the damage brain region.

Within the non-lesion hemisphere there are increased levels of HIF-1α in the non-damaged PT

area.   HIF-1α is involved in cellular responses for protection and repair. The anti-oxidative

response was increased in within the damage of $Mthfr^{+/-}$ mice.

In patients with vascular disease, changes in DNA methylation have been reported (Castro

et al., 2003). Using animal models to understand diseases processes, it has been suggested that

decreases in methylation may promote cell death after stroke (Kalani et al., 2013). Furthermore,

changes in methylation may impact stroke onset and progression (Endres et al., 2000; Kalani et

al., 2013). MTHFR generates methyl groups, and the impact of a MTHFR-deficiency on

methylation after stroke remains unclear; our present study has attempted to fill this gap. Previous

work reported that *Mthfr*$^{+/-}$ mice have increased levels of hypomethylation in the brain (Chen et

al., 2001). MAT2A is a catalytic subunit of methionine adenosyltransferase (MAT), an essential

enzyme for the catalysis of the principle biological methyl donor, SAM from methionine and ATP.

In the present study, we observed increased levels of MAT2A in wild-type mice which is in line

with increased methylation after ischemic stroke that has previously been reported (Endres et al.,

2000; Kalani et al., 2013). Interestingly, lower levels of MAT2A may lead to reduced levels of

SAM, the global methyl donor (Frau et al., 2012), however this requires further confirmation.

HIF-1α is a master regulator of cell response to hypoxia, since it leads to the expression of

several genes involved in the adaptation of oxygen availability and angiogenesis (Jiang et al., 2001;

Cunningham et al., 2012). In the kidney, HIF-1α stabilizes cells after ischemia and interference

with HIF-1α promotes tissue damage and dysfunction (Conde et al., 2012). In the present study,

we observed increased levels of HIF-1α within the non-PT hemisphere in *Mthfr*$^{+/-}$ mice. These

results support our previous work, in which we report increased HIF-1α levels in primary astrocyte

cultures from *Mthfr*$^{+/-}$ and *Mthfr*$^{-/-}$ embryos after a hypoxia treatment (Jadavji et al., 2018).

Increased levels of HIF-1α in the non-PT hemisphere may be another adaptive response to

compensate for damage in the other hemisphere.

In the brain, both ischemic stroke (Siti et al., 2015) and increased levels of homocysteine

(Hoffman, 2011) result in oxidative stress. The anti-oxidative response is important in regulating

the impact of oxidative stress in brain tissue especially after damage, during neurodegeneration

(Zhang and Butterfield, 2017). Our previous work has shown that dietary supplementation of one-

carbon metabolites including folic acid (vitamin B9), choline, vitamin B12, and riboflavin (vitamin

B2) increase the anti-oxidative response in brain tissue after ischemic stroke (Jadavji 2017).

Targeting anti-oxidants after stroke in patients is complicated (Siti et al., 2015), but increasing

activation of Nrf2 could be a potential target (Zhang et al., 2017). Additionally, the combination

of anti-oxidant through diet and pharmaceutical therapies may be beneficial for stroke-affected

patients.

Together, the data from this study provides insight on potential mechanisms that enhance

stroke vulnerability in *Mthfr*$^{+/-}$ mice. Ischemic stroke in MTHFR-deficient animals changes

cellular responses in brain tissue, making *Mthfr*$^{+/-}$ mice more vulnerable to damage. Further

investigation is required to dissect the impact of changes to SAM levels and consequential global

genetic silencing processes in MTHFR-deficiency that contribute to ischemic stroke outcomes.

Acknowledgements

This research was supported by Natural Sciences and Engineering Research Council of Canada

(NMJ and PDS) and Midwestern University start-up funds (NMJ).

267 **Figure & Captions**

271 **Figure 1.** Behavioral function in adult *Mthfr*$^{+/+}$ and *Mthfr*$^{+/-}$ animals 4 weeks after PT damage.

272 Neurological testing (A) and forepaw asymmetry (B) scores in *Mthfr*$^{+/+}$ and *Mthfr*$^{+/-}$ mice. Bars

273 represent mean $\pm$ SEM of 7 animals per group. ** $p < 0.01$ un-paired t-test between *Mthfr*$^{+/+}$ and

274 *Mthfr*$^{+/-}$ mice.

Figure 2. The impact of MTHFR-deficiency on damage volume 4-weeks after PT damage to the

sensorimotor cortex. Representative MRI T2-weighted images from *Mthfr*$^{+/+}$ (A) and *Mthfr*$^{+/-}$ (B)

mice. Quantification of lesion volume (C). Bars represent mean $\pm$ SEM of 4 animals per group. *

p < 0.05 un-paired t-test between *Mthfr*$^{+/+}$ and *Mthfr*$^{+/-}$ mice.

Figure 3. The impact of MTHFR-deficiency after ischemic damage on cellular responses to

ischemia. Protein levels of methionine adenosyltransferase 2A (MAT2A) in PT hemisphere (A),

and non-PT hemisphere levels of hypoxia-inducible factor 1-alpha (HIF-1α) (B) in *Mthfr*^{+/+} and

Mthfr^{+/-} animals 4-weeks after PT damage. Bars represent mean ± SEM of 7 mice per group. The

panels below the graphs depict a representative Western Blot. *p<0.05 and **p<0.01 indicate

Tukey's pairwise comparison between indicated groups.

Figure 4. Antioxidant activity within ischemic area of MTHFR-deficient mice. Representative

images of nuclear factor (erythroid-derived 2)-like 2 (Nrf2), (A) and (B). Quantification of Nrf2,

neuronal nuclei (NeuN), and DAPI (4′,6-diamidino-2-phenylindole) positive cells (C).

Representative superoxide dismutase 2 (SOD2) images, (D) and (E). Quantification of SOD2,

NeuN and DAPI positive cells within damage area (F). Bars represent mean $\pm$ SEM of 4-5 mice

per group. *p<0.05 and **p<0.01 indicate unpaired t-test comparison between indicated groups.

References

Azad MAK, Huang P, Liu G, Ren W, Teklebrh T, Yan W, Zhou X, Yin Y (2017)
Hyperhomocysteinemia and cardiovascular disease in animal model. Amino Acids
Available at: http://link.springer.com/10.1007/s00726-017-2503-5.

Bederson JB, Pitts LH, Tsuji M, Nishimura MC, Davis RL, Bartkowski H (1986) Rat middle
cerebral artery occlusion: evaluation of the model and development of a neurologic
examination. Stroke 17:472–476 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/3715945 [Accessed October 7, 2014].

Bottiglieri T, Reynolds EH, Laundy M (2000) Folate in CSF and age. J Neurol Neurosurg
Psychiatry 69:562 Available at: http://www.ncbi.nlm.nih.gov/pubmed/11183038 [Accessed
July 13, 2018].

Casas JP, Bautista LE, Smeeth L, Sharma P, Hingorani AD (2005) Homocysteine and stroke:
Evidence on a causal link from mendelian randomisation. Lancet 365:224–232.

Castro R, Rivera I, Blom HJ, Jakobs C, Tavares de Almeida I (2006) Homocysteine metabolism,
hyperhomocysteinaemia and vascular disease: an overview. J Inherit Metab Dis 29:3–20
Available at: http://www.ncbi.nlm.nih.gov/pubmed/16601863 [Accessed June 25, 2014].

Castro R, Rivera I, Struys EA, Jansen EEW, Ravasco P, Camilo ME, Blom HJ, Jakobs C, De
Almeida IT (2003) Increased homocysteine and S-adenosylhomocysteine concentrations
and DNA hypomethylation in vascular disease. Clin Chem 49:1292–1296.

Chen Z, Karaplis AC, Ackerman SL, Pogribny IP, Melnyk S, Lussier-Cacan S, Chen MF, Pai A,
John SW, Smith RS, Bottiglieri T, Bagley P, Selhub J, Rudnicki MA, James SJ, Rozen R
(2001) Mice deficient in methylenetetrahydrofolate reductase exhibit
hyperhomocysteinemia and decreased methylation capacity, with neuropathology and aortic
lipid deposition. Hum Mol Genet 10:433–443 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/11181567.

Conde E, Alegre L, Blanco-Sánchez I, Sáenz-Morales D, Aguado-Fraile E, Ponte B, Ramos E,
Sáiz A, Jiménez C, Ordoñez A, López-Cabrera M, del Peso L, de Landázuri MO, Liaño F,
Selgas R, Sanchez-Tomero JA, García-Bermejo ML (2012) Hypoxia inducible factor 1-
alpha (hif-1 alpha) is induced during reperfusion after renal ischemia and is critical for
proximal tubule cell survival. PLoS One 7.

Cronin S, Furie KL, Kelly PJ (2005) Dose-Related Association of MTHFR 677T Allele With
Risk of Ischemic Stroke: Evidence From a Cumulative Meta-Analysis. Stroke 36:1581–
1587 Available at:
http://stroke.ahajournals.org/cgi/doi/10.1161/01.STR.0000169946.31639.af.

Cunningham LA, Candelario K, Li L (2012) Roles for HIF-1α in neural stem cell function and
the regenerative response to stroke. Behav Brain Res 227:410–417.

Durga J, van Boxtel MPJ, Schouten EG, Kok FJ, Jolles J, Katan MB, Verhoef P (2007) Effect of
3-year folic acid supplementation on cognitive function in older adults in the FACIT trial: a
randomised, double blind, controlled trial. Lancet (London, England) 369:208–216
Available at: http://www.ncbi.nlm.nih.gov/pubmed/17240287 [Accessed October 18, 2017].

Endres M, Meisel a, Biniszkiewicz D, Namura S, Prass K, Ruscher K, Lipski a, Jaenisch R, Moskowitz M a, Dirnagl U (2000) DNA methyltransferase contributes to delayed ischemic brain injury. J Neurosci 20:3175–3181.

Frau M, Tomasi ML, Simile MM, Demartis MI, Salis F, Latte G, Calvisi DF, Seddaiu MA, Daino L, Feo CF, Brozzetti S, Solinas G, Yamashita S, Ushijima T, Feo F, Pascale RM (2012) Role of transcriptional and posttranscriptional regulation of methionine adenosyltransferases in liver cancer progression. Hepatology 56:165–175.

Frosst P, Blom HJ, Milos R, Goyette P, Sheppard CA, Matthews RG, Boers GJH, den Heijer M, Kluijtmans LAJ, van den Heuvel LP, Rozen R (1995) A candidate genetic risk factor for vascular disease: a common mutation in methylenetetrahydrofolate reductase. Nat Genet 10:196–201.

Hoffman M (2011) Hypothesis: hyperhomocysteinemia is an indicator of oxidant stress. Med Hypotheses 77:1088–1093 Available at: http://www.ncbi.nlm.nih.gov/pubmed/21963358 [Accessed September 8, 2013].

Jadavji N, Emmerson JT, Shanmugalingam U, MacFarlane AJ, Willmore WG, Smith PD (2018) A genetic deficiency in folic acid metabolism impairs recovery after ischemic stroke. Exp Neurol 309:14–22.

Jadavji NM, Deng L, Leclerc D, Malysheva O, Bedell BJ, Caudill MA, Rozen R (2012) Severe methylenetetrahydrofolate reductase deficiency in mice results in behavioral anomalies with morphological and biochemical changes in hippocampus. Mol Genet Metab 106:149–159 Available at: http://www.ncbi.nlm.nih.gov/pubmed/22521626 [Accessed August 30, 2013].

Jadavji NM, Emmerson J, Willmore WG, MacFarlane AJ, Smith P (2017) B-vitamin and choline supplementation increases neuroplasticity and recovery after stroke. Neurobiol Dis 103:89–100.

Jadavji NM, Mosnier H, Kelly E, Lawrence K, Cruickshank S, Stacey S, McCall A, Dhatt S, Arning E, Bottiglieri T, Smith PD (2019) One-carbon metabolism supplementation improves outcome after stroke in aged male MTHFR-deficient mice. Neurobiol Dis:104613 Available at: http://www.ncbi.nlm.nih.gov/pubmed/31525435 [Accessed September 25, 2019].

Jiang BH, Jiang GQ, Zheng JZ, Lu ZM, Hunter T, Vogt PK (2001) Phosphatidylinositol 3-kinase signaling controls levels of hypoxia-inducible factor. Cell Growth Differ 12:363–369.

Kalani A, Kamat PK, Tyagi SC, Tyagi N (2013) Synergy of homocysteine, MicroRNA, and epigenetics: A novel therapeutic approach for stroke. Mol Neurobiol 48:157–168.

Katan M, Luft A (2018) Global Burden of Stroke. Semin Neurol 38:208–211.

Katsimpardi L, Litterman NK, Schein PA, Miller CM, Loffredo FS, Wojtkiewicz GR, Chen JW, Lee RT, Wagers AJ, Rubin LL (2014) Vascular and neurogenic rejuvenation of the aging mouse brain by young systemic factors. Science 344:630–634 Available at: http://www.sciencemag.org/cgi/doi/10.1126/science.1251141 [Accessed April 10, 2017].

Lee J-K, Kim J-E, Sivula M, Strittmatter SM (2004) Nogo receptor antagonism promotes stroke recovery by enhancing axonal plasticity. J Neurosci 24:6209–6217.

Lehotský J, Tothová B, Kovalská M, Dobrota D, Beňová A, Kalenská D, Kaplán P (2016) Role
of Homocysteine in the Ischemic Stroke and Development of Ischemic Tolerance. Front
Neurosci 10:1–16 Available at:
http://journal.frontiersin.org/article/10.3389/fnins.2016.00538/full.

Li A, Shi Y, Xu L, Zhang Y, Zhao H, Li Q, Zhao X, Cao X, Zheng H, He Y (2017) A possible
synergistic effect of MTHFR C677T polymorphism on homocysteine level variations
increased risk for ischemic stroke. Medicine (Baltimore) 96:e9300.

Mao X, Han L (2018) The Relationship of Methylenetetrahydrofolate Reductase Gene C677T
Polymorphism and Ischemic Stroke in Chinese Han Population. Ann Clin Lab Sci 48:242–
247 Available at: http://www.ncbi.nlm.nih.gov/pubmed/29678854 [Accessed April 30,
2018].

Mozaffarian D et al. (2015) Heart Disease and Stroke Statistics-2016 Update: A Report From the
American Heart Association. Circulation 133:e38-360.

Schneider JA, Rees DC, Liu YT, Clegg JB (1998) Worldwide distribution of a common
methylenetetrahydrofolate reductase mutation. Am J Hum Genet 62:1258–1260 Available
at:
http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=1377093&tool=pmcentrez&ren
dertype=abstract.

Shlisky J, Bloom DE, Beaudreault AR, Tucker KL, Keller HH, Freund-levi Y (2017) Nutritional
Considerations for Healthy Aging and Reduction in Age-Related Chronic Disease. Adv
Nutr 8:17–26.

Siti HN, Kamisah Y, Kamsiah J (2015) The role of oxidative stress, antioxidants and vascular
inflammation in cardiovascular disease (a review). Vascul Pharmacol 71:40–56.

Song Y, Li B, Wang C, Wang P, Gao X, Liu G (2016) Association between 5,10-
Methylenetetrahydrofolate Reductase C677T Gene Polymorphism and Risk of Ischemic
Stroke: A Meta-analysis. J Stroke Cerebrovasc Dis:1–9 Available at:
http://linkinghub.elsevier.com/retrieve/pii/S1052305715006667.

Theoret JKJK, Jadavji NMNM, Zhang M, Smith PDPD (2015) Granulocyte macrophage colony
stimulating factor treatment results in recovery of motor function after white matter damage
in mice. Eur J Neurosci 43:1–8 Available at:
http://www.ncbi.nlm.nih.gov/pubmed/26474338.

Zhang B, Dong J, Chen Y, Liu Y, Huang S, Zhong X, Cheng Y, Wang Z (2017) Nrf2 mediates
the protective effects of homocysteine by increasing the levels of GSH content in HepG2
cells. Mol Med Rep:597–602 Available at: http://www.spandidos-
publications.com/10.3892/mmr.2017.6633.

Zhang J, Butterfield DA (2017) Oxidative stress and neurodegeneration. Brain Res Bull 133:1–3
Available at: https://linkinghub.elsevier.com/retrieve/pii/S0361923017302496 [Accessed
November 18, 2019].

428