Niche-selective inhibition of pathogenic Th17 cells by targeting metabolic redundancy

Lin Wu[1,3,7], Kate E.R. Hollinshead[2], Yuhan Hao[3,4], Christy Au[1,5], Lina Kroehling[1], Charles Ng[1],

Woan-Yu Lin[1], Dayi Li[1], Hernandez Moura Silva[1], Jong Shin[6], Juan J. Lafaille[1,6], Richard

Possemato[6], Michael E. Pacold[2], Thales Papagiannakopoulos[6], Alec C. Kimmelman[2], Rahul

Satija[3,4], Dan R. Littman[1,5,6,7,8]

[1]The Kimmel Center for Biology and Medicine of the Skirball Institute, New York University

School of Medicine, New York, NY, USA; [2]Department of Radiation Oncology and Perlmutter

Cancer Center, New York University School of Medicine, New York, NY, USA; [3]New York

Genome Center, New York, NY, USA; [4]Center for Genomics and Systems Biology, New York

University, New York, NY, USA; [5]Howard Hughes Medical Institute, New York, NY, USA;

[6]Department of Pathology, New York University School of Medicine, New York, NY, USA;

[7]Corresponding Authors; [8]Lead Contact.

Lead Contact: dan.littman@med.nyu.edu

Correspondence: lin.wu@med.nyu.edu and dan.littman@med.nyu.edu

Key words: Autoimmunity, metabolic plasticity, hypoxia, glycolysis, CRISPR, OXPHOS, inflammation, segmented filamentous bacteria, EAE, colitis

Summary

Targeting glycolysis has been considered therapeutically intractable owing to its essential housekeeping role. However, the context-dependent requirement for individual glycolytic steps has not been fully explored. We show that CRISPR-mediated targeting of glycolysis in T cells in mice results in global loss of Th17 cells, whereas deficiency of the glycolytic enzyme glucose phosphate isomerase (Gpi1) selectively eliminates inflammatory encephalitogenic and colitogenic Th17 cells, without substantially affecting homeostatic microbiota-specific Th17 cells. In homeostatic Th17 cells, partial blockade of glycolysis upon *Gpi1* inactivation was compensated by pentose phosphate pathway flux and increased mitochondrial respiration. In contrast, inflammatory Th17 cells experience a hypoxic microenvironment known to limit mitochondrial respiration, which is incompatible with loss of *Gpi1*. Our study suggests that inhibiting glycolysis by targeting Gpi1 could be an effective therapeutic strategy with minimum toxicity for Th17-mediated autoimmune diseases, and, more generally, that metabolic redundancies can be exploited for selective targeting of disease processes.

Introduction

Cellular metabolism is a dynamic process that supports all aspects of the cell's activities. It is orchestrated by more than 2000 metabolic enzymes organized into pathways that are specialized for processing and producing distinct metabolites. Multiple metabolites are shared by different pathways, serving as nodes in a complex network with many redundant elements. A well-appreciated example of metabolic plasticity is the generation of ATP by either glycolysis or by mitochondrial oxidative phosphorylation (OXPHOS), making the two processes partially redundant despite having other non-redundant functions (O'Neill et al., 2016). Here, we explore metabolic plasticity in the context of T cell function and microenvironment, and the consequence of limiting that plasticity by inhibiting specific metabolic enzymes.

Th17 cells are a subset of CD4$^+$ T cells whose differentiation is governed by the transcription factor RORγt, which regulates the expression of the signature IL-17 cytokines (Ivanov et al., 2006). Under homeostatic conditions, Th17 cells typically reside at mucosal surfaces where they provide protection from pathogenic bacteria and fungi and also regulate the composition of the microbiota (Kumar et al., 2016; Mao et al., 2018; Milner and Holland, 2013; Okada et al., 2015; Puel et al., 2011). However, under conditions that favor inflammatory processes, Th17 cells can adopt a pro-inflammatory program that promotes autoimmune diseases, including inflammatory bowel disease (Hue et al., 2006; Kullberg et al., 2006), psoriasis (Piskin et al., 2006; Zheng et al., 2007) and diverse forms of arthritis (Hirota et al., 2007; Murphy et al., 2003; Nakae et al., 2003a; Nakae et al., 2003b). This program is dependent on IL-23 and is typically marked by the additional expression of T-bet and IFN-γ (Ahern et al., 2010; Cua et al., 2003; Hirota et al., 2011; Morrison et al., 2013). The Th17 pathway has been targeted with neutralizing antibodies specific for IL-17A or IL-23 to effectively treat psoriasis,

ulcerative colitis, Crohn's disease, and some forms of arthritis (Fotiadou et al., 2018; Hanzel and

D'Haens, 2020; Langley et al., 2018; Moschen et al., 2019; Pariser et al., 2018; Tahir, 2018;

Wang et al., 2017), but these therapies inevitably expose patients to potential fungal and bacterial

infections, as they also inhibit homeostatic Th17 cell functions. In this study, we aimed to

identify genes and pathways that are required for the function of inflammatory pathogenic Th17

cells, but are dispensable for homeostatic Th17 cells, which may enable the selective therapeutic

targeting of pathogenic Th17 cells.

T cell activation leads to extensive clonal expansion, demanding a large amount of energy

and biomass production (O'Neill et al., 2016; Olenchock et al., 2017). Glycolysis is central both

in generating ATP and providing building blocks for macromolecular biosynthesis (Zhu and

Thompson, 2019). Glycolysis catabolizes glucose into pyruvate through ten enzymatic reactions.

In normoxic environments, pyruvate is typically transported into the mitochondria to be

processed by the tricarboxylic acid cycle, driving OXPHOS for ATP production. In hypoxic

environments, OXPHOS is suppressed and glycolysis is enhanced, generating lactate from

pyruvate in order to regenerate nicotinamide adenine dinucleotide (NAD^+) needed to support

ongoing glycolytic flux (Birsoy et al., 2015; Semenza, 2014; Sullivan et al., 2015). In highly

proliferating cells, such as cancer cells and activated T cells, pyruvate can be converted into

lactate in the presence of oxygen, a phenomenon termed aerobic glycolysis, or the "Warburg

effect" (MacIver et al., 2013; Vander Heiden et al., 2009; Vander Heiden and DeBerardinis,

2017; Warburg et al., 1958; Zhu and Thompson, 2019). Although inhibiting glycolysis is an

effective method of blocking T cell activation (Macintyre et al., 2014; Peng et al., 2016; Shi et

al., 2011), inhibitors such as 2-Deoxy-D-glucose (2DG) have limited application in patients, due

to toxic side effects as a result of the housekeeping function of glycolysis in multiple cell types

(Raez et al., 2013).

Here, we systematically interrogate the requirement for individual glycolytic reactions in

Th17 cell models of inflammation and homeostatic function. We found that the glycolysis gene

Gpi1 is unique in that it is selectively required by inflammatory pathogenic Th17 but not

homeostatic Th17 cells. All other tested glycolysis genes were essential for functions of both

Th17 cell types. Our mechanistic study revealed that upon Gpi1 loss during homeostasis, pentose

phosphate pathway (PPP) activity was sufficient to maintain basal glycolytic flux for biomass

generation, while increased mitochondrial respiration compensated for reduced glycolytic flux.

In contrast, Th17 cells in the inflammation models are confined to hypoxic environments and

hence were unable to increase respiration to compensate for reduced glycolytic flux. This

metabolic stress led to the selective elimination of Gpi1-deficient Th17 cells in inflammatory

settings but not in healthy intestinal lamina propria. Overall, our study reveals a context-

dependent metabolic requirement that can be targeted to eliminate pathogenic Th17 cells. As

Gpi1 blockade can be better tolerated than inhibition of other glycolysis components, it is a

potentially attractive target for clinical use.

Results

Inflammatory Th17 cells have higher expression of glycolysis pathway genes than

commensal bacteria-induced homeostatic Th17 cells

Commensal SFB induce the generation of homeostatic Th17 cells in the small intestine lamina

propria (SILP) (Ivanov et al., 2009), whereas immunization with myelin oligodendrocyte

glycoprotein peptide (MOG) in complete Freund's adjuvant (CFA) induces pathogenic Th17

cells in the spinal cord (SC), resulting in experimental autoimmune encephalomyelitis (EAE)

(Cua et al., 2003). Th17 cell pathogenicity requires expression of IL-23R (Ahern et al., 2010;

McGeachy et al., 2009). To identify genes involved in the pathogenic but not the homeostatic

Th17 cell program, we reconstituted Rag1$^{-/-}$ mice with isotype-marked $Il23r$-sufficient

($Il23r^{GFP/+}$) and $Il23r$ knockout ($Il23r^{GFP/GFP}$) bone marrow cells, and sorted CD4$^+$ T cells of both

genetic backgrounds from the SILP of SFB-colonized mice and the SC of EAE mice for single

cell RNA sequencing (scRNAseq) analysis (Figure S1A). We identified 4 clusters of CD4$^+$ T

cells in the SC of EAE mice and 5 clusters of CD4$^+$ T cells in ileum lamina propria of SFB-

colonized mice (Figure S1B,C)). Clusters were annotated based on their signature genes, i.e.

Th17 cells ($Il17a$), Th1 cells ($Ifng$), Treg cells ($Foxp3$), and Tcf7$^+$ cells ($Tcf7$) (Figure S1D,E).

Importantly, two of the EAE clusters consist mostly of IL-23R-sufficient T cells (Figures S1F-

H), indicating that these populations are likely pathogenic. One of them was annotated as Th17

and the other as Th1* according to earlier publications describing such cells as $Il23r$-dependent

IFN-γ producers (Hirota et al., 2011; Okada et al., 2015), and they were validated using flow

cytometry (Figure S1I). Importantly, pathway enrichment analysis revealed glycolysis as the top

pathway upregulated in both EAE Th17 and Th1* compared to the SFB-induced Th17 cells

(Figures S1J,K).

We selected a subset of genes that were differentially expressed in the pathogenic Th17/Th1*

cells (EAE model) versus the homeostatic Th17 cells (SFB model) for pilot functional studies

(Figures S1L,M). We developed a CRISPR- based T cell transfer EAE system to evaluate the

roles of the selected genes in pathogenicity (Figures S2A-C; see Methods). Among 12 genes

tested, only triosephosphate isomerase 1 ($Tpi1$), encoding an enzyme in the glycolysis pathway,

was required for disease onset following transfer of in vitro differentiated MOG-specific 2D2

TCR transgenic T cells (Figure S2D). This result was confirmed by analyzing $Tpi1^{f/f}Cd4^{cre}$ mice,

which, unlike Tpi1$^{+/+}$ CD4^{cre} littermates, were completely resistant to EAE induction (Figure

1A). This finding prompted us to direct our focus on the glycolysis pathway. The relatively

sparse scRNAseq data detected the transcripts of only a few glycolysis pathway genes. We

therefore compared bulk RNAseq data of the two major pathogenic populations (Th1*: Klrc1^{+}

Foxp3^{RFP-}, Th17: IL17a^{GFP+} Klrc1^{-} Foxp3^{RFP-}) and the Treg cells (Foxp3^{RFP+} IL17a^{GFP-}) from the

spinal cord of EAE model mice and of the Th17 (IL17a^{GFP+} Foxp3^{RFP-}) and Treg (IL17a^{GFP-}

Foxp3^{RFP+}) cells from the SILP of the SFB model (Figures S2E, F). We found that almost every

glycolysis pathway gene was expressed at a higher level in the EAE model than in the SILP

CD4^{+} T cells, irrespective of whether cells were pathogenic or Treg (Figure 1B), suggesting that

most CD4^{+} T cells have more active glycoysis in the inflamed spinal cord microenvironment

than in the SILP at homeostasis.

Despite these differences in gene expression, Tpi1 was also indispensable for the SFB-

dependent induction of homeostatic Th17 cells. We analyzed CD4^{+} T cell profiles in chimeric

Rag1 KO mice reconstituted with a 1:1 mix of bone marrow cells from $Tpi1^{+/+}CD4^{cre}$ CD45.1/2

and $Tpi1^{f/f}CD4^{cre}$ CD45.2/2 donors (Figure S3A). Tpi1 deficiency led to 33-fold reduction of

total CD4^{+} T cells in the SILP and to 1.4-fold and 1.7-fold reduction in the mLN and the

peripheral blood, respectively (Figure 1D). Moreover, RORγt, Foxp3, IL-17a, and IFN-γ were all

severely reduced in mutant T cells relative to the WT cells in the SILP and mLN (Figures

S3A,B). The same pattern was observed in the EAE model, using mixed bone marrow chimeric

mice (Figures 1E, and S3C), indicating that Tpi1 is essential for the differentiation of all CD4^{+} T

cell subsets.

Gpi1 is selectively required by inflammatory but not homeostatic Th17 cells

The higher glycolytic activity of the inflammatory Th17 cells suggested that they may be more

sensitive than the homeostatic Th17 cells to inhibition of glycolysis, which might be achieved by

inactivation of *Tpi1, Gpi1 or Ldha* (explained in figure S4A). However, neither cell type is likely

to survive a complete block in glycolysis by *Gapdh* knockout. To test the hypothesis, we

assessed the ability of co-tranferred control and targeted TCR transgenic T cells to differentiate

into Th17 cells in models of homeostasis (7B8 T cells in SFB-colonized mice (Yang et al.,

2014)) and inflammatory disease (2D2 T cells for EAE (Bettelli et al., 2003) and HH7.2 T cells

for *Helicobacter*-dependent transfer colitis (Xu et al., 2018)) (Figure 2A). The genes of interest

(indicated in Figure S4A) or the control gene *Olfr2* were inactivated by guide RNA

electroporation of naïve isotype-marked CD4$^+$ TCR and Cas9 transgenic T cells (Platt et al.,

2014), which were then transferred in equal numbers into recipient mice (Figures 2A, S4B).

Targeting was performed with sgRNAs that achieved the highest in vitro knockout efficiency

(~80-90%) (Figures S4C-E), which corresponded to the frequency of targeted cells at 4 days

after electroporation and transfer into mice, as assessed by targeting of GFP (Figure S4B). This

strategy also allowed for efficient double gene KO (Figure S4F).

In both SFB and EAE T cell transfer experiments, there were roughly 9-fold fewer *Tpi1*-

targeted than co-transferred control Th17 cells in the SILP or the SC, respectively (Figures 2B-

D), consistent with the *in vitro* KO efficiency (Figure S4C), suggesting that the majority of the

Tpi1 KO cells were eliminated in the total T cell pool. The remaining cells expressed RORγt and

IL-17a at a similar level to control populations (Figures S5A-D), which was not observed in the

previous *Tpi1* bone marrow-reconstituted mice (Figures S3B), suggesting that they were not

targeted. These results indicated that the electroporation transfer system could be successfully

used to evaluate T cell gene functions *in vivo*.

Similar to the *Tpi1* KO, cells deficient for *Gapdh* were eliminated in all three models tested

(Figures 2B-E). Inactivation of *Ldha* resulted in loss of the cells in the EAE model, and a less

striking, but still substantial, cell number reduction in the homeostatic SFB model (Figures 2B-

E). In contrast, although *Gpi1* ablation reduced cell number by 75% in the inflammatory EAE

and colitis models, it had no significant effect on cell number in the homeostatic Th17 cell model

(Figures 2B-E). Consistent with the reduction in cell number, *Gpi1*-mutant 2D2 cells were

unable to induce EAE (Figure 2F), suggesting that Gpi1 is essential for pathogenicity. We also

investigated Th17 cell differentiation and cytokine production by *Gpi1*-deficient cells in the SFB

model, and found no difference from co-transferred control cells in terms of RORγt/Foxp3

expression (Figures 2G) or IL-17a production, as demonstrated using 7B8 cells from mice bred

to the *Il17a$^{GFP/+}$* reporter strain (Figure 2H). *In vitro* restimulation further confirmed that

transferred Gpi1-targeted 7B8 cells were indeed Gpi1 deficient (Figures S5C, D). Collectively,

these data suggest that Gpi1 is selectively required by the inflammatory encephalitogenic and

colitogenic Th17 cells, but not by homeostatic SFB-induced Th17 cells. In contrast, Gapdh,

Tpi1 and Ldha are required for the accumulation of both types of Th17 cells.

PPP compensates for Gpi1 deficiency in the homeostatic SFB model

To explain why Gpi1 is dispensable in the homeostatic SFB model while other glycolysis genes

are not, we hypothesized that the PPP shunt might maintain some glycolytic activity in the *Gpi1*

KO cells and thus compensate for Gpi1 deficiency (Figure S4A). We tested the hypothesis with

in vitro isotope tracing. Irrespective of the non-pathogenic (npTh17, cultured in IL-6 and TGF-

β (Veldhoen et al., 2006)) or pathogenic (pTh17, cultured in IL-6, IL-1β, and IL-23 (Ghoreschi

et al., 2010)) conditions used, *in vitro* differentiated Th17 cells displayed less of a growth defect

(25% reduction) following targeting of *Gpi1* than upon inactivation of *Gapdh* (70% reduction)

(Figure 3A). This result resembles the different sensitivities observed in the homeostatic SFB

model, suggesting that the cytokine milieu of the pathogenic and homeostatic Th17 cells does

not account for the phenotypes of the *Gpi1* mutant mice. We incubated np/p Th17 cells with

$^{13}C_{1,2}$-glucose and analyzed downstream ^{13}C label incorporation. The $^{13}C_{1,2}$-glucose tracer is

catabolized through glycolysis via Gpi1, producing pyruvate or lactate containing two heavy

carbons (M2), while Gpi1-independent shunting through the PPP produces pyruvate and lactate

containing one heavy carbon (M1) (Figure 3B) (Lee et al., 1998). Consistent with our hypothesis,

we observed ~ 3-fold increase in relative PPP activity in the *Gpi1* KO Th17 cells compared to

the *Olfr2* KO control, irrespective of npTh17 or pTh17 condition (Figure 3C), indicating that

Gpi1 deficiency maintains active glycolytic flux through PPP shunting.

To test whether PPP activity compensates for Gpi1 deficiency *in vivo* in the homeostatic SFB

model, we knocked out both *Gpi1* and *G6pdx*, which catalyzes the initial oxidative step in the

PPP, with the CRISPR-electroporation transfer system. Combined deletion of these two genes

would be expected to block glycolysis, similar to *Gapdh* KO. Consistent with this hypothesis, the

Gpi1/G6pdx double KO reduced cell number by about 75%, which is similar to that with the

Gapdh single KO, yet significantly different from that with inactivation of either *Gpi1* or *G6pdx*

alone (Figure 3D). These results suggest that *in vivo* PPP activity maintains viability of the

homeostatic SFB-induced Th17 cells lacking Gpi1.

Increased mitochondrial respiration additionally compensates for Gpi1 deficiency in the

homeostatic SFB model

To determine the impact of Gpi1 deficiency on aerobic glycolysis activity, we quantified lactate

production by GC-MS and found it to be markedly reduced in *Gpi1* KO cells (Figure 4A). A

significant reduction in lactate production may suggest compromised glycolytic ATP production

in *Gpi1* KO cells. As *in vitro* cultured *Gpi1* KO Th17 cells and homeostatic SFB-specific *Gpi1*

KO Th17 cells were largely unaffected in terms of cell number and cytokine production, we

speculate that mitochondrial respiration may provide an alternative source of ATP to

compensate. Indeed, in vitro cultured *Gpi1* KO cells displayed higher ATP-linked respiration

rate than *Olfr2* control cells, irrespective of npTh17 or pTh17 cell culture condition (Figure

4B,C). Furthermore, upon Antimycin A (Complex III inhibitor) treatment, *Gpi1* KO cells were

decreased in cell number, similar to *Gapdh*-deficient cells (Figure 4D), consistent with increased

mitochondrial respiration compensating for Gpi1 deficiency in the *in vitro* cell culture system.

    To determine whether increased respiration rescues Gpi1 deficiency *in vivo* in the

homeostatic SFB model, we targeted *Gpi1* in 7B8 T cells along with genes responsible for the

oxidation of three major substrates feeding the TCA cycle: *Pdha1* (pyruvate), *Gls1* (glutamine),

and *Hadhb* (fatty acids). Following co-transfer of control cells and cells lacking either *Pdha1*,

Gls1, or *Hadhb*, there was no defect in Th17 cell number, RORγt expression, or cytokine

production (data not shown). However, only double knockout of *Gpi1* and *Pdha1* led to a

significant reduction in cell number, similar to that with the *Gapdh* KO cells (Figure 4E),

suggesting that mitochondrial respiration through pyruvate oxidation compensates for Gpi1

deficiency in homeostatic Th17 cell differentiation induced by SFB colonization.

PPP utilization favors biosynthetic metabolite synthesis in Gpi1-deficient cells

Compensation for *Gpi1* deletion by the combination of the PPP and mitochondrial respiration

suggests the existence of a partial metabolic redundancy for Gpi1. To understand the

compensatory mechanism, we performed kinetic flux profiling, employing GC-MS to measure

the passage of isotope label from $^{13}C_6$-glucose into downstream metabolites (Yuan et al., 2008).

The resulting kinetic data, along with metabolite abundance, was used to quantify metabolic flux

in npTh17 cells prepared from *Gpi1* KO, *Olfr2* KO and cells treated with koningic acid (KA), to

inhibit Gapdh (Liberti et al., 2017), since more than 50% of cells in the *Gapdh* KO culture were

WT escapees (Figure S4C)).

Loss of Gpi1 resulted in reduction of glucose uptake (Figure 5A) and in substantially

decreased transmission of isotopic label to intracellular pyruvate and lactate, whose abundance

was also reduced (Figures 5C,D,I). Interestingly, a lower lactate production/glucose

consumption ratio was observed in *Gpi1* KO than control cells (Figure 5B), suggesting that

Gpi1-deficient cells may be more inclined to use glucose carbons for the synthesis of

biosynthetic metabolites, rather than lactate production. Indeed, the labeling rate and total

abundance of the biosynthetic metabolites alanine, serine, glycine, and citrate either did not

change or were slightly increased in the *Gpi1* KO cells (Figures 5E-I). The flux of pyruvate,

lactate, alanine, serine, and glycine in the Gpi1-deficient cells, as compared to *Olfr2* KO control

(Figure 5J), was consistent with the PPP supporting the branching pathways of glycolysis

important for biosynthetic metabolite production, despite a reduced glycolysis rate. In contrast,

Gapdh inhibition by KA treatment almost completely prevented the synthesis of downstream

metabolites (Figures 5C-H) and lactate production (Figure 5K). However, it did not induce an

increase in OCR (Figures 5K, L), possibly due to the severely limited production of pyruvate.

Taken together, we conclude that the reduced amount of glucose metabolized via the PPP by

Gpi1-deficient cells was sufficient to support the production of glycolytic intermediates and

maintain pyruvate oxidation. *Gpi1*-deficient cells increased their mitochondrial respiration to

compensate for the loss of glycolytic flux. In contrast, glycolytic blockade by Gapdh inhibition

completely prevented carbon flux needed for key biosynthetic metabolites, which is incompatible

with cell viability.

Gpi1 is essential in the hypoxic setting of Th17-mediated inflammation

To understand why inflammatory Th17 cells are particularly sensitive to Gpi1 deficiency, we

reasoned that metabolic compensation (either through PPP or mitochondrial respiration), which

occurs in the homeostatic model, may be restricted in inflammatory Th17 cells. *G6pdx* KO in the

EAE model resulted in a 70% reduction in 2D2 cell number in the spinal cord (Figures S6A,B),

suggesting that PPP flux is still active in the setting of inflammation. On the other hand, a

number of reports have shown that inflamed tissues, including the spinal cord in EAE and the

LILP in colitis, are associated with low oxygenation (hypoxia) (Davies et al., 2013; Johnson et

al., 2016; Karhausen et al., 2004; Naughton et al., 1993; Ng et al., 2010; Peters et al., 2004; Van

Welden et al., 2017; Yang and Dunn, 2015). This suggests that impaired mitochondrial

respiration might occur which would result in an inability to provide sufficient ATP to

compensate for energy loss due to Gpi1 deficiency in these tissues. To test this hypothesis, we

cultured Th17 cells in normoxic (20% O_2) and mild hypoxic (3% O_2) conditions, and found that

Gpi1 KO Th17 cells in the hypoxic state, irrespective of npTh17 or pTh17 differentiation

protocols, displayed a growth defect similar to that occurring with Gapdh deficiency (Figure

6A). Furthermore, Olfr2 control npTh17 cells displayed about a 2-fold increase in lactate

production in hypoxic compared to normoxic conditions (Figure S6C), indicating that hypoxic

cells rely more on glycolysis to generate ATP. *Gpi1* KO cells had substantially decreased lactate

production in both normoxic and hypoxic conditions (Figure S6C), but there was reduced

intracellular ATP only with hypoxia (Figure S6D), suggesting that restrained mitochondrial

respiration in hypoxia cannot compensate for Gpi1 inactivation, and hence leads to energy crisis.

We next investigated whether the inflamed tissue in the EAE model differs from the

homeostatic tissue in the SFB model with regard to oxygen availability. To address this, we

performed I.V. injection of pimonidazole, an indicator of O_2 levels, into mice with EAE or with

SFB colonization, and found that $CD4^+$ T cells in the EAE model spinal cord had two-fold

higher pimonidazole staining than in the dLN and ~2.7-fold compared to $CD4^+$ T cells in SILP

and mLN (Figure 6B). This result suggests that $CD4^+$ T cells in the spinal cord of the EAE

model experience greater oxygen deprivation than in other tissues examined. The increased

labeling in the spinal cord was observed in all subsets of $CD4^+$ T cells (Figures S6E,F),

indicating that decreased oxygen availability is likely a tissue feature.

Hif1α is an oxygen sensor that, upon activation by hypoxia, initiates transcription of genes,

including glycolysis genes, to allow cells to adapt in poorly oxygenated tissue (Corcoran and

O'Neill, 2016; Lee et al., 2020; Semenza, 2013). To further examine the oxygenation level of

SILP and SC, we investigated the effect of *Hif1a* deficiency in the different models of $CD4^+$ T

cell differentiation. Using chimeric *Rag1* KO mice reconstituted with equal numbers of

Hif1a$^{+/+}$CD4^{cre} and *Hif1a$^{f/f}$CD4^{cre}* bone marrow cells, there was ~ 40% reduction of Hif1α-

deficient $CD4^+$ T cells in the spinal cord of mice with EAE, but little difference in the SILP

(Figures 6C-E). Furthermore, in the SILP of SFB-colonized mice, there was no significant

difference in RORγt and Foxp3 expression or IL-17a and IFN-γ production between WT and KO

CD4$^+$ T cells (Figures S6G,H). In contrast, in the spinal cord of the EAE mice, there was a slight

decrease in the Th17 cell fractions and increase in the Foxp3$^+$ fraction in the *Hif1a* KO compared

to WT CD4$^+$ T cells (Figures S6G, H). Overall, these data suggest that Hif1α is functionally

important for the pathogenic Th17 cells (as well as Treg cells) in the EAE model, but not for

homeostatic SILP Th17 cells, reinforcing the conclusion of the pimonidazole experiment that T

cells in the inflamed spinal cord experience greater oxygen deprivation than those in the healthy

small intestine lamina propria.

To address when and how Gpi1 deficiency impairs inflammatory Th17 cell differentiation

and/or function, we performed a time course experiment to assess the kinetics of the co-

transferred *Olfr2* KO control and *Gpi1* KO 2D2 cells in the EAE model. *Gpi1* KO T cells, like

Olfr2 KO control cells, were maintained in the dLN until day 13, the last time point when

transferred 2D2 cells were still detectable in the dLN (Figure 6E). However, at 13 days post

immunization, *Gpi1* mutant cell number was reduced by 50% compared to control T cells in the

spinal cord (Figure 6E). This finding is consistent with the previous observation that the spinal

cord, but not the dLN, is hypoxic. Accordingly, we observed reduced proliferation of *Gpi1* KO

2D2 cells in the spinal cord but not in the dLN (Figures 6F and S6I). Importantly, 2D2 cells in

the dLN of mice with EAE proliferated at a markedly faster rate than 7B8 cells in the mLN of

the SFB model (Figures S6J,K). This demonstrates that Gpi1-deficient 2D2 cells can maintain

the higher demand for energy and biomass associated with rapid proliferation and emphasizes

the role of the SC hypoxic environment in the T cell's dependence on Gpi1. We also examined

Th17 cell differentiation/cytokine production, by staining for RORγt/Foxp3 and IL-17a,

apoptosis, by staining for the active form of caspase 3, and potential for migration, by staining

for Ccr6, and observed no differences between *Gpi1*-deficient and co-transferred control *Olfr2*

KO 2D2 T cells (Figures S6L-P). We conclude that Th17 cells increase glycolysis activity to

adapt to the hypoxic environment in the EAE spinal cord, and the restrained mitochondrial

respiration in this setting cannot compensate for the loss of glycolytic ATP production upon

Gpi1 inactivation, leading to energy crisis and cell elimination.

Discussion

Our results demonstrate the remarkable plasticity of cellular metabolism owing to redundant

components in the network, and highlight that plasticity can vary based on microenvironmental

factors. Due to the unique position of Gpi1 in the glycolysis pathway, its deficiency was

compensated by the combination of two metabolic pathways – the PPP and mitochondrial

respiration. The PPP maintained glycolytic activity, albeit at a reduced level, to fully support

biosynthetic precursor synthesis through branching pathways. Pyruvate, which is also produced

by way of the PPP, increased its flux into the TCA cycle, elevating ATP production from

mitochondrial respiration, thus compensating for loss of ATP production from glycolysis.

Therefore, upon Gpi1 inactivation, homeostatic SFB-induced Th17 cells reprogrammed their

metabolism, retaining the ability to provide adequate biomass and energy to support cell

function. In contrast, limitation of mitochondrial respiration due to low oxygen availability in the

inflamed EAE spinal cord rendered Gpi1-deficient cells unable to generate sufficient ATP

molecules either through glycolysis or mitochondrial respiration, resulting in energy crisis and

cell elimination. Inactivation of Gapdh, whose function, unlike that of Gpi1, cannot be

compensated by other metabolic pathways, completely blocked glycolysis and hence led to

elimination of both homeostatic and inflammatory Th17 cells. Our results suggest not only that

362 suppressing glycolysis by Gpi1 inhibition could be a well-tolerated treatment for hypoxia-related

363 autoimmune diseases, but also that metabolic redundancies can be exploited for targeting disease

364 processes in selected tissues.

366 **Metabolic requirements for CD4$^+$ T cell activation and differentiation**

367 We found that CD4$^+$ T cells (pathogenic or Treg) in the EAE model had greater expression of

368 glycolytic genes than homeostatic CD4$^+$ T cells, which is consistent with the recent

369 demonstration that *Citrobacter*-induced inflammatory Th17 cells were more glycolytic than

370 SFB-induced homeostatic Th17 cells (Omenetti et al., 2019). Our data suggest that T cells adapt

371 to the hypoxic environment in the inflamed spinal cord by up-regulating glycolysis, most likely

372 by hypoxia-stabilized Hif1α.

373  In spite of the difference in glycolytic activity, blockade of glycolysis by *Gapdh* KO is

374 incompatible with Th17 cell expansion and differentiation in either homeostatic or inflammatory

375 models. However, the finding that Gpi1-deficient 2D2 T cells proliferated normally in the dLN

376 of the EAE model suggests that T cell activation and differentiation *in vivo* do not necessarily

377 need a fully functional glycolytic pathway. With the PPP providing sufficient metabolites for

378 biomass production and mitochondria generating greater amounts of ATP, T cell proliferation

379 and execution of the non-pathogenic Th17 program *in vivo* can be largely maintained in the *Gpi1*

380 KO cells. Our data highlight the complexity of the glycolysis pathway and suggest that not every

381 enzyme is equally important to glycolytic activity and cell function.

382  Treg cell differentiation has been proposed to depend on OXPHOS (Kullberg et al., 2006;

383 Macintyre et al., 2014; MacIver et al., 2013; Michalek et al., 2011), and was promoted by

384 glycolysis inhibition *in vitro* with 2DG (Shi et al., 2011). Our *in vivo* genetic data however

support an indispensable role of glycolysis in Treg cell differentiation, as demonstrated by

complete loss of the intestinal and SC Foxp3[+] T cells in mice deficient for *Tpi1* in T cells.

One intriguing finding that we have not characterized in detail is the dispensable role of

Pdha1, Gls1, and Hadhb in SFB-induced homeostatic Th17 cell differentiation. While OXPHOS

is essential for SFB-specific Th17 cells (data not shown), their ability to differentiate in the

absence of genes controlling metabolite fueling of the TCA cycle suggests potential

redundancies. Furthermore, the discrepancy between the in vivo and in vitro function of

Pdha1(as shown in this study) and Gls1 (by comparing the in vivo data in this study with

reported in vitro data (Johnson et al., 2018)) suggests that the microenvironment is critical for T

cell metabolism and gene function, as does a recent report showing distinct metabolism profiles

of in vivo and in vitro activated T cells (Ma et al., 2019).

Gpi1 as a drug target for hypoxia-related diseases

Our study reveals that Gpi1 may be a good drug target for Th17-mediated autoimmune diseases.

The sensitivity to Gpi1 deficiency is dependent on whether cells are competent to increase

mitochondrial respiration. Therefore, other disease processes involving hypoxia-related

pathologies may also be susceptible to Gpi1 inhibition. A recent report showed that 1% O_2

almost completely inhibited *in vitro* growth of *Gpi1*-mutant tumor cell lines, but the proliferation

defect was much milder when cells were cultured in normoxic condition (de Padua et al., 2017).

This suggests that hypoxia-mediated sensitivity to Gpi1 deficiency can be a general

phenomenon, from Th17 cells to tumor cells, which may extend the deployment of Gpi1

inhibition to a broader range of diseases.

Sensitivity to inhibition of both Hif1α and Gpi1 is dependent on oxygen availability. In the

mixed bone marrow chimera experiments with the EAE model, there was less than a 2-fold

reduction in *Hif1a* mutant compared to wild-type spinal cord CD4$^+$ T cells, whereas *Gpi1*-

targeted T cells were reduced 5-fold compared to wild-type cells in the transfer model of EAE.

Hif1a inactivation efficiency was almost complete (data not shown), while *Gpi1* KO efficiency

was roughly 80%, which suggests that Gpi1 deficiency resulted in a more severe defect than

Hif1a deficiency when the T cells were in a hypoxic environment. This is not unexpected, as

Gpi1 itself participates in the glycolysis pathway while Hif1α regulates the transcription of genes

in the pathway. Although EAE was attenuated following *Hif1a* inactivation in CD4$^+$ T cells

(Dang et al., 2011; Shi et al., 2011), our results suggest that Gpi1 inhibition may be at least as

effective a means of blocking cell function in hypoxic microenvironments.

While our data suggest that hypoxia is an essential factor that restains proliferation of Gpi1-

deficient Th17 cells in the spinal cord, we cannot exclude the possibility that other

environmental factors also contribute to the phenotype, as oxygenation level is certainly not the

only difference between the inflamed spinal cord and the healthy small intestine lamina propria.

Gpi1 targeting can be tolerated

Glycolysis is a universal metabolic pathway whose blockade by inhibitors such as 2DG can yield

adverse side effects even at dosages too low to control tumor growth (Raez et al., 2013). 2DG

inhibits the first three steps of glycolysis, an effect that is recapitulated by Gapdh inhibition,

given that no alternative pathway exists for glucose catabolism. Although glycolysis is essential

for T cell activation, treating diseases caused by autoimmune T cells with drugs that fully inhibit

the pathway, e.g. 2DG or KA, would also likely be too toxic at potentially efficacious dosages.

In light of the results presented here, Gpi1 may represent a better drug target than other

glycolysis gene products for treating Th17 cell-mediated autoimmune diseases, as targeting Gpi1

may be well tolerated by most cells and tissues, thus causing minimal side effects.

This hypothesis is supported by genetic data obtained from patients with glycolysis gene

deficiencies. With few exceptions, patients deficient for *Tpi1* or *Pgk1* have combined symptoms

of anemia, mental retardation and muscle weakness [Online Mendelian Inheritance in Man

(OMIM) entry 190450, 311800] (Schneider, 2000). Mutations in glycolysis gene products that

have redundant isozymes can result in dysfunction of cell types in which only the mutated

isozyme gets expressed, such that *Pgam2* deficiency results in muscle breakdown and *Pklr*

deficiency causes anemia (Zanella et al., 2005). The human genetic data are thus consistent with

the prediction that inhibition of glycolysis enzymes would result in serious side effects.

However, the vast majority of patients with *Gpi1* mutations, which can result in preservation of

less than 20% enzymatic activity, have mild to severe anemia that is treatable, with no

neurological or muscle developmental defects (Baronciani et al., 1996; Kanno et al., 1996;

Kugler et al., 1998; McMullin, 1999; Zaidi et al., 2017) [OMIM entry 172400]. These clinical

data indicate that neurons and muscle cells can afford losing most of their Gpi1 activity even

though they require glycolysis, and hence they behave similarly to homeostatic Th17 cells.

Collectively, these patient data strongly support our proposal that Gpi1 can be a therapeutic

target.

Selective metabolic redundancies permit selective cell inhibition

Metabolic redundancy has been demonstrated in many biological systems, from bacteria to

cancer cell lines (Bulcha et al., 2019; Deutscher et al., 2006; Horlbeck et al., 2018; Segre et al.,

2005; Thiele et al., 2005; Zhao et al., 2018). Our study demonstrates compensatory metabolic

pathways in CD4$^+$ T cells *in vivo*, and highlights that such redundancy can be selective,

depending on the cellular microenvironment. Indeed, specific inhibition of pathogenic cells can

be achieved by metabolic targeting of selective redundant components. Therefore,

characterization of the metabolic redundancy of cells in microenvironments with varying oxygen

level, pH, abundance of extracellular metabolites, redox status, cytokine milieu, or temperature,

may provide opportunities for metabolic targeting of cells in selected tissue microenvironments

for therapeutic purposes.

Acknowledgements: We thank the NYU Langone Genome Technology Center for expert

library preparation and sequencing. This shared resource is partially supported by the Cancer

Center Support Grant P30CA016087 at the Laura and Isaac Perlmutter Cancer Center. We thank

Drew R. Jones and Rebecca E. Rose in the NYU Metabolomics Laboratory for helpful

discussion and aid with LC-MS data generation and analysis. We thank S.Y. Kim in the NYU

Rodent Genetic Engineering Laboratory (RGEL) for generating the Tpi1 conditional KO mice.

We thank Anne R. Bresnick (Department of Biochemistry, Albert Einstein College of Medicine)

for sharing the S100a4 mutant mice. This work was supported by National Multiple Sclerosis

Society Fellowship FG 2089-A-1 (L.W.), by Immunology and Inflammation training grant

T32AI100853 (C.N.), by NIH grant R01AI121436 (D.R.L.), by the Howard Hughes Medical

Institute (D.R.L.), and the Helen and Martin Kimmel Center for Biology and Medicine (D.R.L.)

Author Contributions: L.W. and D.R.L. conceived the project, and wrote the manuscript. L.W.

designed, performed experiments, and analyzed the data. L.W. and K.E.R.H. designed,

performed the GC-MS flux, tracing, and seahorse experiments, analyzed the data. Y.H. and R.S.

analyzed the scRNAseq data. L.K. analyzed the bulk RNAseq data. C.A., W.L., D.L., and

H.M.S. helped with mouse experiments. C.N. and W.L. helped with the development of the

CRISPR/Cas9 transfection method. S.J. and R.P. helped with GC-MS experiments. K.E.R.H.,

J.J.L., R.P., M.E.P., T.Y.P., and A.C.K. assisted with the analysis and interpretation of the

metabolic data. K.E.R.H., H.M.S., S.J., J.J.L., R.P., M.E.P., T.Y.P., A.C.K., and R.S. edited the

manuscript. D.R.L. supervised the work.

Declaration of Interests: D.R.L. consults and has equity interest in Chemocentryx, Vedanta,

and Pfizer Pharmaceuticals. The NYU School of Medicine has filed a provisional patent

application related to this work.

Main figure titles and legends

Figure 1. Encephalitogenic Th17 cells have higher glycolysis pathway gene expression than

homeostatic SFB-induced Th17 cells.

(A) Clinical disease course of MOG-CFA-induced EAE in $Tpi1^{f/f}Cd4^{cre}$ (n=7) and $Tpi1^{+/+}Cd4^{cre}$

littermate control mice (n=8). The experiment was repeated twice with the same

conclusion. P values were determined using two-way ANOVA, * (P<0.05), ****

(P<0.0001).

(B) Heatmap of glycolysis pathway genes expressed in spinal cord Th1[*], Th17, and Treg cells in

mice with EAE (day 15, score 5) compared to Th17 and Treg cells in the SILP of SFB-

colonized mice. Genes are arranged according to the fold change between EAE Th17 and

SFB Th17.

(C) Experimental design of the Tpi1 bone marrow reconstitution experiment. Bone marrow cells

from *Tpi1*$^{+/+}$ *Cd4*^{cre} CD45.1/2 and *Tpi1*$^{f/f}$*Cd4*^{cre} CD45.2/2 donor mice were transferred into

lethally irradiated CD45.1/1 recipients. Peripheral blood was collected two months later for

reconstitution analysis. dLN and SC of the EAE model, and mLN and SILP of the SFB

model were dissected at day 15 post EAE induction or SFB colonization.

(D,E) Cell number analysis of *Tpi1* WT and KO CD4$^+$ T cells in the SFB model (D) and the

EAE model (E). Left panels, representative FACS plots showing the frequencies of the

CD45.1/2 and the CD45.2/2 B cells among all B cells in peripheral blood and of the

CD45.1/2 and the CD45.2/2 CD4$^+$ T cells among all CD4$^+$ T cells in peripheral blood, mLN

or dLN, and SILP of SFB colonized mice or SC of EAE mice. Middle panels, compilation of

the cell number ratio of CD45.2/2 to CD45.1/2. Right, normalized KO/WT total CD4$^+$ T cell

number ratio in each tissue. The cell number ratio of peripheral B cells was set as 1. P values

were determined using paired t tests.

See also Figures S1,2,3.

Figure 2. Gpi1 is selectively required by inflammatory encephalitogenic or colitogenic Th17

cells, but not by homeostatic SFB-induced Th17 cells.

(A) Experimental setup. TCR transgenic Cas9-expressing naïve CD4$^+$ T cells were electroporated

with guide amplicons, and co-transferred with *Olfr2* amplicon-electroporated control cells

into recipient mice that were immunized for EAE induction (EAE model), or had been

colonized with SFB (SFB model) or *Helicobacter hepaticus* (Colitis model).

(B) Representative FACS plots showing the frequencies of *Olfr2* KO cells and the co-transferred

glycolytic gene KO cells among total CD4$^+$ T cells in each model.

(C-E) Compilation of cell number ratio of gene KO group to the co-transferred *Olfr2* control for

each model, as shown in panel B. The ratio was normalized to *Olfr2*/*Olfr2* co-transfer

(*Olfr2*/*Olfr2* cell number ratio was set to 1). Three independent experiments were performed

with the same conclusion.

(F) Clinical disease course of EAE in *Rag1* KO mice receiving *Olfr2* KO 2D2 cells or *Gpi1* KO

2D2 cells. Experiment was conducted as illustrated in Figure S2C. n=5 mice/group. The

experiment was repeated twice with the same conclusion.

(G) Left, representative FACS plot showing RORγt and Foxp3 expression of co-transferred

targeted 7B8 cells isolated from the SILP of recipient SFB-colonized mice at day15. Right,

ratio of the percentage of RORγt$^+$ Foxp3$^-$ in *Gpi1*- or *Olfr2*-targeted versus co-transferred

Olfr2 control cells in each recipient.

(H) Left, representative FACS histogram showing the overlay of IL17a-GFP expression of co-

transferred *Olfr2* KO and *Gpi1* KO 7B8 cells isolated from the SILP of SFB-colonized mice

at day15. Right, ratio of the percentage of IL-17a$^+$ cells in *Gpi1*- or *Olfr2*-targeted versus co-

transferred *Olfr2* control cells in each recipient.

P values were determined using t tests.

See also Figures S4,5.

Figure 3. PPP activity rescues *Gpi1* deficiency in the homeostatic SFB-induced Th17 cells

(A) Relative number of Th17 cells cultured for 120 h *in vitro*. Naïve Cas9-expressing CD4$^+$ T

cells were electroporated with corresponding guide-amplicons and cultured in both npTh17

condition and pTh17 condition. Cell number was normalized to *Olfr2* KO group (*Olfr2* cell

number was set as 100). N=3, data are representative of three independent experiments.

(B) Schematic of $^{13}C_{1,2}$-glucose labeling into downstream metabolites. ^{13}C is labeled as filled

circle. Catalytic reactions of PPP are labeled as red arrows, while those of glycolysis are in

black.

(C) Relative PPP activity of *in vitro* cultured Th17 cells. Relative PPP activity from $^{13}C_{1,2}$-

glucose was determined using the following equation: PPP= M1/ (M1+M2), where M1 is the

fraction of $^{13}C_{1,2}$-glucose derived from the PPP and M2 is the fraction of $^{13}C_{1,2}$-glucose

derived from glycolysis.

(D) 7B8 Cas9 transgenic naïve $CD4^+$ T cells were electroporated with a mixture of two guide

amplicons (for double KO) before being transferred into recipient mice. Left, representative

FACS plots showing the frequencies of co-transferred *Olfr2+Olfr2* control $CD4^+$ T cells and

the indicated gene KO $CD4^+$ T cells in the SILP of the SFB model. Right, compilation of cell

number ratios of the indicated gene KO combinations to the co-transferred *Olfr2+Olfr2*

control. The ratio was normalized to *Olfr2+Olfr2/Olfr2+Olfr2* co-transfer. Two independent

experiments were performed with same conclusion.

P values were determined using t tests.

Figure 4. Mitochondrial respiration compensates for Gpi1 deficiency in the homeostatic

SFB Th17 cells

(A) Lactate secretion of *in vitro* cultured *Olfr2* and *Gpi1* KO np/p Th17 cells. Cells were cultured

as in Figure 3A. At 96 h cells were re-plated in fresh RPMI medium with 10% dialyzed FCS,

and supernatants were collected 12 h later for lactate quantification by GC-MS.

(B) Seahorse experiment showing the oxygen consumption rate (OCR) of *in vitro* cultured Th17

cells at baseline and in response to oligomycin (Oligo), carbonyl cyanide 4-

(trifluoromethoxy) phenylhydrazone (FCCP), and rotenone plus antimycin (R + A). 10

replicates were used for the npTh17 *Olfr2* and *Gpi1* targeted cells, 7 replicates for the pTh17

Olfr2, and 6 replicates for the pTh17 *Gpi1* sets of targeted cells.

(C) ATP linked respiration rate quantified based on experiments in panel (B). Representative

data from three independent experiments.

(D) Relative number of Th17 cells cultured for 120 h *in vitro* (as in Figure 3A), in the presence or

absence of 10 nM antimycin A. Cell number was normalized to control (*Olfr2*-targeted cell

number was set as 100). n=3, data are representative of two independent experiments.

(E) 7B8 Cas9 naïve CD4$^+$ T cells were electroporated with a mixture of two guide amplicons (for

double KO) before transfer into recipient mice. Left, representative FACS plots showing the

frequencies of co-transferred *Olfr2+Olfr2* control and indicated gene KO CD4$^+$ T cells in the

SILP of the SFB model at day 15. Right, compilation of cell number ratios of glycolysis gene

KO group to the co-transferred *Olfr2+Olfr2* control. The ratio was normalized to

Olfr2+Olfr2/Olfr2+Olfr2 co-transfer. Three independent experiments were performed with

same conclusion.

P values were determined using t tests.

Figure 5. PPP supports normal biomass synthesis in the *Gpi1*-deficient npTh17 cells

Olfr2- or *Gpi1*-targeted npTh17 cells (cultured for 96 h as in Figure 3A) were collected for

further metabolic analysis.

(A, B) Cells were re-plated in U-^{13}C$_6$-glucose tracing medium to measure (A) glucose

consumption rate, and (B) ratio of released lactate to consumed glucose by LC-MS.

(C-H) Cells were re-plated in U-$^{13}C_6$-glucose tracing medium, and cell pellets were collected at

different time points to measure ^{13}C incorporation kinetics of pyruvate (C), lactate (D), serine

(E), glycine (F), alanine (G) and citrate (H) in *Olfr2* KO, *Gpi1* KO, or koningic acid (KA)-

treated npTh17 cells by GC-MS.

(I) Intracellular abundance of pyruvate, lactate, alanine, serine, glycine and citrate in targeted

npTh17 cells, as determined by GC-MS. Raw ion counts were normalized to extraction

efficiency and cell number. The abundance is shown relative to the *Olfr2* sample set as 100.

(J) Quantification of the production flux of serine, glycine, alanine, pyruvate and lactate based on

the ^{13}C incorporation kinetics and intracellular abundance of each metabolite. Three

replicates for each sample.

(K) OCR (top) and Extracellular Acidification Rate (ECAR) (bottom) of targeted or KA-treated

npTh17 cells at baseline and in response to oligomycin (Oligo), carbonyl cyanide 4-

(trifluoromethoxy) phenylhydrazone (FCCP), and rotenone plus antimycin (R + A). 10

replicates for each condition.

(L) ATP-linked respiration rate, quantified based on panel K. Data are representative of two

independent experiments.

P values were determined using t tests.

Figure 6. Hypoxia in the inflamed spinal cord of the EAE model restrains mitochondrial

respiration, leading to the elimination of Gpi1-deficient Th17 cells

(A) Relative number of gRNA-targeted Th17 cells cultured for 120 h *in vitro* in either normoxic

(20% O_2) or hypoxic (3% O_2) conditions. Cell number was normalized to *Olfr2* KO (*Olfr2*

cell number was set as 100).

(B) Representative FACS histogram showing pimonidazole labeling of total CD4$^+$ T cells

isolated from the SC and the dLN at day 15 after induction of EAE and from the SILP and

the mLN of SFB-colonized mice.

(C) Experimental design of the Hif1a bone marrow reconstitution experiment. *Rag1* KO mice

were lethally irradiated and reconstituted with equal numbers of bone marrow cells from

Hif1a$^{+/+}$ Cd4^{cre} CD45.1/2 and *Hif1a*$^{f/f}$ *Cd4*^{cre} CD45.1/1 donors. CD4$^+$ T cells were examined

2-months later in peripheral blood (PB), after which mice were either gavaged with SFB or

immunized for EAE induction, and CD4$^+$ T cells from SC or SILP from each model were

isolated at day15 for examination.

(D) Top, representative FACS plots showing the frequencies of *Hif1a$^{+/+}$ Cd4*^{cre} and *Hif1a*$^{f/f}$*Cd4*^{cre}

CD4$^+$ T cells in the peripheral blood and the SILP or the SC of the SFB and the EAE models,

respectively. Bottom, compilation of the KO/WT total CD4$^+$ cell number ratio obtained in

the PB and the SILP or the SC of the SFB and EAE models. Two experiments were

performed with the same conclusion.

(E) Time-course of 2D2 cas9 co-transfer experiment. Left, representative FACS plot showing the

frequencies of the *Olfr2* control and the co-transferred *Olfr2* or *Gpi1* KO 2D2 cells in total

CD4$^+$ T cells isolated from the dLN or the SC at different time points of EAE. Right,

compilation of the KO/Olfr2 control cell number ratio in the dLN or the SC. Data are

representative of two independent experiments.

(F) EDU in vivo labeling of co-transferred 2D2 cells in the EAE model. Left, representative

FACS plots showing the frequencies of EDU$^+$ among co-transferred *Olfr2* control and *Gpi1*

KO 2D2 cells in the SC at the onset of EAE (day10). Middle, compilation of the percentage

EDU$^+$ cells in the *Olfr2* control and the *Gpi1* KO cells. Right, the *Gpi1* KO/*Olfr2* control

ratios of percentage EDU$^+$ cells. Two experiments were performed with the same conclusion.

P values were determined using t tests. Panel D, F, and Day13 dLN and SC comparison in panel

E were analyzed with paired t tests.

See also Figure S6.

Supplemental figure titles and legends

Figure S1. Single cell RNAseq analysis of ex vivo CD4$^+$ T cells from spinal cord in the EAE

model and the SILP of SFB-colonized mice, related to Figure 1

(A) Experimental design of the scRNAseq analysis. Lethally irradiated *Rag1* KO mice were

reconstituted with equal numbers of bone marrow cells from isotype-marked *Il23r*$^{Gfp/+}$ and

Il23r$^{Gfp/Gfp}$ donors. The CD4$^+$ T cells of both *Il23r* het and KO background in the SC and

SILP of each model were sorted for scRNAseq analysis.

(B, C) UMAP projections of the scRNAseq data showing the five CD4$^+$ T cell sub-populations in

the SILP of SFB colonized mice and the four sub-populations in the SC of EAE mice.

(D, E) Violin plots showing the expression level of selected marker genes for each cluster in the

SFB model (D) and the EAE model (E).

(F, G) Distribution of *Il23r*$^{Gfp/+}$ and *Il23r*$^{Gfp/Gfp}$ cells in the UMAP projection of SILP CD4$^+$ T

cells (F) and EAE SC CD4$^+$ T cells (G).

(H) Pie chart showing the percentage of *Il23r*^{het} and *Il23r*^{KO} cells in each cluster as in panels D-

G. The total number of *Il23r*$^{Gfp/+}$ cells is scaled according to that of *Il23r*$^{Gfp/Gfp}$ cells for the

analysis (assuming same number of heterozygous and homozygous cells were analyzed).

The actual cell number of each cluster is indicated beneath the corresponding chart.

(I) Validation of marker gene expression in EAE clusters shown in panel E. Top, Klrc1 and

Klrd1 are co-expressed on cells that mainly produce IFN-γ (left), but are not expressed on

Treg cells (right), and are thus considered as Th1* (IFN-γ producing pathogenic Th17)

markers. Bottom left, Treg cells express less Rankl (encoded by *Tnfsf11*). Bottom right,

majority of the IFN-γ- and IL-17a-producing cells are Rankl$^+$.

(J, K) Pathway enrichment analysis showing the top 2 pathways activated in EAE-associated

Th17 (J) or Th1* (K) cells compared to SFB-induced Th17 cells.

(L, M) Volcano plots based on the scRNAseq analysis showing the differentially expressed

genes in EAE-associated Th17 (L) or Th1* cells (M) compared to SFB-induced Th17 cells.

Genes selected for functional screening are highlighted.

Figure S2. Tpi1 is essential for EAE-associated Th17 cell pathogenicity, related to Figure 1

and STAR Methods

(A) Schematic of the PSIN retroviral vector encoding the Thy1.1 reporter and two gRNAs.

(B) Knockout efficiency with single compared to double gRNA vectors. In vitro cultured CD4$^+$ T

cells were infected with the retrovirus containing a single or two gRNAs targeting *Cd4,* and

Cd4 expression was evaluated 72 h later.

(C) Experimental design for functional screening of pathogenic genes in the transfer EAE model.

Naïve 2D2 Cas9 CD4$^+$ T cells were activated in vitro, infected with the double gRNA

retrovirus, and differentiated into Th17 cells. Thy1.1 high cells were purified and transferred

into *Rag1* KO mice (50k cells/mouse), which were immunized with MOG/CFA at the same

time. EAE progression was monitored to evaluate the requirement for each gene in

pathogenesis.

(D) Clinical disease course of EAE in *Rag1* KO mice receiving control *(Olfr2* KO) 2D2 cells or

candidate gene KO 2D2 cells, and in MOG-CFA-immunized *S100a4* KO and WT littermate

control mice (6 pairs). The 2D2 transfer system was validated by failure of 2D2 cells

deficient in *Bhlhe40*, a gene known to be essential for T cell pathogenicity, to induce EAE.

At least 5 mice were used for each group, and each experiment was repeated at least twice.

(E, F) Sorting strategy for bulk RNAseq analysis of EAE-associated Th17, Th1*, Treg (E)

and SFB-induced Th17 and Treg cells (F).

Figure S3. *Tpi1* is essential for the differentiation of all CD4+ T cells, related to Figure 1

(A, B) Impaired RORγt, Foxp3, IL-17a and IFN-γ expression by *Tpi1* mutant compared to WT

CD4+ T cells in the mLN (A) and the SILP (B) of reconstituted mixed bone marrow chimeric

mice colonized with SFB. Left panels, representative FACS plots showing RORγt/Foxp3 and

IL-17a/IFN-γ expression in paired *Tpi1* WT (red) and KO (blue) CD4+ T cells in each mouse.

Middle panels, frequencies of each CD4+ T cell subtype among the *Tpi1* WT and *Tpi1* KO

cells. Right panels, *Tpi1* KO/*Tpi1* WT ratios for each CD4+ T cell subtype in mLN and SILP.

(C) Impaired RORγt, Foxp3, IL-17a and IFN-γ expression in *Tpi1* mutant compared to WT CD4+

T cells in the dLN of mixed bone marrow reconstituted mice, at day 15 of MOG-induced

EAE. The number of *Tpi1* KO CD4+ T cells obtained from SC was too low for a meaningful

subtype analysis, and hence is not shown. Panels are organized as in (B).

Figure S4. Electroporation-based CRISPR-mediated targeting of metabolic genes in naïve

CD4+ T cells, related to Figure 2 and STAR Methods

(A) Map of metabolic pathways, showing relevant genes and metabolites in glycolysis, PPP and

the TCA cycle. The biosynthetic processes from glycolysis metabolites are labeled in green.

Genes that were targeted using CRISPR in this study are labeled in blue. Gpi1, Tpi1 and

Ldha are the three enzymes that are expected to not be essential for glycolysis: (1) The

reaction catalyzed by Gpi1 (glucose-6-phosphate to fructose-1,6-bisphosphate) can also be

achieved by the PPP, making Gpi1 possibly redundant for this reaction; (2) Tpi1 controls the

inter conversion of glyceraldehyde-3-phosphate and dihydroxyacetone phosphate, the two 3-

carbon molecules generated by degrading fructose-1,6-phosphate. Loss of Tpi1 should lead

to the loss of 50%, but not all of the glycolytic activity, and thus may be not lethal; (3) Ldha

controls the conversion of pyruvate to lactic acid, and the regeneration of NAD^+. However,

pyruvate could still enter the TCA cycle in Ldha-deficient cells, for energy production and

biomass synthesis.

(B) Schematic of gene targeting in naïve T cells. Left, PCR amplicons encoding gRNAs were

electroporated into Cas9-expressing naïve $CD4^+$ T cells, which were subsequently cultured in

vitro or transferred into recipient mice. Right, *in vivo* KO efficiency can be assessed by

measuring the KO efficiency of the in vitro cultured Th17 cells. Cas9-expressing 7B8 naïve

$CD4^+$ T cells were electroporated with *Gfp* guide amplicons to target the Gfp that is co-

expressed with Cas9 at the Rosa26 locus, and then cultured *in vitro* in Th17 conditions

(IL6+TGF-β), or transferred into SFB-colonized mice. Gfp expression was examined 4 days

later in the in vitro cultured Th17 cells (top) and the ex vivo cells isolated from the mLN

(bottom).

(C-E) Evaluation of the knockout efficiency of guides used in this study with in vitro Th17 cell

culture. 8000 electroporated naïve $CD4^+$ T cells were seeded in each well at the beginning of

Th17 cell culture. Because gene function is related to cell growth and differentiation, we

estimated the KO efficiency based on three variable factors at 120 h post electroporation:

protein expression level (C), live cell number (D), and IL-17a production (E). Taking *Gapdh*

targeting as an example, western blot data shows Gapdh protein expression decreased only

~40%. However, the cell number was reduced ~70%, and IL-17a production by live cells was

further reduced more than 50% compared to the *Olfr2* control. Taken together, we estimate

that the Gapdh guide achieved more than 80% knockout *in vitro*. Data are representative of

three independent experiments.

(F) Efficiency of gene inactivation following electroporation of naïve Cas9 CD4$^+$ T cells with a

mixture of two guide amplicons targeting *Cd4* and *Cd45* in cultured Th17 cells at 96 h.

P values were determined using t tests.

Figure S5. Differentiation of glycolysis gene-targeted 7B8 T cells in the SILP of SFB-

colonized mice, related to Figure 2

(A) Representative FACS plots showing the expression of RORγt and Foxp3 in the *Olfr2* KO

control cells and the co-transferred *Olfr2* control or glycolysis gene KO cells in the SILP at

day 15 post transfer.

(B) Left, summary of the Th17 frequencies (RORγt$^+$ Foxp3$^-$) in the *Olfr2* KO control cells

(green) and the co-transferred *Olfr2* control or glycolysis gene KO cells (red), as in panel

(A). Right, the KO/Olfr2 ratios of Th17 frequencies.

(C) Representative FACS plots showing IL-17a expression in *Olfr2* control and co-transferred

Olfr2 control or glycolysis gene KO Th17 cells (RORγt$^+$ Foxp3$^-$ cells) isolated from the SILP

at day 15 post transfer and following *in vitro* PMA/Ionomycin re-stimulation.

(D) Left, summary of IL-17a expression based on results in panel (C). Right, KO/*Olfr2* ratios of

IL-17a expression. There was consistent reduction of IL-17a in *Gpi1-* or *Ldha*-targeted, but

not in *Olfr2*-targeted 7B8 cells isolated from the SILP and re-stimulated in vitro. While the

reduced cytokine production observed in the *in vitro* restimulation assay supports the

maintainence of Gpi1-deficient cells in the SILP, it is not necessarily contradictory to our

previous claim that Gpi1-deficient 7B8 cells produce normal amounts of IL-17a (based on

Il17a$^{GFP/+}$ assay (Figure 2H))*,* as *in vitro* restimulation was conducted in an artificial

microenvironment with strong stimuli, wheareas *Il17a$^{GFP/+}$* reporter cells report the true

cytokine production in real physiological conditions.

P values were determined using t tests.

Figure S6. Hypoxic microenvironment in the EAE spinal cord impairs proliferation of

***Gpi1*-deficient Th17 cells, related to Figure 6**

(A, B) The PPP contributes to Th17 cell accumulation in the SC in EAE. (A) Representative

FACS plots showing the frequencies of co-transferred *Olfr2* KO and *Olfr2* control or *G6pdx*

KO 2D2 T cells in the EAE SC at day 15. (B) Compilation of KO/*Olfr2* cell number ratios as

shown in panel A.

(C) Lactate secretion of *in vitro* cultured *Olfr2* and *Gpi1* KO npTh17 cells in normoxia and

hypoxia. Cells were cultured as in Figure 3A in normoxia. At 96 h cells were re-plated in

new wells with fresh npTh17 medium in normoxia and hypoxia, and supernatants were

collected 12 h later for lactate quantification by GC-MS.

(D) Relative intracellular ATP abundance of in vitro cultured Olfr2 and Gpi1 KO npTh17 cells

in normoxia and hypoxia. Cells were cultured as in Figure 3A in normoxic condition. At 96 h

cells were re-plated in new wells with fresh npTh17 medium in normoxic and hypoxic

condition, and ATP level was measured 12 h later.

(E, F) Pimonidazole staining of different CD4$^+$ T cell subsets in the SFB and EAE models. (E)

Representative FACS plots showing pimonidazole staining of each subtype of CD4$^+$ T cell in

the mLN and the SILP of SFB-colonized healthy mice and in the dLN and the SC of EAE

mice. Treg (Foxp3$^+$ RORγt$^-$ Tbet$^-$), Th17 (RORγt$^+$ Foxp3$^-$Tbet$^-$), Th1 (Tbet$^+$ RORγt$^-$ Foxp3$^-$),

Triple$^-$ (RORγt$^-$ Tbet$^-$ Foxp3$^-$). (F) Summarized pimonidazole staining based on panel C.

(G) Transcription factor staining in CD4$^+$ T cells from *Hif1a* WT and KO mice colonized with

SFB (top, cells from SILP)) or subjected to EAE (bottom, from SC at day15). Left,

representative FACS plots showing the frequencies of Th17 cells (RORγt$^+$ Foxp3$^-$) and Treg

cells (Foxp3$^+$ RORγt$^-$) among total *Hif1a* WT and KO CD4$^+$ T cells; Right, summary of

transcription factor staining from two independent experiments.

(H) Cytokine production in CD4$^+$ T cells from *Hif1a* WT and KO CD4 T cells in the SFB (top)

and EAE (bottom) models after in vitro PMA/ionomycin restimulation. Left, representative

plots showing the frequencies of IL-17a producing cells among RORγt$^+$ Foxp3$^-$ *Hif1a* WT or

KO cells in the SILP of SFB-colonized mice (top) and frequencies of IL-17a, IL-17a/IFN-γ,

and IFN-γ producing cells among Foxp3$^-$ *Hif1a* WT or KO cells in the SC at day 15 after

EAE induction (bottom); Right, summary of cytokine production from two independent

experiments. .

(I) Left, representative plots showing EDU incorporation in co-transferred *Olfr2* and *Olfr2*

control or *Gpi1* KO 2D2 cells at different days post transfer in the dLNs. Right, compiled

data of one representative experiment. Two independent experiments were performed with

the same conclusion.

(J, K) Proliferation of dLN 2D2 cells in the EAE model and mLN 7B8 cells in SFB-colonized

mice. Naïve 7B8 or 2D2 CD4$^+$ T cells were labeled with CFSE and 100K cells were

transferred into recipient mice that had been previously colonized with SFB or were

immunized with MOG/CFA immediately after cell transfer. The mLN of the 7B8 model and

dLN of the EAE model were collected at different time points for cell division analysis. (J)

Left, representative FACS plot showing CFSE labeling of the 7B8 and 2D2 cells at different

time points. Right, pie charts showing the frequencies of cells with different numbers of

divisions. Division number was labeled in different colors. (K) Total cell number count of

mLN 7B8 cells and dLN 2D2 cells at different time points. For each time point, 4-10 mice

were used for each group. Two independent experiments were performed with the same

conclusion.

(L, M) IL-17a production (L) and RORγt/Foxp3 expression (M) in 2D2 cells from dLN at day 10

in the EAE model. Left, representative FACS plots showing the overlay of IL-17a expression

(L) or RORγt/Foxp3 expression (M) in *Olfr2* control and co-transferred *Olfr2* KO or *Gpi1*

KO 2D2 cells. Middle, compilation of frequencies of IL-17a$^+$ (L) or RORγt$^+$Foxp3$^-$ cells (M)

among *Olfr2* control and co-transferred *Olfr2* KO or *Gpi1* KO 2D2 cells. Right, the

KO/Olfr2 ratios of the IL-17a$^+$ frequencies (L) and RORγt$^+$Foxp3$^-$ frequencies (M).

(N, O) Left, compiled data showing the frequencies of Ccr6$^+$ cells among co-transferred *Olfr2*

and *Olfr2* control or *Gpi1* KO 2D2 cells in the dLN at day 9 (N) or in the SC at day 13 (O)

post transfer and EAE induction. Right, the KO/Olfr2 ratios of the Ccr6$^+$ frequencies. Two

independent experiments were performed with the same conclusion.

(P) Compiled data showing the frequencies of active Caspase3$^+$ cells among co-transferred *Olfr2*

and *Olfr2* control or *Gpi1* KO 2D2 cells in the SC at day 10 post transfer and EAE induction.

n=10.

P values of panel B, C, D, K, L, M, N, and O were determined using t tests. P values of panel G,

H, and P, were determined using paired t tests.

STAR Methods

RESOURCE AVAILABILITY

LEAD CONTACT AND MATERIALS AVAILABILITY

826   Further information and requests for resources and reagents should be directed to and will be

827   fulfilled by the Lead Contact, Dan R. Littman (Dan.Littman@med.nyu.edu).

Materials Availability

830   This study did not generate new unique reagents

Data and Code Availability

833   The bulk RNAseq data and the scRNA seq data supporting the current study have been deposited

834   in GEO (accession number GSE141006).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mouse Strains

C57BL/6 mice were obtained from Taconic Farms or the Jackson Laboratory. All transgenic

animals were bred and maintained in the animal facility of the Skirball Institute (NYU School of

Medicine) in specific-pathogen-free (SPF) conditions. *Il-23r^{GFP}* mice (Awasthi et al., 2009) were

provided by M. Oukka. *S100a4* knockout mice (Li et al., 2010) were provided by Dr. Anne R.

Bresnick. *Cas9^{Tg}* mice (Platt et al., 2014) were obtained from Jackson Laboratory (JAX;

B6J.129(Cg)-Gt(ROSA)26Sortm1.1(CAG-cas9*,-EGFP)Fezh/J), and maintained on the Cd45.1

background (JAX; B6. SJL-Ptprca Pepcb/BoyJ). SFB-specific TCRTg (7B8) mice (Yang et al.,

2014), MOG-specific TCRTg (2D2) mice (Bettelli et al., 2003), and *Helicobacter hepaticus*-

specific TCRTg (HH7-2) mice (Xu et al., 2018) were previously described. They were crossed to

the Cas9^{Tg} Cd45.1/1 mice to generate 7B8 *Cas9^{Tg}*, 2D2 *Cas9^{Tg}*, and HH7-2 *Cas9^{Tg}* mice with

either Cd45.1/1 or Cd45.1/2 congenic markers. 7B8 Cas9 CD45.1/2 mice were further crossed

with *IL17a$^{GFP/GFP}$* reporter mice (JAX; C57BL/6-Il17atm1Bcgen/J) to generate 7B8 *Cas9*

IL17a$^{GFP/+}$ Cd45.1/2 or Cd45.2/2 mice. Female *Foxp3$^{RFP/RFP}$* reporter mice (Jax; C57BL/6-

Foxp3^{tm1Flv}/J) were crossed with male *IL17a$^{GFP/GFP}$* reporter mice to generate *IL17a$^{GFP/+}$*

Foxp3^{RFP} male mice for bulk RNAseq. *Hif1a$^{f/f}$* mice (Jax; B6.129-Hif1a^{tm3Rsjo}/J) were crossed

with *CD4^{cre}* mice (Jax; B6.Cg-Tg(Cd4-cre)1Cwi/BfluJ) and CD45.1/1 mice to generate *Hif1a$^{+/+}$*

CD4^{cre} CD45.1/1 and *Hif1a$^{f/f}$ CD4^{cre}* CD45.2/2 littermates. *Tpi1$^{f/+}$* mice generated by crossing

Tpi1$^{tm1a(EUCOMM)Wtsi}$ mice (Wellcome Trust Sanger Institute; B6Brd;B6N-Tyr^{c-Brd}

Tpi1$^{tm1a(EUCOMM)Wtsi}$/WtsiCnbc) with a FLP deleter strain (Jax; B6.129S4-

Gt(ROSA)26Sor$^{tm1(FLP1)Dym}$/RainJ) were further crossed with CD4^{cre} and CD45.1/1 strains to

generate *Tpi1$^{+/+}$Cd4^{cre}* and *Tpi1$^{f/f}$CD4^{cre}* mice with different congenic markers.

METHOD DETAILS

Flow cytometry

For transcription factor and cytokine staining of *ex vivo* cells, single cell suspensions were

incubated for 3h in RPMI with 10% FBS, phorbol 12-myristate 13-acetate (PMA) (50 ng/ml;

Sigma), ionomycin (500 ng/ml;Sigma) and GolgiStop (BD). Cells were then resuspended with

surface-staining antibodies in HEPES-buffered HBSS. Staining was performed for 20-30 min on

ice. Surface-stained cells were washed and resuspended in live/dead fixable blue (ThermoFisher)

for 5 min prior to fixation. Cells were treated using the FoxP3 staining buffer set from

eBioscience according to the manufacturer's protocol. Intracellular stains were prepared in 1X

eBioscience permwash buffer containing anti-CD16/anti-CD32, normal mouse IgG, and normal

rat IgG Staining was performed for 30-60 min on ice.

For cytokine analysis of *in vitro* cultured Th17 cells, cells were incubated in the restimulation

buffer for 3 h. After surface and live/dead staining, cells were treated using the

Cytofix/Cytoperm buffer set from BD Biosciences according to the manufacturer's protocol.

Intracellular stains were prepared in BD permwash in the same manner used for transcription

factor staining. Flow cytometric analysis was performed on an LSR II (BD Biosciences) or an

Aria II (BD Biosciences) and analyzed using FlowJo software (Tree Star).

Retroviral transduction and 2D2 TCRtg Cas9^{Tg} T cell-mediated EAE

The protocol described in this section was only used for experiments shown in Figure 2F and

Figure S2A-D. sgRNAs were designed using the Crispr guide design software

(http://crispr.mit.edu/). sgRNA encoding sequences were cloned into the retroviral sgRNA

expression vector pSIN-U6-sgRNAEF1as-Thy1.1-P2A-Neo that has been reported before (Ng et

al., 2019). To make the double sgRNA vector for experiments shown in Figures S2A-D, the

sgRNA transcription cassette, ranging from the U6 promoter to the U6 terminator, was amplified

(primer sequences can be found in the Key Resources Table) and cloned into the pSIN vector

after the first sgRNA encoding cassette. For the EAE experiment shown in Figure 2F, only one

sgRNA was used for each virus. Retroviruses were packaged in PlatE cells by transient

transfection using TransIT 293 (Mirus Bio).

Tissue culture plates were coated with polyclonal goat anti-hamster IgG (MP Biomedicals) at

37 °C for at least 3 h and washed 3× with PBS before cell plating. FACS-sorted

$CD4^+CD8^-CD25^-CD62L^+CD44^{low}$ naïve 2D2 TCRtg $Cas9^{Tg}$ T cells were seeded for 24 h in T

cell medium (RPMI supplemented with 10% FCS, 2mM β-mercaptoethanol, 2mM glutamine),

along with anti-CD3 (BioXcell, clone 145-2C11, 0.25μg/ml) and anti-CD28 (BioXcell, clone

37.5.1, 1μg/ml) antibodies. Cells were then transduced with the pSIN virus by spin infection

at 1200 × g at 30 °C for 90 min in the presence of 10μg/ml polybrene (Sigma). Viral

supernatants were removed 12 h later and replaced with fresh medium containing anti-

CD3/anti-CD28, 20ng/ml IL-6 and 0.3ng/ml TGF-β. Cells were lifted off 48 h later and

supplemented with IL-2 for another 48hours. $Thy1.1^{high}$ cells were then sorted with the ARIA

and i.v. injected into $Rag1^{-/-}$ mice (50k cells/mouse). Mice were then immunized

subcutaneously on day 0 with 100μg of MOG_{35-55} peptide, emulsified in CFA (Complete

Freund's Adjuvant supplemented with 2mg/mL *Mycobacterium tuberculosis* H37Ra), and

injected i.p. with 200 ng pertussis toxin (Calbiochem). The EAE scoring system was as follows:

0-no disease, 1-Partially limp tail; 2-Paralyzed tail; 3-Hind limb paresis, uncoordinated

movement; 4-One hind limb paralyzed; 5-Both hind limbs paralyzed; 6-Hind limbs paralyzed,

weakness in forelimbs; 7-Hind limbs paralyzed, one forelimb paralyzed; 8-Hind limbs paralyzed,

both forelimbs paralyzed; 9-Moribund; 10-Death.

sgRNA guide amplicon electroporation of naïve CD4$^+$ T cells

sgRNA-encoding sequences were cloned into the pSIN vector (single sgRNA in one vector).

The sgRNA encoding cassette, ranging from the U6 promoter to the U6 terminator, was

amplified with 34 cycles of PCR with Phusion (NEB), and purified with Wizard SV Gel and

PCR Clean-up System (Promega), and concentrated with a standard ethanol DNA

precipitation protocol. Amplicons were resuspended in a small volume of H_2O and kept at -

80 °C until use. Amplicon concentration was measured with Qubit dsDNA HS assay kit

(Thermofisher)

The Amaxa P3 Primary Cell 4D-Nucleofector X kit was used for T cell electroporation. 2

million FACS-sorted naïve CD4$^+$ $Cas9^{Tg}$ cells were pelleted and resuspended in 100ul P3

primary cell solution supplemented with 0.56ug sgRNA amplicon (for single gene knockout),

or equal amounts of sgRNA amplicons (0.56ug each for targeting two genes simultaneously in

Figures 3D, and 4E). For GFP KO (Figure S4B), a mixture of three pSIN vectors at equal

amount containing different sgRNA-encoding sequences (all targeting GFP) was used as

template for PCR amplifaction, and 0.56ug of amplicon was used for electroporation. Cell

suspensions were transferred into a Nucleocuvette Vessel for electroporation in the

Nucleofector X unit, using program "T cell. Mouse. Unstim.". Cells were then rested in 500 ul

mouse T cell nucleofector medium (Lonza) for 30 min before in vitro culture or transfer into

mice.

In vitro cell culture – 96 well flat-bottom tissue culture plates (Corning) were coated with

polyclonal goat anti-hamster IgG (MP Biomedicals) at 37 °C for at least 3 h and washed 3×

with PBS before cell plating. Electroporated cells were plated into each well containing 200ul

T cell culture medium (glucose-free RPMI (ThermoFisher) supplemented with 10% FCS

(Atlanta Biologicals), 4g/L glucose, 4mM glutamine, 50uM β-mercaptoethanol, 0.25μg/ml

anti-CD3 (BioXcell, clone 145-2C11) and 1μg/ml anti-CD28 (BioXcell, clone 37.5.1)

antibodies). For npTh17 culture, 20ng/ml IL-6 and 0.3ng/ml TGF- β were further added into

the T cell medium. For pTh17 cell culture, 20ng/ml IL-6, 20ng/ml IL-23, and 20ng/ml IL-1β

were further added into the T cell medium. Cells were then cultured for 96 h or 120 h in

normoxia or hypoxia (3% oxygen) before metabolic experiments or cell number

counting/western/FACS profiling, respectively.

T cell transfer into mice – TCR transgenic $Cas9^{Tg}$ naïve CD4$^+$ T cells (7B8 TCRtg for SFB

model, 2D2 TCRtg for EAE model, and HH7-2 TCRtg for colitis model) with different

isotype markers were electroporated at the same time. Equal numbers of isotype-distinct

electroporated resting cells were mixed, pelleted and resuspended in dPBS supplemented with

1% filtered FBS before retro-orbital injection into recipient mice (100,000 cells in total for

each mouse). The transferred cells were isolated from SILP of the SFB model, SC of the EAE

model, and the LILP of the colitis model at day 15 post transfer, unless otherwise indicated.

SFB model

C57BL/6 mice were purchased from Jackson Laboratory and maintained in SFB-free SPF cages.

Fresh feces of SFB mono-colonized mice maintained in our gnotobiotic animal facility were

collected and frozen at -80 °C until use. Fecal pellets were meshed through a 100uM strainer

(Corning) in cold sterile dPBS (Hyclone). 300 μl feces suspension, equal to the amount of 1/3

pellet, was administered to each mouse by oral gavage. Mice were inoculated one more time at 24

h. For the transfer experiment, electroporated cells were injected 96 h after the initial gavage.

Colitis model

H. hepaticus was provided by J. Fox (MIT) and grown on blood agar plates (TSA with 5% sheep

blood, Thermo Fisher) as previously described (Xu et al., 2018). Briefly, $Rag1^{-/-}$ mice were

colonized with *H. hepaticus* by oral gavage on days 0 and 4 of the experiment. At day 7,

electroporated naïve HH7-2 TCRtg $Cas9^{Tg}$ CD4$^+$ T cells were adoptively transferred into the *H.*

hepaticus-colonized recipients. In order to confirm *H. hepaticus* colonization, *H. hepaticus*-

specific 16S primers were used on DNA extracted from fecal pellets. Universal 16S were

quantified simultaneously to normalize *H. hepaticus* colonization of each sample.

EAE model

Right after the transfer of electroporated 2D2 $Cas9^{Tg}$ cells, mice were immunized subcutaneously

on day 0 with 100μg of MOG$_{35-55}$ peptide, emulsified in CFA (Complete Freund's Adjuvant

supplemented with 2mg/mL Mycobacterium tuberculosis H37Ra), and injected i.p. on days 0

and 2 with 200 ng pertussis toxin (Calbiochem).

Isolation of lymphocytes from lamina propria, spinal cord, and lymph nodes

For isolating mononuclear cells from the lamina propria, the intestine (small and/or large) was

removed immediately after euthanasia, carefully stripped of mesenteric fat and Peyer's

patches/cecal patch, sliced longitudinally and vigorously washed in cold HEPES buffered

(25mM), divalent cation-free HBSS to remove all fecal traces. The tissue was cut into 1-inch

fragments and placed in a 50ml conical containing 10ml of HEPES buffered (25mM), divalent

cation-free HBSS and 1 mM of fresh DTT. The conical was placed in a bacterial shaker set to

37 °C and 200rpm for 10 minutes. After 45 seconds of vigorously shaking the conical by hand,

the tissue was moved to a fresh conical containing 10ml of HEPES buffered (25mM), divalent

cation-free HBSS and 5 mM of EDTA. The conical was placed in a bacterial shaker set to 37 °C

and 200rpm for 10 minutes. After 45 seconds of vigorously shaking the conical by hand, the

EDTA wash was repeated once more in order to completely remove epithelial cells. The tissue

was minced and digested in 5-7ml of 10% FBS-supplemented RPMI containing collagenase (1

mg/ml collagenaseD; Roche), DNase I (100 µg/ml; Sigma), dispase (0.05 U/ml; Worthington)

and subjected to constant shaking at 155rpm, 37 °C for 35 min (small intestine) or 55 min (large

intestine). Digested tissue was vigorously shaken by hand for 2 min before adding 2 volumes of

media and subsequently passed through a 70 µm cell strainer. The tissue was spun down and

resuspended in 40% buffered percoll solution, which was then aliquoted into a 15ml conical. An

equal volume of 80% buffered percoll solution was underlaid to create a sharp interface. The

tube was spun at 2200rpm for 22 min at 22 °C to enrich for live mononuclear cells. Lamina

propria (LP) lymphocytes were collected from the interface and washed once prior to staining.

For isolating mononuclear cells from spinal cords during EAE, spinal cords were mechanically

disrupted and dissociated in RPMI containing collagenase (1 mg/ml collagenaseD; Roche),

DNase I (100 µg/ml; Sigma) and 10% FBS at 37 °C for 30 min. Leukocytes were collected at the

interface of a 40%/80% Percoll gradient (GE Healthcare).

For isolating mononuclear cells from lymph nodes, mesenteric lymph nodes (for the SFB

model), and draining lymph nodes (Inguinal, axillary, and brachial lymph nodes for the EAE

model) were mechanically dissected and minced in RPMI containing 10% FBS and 25mM

HEPES through 40uM strainer. Cells were collected for further analysis.

1003

Generation of bone marrow chimeric reconstituted mice

Bone marrow mononuclear cells were isolated from donor mice by flushing the long bones. To

generate $Tpi1^{WT}/Tpi1^{KO}$ chimeric reconstituted mice, $Tpi1^{+/+} Cd4^{Cre}$ CD45.1/2 and $Tpi1^{f/f} Cd4^{Cre}$

CD45.2/2 mice were used as donors. To generate $Hif1a^{WT}/Hif1a^{KO}$ chimeric reconstituted mice,

$Hif1a^{+/+} Cd4^{Cre}$ CD45.1/2 and $Hif1a^{f/f} Cd4^{Cre}$ CD45.1/1 mice were used as donors. To generate

$Il23r^{het}/Il23r^{KO}$ chimeric reconstituted mice, $Il23r^{Gfp/+}$ CD45.1/2 and $Il23r^{Gfp/Gfp}$ CD45.1/1 mice

were used as donors. Red blood cells were lysed with ACK Lysing Buffer, and CD4/CD8 T

cells were labeled with CD4/CD8 magnetic microbeads and depleted with a Miltenyi LD

column. The remaining cells were resuspended in PBS for injection in at a 1:1 ratio (WT: KO).

4×10^6 cells were injected intravenously into 6 week old mice (CD45.1/1 in Tpi1 experiment;

Rag1$^{-/-}$ in Hif1a experiment) that were irradiated 4h before reconstitution using 1000 rads/mouse

(2x500rads, at an interval of 3h, at X-RAD 320 X-Ray Irradiator). Peripheral blood samples

were collected and analyzed by FACS 8 weeks later to check for reconstitution, after which SFB

colonization or EAE induction were performed.

1018

Pimonidazole HCl labeling

Pimonidazole HCl (hypoxyprobe) was retro orbitally injected into mice at the dose of 100mg/kg

under general anesthesia (Ketamine 100mg/Kg, Xylazine 15mg/Kg). 3 h later, tissues were

dissected and processed to get the mononuclear cell fraction. After surface/live/dead staining,

cells were treated using the FoxP3 staining buffer set from eBioscience according to the

manufacturer's protocol. Intracellular staining with anti-RORγt, Foxp3, Tbet, and biotin-

pimonidazole adduct antibodies were performed for 60min on ice in 1X eBioscience permwash

buffer containing normal mouse IgG (50μg/ml), and normal rat IgG (50μg/ml). The

pimonidazole labeling was visualized with streptavidin APC staining in the same buffer.

1028

EdU incorporation

Two doses of EdU (50mg/kg for each) were i.p. injected into MOG/CFA-immunized mice in 12

h intervals at days 4, 6, 8, 9, and 12 post immunization. Draining lymph nodes and/or spinal

cords were dissected 12 h post the second injection. Detection of EdU incorporation into the

DNA was performed with EdU-click 647 Kit (Baseclick) according to the manufacturer's

instructions. In brief, surface-stained cells were fixed in 100μl fixative solution, and

permeabilized in 100μl permeabilization buffer. Cells were pelleted and incubated for 30min in

220μl EdU reaction mixture containing 20μl permeabilization buffer, 20μl buffer additive, 1μl

dye azide, 4μl catalyst solution, and 175μl dPBS, before 3x wash and FACS analysis.

1038

CFSE labeling

Sorted naïve 7B8 or 2D2 CD45.1/1 CD4 T cells were stained with CFSE cell proliferation kit

(Life Technology). Labeled cells were administered into each congenic CD45.2/2 recipient

mouse by retro-orbital injection. Mice receiving 7B8 cells were gavaged with SFB mono-feces

twice on days -4 and -3. Mice receiving 2D2 cells were immunized for EAE induction

immediately after cell transfer. mLNs of the SFB-colonized mice were collected at 72h and

120h, and dLN of 2D2-injected mice were collected at 72h, 96h and 120h post transfer for cell

division analysis.

1047

Bulk RNA sequencing

Total RNA from sorted target cell populations was isolated using TRIzol LS (Invitrogen)

followed by DNase I (Qiagen) treatment and cleanup with RNeasy Plus Micro kit (Qiagen;

74034). RNA quality was assessed using Pico Bioanalyser chips (Agilent). RNASeq libraries

were prepared using the Clontech Smart-Seq Ultra low RNA kit (Takara Bio) for cDNA

preparation, starting with 550 pg of total RNA, and 13 cycles of PCR for cDNA amplification.

The SMARTer Thruplex DNA-Seq kit (Takara Bio) was used for library prep, with 7 cycles of

PCR amplification, following the manufacturer's protocol. The amplified library was purified

using AMPure beads (Beckman Coulter), quantified by Qubit and QPCR, and visualized in an

Agilent Bioanalyzer. The libraries were pooled equimolarly, and run on a HiSeq 4000, as paired

end reads, 50 nucleotides in length.

1059

Single cell RNAseq

Libraries from isolated single cells were generated based on the Smart-seq2 protocol (Picelli et

al., 2014) with the following modifications. In brief, RNA from sorted single cells was used as

template for oligo-dT primed reverse transcription with Superscript II Reverse Transcriptase

(Thermo Fisher), and the cDNA library was generated by 20 cycle PCR amplification with IS

PCR primer using KAPA HiFi HotStart ReadyMix (KAPA Biosystems) followed by Agencourt

AMPure XP bead purification (Beckman Coulter) as described. Libraries were tagmented using

the Nextera XT Library Prep kit (Illumina) with custom barcode adapters (sequences available

upon request). Libraries with unique barcodes were combined and sequenced on a HiSeq2500 in

RapidRun mode, with either 2x50 or 1x50 nt reads.

1070

Stable-isotope tracing, metabolite secretion and kinetic flux profiling by GC-MS

In vitro cultured np/pTh17 cells were collected at 96h, washed twice in glucose-free RPMI

(ThermoFisher, 11879020) containing 10% dialyzed FBS (ThermoFisher), and re-plated in anti-

hamster IgG-coated 96-well plates at 60,000/well in np/pTh17 medium (glucose-free RPMI

containing 10% dialyzed FBS, 2mM β-mercaptoethanol, 4mM glutamine, anti-CD3, anti-

CD28, and 4g/L isotype labeled glucose). For KA-treated samples, *Olfr2* KO cells were

previously treated with 30 µM KA before collection and re-plating. The re-plated cells were

cultured in Th17 medium supplemented with 30 µM KA.

1079

To measure relative PPP activity, $^{13}C_{1,2}$-glucose (Cambridge Isotope Laboratories) was added

into the Th17 medium. Re-plated cells were further cultured 12 h before metabolite extraction

and GC-MS analysis. After a brief wash with 0.9% ice-cold saline solution to remove media

contamination, cellular metabolites were extracted using a pre-chilled

methanol/water/chloroform extraction method, as previously described (Metallo et al., 2011).

Aqueous and inorganic layers were separated by cold centrifugation for 15 min. The aqueous

layer was transferred to sample vials (Agilent 5190-2243) and evaporated to dryness using a

SpeedVac (Savant Thermo SPD111V).

1088

1089   For lactate secretion analysis, re-plated cells were cultured in normoxia or hypoxia for 12 h.

1090   media was collected and centrifuged at 1,000 x g to pellet cellular debris. 5 µL of supernatant

1091   was extracted in 80% ice-cold methanol mixture containing isotope-labeled internal standards

1092   and evaporated to dryness.

1093

1094   Metabolite abundance $[X_m]$ was determined using the following equation:

1095

1096 $$[X_m] = \frac{X_{std} - \%M0_x}{100 - \%M0_x}$$

1097

1098   where X_{std} is the molar amount of isotope-labeled standard (e.g. 5 nmol lactate) and $\%M0_x$ is the

1099   relative abundance of unlabeled metabolite X corrected for natural isotope abundance.

1100

1101   To quantify the metabolite secretion flux ($Flux_m$), the molar change in extracellular metabolites

1102   was determined by calculating the difference between conditioned and fresh media and

1103   normalizing to changes in cell density:

1104

1105 $$Flux_m = \frac{X_f - X_i}{\int_0^f X_0\, e^{kt}}$$

1106

1107   where X_f and X_i are the final and initial molar amounts of metabolite, respectively, X_0 is the

1108   initial cell density, k is the exponential growth rate (hr^{-1}), and t is the media conditioning time.

1109

1110   For kinetic flux profiling, npTh17 medium was supplemented with 4g/L $^{13}C_6$-glucose

1111   (Cambridge Isotope Laboratories). Cells were collected at various time points. Unlabeled

1112   metabolite (M0) versus time (t) were plotted and the data were fitted to an equation as previously

1113   described (Yuan et al., 2008) to quantify the metabolite turnover rate (t^{-1}):

1114

1115 $$Y = (1 - \alpha)^* \exp(-k^*t) + \alpha$$

1116

1117   where α is the percentage of unlabeled (M0) metabolite at steady-state, k is the metabolite

1118   turnover rate and t is time.

1119

1120   Metabolite flux was quantified by multiplying the decay rate by the intracellular metabolite

1121   abundance.

1122

1123   Metabolite derivatization using MOX-tBDMS was conducted as previously described (Lewis et

1124   al., 2014). Derivatized samples were analyzed by GC-MS using a DB-35MS column (30m x

1125   0.25mm i.d. x 0.25 μm) installed in an Agilent 7890B gas chromatograph (GC) interfaced with

1126   an Agilent 5977B mass spectrometer as previously described (Grassian et al., 2014) and

1127   corrected for natural abundance using in-house algorithms adapted from (Fernandez et al., 1996).

1128

Intracellular ATP measurement

npTh17 cells were cultured in vitro in normoxic condition for 96 h before being collected and

washed twice in glucose-free RPMI (ThermoFisher) containing 10% FBS (Atlanta Biologicals),

and re-plated in anti-hamster IgG-coated 96-well plates at 20,000/well in 200μl npTh17 medium

(glucose-free RPMI containing 10% dialyzed FBS, 2mM β-mercaptoethanol, 4mM glutamine,

anti-CD3, anti-CD28, and 4g/L glucose). 12 hours later, cell numbers were determined in

replicate wells, and ATP measurements were made using CellTiter-Glo® Luminescent Cell

Viability Assay kit (Promega) according to instructions.

1137

LC-MS analysis

For glucose consumption analysis, supernatants of npTh17 cells, or cell free control, were

collected at 24h post cell re-plating. Metabolites were extracted for LC-MS analysis to measure

glucose and lactate quantities. Glucose consumption rate and lactate production rate was

quantified on a relative basis by calculating the ion count difference between conditioned and

fresh media and normalizing to changes in cell density.

1144

Samples were subjected to an LC-MS analysis to detect and quantify known peaks. A metabolite

extraction was carried out on each sample with a previously described method (Pacold et al.,

2016). In brief, samples were extracted by mixing 10 µL of sample with 490 µL of extraction

buffer (81.63% methanol (Fisher Scientific) and 500 nM metabolomics amino acid mix standard

(Cambridge Isotope Laboratories, Inc.)) in 2.0mL screw cap vials containing ~100 µL of

disruption beads (Research Products International, Mount Prospect, IL). Each was homogenized

for 10 cycles on a bead blaster homogenizer (Benchmark Scientific, Edison, NJ). Cycling

consisted of a 30 sec homogenization time at 6 m/s followed by a 30 sec pause. Samples were

subsequently spun at 21,000 g for 3 min at 4 °C. A set volume of each (360 µL) was transferred

to a 1.5 mL tube and dried down by speedvac (Thermo Fisher, Waltham, MA). Samples were

reconstituted in 40 µL of Optima LC/MS grade water (Fisher Scientific, Waltham, MA).

Samples were sonicated for 2 mins, then spun at 21,000g for 3min at 4°C. Twenty microliters

were transferred to LC vials containing glass inserts for analysis.

1158

The LC column was a MilliporeTM ZIC-pHILIC (2.1 x150 mm, 5 μm) coupled to a Dionex

Ultimate 3000TM system and the column oven temperature was set to 25 °C for the gradient

elution. A flow rate of 100 μL/min was used with the following buffers; A) 10 mM ammonium

carbonate in water, pH 9.0, and B) neat acetonitrile. The gradient profile was as follows; 80-

20%B (0-30 min), 20-80%B (30-31 min), 80-80%B (31-42 min). Injection volume was set to 1

μL for all analyses (42 min total run time per injection).

1165

MS analyses were carried out by coupling the LC system to a Thermo Q Exactive HFTM mass

spectrometer operating in heated electrospray ionization mode (HESI). Method duration was 30

min with a polarity switching data-dependent Top 5 method for both positive and negative

modes. Spray voltage for both positive and negative modes was 3.5kV and capillary temperature

was set to 320°C with a sheath gas rate of 35, aux gas of 10, and max spray current of 100 μA.

The full MS scan for both polarities utilized 120,000 resolution with an AGC target of 3e6 and a

maximum IT of 100 ms, and the scan range was from 67-1000 m/z. Tandem MS spectra for both

positive and negative mode used a resolution of 15,000, AGC target of 1e5, maximum IT of 50

ms, isolation window of 0.4 m/z, isolation offset of 0.1 m/z, fixed first mass of 50 m/z, and 3-

way multiplexed normalized collision energies (nCE) of 10, 35, 80. The minimum AGC target

was 1e4 with an intensity threshold of 2e5. All data were acquired in profile mode.

1177

Seahorse Analysis

OCR and ECAR measurements were performed using a Seahorse XFe96 analyzer (Seahorse

Biosciences). Seahorse XFe96 plates were coated with 0.56ug Cell-Tak (Corning) in 24µl

coating buffer (8µl H_2O + 16µl 0.1M $NaHCO_3$ (PH 8.0)) overnight at 4 °C, and washed 3x with

H_2O before use. *In vitro* cultured Th17 cells were collected at 96 h, washed twice in Seahorse

RPMI media PH7.4 (Agilent, 103576-100) and seeded onto the coated plates at 100,000 per well.

To measure the OCR and ECAR of KA-treated cells, 30µM KA (Santa Cruz) was added into

Olfr2 control Th17 cell samples at the beginning of the Seahorse measurement. Respiratory rates

were measured in RPMI Seahorse media, pH7.4 (Agilent) supplemented with 10 mM glucose

(Agilent) and 2 mM glutamine (Agilent) in response to sequential injections of oligomycin (1

µM) , FCCP (0.5 µM) and antimycin/rotenone (1 µM) (all Cayman Chemicals, Ann Arbor, MI,

USA).

1190

Western blotting

Whole cell extracts were fractionated by SDS-PAGE and transferred to nitrocellulose membrane

by wet transfer for 4 h on ice at 60 V. Blots were blocked in TBS blocking buffer (Licor) and

then stained overnight with primary antibody at 4°C. The next day, membranes were washed

three times in TBST (TBS and 0.1% Tween-20), stained with fluorescently conjugated secondary

antibodies (Licor) at 1:10,000 dilution in TBS blocking buffer for 1 h at room temperature, and

then imaged in the 680- and 800-nm channels. Antibodies used: Gapdh (CST), Gpi1 (Thermo

Fisher), Ldha (CST), Pdha1 (Abcam), Tpi1 (Abcam), G6pdx (Abcam), α-tubulin (Santa Cruz).

1199

1200 **QUANTIFICATION AND STATISTICAL ANALYSIS**

1201 **Bulk RNAseq analysis**

1202   Raw reads from bulk RNA-seq were aligned to the mm10 genome using STAR v 2.6.1d with

1203   the quantmode parameter to get read counts. DESeq2 v 1.22.2 was used for differential

1204   expression analysis. Heatmaps were produced using gplots v 3.0.1.1.

1205

Pre-processing of scRNA-seq data

The single cell RNA raw sequence reads were aligned to the UCSC Mus musculus genome,

mm10, using Kallisto (Bray et al., 2016). The mean number of reads per cell was 241,500. Gene

expression values were quantified into Transcripts Per Kilobase Million (TPM). We removed

cells which had fewer than 800 detected genes or more than 3,000 genes. After filtering, the

mean number of detected genes per cell were 1,765. Then, for the SFB and EAE dataset, we

normalized the gene matrix by SCTransform (Hafemeister and Satija, 2019) regression with

percent of mitochondria reads separately.

1214

Visualization and clustering

To visualize the EAE and SFB data, we performed principal component analysis (PCA) for

dimensionality reduction, and further reduced PCA dimensions into two dimensional space using

UMAP (Becht et al., 2018; McInnes and Healy, 2018). UMAP matrix was also used to build the

weighted shared nearest neighbor graph and clustering of single cells was performed by the

original Louvain algorithm. All the above analysis was performed using Seurat v3 R package

(Butler et al., 2018; Stuart et al., 2019). However, we noticed there was one cluster of cells with

a much lower number of features in which over 60% of cells had fewer features than the average.

Thus, it was removed from the analysis.

A table of average gene expression for each cluster, and a table of log2 Fold Change values with

p-values between each sample were calculated with Seurat. Seurat calculates differential

expression based on the non-parameteric Wilcoxon rank sum test. Log fold change is calculated

using the average expression between two groups. These values were used in Figure S1 J. and K.

with IPA for a pathway analysis, and in Figure S1 L. to make volcano plots.

1229

Statistical analysis

Differences between groups were calculated using the unpaired two-sided Welch's t-test.

Difference between two groups of co-transferred cells, or in bone marrow chimera experiments,

were calculated using the paired two-tailed t-test. Differences between EAE clinical score was

calculated using two-way ANOVA for repeated measures with Bonferroni correction. For RNA-

seq analysis, differential expression was tested by DESeq2 by use of a negative binomial

generalized linear model. We treated less than 0.05 of p value as significant differences. $*p <$

0.05, $**p < 0.01$, $***p < 0.001$, and $****p < 0.0001$. Details regarding number of replicates and

the definition of center/error bars can be found in figure legends.

1239

1240 **References**

1241

Ahern, P.P., Schiering, C., Buonocore, S., McGeachy, M.J., Cua, D.J., Maloy, K.J., and Powrie,

F. (2010). Interleukin-23 Drives Intestinal Inflammation through Direct Activity on T Cells.

Immunity *33*, 279-288.

Awasthi, A., Riol-Blanco, L., Jager, A., Korn, T., Pot, C., Galileos, G., Bettelli, E., Kuchroo,

V.K., and Oukka, M. (2009). Cutting edge: IL-23 receptor gfp reporter mice reveal distinct

populations of IL-17-producing cells. J Immunol *182*, 5904-5908.

Baronciani, L., Zanella, A., Bianchi, P., Zappa, M., Alfinito, F., Iolascon, A., Tannoia, N.,

Beutler, E., and Sirchia, G. (1996). Study of the molecular defects in glucose phosphate

isomerase-deficient patients affected by chronic hemolytic anemia. Blood *88*, 2306-2310.

Becht, E., McInnes, L., Healy, J., Dutertre, C.A., Kwok, I.W.H., Ng, L.G., Ginhoux, F., and

Newell, E.W. (2018). Dimensionality reduction for visualizing single-cell data using UMAP.

Nat Biotechnol.

Bettelli, E., Pagany, M., Weiner, H.L., Linington, C., Sobel, R.A., and Kuchroo, V.K. (2003).

Myelin oligodendrocyte glycoprotein-specific T cell receptor transgenic mice develop

spontaneous autoimmune optic neuritis. J Exp Med *197*, 1073-1081.

Birsoy, K., Wang, T., Chen, W.W., Freinkman, E., Abu-Remaileh, M., and Sabatini, D.M.

(2015). An Essential Role of the Mitochondrial Electron Transport Chain in Cell Proliferation

Is to Enable Aspartate Synthesis. Cell *162*, 540-551.

Bray, N.L., Pimentel, H., Melsted, P., and Pachter, L. (2016). Near-optimal probabilistic RNA-

seq quantification. Nat Biotechnol *34*, 525-527.

Bulcha, J.T., Giese, G.E., Ali, M.Z., Lee, Y.U., Walker, M.D., Holdorf, A.D., Yilmaz, L.S.,

Brewster, R.C., and Walhout, A.J.M. (2019). A Persistence Detector for Metabolic Network

Rewiring in an Animal. Cell Rep *26*, 460-468 e464.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-cell

transcriptomic data across different conditions, technologies, and species. Nat Biotechnol *36*,

411-420.

Corcoran, S.E., and O'Neill, L.A. (2016). HIF1alpha and metabolic reprogramming in

inflammation. J Clin Invest *126*, 3699-3707.

Cua, D.J., Sherlock, J., Chen, Y., Murphy, C.A., Joyce, B., Seymour, B., Lucian, L., To, W.,

Kwan, S., Churakova, T.*, et al.* (2003). Interleukin-23 rather than interleukin-12 is the critical

cytokine for autoimmune inflammation of the brain. Nature *421*, 744-748.

Dang, E.V., Barbi, J., Yang, H.Y., Jinasena, D., Yu, H., Zheng, Y., Bordman, Z., Fu, J., Kim, Y.,

Yen, H.R.*, et al.* (2011). Control of T(H)17/T(reg) balance by hypoxia-inducible factor 1. Cell

146, 772-784.

Davies, A.L., Desai, R.A., Bloomfield, P.S., McIntosh, P.R., Chapple, K.J., Linington, C.,

Fairless, R., Diem, R., Kasti, M., Murphy, M.P.*, et al.* (2013). Neurological deficits caused by

tissue hypoxia in neuroinflammatory disease. Ann Neurol *74*, 815-825.

de Padua, M.C., Delodi, G., Vucetic, M., Durivault, J., Vial, V., Bayer, P., Noleto, G.R., Mazure,

N.M., Zdralevic, M., and Pouyssegur, J. (2017). Disrupting glucose-6-phosphate isomerase

fully suppresses the "Warburg effect" and activates OXPHOS with minimal impact on tumor

growth except in hypoxia. Oncotarget *8*, 87623-87637.

Deutscher, D., Meilijson, I., Kupiec, M., and Ruppin, E. (2006). Multiple knockout analysis of

genetic robustness in the yeast metabolic network. Nat Genet *38*, 993-998.

Fernandez, C.A., DesRosiers, C., Previs, S.F., David, F., and Brunengraber, H. (1996).

Correction of C-13 mass isotopomer distributions for natural stable isotope abundance. J Mass

Spectrom *31*, 255-262.

Fotiadou, C., Lazaridou, E., Sotiriou, E., and Ioannides, D. (2018). Targeting IL-23 in psoriasis:

current perspectives. Psoriasis (Auckl) *8*, 1-5.

Ghoreschi, K., Laurence, A., Yang, X.P., Tato, C.M., McGeachy, M.J., Konkel, J.E., Ramos,

H.L., Wei, L., Davidson, T.S., Bouladoux, N.*, et al.* (2010). Generation of pathogenic T(H)17

cells in the absence of TGF-beta signalling. Nature *467*, 967-971.

Grassian, A.R., Parker, S.J., Davidson, S.M., Divakaruni, A.S., Green, C.R., Zhang, X., Slocum,

K.L., Pu, M., Lin, F., Vickers, C.*, et al.* (2014). IDH1 mutations alter citric acid cycle

metabolism and increase dependence on oxidative mitochondrial metabolism. Cancer Res *74*,

3317-3331.

Hafemeister, C., and Satija, R. (2019). Normalization and variance stabilization of single-cell

RNA-seq data using regularized negative binomial regression. bioRxiv, 576827.

Hanzel, J., and D'Haens, G.R. (2020). Anti-interleukin-23 agents for the treatment of ulcerative

colitis. Expert Opin Biol Ther *20*, 399-406.

Hirota, K., Duarte, J.H., Veldhoen, M., Hornsby, E., Li, Y., Cua, D.J., Ahlfors, H., Wilhelm, C.,

Tolaini, M., Menzel, U.*, et al.* (2011). Fate mapping of IL-17-producing T cells in

inflammatory responses. Nat Immunol *12*, 255-263.

Hirota, K., Hashimoto, M., Yoshitomi, H., Tanaka, S., Nomura, T., Yamaguchi, T., Iwakura, Y.,

Sakaguchi, N., and Sakaguchi, S. (2007). T cell self-reactivity forms a cytokine milieu for

spontaneous development of IL-17(+) Th cells that cause autoimmune arthritis. Journal of

Experimental Medicine *204*, 41-47.

Horlbeck, M.A., Xu, A., Wang, M., Bennett, N.K., Park, C.Y., Bogdanoff, D., Adamson, B.,

Chow, E.D., Kampmann, M., Peterson, T.R.*, et al.* (2018). Mapping the Genetic Landscape of

Human Cells. Cell *174*, 953-967 e922.

Hue, S., Ahern, P., Buonocore, S., Kullberg, M.C., Cua, D.J., McKenzie, B.S., Powrie, F., and

Maloy, K.J. (2006). Interleukin-23 drives innate and T cell-mediated intestinal inflammation.

J Exp Med *203*, 2473-2483.

Ivanov, II, Atarashi, K., Manel, N., Brodie, E.L., Shima, T., Karaoz, U., Wei, D., Goldfarb, K.C.,

Santee, C.A., Lynch, S.V.*, et al.* (2009). Induction of intestinal Th17 cells by segmented

filamentous bacteria. Cell *139*, 485-498.

Ivanov, II, McKenzie, B.S., Zhou, L., Tadokoro, C.E., Lepelley, A., Lafaille, J.J., Cua, D.J., and

Littman, D.R. (2006). The orphan nuclear receptor RORgammat directs the differentiation

program of proinflammatory IL-17+ T helper cells. Cell *126*, 1121-1133.

Johnson, M.O., Wolf, M.M., Madden, M.Z., Andrejeva, G., Sugiura, A., Contreras, D.C.,

Maseda, D., Liberti, M.V., Paz, K., Kishton, R.J.*, et al.* (2018). Distinct Regulation of Th17

and Th1 Cell Differentiation by Glutaminase-Dependent Metabolism. Cell *175*, 1780-1795

e1719.

Johnson, T.W., Wu, Y., Nathoo, N., Rogers, J.A., Wee Yong, V., and Dunn, J.F. (2016). Gray

Matter Hypoxia in the Brain of the Experimental Autoimmune Encephalomyelitis Model of

Multiple Sclerosis. PLoS One *11*, e0167196.

Kanno, H., Fujii, H., Hirono, A., Ishida, Y., Ohga, S., Fukumoto, Y., Matsuzawa, K., Ogawa, S.,

and Miwa, S. (1996). Molecular analysis of glucose phosphate isomerase deficiency

associated with hereditary hemolytic anemia. Blood *88*, 2321-2325.

Karhausen, J., Furuta, G.T., Tomaszewski, J.E., Johnson, R.S., Colgan, S.P., and Haase, V.H.

(2004). Epithelial hypoxia-inducible factor-1 is protective in murine experimental colitis. J

Clin Invest *114*, 1098-1106.

Kugler, W., Breme, K., Laspe, P., Muirhead, H., Davies, C., Winkler, H., Schroter, W., and

Lakomek, M. (1998). Molecular basis of neurological dysfunction coupled with haemolytic

anaemia in human glucose-6-phosphate isomerase (GPI) deficiency. Hum Genet *103*, 450-

454.

Kullberg, M.C., Jankovic, D., Feng, C.G., Hue, S., Gorelick, P.L., McKenzie, B.S., Cua, D.J.,

Powrie, F., Cheever, A.W., Maloy, K.J.*, et al.* (2006). IL-23 plays a key role in Helicobacter

hepaticus-induced T cell-dependent colitis. J Exp Med *203*, 2485-2494.

Kumar, P., Monin, L., Castillo, P., Elsegeiny, W., Horne, W., Eddens, T., Vikram, A., Good, M.,

Schoenborn, A.A., Bibby, K.*, et al.* (2016). Intestinal Interleukin-17 Receptor Signaling

Mediates Reciprocal Control of the Gut Microbiota and Autoimmune Inflammation.

Immunity *44*, 659-671.

Langley, R.G., Tsai, T.F., Flavin, S., Song, M., Randazzo, B., Wasfi, Y., Jiang, J., Li, S., and

Puig, L. (2018). Efficacy and safety of guselkumab in patients with psoriasis who have an

inadequate response to ustekinumab: results of the randomized, double-blind, phase III

NAVIGATE trial. Br J Dermatol *178*, 114-123.

Lee, P., Chandel, N.S., and Simon, M.C. (2020). Cellular adaptation to hypoxia through hypoxia

inducible factors and beyond. Nat Rev Mol Cell Biol.

Lee, W.N., Boros, L.G., Puigjaner, J., Bassilian, S., Lim, S., and Cascante, M. (1998). Mass

isotopomer study of the nonoxidative pathways of the pentose cycle with [1,2-13C2]glucose.

Am J Physiol *274*, E843-851.

Lewis, C.A., Parker, S.J., Fiske, B.P., McCloskey, D., Gui, D.Y., Green, C.R., Vokes, N.I., Feist,

A.M., Vander Heiden, M.G., and Metallo, C.M. (2014). Tracing compartmentalized NADPH

metabolism in the cytosol and mitochondria of mammalian cells. Mol Cell *55*, 253-263.

Li, Z.H., Dulyaninova, N.G., House, R.P., Almo, S.C., and Bresnick, A.R. (2010). S100A4

regulates macrophage chemotaxis. Mol Biol Cell *21*, 2598-2610.

Liberti, M.V., Dai, Z.W., Wardell, S.E., Baccile, J.A., Liu, X.J., Gao, X., Baldi, R.,

Mehrmohamadi, M., Johnson, M.O., Madhukar, N.S.*, et al.* (2017). A Predictive Model for

Selective Targeting of the Warburg Effect through GAPDH Inhibition with a Natural Product.

Cell Metab *26*, 648-+.

Ma, E.H., Verway, M.J., Johnson, R.M., Roy, D.G., Steadman, M., Hayes, S., Williams, K.S.,

Sheldon, R.D., Samborska, B., Kosinski, P.A.*, et al.* (2019). Metabolic Profiling Using Stable

Isotope Tracing Reveals Distinct Patterns of Glucose Utilization by Physiologically Activated

CD8(+) T Cells. Immunity *51*, 856-870 e855.

Macintyre, A.N., Gerriets, V.A., Nichols, A.G., Michalek, R.D., Rudolph, M.C., Deoliveira, D.,

Anderson, S.M., Abel, E.D., Chen, B.J., Hale, L.P.*, et al.* (2014). The glucose transporter

Glut1 is selectively essential for CD4 T cell activation and effector function. Cell Metab *20*,

61-72.

MacIver, N.J., Michalek, R.D., and Rathmell, J.C. (2013). Metabolic regulation of T

lymphocytes. Annu Rev Immunol *31*, 259-283.

Mao, K., Baptista, A.P., Tamoutounour, S., Zhuang, L., Bouladoux, N., Martins, A.J., Huang, Y.,

Gerner, M.Y., Belkaid, Y., and Germain, R.N. (2018). Innate and adaptive lymphocytes

sequentially shape the gut microbiota and lipid metabolism. Nature *554*, 255-259.

McGeachy, M.J., Chen, Y., Tato, C.M., Laurence, A., Joyce-Shaikh, B., Blumenschein, W.M.,

McClanahan, T.K., O'Shea, J.J., and Cua, D.J. (2009). The interleukin 23 receptor is essential

for the terminal differentiation of interleukin 17-producing effector T helper cells in vivo. Nat

Immunol *10*, 314-324.

McInnes, L., and Healy, J. (2018). UMAP: Uniform Manifold Approximation and Projection for

Dimension Reduction. ArXiv *abs/1802.03426*.

McMullin, M.F. (1999). The molecular basis of disorders of red cell enzymes. J Clin Pathol *52*,

241-244.

Metallo, C.M., Gameiro, P.A., Bell, E.L., Mattaini, K.R., Yang, J., Hiller, K., Jewell, C.M.,

Johnson, Z.R., Irvine, D.J., Guarente, L.*, et al.* (2011). Reductive glutamine metabolism by

IDH1 mediates lipogenesis under hypoxia. Nature *481*, 380-384.

Michalek, R.D., Gerriets, V.A., Jacobs, S.R., Macintyre, A.N., MacIver, N.J., Mason, E.F.,

Sullivan, S.A., Nichols, A.G., and Rathmell, J.C. (2011). Cutting edge: distinct glycolytic and

lipid oxidative metabolic programs are essential for effector and regulatory CD4+ T cell

subsets. J Immunol *186*, 3299-3303.

Milner, J.D., and Holland, S.M. (2013). The cup runneth over: lessons from the ever-expanding

pool of primary immunodeficiency diseases. Nat Rev Immunol *13*, 635-648.

Morrison, P.J., Bending, D., Fouser, L.A., Wright, J.F., Stockinger, B., Cooke, A., and Kullberg,

M.C. (2013). Th17-cell plasticity in Helicobacter hepaticus-induced intestinal inflammation.

Mucosal Immunol *6*, 1143-1156.

Moschen, A.R., Tilg, H., and Raine, T. (2019). IL-12, IL-23 and IL-17 in IBD: immunobiology

and therapeutic targeting. Nat Rev Gastroenterol Hepatol *16*, 185-196.

Murphy, C.A., Langrish, C.L., Chen, Y., Blumenschein, W., McClanahan, T., Kastelein, R.A.,

Sedgwick, J.D., and Cua, D.J. (2003). Divergent pro- and antiinflammatory roles for IL-23

and IL-12 in joint autoimmune inflammation. J Exp Med *198*, 1951-1957.

Nakae, S., Nambu, A., Sudo, K., and Iwakura, Y. (2003a). Suppression of immune induction of

collagen-induced arthritis in IL-17-deficient mice. J Immunol *171*, 6173-6177.

Nakae, S., Saijo, S., Horai, R., Sudo, K., Mori, S., and Iwakura, Y. (2003b). IL-17 production

from activated T cells is required for the spontaneous development of destructive arthritis in

mice deficient in IL-1 receptor antagonist. P Natl Acad Sci USA *100*, 5986-5990.

Naughton, D.P., Haywood, R., Blake, D.R., Edmonds, S., Hawkes, G.E., and Grootveld, M.

(1993). A comparative evaluation of the metabolic profiles of normal and inflammatory knee-

joint synovial fluids by high resolution proton NMR spectroscopy. FEBS Lett *332*, 221-225.

Ng, C., Aichinger, M., Nguyen, T., Au, C., Najar, T., Wu, L., Mesa, K.R., Liao, W., Quivy, J.P.,

Hubert, B.*, et al.* (2019). The histone chaperone CAF-1 cooperates with the DNA

methyltransferases to maintain Cd4 silencing in cytotoxic T cells. Genes Dev *33*, 669-683.

Ng, C.T., Biniecka, M., Kennedy, A., McCormick, J., Fitzgerald, O., Bresnihan, B., Buggy, D.,

Taylor, C.T., O'Sullivan, J., Fearon, U.*, et al.* (2010). Synovial tissue hypoxia and

inflammation in vivo. Ann Rheum Dis *69*, 1389-1395.

O'Neill, L.A., Kishton, R.J., and Rathmell, J. (2016). A guide to immunometabolism for

immunologists. Nat Rev Immunol *16*, 553-565.

Okada, S., Markle, J.G., Deenick, E.K., Mele, F., Averbuch, D., Lagos, M., Alzahrani, M., Al-

Muhsen, S., Halwani, R., Ma, C.S.*, et al.* (2015). IMMUNODEFICIENCIES. Impairment of

immunity to Candida and Mycobacterium in humans with bi-allelic RORC mutations. Science

349, 606-613.

Olenchock, B.A., Rathmell, J.C., and Vander Heiden, M.G. (2017). Biochemical Underpinnings

of Immune Cell Metabolic Phenotypes. Immunity *46*, 703-713.

Omenetti, S., Bussi, C., Metidji, A., Iseppon, A., Lee, S., Tolaini, M., Li, Y., Kelly, G.,

Chakravarty, P., Shoaie, S.*, et al.* (2019). The Intestine Harbors Functionally Distinct

Homeostatic Tissue-Resident and Inflammatory Th17 Cells. Immunity *51*, 77-89 e76.

Pacold, M.E., Brimacombe, K.R., Chan, S.H., Rohde, J.M., Lewis, C.A., Swier, L.J., Possemato,

R., Chen, W.W., Sullivan, L.B., Fiske, B.P*., et al.* (2016). A PHGDH inhibitor reveals

coordination of serine synthesis and one-carbon unit fate. Nat Chem Biol *12*, 452-458.

Pariser, D., Frankel, E., Schlessinger, J., Poulin, Y., Vender, R., Langley, R.G., Meng, X.,

Guana, A., and Nyirady, J. (2018). Efficacy of Secukinumab in the Treatment of Moderate to

Severe Plaque Psoriasis in the North American Subgroup of Patients: Pooled Analysis of Four

Phase 3 Studies. Dermatol Ther (Heidelb) *8*, 17-32.

Peng, M., Yin, N., Chhangawala, S., Xu, K., Leslie, C.S., and Li, M.O. (2016). Aerobic

glycolysis promotes T helper 1 cell differentiation through an epigenetic mechanism. Science

354, 481-484.

Peters, C.L., Morris, C.J., Mapp, P.I., Blake, D.R., Lewis, C.E., and Winrow, V.R. (2004). The

transcription factors hypoxia-inducible factor 1alpha and Ets-1 colocalize in the hypoxic

synovium of inflamed joints in adjuvant-induced arthritis. Arthritis Rheum *50*, 291-296.

Picelli, S., Faridani, O.R., Bjorklund, A.K., Winberg, G., Sagasser, S., and Sandberg, R. (2014).

Full-length RNA-seq from single cells using Smart-seq2. Nat Protoc *9*, 171-181.

Piskin, G., Sylva-Steenland, R.M., Bos, J.D., and Teunissen, M.B. (2006). In vitro and in situ

expression of IL-23 by keratinocytes in healthy skin and psoriasis lesions: enhanced

expression in psoriatic skin. J Immunol *176*, 1908-1915.

Platt, R.J., Chen, S., Zhou, Y., Yim, M.J., Swiech, L., Kempton, H.R., Dahlman, J.E., Parnas, O.,

Eisenhaure, T.M., Jovanovic, M*., et al.* (2014). CRISPR-Cas9 knockin mice for genome

editing and cancer modeling. Cell *159*, 440-455.

Puel, A., Cypowyj, S., Bustamante, J., Wright, J.F., Liu, L., Lim, H.K., Migaud, M., Israel, L.,

Chrabieh, M., Audry, M.*, et al.* (2011). Chronic mucocutaneous candidiasis in humans with

inborn errors of interleukin-17 immunity. Science *332*, 65-68.

Raez, L.E., Papadopoulos, K., Ricart, A.D., Chiorean, E.G., Dipaola, R.S., Stein, M.N., Rocha

Lima, C.M., Schlesselman, J.J., Tolba, K., Langmuir, V.K.*, et al.* (2013). A phase I dose-

escalation trial of 2-deoxy-D-glucose alone or combined with docetaxel in patients with

advanced solid tumors. Cancer Chemother Pharmacol *71*, 523-530.

Schneider, A.S. (2000). Triosephosphate isomerase deficiency: historical perspectives and

molecular aspects. Best Pract Res Cl Ha *13*, 119-140.

Segre, D., Deluna, A., Church, G.M., and Kishony, R. (2005). Modular epistasis in yeast

metabolism. Nat Genet *37*, 77-83.

Semenza, G.L. (2013). HIF-1 mediates metabolic responses to intratumoral hypoxia and

oncogenic mutations. J Clin Invest *123*, 3664-3671.

Semenza, G.L. (2014). Oxygen sensing, hypoxia-inducible factors, and disease pathophysiology.

Annu Rev Pathol *9*, 47-71.

Shi, L.Z., Wang, R., Huang, G., Vogel, P., Neale, G., Green, D.R., and Chi, H. (2011).

HIF1alpha-dependent glycolytic pathway orchestrates a metabolic checkpoint for the

differentiation of TH17 and Treg cells. J Exp Med *208*, 1367-1376.

Stuart, T., Butler, A., Hoffman, P., Hafemeister, C., Papalexi, E., Mauck, W.M., 3rd, Hao, Y.,

Stoeckius, M., Smibert, P., and Satija, R. (2019). Comprehensive Integration of Single-Cell

Data. Cell *177*, 1888-1902 e1821.

Sullivan, L.B., Gui, D.Y., Hosios, A.M., Bush, L.N., Freinkman, E., and Vander Heiden, M.G.

(2015). Supporting Aspartate Biosynthesis Is an Essential Function of Respiration in

Proliferating Cells. Cell *162*, 552-563.

Tahir, H. (2018). Therapies in ankylosing spondylitis-from clinical trials to clinical practice.

Rheumatology (Oxford) *57*, vi23-vi28.

Thiele, I., Vo, T.D., Price, N.D., and Palsson, B.O. (2005). Expanded metabolic reconstruction of

Helicobacter pylori (iIT341 GSM/GPR): an in silico genome-scale characterization of single-

and double-deletion mutants. J Bacteriol *187*, 5818-5830.

Van Welden, S., Selfridge, A.C., and Hindryckx, P. (2017). Intestinal hypoxia and hypoxia-

induced signalling as therapeutic targets for IBD. Nat Rev Gastroenterol Hepatol *14*, 596-611.

Vander Heiden, M.G., Cantley, L.C., and Thompson, C.B. (2009). Understanding the Warburg

effect: the metabolic requirements of cell proliferation. Science *324*, 1029-1033.

Vander Heiden, M.G., and DeBerardinis, R.J. (2017). Understanding the Intersections between

Metabolism and Cancer Biology. Cell *168*, 657-669.

Veldhoen, M., Hocking, R.J., Atkins, C.J., Locksley, R.M., and Stockinger, B. (2006). TGFbeta

in the context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-

producing T cells. Immunity *24*, 179-189.

Wang, E.A., Suzuki, E., Maverakis, E., and Adamopoulos, I.E. (2017). Targeting IL-17 in

psoriatic arthritis. Eur J Rheumatol *4*, 272-277.

Warburg, O., Gawehn, K., and Geissler, A.W. (1958). [Metabolism of leukocytes]. Z

Naturforsch B *13B*, 515-516.

Xu, M., Pokrovskii, M., Ding, Y., Yi, R., Au, C., Harrison, O.J., Galan, C., Belkaid, Y.,

Bonneau, R., and Littman, D.R. (2018). c-MAF-dependent regulatory T cells mediate

immunological tolerance to a gut pathobiont. Nature *554*, 373-377.

Yang, R., and Dunn, J.F. (2015). Reduced cortical microvascular oxygenation in multiple

sclerosis: a blinded, case-controlled study using a novel quantitative near-infrared

spectroscopy method. Sci Rep *5*, 16477.

Yang, Y., Torchinsky, M.B., Gobert, M., Xiong, H., Xu, M., Linehan, J.L., Alonzo, F., Ng, C.,

Chen, A., Lin, X.*, et al.* (2014). Focused specificity of intestinal TH17 cells towards

commensal bacterial antigens. Nature *510*, 152-156.

Yuan, J., Bennett, B.D., and Rabinowitz, J.D. (2008). Kinetic flux profiling for quantitation of

cellular metabolic fluxes. Nat Protoc *3*, 1328-1340.

Zaidi, A.U., Kedar, P., Koduri, P.R., Goyette, G.W., Buck, S., Paglia, D.E., and Ravindranath, Y.

(2017). Glucose phosphate isomerase (GPI) Tadikonda: Characterization of a novel

Pro340Ser mutation. Pediatr Hemat Oncol *34*, 449-454.

Zanella, A., Fermo, E., Bianchi, P., and Valentini, G. (2005). Red cell pyruvate kinase

deficiency: molecular and clinical aspects. Br J Haematol *130*, 11-25.

Zhao, D., Badur, M.G., Luebeck, J., Magana, J.H., Birmingham, A., Sasik, R., Ahn, C.S., Ideker,

T., Metallo, C.M., and Mali, P. (2018). Combinatorial CRISPR-Cas9 Metabolic Screens

Reveal Critical Redox Control Points Dependent on the KEAP1-NRF2 Regulatory Axis. Mol

Cell *69*, 699-708 e697.

Zheng, Y., Danilenko, D.M., Valdez, P., Kasman, I., Eastham-Anderson, J., Wu, J., and Ouyang,

W. (2007). Interleukin-22, a T(H)17 cytokine, mediates IL-23-induced dermal inflammation

and acanthosis. Nature *445*, 648-651.

Zhu, J., and Thompson, C.B. (2019). Metabolic regulation of cell growth and proliferation. Nat

Rev Mol Cell Biol *20*, 436-450.

1513