Transgenic mice expressing human alpha-synuclein in noradrenergic neurons develop locus coeruleus pathology and non-motor features of Parkinson's disease

Butkovich LM[1], Houser MC[1], Chalermpalanupap T[2], Porter-Stransky KA[2,3], Eidson LN[1], De Sousa Rodrigues ME[1], Oliver DL[1], Kelly SD[1], Chang J[1], Bengoa-Vergniory N[4], Wade-Martins R[4], Weinshenker D[2], Tansey MG[5]

[1] Department of Physiology, Emory School of Medicine, 30322
[2] Department of Human Genetics, Emory School of Medicine, 30322
[3] Department of Biomedical Sciences, Western Michigan University Homer Stryker M.D. School of Medicine, 49008
[4] Oxford Parkinson's Disease Centre, Department of Physiology, Anatomy and Genetics, University of Oxford, UK OX1 3QX
[5] Department of Neuroscience and Center for Translational Research in Neurodegenerative Disease, University of Florida College of Medicine, 32610

Contact:
Malú G Tansey, PhD
Department of Neuroscience
University of Florida College of Medicine
1149 Newell Drive, L5-101B
Gainesville, FL 32610
mgtansey@ufl.edu

Key words: Parkinson's disease, α-synuclein, locus coeruleus, aging, norepinephrine

Acknowledgements
This work was supported by NIH/NINDS 5F31 NS098673 (LB), NIH/NINDS F32 NS098615 (KP), NIH/NINDS 1R01 NS102306 (DW), NIH/NIA 1RF1 AG047667 (DW), NIH/NIA 1R01AG061175 (DW), NIH/NIA 1R01 AG057247 (MT), NIH/NINDS 5R01 NS092122 (MT), and NIH/NIA 3RF1 AG051514-01 (MT). This research project was supported in part by the Emory University Integrated Cellular Imaging Microscopy Core, by the Rodent Behavioral Core (RBC), which is subsidized by the Emory University School of Medicine and is one of the Emory Integrated Core Facilities. Additional support was provided by the Emory Neuroscience NINDS Core Facilities P30NS055077. Further support was provided by the Georgia Clinical & Translational Science Alliance of the National Institutes of Health under Award Number UL1TR002378. This study was supported in part by the Emory HPLC Bioanalytical Core (EHBC), which was supported by the Department of Pharmacology, Emory University School of Medicine and the Georgia Clinical & Translational Science Alliance of the National Institutes of Health under Award Number UL1TR002378. The content is solely the responsibility of the authors and does not necessarily reflect the official views of the National Institutes of Health.

Word count: Abstract =175; Introduction =603; Discussion =1471

Abstract

Degeneration of locus coeruleus (LC) neurons and dysregulation of noradrenergic signaling are ubiquitous features of Parkinson's disease (PD). The LC is among the first brain regions affected by α-synuclein (asyn) pathology, yet how asyn affects these neurons remains unclear. LC-derived norepinephrine (NE) can stimulate neuroprotective mechanisms and modulate immune cells, while dysregulation of NE neurotransmission may exacerbate disease progression, particularly non-motor symptoms, and contribute to the chronic neuroinflammation associated with PD pathology. Although transgenic mice overexpressing asyn have previously been developed, transgene expression is usually driven by pan-neuronal promoters and thus has not been selectively targeted to LC neurons. Here we report a novel transgenic mouse expressing human wild-type asyn under control of the noradrenergic-specific dopamine β-hydroxylase promoter. These mice developed asyn aggregates in LC neurons, alterations in hippocampal and LC microglial abundance, upregulated GFAP expression, degeneration of LC fibers, decreased striatal dopamine metabolism, and age-dependent behaviors reminiscent of non-motor symptoms of PD. These mice provide novel insights into how asyn pathology affects LC neurons and how LC dysfunction may contribute to early PD pathophysiology.

Significance statement

α-synuclein (asyn) pathology and loss of neurons in the locus coeruleus (LC) are two of the most ubiquitous neuropathologic features of Parkinson's disease (PD). Dysregulated NE neurotransmission is associated with the non-motor symptoms of PD including sleep disturbance, emotional changes such as anxiety and depression, and cognitive decline. Importantly, loss of central NE may contribute to the chronic inflammation in, and

progression of, PD. We have generated a novel transgenic mouse expressing human

wild-type asyn in LC neurons to investigate how increased asyn expression affects the

function of the central noradrenergic transmission and associated behaviors. We report

cytotoxic effects of asyn overexpression associated with astrogliosis in the LC and LC

target regions, degeneration of LC fibers with disruptions in striatal dopamine (DA)

metabolism, and age-dependent alterations in non-motor behaviors.

Introduction

Brain regions affected in Parkinson's disease (PD) exhibit proteinaceous aggregates

(known as Lewy bodies) primarily composed of α-synuclein (asyn), chronic inflammation,

and neuron loss (den Hartog and Bethlem, 1960; Spillantini et al., 1997; Tansey and

Goldberg, 2010). Striatal dopamine (DA) and DA transporter (DAT) are deficient in PD,

but other neurotransmitter systems are affected including acetylcholine, serotonin, and

norepinephrine (NE) (Gonera et al., 1997; Abbott et al., 2005; Ross et al., 2008).

The locus coeruleus (LC) is among the first brain regions affected in PD. The LC is a

bilateral pontine nucleus at the lateral floor of the fourth ventricle, and is the main source

of NE for the central nervous system (CNS) (Iversen et al., 1983; Mann et al., 1983; Braak

et al., 2001). Asyn aggregates and neuronal degeneration in the LC are ubiquitous

features of PD and are associated with non-motor symptoms including sleep disorders,

mood disturbances, and cognitive deficits (Iversen et al., 1983; Chui et al., 1986; German

et al., 1992; Braak et al., 2001; Zarow et al., 2003; Weinshenker, 2018). Imaging and

histological studies show a progressive loss of central NE, noradrenergic neurons, and

accumulation of asyn pathology in the LC in PD (Halliday et al., 1990; German et al.,

1992; Brunnstrom et al., 2011; Pifl et al., 2012; Keren et al., 2015) which may exacerbate

degeneration of DA neurons in the midbrain substantia nigra pars compacta (SNpc)

(Zarow et al., 2003; Chen et al., 2014).

Depletion of LC-NE exacerbates 6-hydroxydopamine (6-OHDA) or 1-methyl-4phenyl-

1,2,3,6-tetrahydropyridine (MPTP)-induced nigrostriatal pathology in rodents and

primates (Mavridis et al., 1991; Srinivasan and Schmidt, 2003; Rommelfanger et al.,

2007), while increasing extracellular NE is protective (Kilbourn et al., 1998;

Rommelfanger et al., 2004; Kreiner et al., 2019). Furthermore, lesioning LC neurons

induces inflammation, and dysregulated NE neurotransmission may contribute to the

chronic inflammation seen in PD (Kim and Joh, 2006; Tansey and Goldberg, 2010; Yao

et al., 2015; Bharani et al., 2017; Song et al., 2018).

The initiating event in asyn aggregation in sporadic PD is unclear, but a candidate

mechanisms include environmental exposures, and increased expression of asyn, as

individuals with a multiplication mutation in the gene encoding asyn (*SNCA*) develop

autosomal dominantly-inherited PD (Singleton et al., 2003; Chartier-Harlin et al., 2004;

Ferese et al., 2015). Age is the primary risk factor for PD, and rodent models of asyn

overexpression develop age-dependent asyn aggregates and PD-like behavioral

abnormalities (Masliah et al., 2000; Giasson et al., 2002; Hansen et al., 2013). However,

in most of these models, transgene expression is driven by a pan-neuronal promoter with

asyn overexpression in multiple regions of the CNS, with only one model reporting LC

pathology (Masliah et al., 2000; Giasson et al., 2002; Maskri et al., 2004; Schell et al.,

2009; Koprich et al., 2010; Sotiriou et al., 2010; Delenclos et al., 2017). Viral-mediated

expression has been used to target asyn overexpression to specific brain regions

(Baekelandt et al., 2002; Kirik et al., 2002; Delenclos et al., 2017; Ip et al., 2017; Niu et

al., 2018). Notably, viral overexpression of a familial PD mutant asyn in LC neurons

resulted in asyn aggregation, inflammation, and degeneration of LC neurons (Henrich et

al., 2018). To investigate specifically how pathology induced by wild-type asyn affects

noradrenergic neurons in the LC in an aging organism, we targeted expression of human

wild-type asyn to LC neurons under control of the noradrenergic/adrenergic-specific

dopamine β-hydroxylase (DBH) promoter using bacterial artificial chromosome (BAC)

transgenesis. To determine the molecular, cellular, and behavioral age-dependent

consequences of increased asyn expression in LC neurons, 3-, 14-, and 24-month (mo)

old DBH-hSNCA transgenic (Tg) mice and non-transgenic (nTg) littermate controls were

examined.

Methods

Experimental design. To determine how expression of human asyn in LC neurons

affects neuronal health and LC-NE-associated behaviors, mice were aged undisturbed

until the time of behavioral testing or sacrifice at ages 3-, 14-, and 24-mos.

Generation of the DBH-hSNCA mouse model. Male and female mice expressing

human wild-type α-synuclein (DBH-hSNCA) were engineered using a commercially

available human bacterial artificial chromosomal (BAC) RP11-746P3 (Cubells et al.,

2016) encompassing the *DBH* gene. The wild-type *hSNCA* cDNA open reading frame

(400bp) was targeted to the translational start site of *DBH* by standard BAC

recombineering methods by the University of North Carolina – Chapel Hill Molecular

Neuroscience Core (currently Animal Model Core). The BAC construct was injected into

C57BL/6N pronuclei by the Emory University Mouse Transgenic and Gene Targeting

Core Facility (http://www.cores.emory.edu/tmc/index.html), transgene expression in

founder pups was determined by PCR, and breeding lines were established. Mice

carrying the hSNCA sequence were crossed with wild-type C57Bl/6N mice (Charles

River) to establish the hemizygous transgenic DBH-hSNCA line.

To improve efficiency and accuracy of LC tissue isolation for western blot and mRNA

analysis, DBH-hSNCA mice were crossed with the TH-EGFP reporter mouse expressing

enhanced green fluorescent protein (EGFP) under the tyrosine hydroxylase (TH)

promotor (Sawamoto et al., 2001).

Animals. Male and female DBH-hSNCA mice were maintained on a C57Bl/6 background.

Mice were group housed (maximum 5 mice per cage) until two weeks prior to the start of

behavioral testing, when they were singly housed until euthanized. Animals were

maintained on a 12/12h light/dark cycle with access to standard rodent chow and water

ad libitum. Hemizygous animals served as experimental mice, with non-transgenic

littermates as controls. Genotypes were determined by tail snip PCR with two sets of

primers: Forward 5' TGTCCAAGATGGACCAGACTC 3' Reverse 3'

ACTGGTCTGAGGCAGGGAGCA 5'; Set Forward 5' GCCCTCAGTCTACTTGCGGGA 3'

Reverse 3' GCGAGAGCATCATAGGGAGT 5'. Experimental procedures involving use of

animals were performed in accordance with the NIH Guidelines for Animal Care and Use

and approved by the Institutional Animal Care and Use Committee at Emory University

School of Medicine.

Sleep latency test. Latency to fall asleep was quantified as the duration of time following

gentle handling until their first sleep bout, which was defined as sleeping continuously for

2 min, and for a total of 75% of the 10-min period that began at sleep onset (Hunsley and

Palmiter, 2004). Sleep testing began at 9 AM, 2 h into the light cycle when internal

pressure to sleep is high. The sessions were video recorded and scored by

an experienced observer blind to the genotype. We have validated this behavioral sleep

scoring method with EEG (Porter-Stransky et al., 2019).

Marble burying test. Marble burying was conducted as previously described (de Sousa

Rodrigues et al., 2017) to determine whether expression of human asyn in LC neurons

promotes anxiety-like behavior. Mice were placed in a plastic tub (50.5 x 39.4 x 19.7 cm)

containing 5 inches of lightly pressed bedding. Twenty marbles of uniform size and color

were placed in 5 rows of 4 marbles each on top of the bedding. Mice were placed in the

containers and allowed to roam freely for 30 min. At the end of testing, the mice were

placed back in home cages, and the number of marbles buried at least two-thirds of their

height were counted. Marble burying was conducted 2 weeks after sleep latency testing.

Open Field testing. In the open field test, a mouse that spends less time in or hesitates

to re-enter the open center of the testing chamber is considered to be exhibiting anxiety-

like behavior (Britton and Britton, 1981). During the light phase of the light/dark cycle,

mice were acclimated to a dark testing room under red light for 1 h before testing. Mice

were placed into the open field (45 cm X 45 cm square box) and allowed to move freely

for 10 min. Distance, velocity, center, and border statistics were measured using

Noldus/Ethovision software. Center was defined as the central 22.5 cm X 22.5 cm. Open

field was conducted 1 week after marble burying.

Circadian locomotion. All testing mice were acclimated to the testing room for 2 d prior

to the experiment. Mice were each placed in a clear Plexiglas (15.75" L, 13.25" L, 7.38"

H) activity cage equipped with infrared photobeams (San Diego Instruments, La Jolla,

CA). Food and water were available *ad libitum* during the 23-h testing period. Ambulations

(consecutive photobeam breaks) were recorded by PAS software. Circadian locomotion

behavior was assessed 2 weeks after open field testing.

Fear conditioning. Fear conditioning training and contextual and cued fear testing is a

test of memory for the association of an aversive stimulus with an environment cue or

context, and was conducted as previously described (Chalermpalanupap et al., 2018)

over 3 consecutive days. Mice were placed in the fear conditioning apparatus (7" W, 7"

D, 12" H, Coulbourn) with metal shock grid floor and allowed to explore the enclosure for

3 min. Following habituation, three conditioned stimulus (CS)-unconditioned stimulus

(US) pairings were presented with a one-min inter-trial interval. The CS was a 20 sec 85

db tone, and the US was a 2 sec 0.5mA footshock (Precision Animal Shocker, Colbourn)

which co-terminated with CS presentation. The contextual test was conducted on the

following day when animals were placed back into the same chamber. On day three, the

animals were placed in a novel compartment and allowed to habituate for 2 min. Following

habituation, the 85 db tone was presented, and the amount of freezing behavior recorded.

No shocks were given during the contextual or cued tests. Fear conditioning was

conducted 1 week after circadian locomotion behavior.

Tissue collection. Animals used in immunohistochemical and high-performance liquid

chromatography (HPLC) analyses were anesthetized by injection of sodium pentobarbital

(Euthasol, Virbac) until unresponsive. Mice were transcardially perfused with phosphate-

buffered saline (PBS; pH 7.4) until exiting blood ran clear. Brain tissue was removed, with

one hemisphere post-fixed in 4% paraformaldehyde for immunohistochemistry, and the

other dissected and flash frozen for HPLC. Animals used for qPCR or western blot

analyses were euthanized by cervical dislocation under isoflurane anesthesia. Tissue was

flash frozen and stored at -80°C until processing.

Immunohistochemistry. Brain tissue was sectioned on a freezing microtome (Leica

SM2010R, Buffalo Grove, IL) at 40μm and stored in cryoprotectant (30% ethylene glycol,

30% sucrose, 13.32mM NaH_2PO_4, 38.74mM Na_2HPO_4, 250μM Polyvinylpyrrolidone)

solution at -20°C until staining. Sections were washed in PBS before blocking in 5%

normal goat serum (Jackson ImmunoResearch 005-000-121; NGS) with 0.05% Triton X-

100 (Sigma #T9284100) in Tris-buffered saline pH 7.4 (TBS) for 1 h at room temperature.

Sections were transferred directly to primary antibody solution containing 1% NGS, 0.05%

Triton-X 100, and antibody at the concentrations described in Table 1 and incubated

overnight at room temperature (05-02 Ms anti-NET, 1:1,000) or 4°C (all other primary

antibodies). Fluorescently conjugated secondary antibodies (described in Table 1) were

diluted in 0.1% NGS with 0.05% Triton X-100, and tissue sections were incubated for 1 h

at room temperature in the dark. Sections were mounted on Superfrost Plus slides (VWR)

and were coverslipped with Vectashield with DAPI (Vector). All immunofluorescent

images were acquired as z-stack images and the file compressed on a Keyence BZ-X700

microscope system (Itasca, IL). The Allen Brain Atlas version 1 (2008) was used to

identify regions of interest (ROIs). One section per mouse containing the dorsal

hippocampus (near bregma -1.995 mm) and one containing the LC (bregma -5.555 mm)

were analyzed for percent immunoreactivity (IR) within a standard ROI. A detection

threshold was set uniformly across images in each analysis, and % IR determined using

the "Measure" feature of ImageJ. Percent IR was calculated as area of IR within the ROI

divided by the total ROI area and multiplied by 100. Quantification of Iba1-positive cells

(microglia) was also analyzed with a standard threshold, ROI, and upper and lower size

limits (pixel^2) using the "Analyze particles" function in ImageJ.

Proximity ligation assay (PLA). Paraffin embedded tissue was rehydrated by

consecutive incubations in Xylene, Histoclear, 100% ethanol, 95% ethanol, 70% ethanol

and H_2O. Samples were then incubated in 10% H_2O_2 in PBS to reduce background and

heated in a microwave in citrate buffer (pH 6.0; Abcam) for antigen retrieval. After antigen

retrieval, samples were processed for immunofluorescence: 1 h RT incubation in 10%

Serum with 0.05% Tween-20 in TBS block, 1 h incubation in primary antibody, TBS with

0.05% Tween-20 (TBS-T) wash. Slides were then incubated for 1 h with secondary

antibodies (Alexa488 Life Technologies), and washed again with TBS-T. For

immunohistochemistry, samples were then coverslipped with FluorSave (Calbiochem).

For PLA, samples were covered in manufacturers PLA blocking solution (Sigma) for 1 h

at 37°C, and then incubated overnight with PLA conjugates (a-syn211; ab80627 Abcam).

On the next day, samples were washed with TBS-T, incubated in ligation solution for 1 h

at 37C°, washed with TBS-T, incubated in amplification solution for 2.5 h at 37C°, washed

with TBS, counterstained with DAPI, and mounted with FluorSave. All PLA reagents were

used as per manufacturer's instructions (Sigma; cat #92008).

PLA puncta were counted blindly from 25 cells per LC section per animal. Two

independent experimental rounds were normalized and averaged in order to provide the

final quantification.

RNA Scope. *In situ* RNA analysis was performed using RNAScope Multiplex Fluorescent

v2 kit (ACD Bio 3231000). Tissue prep and analysis were conducted as described in

manufacturer's protocol. Briefly, following transcardial perfusion with saline, brains were

incubated in 4% paraformaldehyde (PFA) for 24 hours followed by a series of increasing

sucrose concentrations before being frozen in optimal temperature cutting medium

(Sakura) and stored at -80°C until sectioning. Tissue sections (12μm) were collected on

a Leica CM1900 cryostat and mounted on Superfrost Plus slides. To prevent tissue

detachment, slides were dried at 60°C for 30 min and fixed in 4% PFA for 15 min at 4°C

before ethanol dehydration. Tissue was processed as described in ACD Bio protocol

(acdbio.com/technical-support/user-manuals #323100-USM).

Western immunoblotting. Western blots were conducted as previously described (de

Sousa Rodrigues et al., 2017). Flash frozen samples were stored at -80°C until

processing. Protein was isolated from LC samples with RIPA buffer (1% Triton-X 100,

50mM Tris HCL, 0.1% sodium dodecyl sulfate, 150mM NaCL, pH 8.0), or Trizol (Life

Technologies #15596-018). RIPA samples were centrifuged at 12,000 rpm for 20 min at

4°C. Supernatant was transferred to new tube for bicinchoninic acid protein assay (Pierce

Scientific #23225). Trizol samples were resuspended in 1% SDS. Samples were diluted

to 1µg/µl in 4x sample buffer (BioRad #1610747) and boiled at 90°C for 5 min.

Electrophoresis was performed using 12% gels (BioRad #4568046; 5µl) and transferred

to 0.45µm PVDF membrane using Trans-Blot Turbo Transfer System (BioRad). For asyn

immunoblotting, the membrane was fixed in 0.4% PFA for 30 min following transfer. After

a brief wash, blots were incubated in 5% milk blocking buffer (BioRad) for 1 hour at 4°C

before primary antibody overnight at 4°C. Membranes were washed with TBST (0.01%

Tween-20) and incubated in HRP-conjugated secondary antibodies in blocking buffer for

1 hour at room temperature. Images were acquired using Azure Biosystems and analyzed

by ImageStudio Lite software. Protein expression was normalized to total protein (Li-Cor

#926-11015).

High performance liquid chromatography (HPLC).

Monoamines were examined by high performance liquid chromatography with
electrochemical detection as described previously (Song et al., 2012). For HPLC, an ESA
5600A CoulArray detection system, equipped with an ESA Model 584 pump and an ESA
542 refrigerated autosampler was used. Separations were performed using an MD-150 ×
3.2 mm C18 (3 µM) column at 25ºC. The mobile phase consisted of 8% acetonitrile, 75
mM NaH_2PO4, 1.7 mM 1-octanesulfonic acid sodium and 0.025% trimethylamine at pH
2.9. Twenty-five microliters of sample were injected. The samples were eluted
isocratically at 0.4 mL/min and detected using a 6210-electrochemical cell (ESA, Bedford,
MA) equipped with 5020 guard cell. Guard cell potential was set at 475 mV, while
analytical cell potentials were −175, 150, 350 and 425 mV. The analytes were identified
by the matching criteria of retention time and sensor ratio measures to known standards
(Sigma Chemical Co., St. Louis MO.) consisting of dopamine, norepinephrine, 3,4-
dihydroxyphenylacetic acid (DOPAC), and 4-Hydroxy-3-methoxyphenylglycol (MHPG).
Compounds were quantified by comparing peak areas to those of standards on the
dominant sensor.

Statistical analysis. Student's t-test was used to assess differences by genotype within
each age group in sleep latency, marble burying, open field, western blot, and all
immunofluorescent analyses. Repeated measures two-way ANOVA was used to analyze
differences by genotype within each age group in fear conditioning, and circadian
locomotor assay followed by Tukey's post hoc test where applicable. Comparisons across
age groups were not conducted, as behavioral assays, HPLC, and immunofluorescence
of each cohort were conducted at separate time points. The analyses were performed

using GraphPad Prism 7 (GraphPad Software, Inc., La Jolla, CA) with a p-value threshold

of <0.05.

Results

Generation of DBH-hSNCA mice

The DBH-hSNCA mouse model was developed using a DBH-BAC construct carrying the

wild type human *SNCA* cDNA open reading frame at the translational start site of *DBH*

(Fig. 1). Transgene integration was confirmed by PCR, and founder mice were bred with

wild type C57BL/6 mice to establish the hemizygous DBH-hSNCA line.

Human *Snca* mRNA is expressed in DBH-hSNCA LC neurons

Fluorescent *in situ* mRNA analysis revealed human *Snca* mRNA in Tg LC neurons (Fig

2), which co-localized with mouse *Th* and mouse *Dbh* mRNA, while human *Snca* mRNA

expression was not detected in nTg LC neurons.

Human asyn is expressed in LC neurons of DBH-hSNCA transgenic mice

Human asyn protein was analyzed by immunofluorescence and western blot. Using an

antibody specific for human asyn (Biolegend 807801), expression of human asyn was

found to co-localize with TH-expressing LC neurons only in brain sections from DBH-

hSNCA Tg mice (Fig. 3A). No human asyn-specific immunofluorescence was detected

in LC neurons of nTg littermates or in SNpc neurons regardless of genotype (Fig. 3A,C).

Human asyn was also detectable specifically in LC protein lysate from Tg tissue by

immunoblot and not in nTg lysates (Fig 3B; n=4). Quantitative western blot analysis of LC

protein using a pan-asyn antibody to detect both human and mouse asyn protein revealed

a significant ~30% increase of total asyn in Tg mice at 3-mo relative to that in nTg

littermates (Fig 3D; $t_{(6)}$=3.156, p=0.0197, n=4). Human *SNCA* mRNA expression in Tg LC

neurons was confirmed by qPCR (Data not shown).

DBH-hSNCA mice exhibit behavioral phenotypes that resemble features of non-

motor PD symptoms

A primary role of the LC-NE system is promoting arousal and wakefulness; LC activity is

highest just prior to, and during wake (Hobson et al., 1975). Sleep disturbances are one

of the most common non-motor PD symptoms, and PD patients with disturbed sleep have

greater asyn pathology in the LC than PD patients without sleep complaints (Kalaitzakis

et al., 2013). Thus, sleep latency was assessed to examine whether features of the

sleep/wake cycle were affected by human asyn expression in the LC. Mice were gently

handled and returned to their home cage, and video recording was scored by an observer

blind to the genotype to determine latency to fall asleep. Our findings indicate that there

was a significant increase in sleep latency in Tg mice at 3-mos, (Fig 4A; $t_{(14)}$=4.36

p=0.0007; n=8) and 14-mos ($t_{(17)}$=2.51, p=0.0225; n=9), indicative of an elevated arousal

state. No differences were observed at 24-mos ($t_{(8)}$=0.821, p=0.4354, n=7-8).

LC neurons are activated by stress exposure, and blocking LC activity abolishes stress-

induced anxiety-like behavior (McCall et al., 2015). In PD, anxiety severity is inversely

correlated with LC function (Remy et al., 2005). To examine anxiety-like behavior, the

behavior DBH-hSNCA mice was assessed in marble burying and open field tests. In the

marble burying assay, there was a significant increase in the number of marbles buried

by Tg mice at 14- (Fig 4B; $t_{(18)}$=2.735, p=0.0136; n=8-12) and 24-mos ($t_{(13)}$=2.212,

p=0.0455, n=7-8). In open field tests, an age-dependent anxiety-like phenotype was also

evident in 14-mo Tg mice, evinced by an increase in the latency to re-enter the center of

the testing field (Fig 4C; $t_{(17)}2.359$, p=0.0305; n=7-12), with no differences in 3-

($t_{(11)}$=0.1988, p=0.8460, n=7-8) or 24-mo old animals ($t_{(11)}$=0.5075, p=0.6219, n=7-8).

Fear conditioning is a measure of hippocampal-dependent (contextual) or -independent

(cued) associative learning (Phillips and LeDoux, 1992), and mice lacking NE exhibit

impaired contextual learning (Murchison et al., 2004). In a standard fear conditioning

paradigm, 14-mo old DBH-hSNCA Tg mice exhibited increased freezing behavior during

the fear training session (Fig 4D center; Interaction $F_{(6, 120)}$=2.735, p=0.0159; n=8-14), as

well as during the contextual test (Fig 4E center; effect of genotype $F_{(1, 20)}$=5.566,

p=0.0286), with no differences in freezing behavior during the cued test (Fig 4F). No

genotype differences were found in the fear training (Fig 4D), contextual test (Fig 4E), or

cued test (Fig 4F) in 3- or 24-mo old mice.

Because disruption in locomotor activity can affect behavioral testing, we evaluated the

locomotion over 24 h and found no genotype differences at 3- ($F_{(1, 12)}$=1.158, p=0.3031;

n=7), 14- ($F_{(1, 11)}$=0.1864, p=0.6743; n=7), or 24-mos ($F_{(1, 13)}$=0.0603, p=0.8099; n=7-8) of

age (Fig. 4F). Initially, all groups had high levels of activity, as would be expected in a

novel environment, which decreased as mice habituated to the test apparatus.

Ambulations increased normally in all genotypes at commencement of the dark phase,

when mice are typically more active, and decreased once the next light cycle began.

However, the 14-mo old Tg mice exhibited significantly more rearing behavior during the

dark phase than the nTg mice (Fig 4G; Interaction $F_{(21, 231)}$=1.911, p=0.0113).

Human asyn forms aggregates in LC neurons at 14-mos of age

To determine whether transgenic expression of human asyn in LC neurons leads to

formation of asyn aggregates, tissue sections containing the LC were analyzed using a

human asyn proximity ligation assay (PLA). In 14-mo old Tg mice (Fig 5), LC neurons

displayed more asyn puncta per field than LC neurons from nTg mice ($t_{(8)}$=2.532,

p=0.0352; n=4-6). Asyn aggregates are fibril-rich and highly phosphorylated at residue

129 (pSer129), and labeling these forms of the protein is commonly used to identify asyn

aggregates (Fujiwara et al., 2002; Wakamatsu et al., 2007; Schell et al., 2009). Therefore,

we immunolabeled tissue sections with an antibody specific for fibril-rich asyn (Syn-F1)

or asyn pSer129 and found no detectable signal (data not shown), suggesting that at 14

mos of age the asyn-immunoreactive puncta are likely to represent intermediate asyn

oligomers rather than mature aggregates.

Human asyn expression in LC neurons impacts striatal dopamine metabolism in

24-mo old mice

Dysregulated catecholamine metabolism and degeneration of catecholaminergic neurons

are well-established features of PD (Iversen et al., 1983; Mann et al., 1983; Hirsch et al.,

1988; Fearnley and Lees, 1991). Therefore, we measured catecholamine levels using

high-performance liquid chromatography (HPLC). Hippocampal and striatal tissue

content of NE, the NE metabolite MHPG, DA, and the DA metabolite DOPAC were

quantified, revealing that NE (Fig. 6A,B) and DA (Fig. 6E,F) were not significantly affected

in the hippocampus or striatum at any age Similarly, the ratio of the major NE metabolite

MHPG to NE was unaffected (Fig, 6C,D). The ratio of the DA metabolite DOPAC to DA

in the hippocampus was unaltered (Fig. 6G), but was significantly reduced in the Tg

striatum at 24-mos (Fig 6H; $t_{(10)}$=3.546, p=0.0046; n=5-7), consistent with decreased DA

turnover.

Human asyn expression does not affect LC neuronal integrity

LC neurons were visualized using NET immunoreactivity (IR). NET is a reliable marker of

LC neurons, and its expression is reduced in PD patients (Remy et al., 2005). Using a

standard ROI, no difference in the percent NET IR was detected between genotypes at

3- (Fig 7; $t_{(17)}$=0.4537, p=0.6558; n=9-10), 14- ($t_{(10)}$=0.8908, p=0.3939; n=5-7), or 24-mos

($t_{(8)}$=0.8069, p=0.4430; n=5-7).

Elevation of tyrosine hydroxylase in the LC of DBH-hSNCA mice

Tyrosine hydroxylase (TH) is the rate-limiting enzyme in NE and DA synthesis, and its

long-term activity depends on its expression levels (Levitt et al., 1965; Haycock, 1993;

Kumer and Vrana, 1996). To determine whether TH expression in the LC is affected by

human asyn, we assessed TH IR in sections from 3-, 14-, and 24-mo old mice. TH IR was

normalized to NET IR to control for potential differences in Bregma level between

sections. TH expression was increased in Tg LC neurons (Fig 8A,B) at both 3-

($t_{(17)}$=2.154, p=0.0459; n=9) and 14-mo ($t_{(10)}$=2.463, p=0.0335; n=5-7) of age. Western

blot analysis from 3-mo old TH-EGFP-expressing LC neurons (Fig 8C,D) confirmed

higher TH expression in Tg animals ($t_{(10)}$=3.837, p=0.0033; n= 5-7).

Human asyn expression in LC neurons affects expression of local inflammatory

markers

A wealth of studies suggest that dysregulated noradrenergic neurotransmission is

associated with inflammation (reviewed by Butkovich et al. 2018), and that dysregulation

of the LC-NE system could contribute to the chronic neuroinflammation observed in PD

(Fujita et al., 1998; Gyoneva and Traynelis, 2013; Johnson et al., 2013; Butkovich et al.,

2018). To determine whether human asyn expression in LC neurons affects the number

of myeloid cells in the brain (both brain-resident microglia and potentially infiltrating

monocytes), we quantified the number of Iba1-positive cells in the LC by

immunofluorescence. At 14-mo, there was a significant decrease in the number of Iba1-

expressing cells in the LC of Tg animals (Fig 9A,B; $t_{(13)}$=2.845, p=0.0138; n=8), with no

changes at other ages. To determine astrocyte activation, we quantified glial fibrillary

acidic protein (GFAP) IR, commonly used as a protein marker of astrogliosis (Eng and

Ghirnikar, 1994), in the LC. At 24-mo, there was a significant increase in astrocytic GFAP

expression in the LC of Tg animals (Fig 9C, D; $t_{(10)}$=2.744, p=0.0207; n=5-7).

Human asyn expression in LC neurons is associated with loss of hippocampal LC

fibers at 24-mos

The LC is the sole source of hippocampal NE, which is necessary for proper memory

formation and retrieval (Devauges and Sara, 1991). Noradrenergic LC fibers express

NET, and PD brain tissue shows substantial LC denervation (Pavese et al., 2011). To

determine whether hippocampal LC projections degenerate in DBH-hSNCA mice, we

examined NET IR in the CA1, CA3, and dentate gyrus regions of the hippocampus. At

24-mos, we found a reduction in LC fibers in the dentate gyrus (Fig 12C,D; $t_{(10)}$=2.974,

p=0.0156; n=5-7), with a trend for reduction in CA1 (Fig 12A, B; $t_{(10)}$=1.899, p=0.0901)

and CA3 (Fig 12E,F; $t_{(10)}$=1.538, p=0.1585). No differences were observed in mice at 3-,

or 14-mos.

Hippocampal astrogliosis and changes in number of hippocampal Iba1-expressing

cells in aged DBH-hSNCA mice

To determine whether degeneration of hippocampal LC projections is associated with

inflammation, GFAP IR was visualized in the CA1, CA3, and dentate gyrus regions of the

hippocampus. At 14-mos, there was a significant increase in GFAP expression in CA1

(Fig 10A,B; $t_{(19)}$=2.723 ,p=0.0135, n=9-12) and CA3 (Fig 10E,F; $t_{(19)}$=2.275 ,p=0.0347),

but not the dentate gyrus (Fig 11C,D; $t_{(19)}$=1.607, p=0.1246), of Tg mice. Similar to what

we observed in the LC, the number of Iba1-expressing cells in CA1 was reduced in Tg

mice at 14-mos (Fig 11; $t_{(16)}$=2.592, p=0.0196, n=9).

Discussion

Based on a wealth of evidence (Mavridis et al., 1991; Srinivasan and Schmidt, 2003; Tong

et al., 2006; Rommelfanger et al., 2007; Yao et al., 2015), we posit that asyn pathology

and degeneration of LC neurons may represent a tipping point in PD progression;

therefore, understanding how asyn accumulation in LC neurons affects their function and

survival may help inform development of new therapeutics for earlier interventions. To

this end, we developed a new BAC transgenic mouse expressing human wild-type asyn

under the control of the noradrenergic-specific DBH.

Human asyn was detectable in Tg LC neurons by immunofluorescence and western blot

at 3-mos. Analysis of total asyn (mouse + human) revealed that the total asyn burden is

increased in Tg LC neurons at 3-mos. While pSer129-positive, or fibrillar asyn aggregates

were not detected in LC neurons at any age, increasing the burden of asyn expression in

LC neurons resulted in the formation of putative oligomeric asyn aggregates. In PD, asyn

aggregates appear prior to neuron degeneration (Chevalier-Larsen and Holzbaur, 2006;

Chu et al., 2012), and while we observed no loss of LC cell bodies, NET-expressing fibers

were reduced in the hippocampus of 24-mo old mice. This finding resembles the pattern

of neuron death observed in PD, with axon terminals degenerating prior to frank cell loss

in the LC (Hornykiewicz, 1998). Interestingly, the selective loss of LC fibers in the dentate

gyrus of DBH-hSNCA mice is reminiscent of what we observed in TgF344-AD rat model

of Alzheimer's disease that accumulates tau pathology in the LC (Rorabaugh et al., 2017).

To determine the functional outcome of asyn aggregates in LC neurons, we measured

catecholamine levels in the hippocampus and striatum. Despite the loss of hippocampal

LC fibers at 24-mos we did not detect changes in hippocampal NE content by HPLC,

which may indicate a compensatory enhancement in NE neurotransmission. We

observed a trend toward decreased striatal NE 24-mos, although it was not statistically

significant. The sparse innervation by NE, it is possible that changes in NE

neurotransmission are detectable in the striatum prior to more densely innervated

regions. Dysregulated DA metabolism is a central feature of PD (Leenders et al., 1990),

and midbrain dopaminergic innervation to the striatum is modulated by LC-NE (Lategan

et al., 1990; Grenhoff et al., 1993; Rommelfanger et al., 2007; Rommelfanger and

Weinshenker, 2007). Studies of catecholamine function in rodents show that enhancing

LC-NE stimulates midbrain DA release in the striatum, whereas LC lesions or NE

deficiency reduce striatal DA release (Lategan et al., 1990; Grenhoff et al., 1993; Schank

et al., 2006). We found that at 24-mos, Tg mice had a reduced striatal ratio of DOPAC:DA,

and a trend toward increased DA, suggesting that human asyn expression in aging LC

neurons causes a reduction in striatal DA turnover. Future studies will determine whether

NE innervation to midbrain DA neurons is reduced in 24-mo DBH-hSNCA mice, and

whether loss of noradrenergic transmission impairs striatal DA release. While no

significant differences were detected in hippocampal or striatal NE content at any age, Tg

LC neurons had increased TH expression at 3- and 14-mos relative to nTg littermates,

suggesting an increased capacity for NE synthesis.

Neuroinflammation is a central feature of PD pathology (McGeer et al., 1988; Gerhard et

al., 2006; Tansey and Goldberg, 2010), with extensive evidence of changes in microglial

activation in brain regions that degenerate in PD (Kim and Joh, 2006; Tansey and

Goldberg, 2010), and we know that microglia and astrocyte functions are modulated by

NE neurotransmission (Fahrig, 1993; Heneka et al., 2010; Bharani et al., 2017).

Unexpectedly, we observed fewer Iba1-expressing cells in the LC and CA1 region of the

hippocampus of DBH-hSNCA mice at 14-mos compared to nTg littermates. It has been

reported that NE can modulate microglial activity, and the functional outcome of microglia

adrenergic receptor activation appears to depend on the physiological context (Fujita et

al., 1998; Gyoneva and Traynelis, 2013; Johnson et al., 2013). At 14-mos, there was an

increase in hippocampal astrocytic GFAP expression. Astrocytes throughout the brain are

activated by NE, and ARs signal transduction stimulates transient calcium release, and

release of inflammatory signaling molecules by astrocytes (Norris and Benveniste, 1993;

Duffy and MacVicar, 1995; Paukert et al., 2014). However, the number GFAP-expressing

astrocytes is inversely correlated to the amount of dopaminergic cell loss in PD and

increased GFAP immunoreactivity in the hippocampus occurred prior to loss of LC

projections (Damier et al., 1993; Eng and Ghirnikar, 1994). Further studies are required

to   determine   whether   increased   GFAP   activity   is   due   to   dysregulated   NE

neurotransmission, or an inflammatory response secondary to degeneration of LC

projections.

Historically, the LC has been implicated in arousal state and stress responses. For

example, LC activity tracks with sleep cycles (with highest firing during wake and

immediately preceding sleep-wake transitions), and chemogenetic or optogenetic

activation of LC neurons increases wakefulness (Carter et al., 2010; Vazey and Aston-

Jones, 2014; Porter-Stransky et al., 2019). Furthermore, the LC is activated by stress,

and stimulation of LC neurons elicits anxiety-like behaviors (Valentino and Van

Bockstaele, 2008; McCall et al., 2015). DBH-hSNCA mice exhibited age-dependent

behavioral phenotypes that peaked at 14-mos and are consistent with LC hyperactivity

and increased NE transmission. Specifically, compared to nTg littermates, DBH-hSNCA

mice displayed increased arousal (as measured by latency to fall asleep), anxiety (as

measured by marble burying and latency to re-enter the center of an open field), and

stress responses (as measured by freezing during fear conditioning training and context

re-exposure). These phenotypes are relevant to multiple non-motor symptoms of PD.

Sleep disturbances are one of the most common complaints of PD patients, and patients

who experience disturbed sleep appear to have greater LC asyn pathology than those

who do not report sleep disturbances (Kalaitzakis et al., 2013). Anxiety is also a common

complaint, as up to 60% of PD patients report experiencing anxiety (Chaudhuri and

Schapira, 2009; Lin et al., 2015). Importantly, DBH-hSNCA mice did not display changes

in the number or speed of ambulations relative to nTg mice at any age examined, ruling

out a general locomotor abnormality that has been observed in more ubiquitous asyn

overexpression mice (Giasson et al., 2002; Fleming et al., 2004; Graham and Sidhu,

2010).

We have previously speculated that asyn pathology promotes LC hyperactivity and non-

motor symptoms during PD progression prior to the degeneration of noradrenergic

neurons later in the disease (Weinshenker, 2018), and the present data support that idea.

Indeed, the behavioral phenotypes we observed were most prominent at 14-mos, while

degeneration of LC fibers was not evident until 24-mos, a time when behavioral

abnormalities abated. There are several potential mechanisms by which increases in

asyn may be affecting the LC-NE system (Fig 13). TH protein was increased in 3- and

14-mo old Tg mice, which is when behavioral phenotypes consistent with NE over-activity

emerged. Because TH is the rate-limiting enzyme in NE production, this may reflect

increased NE synthetic capacity, although the relevance is not clear because we did not

detect differences in tissue NE content. Human wild-type asyn has been reported to

increase in the size, and delay in the closing, of the vesicular fusion pore, allowing more

neurotransmitter to spill into the extracellular space (Larsen et al., 2006; Logan et al.,

2017). In addition, viral-mediated overexpression of A53T mutant asyn in LC neurons

causes increased firing rate (Henrich et al., 2018). Alternatively, if human asyn expression

is negatively impacting LC neuron health as suggested by the presence of asyn

aggregates at 14-mos and the loss of hippocampal LC fibers at 24-mos, there may be

compensatory increases in neuronal activity. Chemical lesion of LC neurons transiently

increases NE neurotransmission by increasing neuronal firing frequency and activity

patterns (Szot et al., 2016). Future studies directly examining LC neuron firing and NE

release will be necessary to identify the mechanism in play here.

Little is known about the potential involvement of other brainstem NE regions in PD. One

study reported increased DBH activity in the A2/C2 of PD patients versus healthy controls

(Molinoff and Axelrod, 1971; Kopp et al., 1982). While generally considered an integrative

region for autonomic functions, A2/C2 neurons are activated by stress and can influence

behavior (Cecchi et al., 2002; Ritter et al., 2003). Future studies are required to determine

if other central noradrenergic regions contribute to the DBH-hSNCA phenotype.

LC neurons are among the first affected in PD, and the features of DBH-hSNCA mice

may represent early pathology and non-motor components of PD that provide insight into

the functional impact of human asyn expression in LC neurons during aging. Additionally,

studies involving exposure to environmental factors that synergize with asyn expression

to influence the risk of PD will likely elucidate the genetic and environmental interactions

that contribute to LC involvement in the pre-clinical pre-motor stages of PD, as well as

the impact of LC degeneration on the trajectory of PD pathogenesis.

References

Abbott RD, Ross GW, White LR, Tanner CM, Masaki KH, Nelson JS, Curb JD, Petrovitch H (2005) Excessive daytime sleepiness and subsequent development of Parkinson disease. Neurology 65:1442-1446.

Baekelandt V, Claeys A, Eggermont K, Lauwers E, De Strooper B, Nuttin B, Debyser Z (2002) Characterization of lentiviral vector-mediated gene transfer in adult mouse brain. Hum Gene Ther 13:841-853.

Bharani KL, Derex R, Granholm AC, Ledreux A (2017) A noradrenergic lesion aggravates the effects of systemic inflammation on the hippocampus of aged rats. PLoS One 12:e0189821.

Braak E, Sandmann-Keil D, Rub U, Gai WP, de Vos RA, Steur EN, Arai K, Braak H (2001) alpha-synuclein immunopositive Parkinson's disease-related inclusion bodies in lower brain stem nuclei. Acta Neuropathol 101:195-201.

Britton DR, Britton KT (1981) A sensitive open field measure of anxiolytic drug activity. Pharmacol Biochem Behav 15:577-582.

Brunnstrom H, Friberg N, Lindberg E, Englund E (2011) Differential degeneration of the locus coeruleus in dementia subtypes. Clin Neuropathol 30:104-110.

Butkovich LM, Houser MC, Tansey MG (2018) alpha-Synuclein and Noradrenergic Modulation of Immune Cells in Parkinson's Disease Pathogenesis. Front Neurosci 12:626.

Carter ME, Yizhar O, Chikahisa S, Nguyen H, Adamantidis A, Nishino S, Deisseroth K, de Lecea L (2010) Tuning arousal with optogenetic modulation of locus coeruleus neurons. Nat Neurosci 13:1526-1533.

Cecchi M, Khoshbouei H, Javors M, Morilak DA (2002) Modulatory effects of norepinephrine in the lateral bed nucleus of the stria terminalis on behavioral and neuroendocrine responses to acute stress. Neuroscience 112:13-21.

Chalermpalanupap T, Schroeder JP, Rorabaugh JM, Liles LC, Lah JJ, Levey AI, Weinshenker D (2018) Locus Coeruleus Ablation Exacerbates Cognitive Deficits, Neuropathology, and Lethality in P301S Tau Transgenic Mice. J Neurosci 38:74-92.

Chartier-Harlin MC, Kachergus J, Roumier C, Mouroux V, Douay X, Lincoln S, Levecque C, Larvor L, Andrieux J, Hulihan M, Waucquier N, Defebvre L, Amouyel P, Farrer M, Destee A (2004) Alpha-synuclein locus duplication as a cause of familial Parkinson's disease. Lancet 364:1167-1169.

Chaudhuri KR, Schapira AH (2009) Non-motor symptoms of Parkinson's disease: dopaminergic pathophysiology and treatment. Lancet Neurol 8:464-474.

Chen X, Huddleston DE, Langley J, Ahn S, Barnum CJ, Factor SA, Levey AI, Hu X (2014) Simultaneous imaging of locus coeruleus and substantia nigra with a quantitative neuromelanin MRI approach. Magn Reson Imaging 32:1301-1306.

Chevalier-Larsen E, Holzbaur EL (2006) Axonal transport and neurodegenerative disease. Biochim Biophys Acta 1762:1094-1108.

Chu Y, Morfini GA, Langhamer LB, He Y, Brady ST, Kordower JH (2012) Alterations in axonal transport motor proteins in sporadic and experimental Parkinson's disease. Brain 135:2058-2073.

Chui HC, Mortimer JA, Slager U, Zarow C, Bondareff W, Webster DD (1986) Pathologic
correlates of dementia in Parkinson's disease. Arch Neurol 43:991-995.
Cubells JF, Schroeder JP, Barrie ES, Manvich DF, Sadee W, Berg T, Mercer K, Stowe TA, Liles LC,
Squires KE, Mezher A, Curtin P, Perdomo DL, Szot P, Weinshenker D (2016) Human
Bacterial Artificial Chromosome (BAC) Transgenesis Fully Rescues Noradrenergic
Function in Dopamine beta-Hydroxylase Knockout Mice. PLoS One 11:e0154864.
Damier P, Hirsch EC, Zhang P, Agid Y, Javoy-Agid F (1993) Glutathione peroxidase, glial cells and
Parkinson's disease. Neuroscience 52:1-6.
de Sousa Rodrigues ME, Bekhbat M, Houser MC, Chang J, Walker DI, Jones DP, Oller do
Nascimento CMP, Barnum CJ, Tansey MG (2017) Chronic psychological stress and high-
fat high-fructose diet disrupt metabolic and inflammatory gene networks in the brain,
liver, and gut and promote behavioral deficits in mice. Brain Behav Immun 59:158-172.
Delenclos M, Faroqi AH, Yue M, Kurti A, Castanedes-Casey M, Rousseau L, Phillips V, Dickson
DW, Fryer JD, McLean PJ (2017) Neonatal AAV delivery of alpha-synuclein induces
pathology in the adult mouse brain. Acta Neuropathol Commun 5:51.
den Hartog JW, Bethlem J (1960) The distribution of Lewy bodies in the central and autonomic
nervous systems in idiopathic paralysis agitans. J Neurol Neurosurg Psychiatry 23:283-
290.
Devauges V, Sara SJ (1991) Memory retrieval enhancement by locus coeruleus stimulation:
evidence for mediation by beta-receptors. Behav Brain Res 43:93-97.
Duffy S, MacVicar BA (1995) Adrenergic calcium signaling in astrocyte networks within the
hippocampal slice. J Neurosci 15:5535-5550.
Eng LF, Ghirnikar RS (1994) GFAP and astrogliosis. Brain Pathol 4:229-237.
Fahrig T (1993) Receptor subtype involved and mechanism of norepinephrine-induced
stimulation of glutamate uptake into primary cultures of rat brain astrocytes. Glia 7:212-
218.
Fearnley JM, Lees AJ (1991) Ageing and Parkinson's disease: substantia nigra regional
selectivity. Brain 114 (Pt 5):2283-2301.
Ferese R, Modugno N, Campopiano R, Santilli M, Zampatti S, Giardina E, Nardone A, Postorivo
D, Fornai F, Novelli G, Romoli E, Ruggieri S, Gambardella S (2015) Four Copies of SNCA
Responsible for Autosomal Dominant Parkinson's Disease in Two Italian Siblings.
Parkinsons Dis 2015:546462.
Fleming SM, Salcedo J, Fernagut PO, Rockenstein E, Masliah E, Levine MS, Chesselet MF (2004)
Early and progressive sensorimotor anomalies in mice overexpressing wild-type human
alpha-synuclein. J Neurosci 24:9434-9440.
Fujita H, Tanaka J, Maeda N, Sakanaka M (1998) Adrenergic agonists suppress the proliferation
of microglia through beta 2-adrenergic receptor. Neurosci Lett 242:37-40.
Fujiwara H, Hasegawa M, Dohmae N, Kawashima A, Masliah E, Goldberg MS, Shen J, Takio K,
Iwatsubo T (2002) alpha-Synuclein is phosphorylated in synucleinopathy lesions. Nat Cell
Biol 4:160-164.
Gerhard A, Pavese N, Hotton G, Turkheimer F, Es M, Hammers A, Eggert K, Oertel W, Banati RB,
Brooks DJ (2006) In vivo imaging of microglial activation with [11C](R)-PK11195 PET in
idiopathic Parkinson's disease. Neurobiol Dis 21:404-412.

German DC, Manaye KF, White CL, 3rd, Woodward DJ, McIntire DD, Smith WK, Kalaria RN, Mann DM (1992) Disease-specific patterns of locus coeruleus cell loss. Ann Neurol 32:667-676.

Giasson BI, Duda JE, Quinn SM, Zhang B, Trojanowski JQ, Lee VM (2002) Neuronal alpha-synucleinopathy with severe movement disorder in mice expressing A53T human alpha-synuclein. Neuron 34:521-533.

Gonera EG, van't Hof M, Berger HJ, van Weel C, Horstink MW (1997) Symptoms and duration of the prodromal phase in Parkinson's disease. Mov Disord 12:871-876.

Graham DR, Sidhu A (2010) Mice expressing the A53T mutant form of human alpha-synuclein exhibit hyperactivity and reduced anxiety-like behavior. J Neurosci Res 88:1777-1783.

Grenhoff J, Nisell M, Ferre S, Aston-Jones G, Svensson TH (1993) Noradrenergic modulation of midbrain dopamine cell firing elicited by stimulation of the locus coeruleus in the rat. J Neural Transm Gen Sect 93:11-25.

Gyoneva S, Traynelis SF (2013) Norepinephrine modulates the motility of resting and activated microglia via different adrenergic receptors. J Biol Chem 288:15291-15302.

Halliday GM, Li YW, Blumbergs PC, Joh TH, Cotton RG, Howe PR, Blessing WW, Geffen LB (1990) Neuropathology of immunohistochemically identified brainstem neurons in Parkinson's disease. Ann Neurol 27:373-385.

Hansen C, Bjorklund T, Petit GH, Lundblad M, Murmu RP, Brundin P, Li JY (2013) A novel alpha-synuclein-GFP mouse model displays progressive motor impairment, olfactory dysfunction and accumulation of alpha-synuclein-GFP. Neurobiol Dis 56:145-155.

Haycock JW (1993) Multiple signaling pathways in bovine chromaffin cells regulate tyrosine hydroxylase phosphorylation at Ser19, Ser31, and Ser40. Neurochem Res 18:15-26.

Heneka MT, Nadrigny F, Regen T, Martinez-Hernandez A, Dumitrescu-Ozimek L, Terwel D, Jardanhazi-Kurutz D, Walter J, Kirchhoff F, Hanisch UK, Kummer MP (2010) Locus ceruleus controls Alzheimer's disease pathology by modulating microglial functions through norepinephrine. Proc Natl Acad Sci U S A 107:6058-6063.

Henrich MT, Geibl FF, Lee B, Chiu WH, Koprich JB, Brotchie JM, Timmermann L, Decher N, Matschke LA, Oertel WH (2018) A53T-alpha-synuclein overexpression in murine locus coeruleus induces Parkinson's disease-like pathology in neurons and glia. Acta Neuropathol Commun 6:39.

Hirsch E, Graybiel AM, Agid YA (1988) Melanized dopaminergic neurons are differentially susceptible to degeneration in Parkinson's disease. Nature 334:345-348.

Hobson JA, McCarley RW, Wyzinski PW (1975) Sleep cycle oscillation: reciprocal discharge by two brainstem neuronal groups. Science 189:55-58.

Hornykiewicz O (1998) Biochemical aspects of Parkinson's disease. Neurology 51:S2-9.

Hunsley MS, Palmiter RD (2004) Altered sleep latency and arousal regulation in mice lacking norepinephrine. Pharmacol Biochem Behav 78:765-773.

Ip CW, Klaus LC, Karikari AA, Visanji NP, Brotchie JM, Lang AE, Volkmann J, Koprich JB (2017) AAV1/2-induced overexpression of A53T-alpha-synuclein in the substantia nigra results in degeneration of the nigrostriatal system with Lewy-like pathology and motor impairment: a new mouse model for Parkinson's disease. Acta Neuropathol Commun 5:11.

Iversen LL, Rossor MN, Reynolds GP, Hills R, Roth M, Mountjoy CQ, Foote SL, Morrison JH,
Bloom FE (1983) Loss of pigmented dopamine-beta-hydroxylase positive cells from locus
coeruleus in senile dementia of Alzheimer's type. Neurosci Lett 39:95-100.
Johnson JD, Zimomra ZR, Stewart LT (2013) Beta-adrenergic receptor activation primes
microglia cytokine production. J Neuroimmunol 254:161-164.
Kalaitzakis ME, Gentleman SM, Pearce RK (2013) Disturbed sleep in Parkinson's disease:
anatomical and pathological correlates. Neuropathol Appl Neurobiol 39:644-653.
Keren NI, Taheri S, Vazey EM, Morgan PS, Granholm AC, Aston-Jones GS, Eckert MA (2015)
Histologic validation of locus coeruleus MRI contrast in post-mortem tissue. Neuroimage
113:235-245.
Kilbourn MR, Sherman P, Abbott LC (1998) Reduced MPTP neurotoxicity in striatum of the
mutant mouse tottering. Synapse 30:205-210.
Kim YS, Joh TH (2006) Microglia, major player in the brain inflammation: their roles in the
pathogenesis of Parkinson's disease. Exp Mol Med 38:333-347.
Kirik D, Rosenblad C, Burger C, Lundberg C, Johansen TE, Muzyczka N, Mandel RJ, Bjorklund A
(2002) Parkinson-like neurodegeneration induced by targeted overexpression of alpha-
synuclein in the nigrostriatal system. J Neurosci 22:2780-2791.
Kopp N, Denoroy L, Tommasi M, Gay N, Chazot G, Renaud B (1982) Increase in noradrenaline-
synthesizing enzyme activity in medulla oblongata in Parkinson's disease. Acta
Neuropathol 56:17-21.
Koprich JB, Johnston TH, Reyes MG, Sun X, Brotchie JM (2010) Expression of human A53T alpha-
synuclein in the rat substantia nigra using a novel AAV1/2 vector produces a rapidly
evolving pathology with protein aggregation, dystrophic neurite architecture and
nigrostriatal degeneration with potential to model the pathology of Parkinson's disease.
Mol Neurodegener 5:43.
Kreiner G, Rafa-Zablocka K, Barut J, Chmielarz P, Kot M, Baginska M, Parlato R, Daniel WA,
Nalepa I (2019) Stimulation of noradrenergic transmission by reboxetine is beneficial for
a mouse model of progressive parkinsonism. Sci Rep 9:5262.
Kumer SC, Vrana KE (1996) Intricate regulation of tyrosine hydroxylase activity and gene
expression. J Neurochem 67:443-462.
Larsen KE, Schmitz Y, Troyer MD, Mosharov E, Dietrich P, Quazi AZ, Savalle M, Nemani V,
Chaudhry FA, Edwards RH, Stefanis L, Sulzer D (2006) Alpha-synuclein overexpression in
PC12 and chromaffin cells impairs catecholamine release by interfering with a late step
in exocytosis. J Neurosci 26:11915-11922.
Lategan AJ, Marien MR, Colpaert FC (1990) Effects of locus coeruleus lesions on the release of
endogenous dopamine in the rat nucleus accumbens and caudate nucleus as
determined by intracerebral microdialysis. Brain Res 523:134-138.
Leenders KL, Salmon EP, Tyrrell P, Perani D, Brooks DJ, Sager H, Jones T, Marsden CD,
Frackowiak RS (1990) The nigrostriatal dopaminergic system assessed in vivo by positron
emission tomography in healthy volunteer subjects and patients with Parkinson's
disease. Arch Neurol 47:1290-1298.
Levitt M, Spector S, Sjoerdsma A, Udenfriend S (1965) Elucidation of the Rate-Limiting Step in
Norepinephrine Biosynthesis in the Perfused Guinea-Pig Heart. J Pharmacol Exp Ther
148:1-8.

Lin CH, Lin JW, Liu YC, Chang CH, Wu RM (2015) Risk of Parkinson's disease following anxiety
disorders: a nationwide population-based cohort study. Eur J Neurol 22:1280-1287.

Logan T, Bendor J, Toupin C, Thorn K, Edwards RH (2017) alpha-Synuclein promotes dilation of
the exocytotic fusion pore. Nat Neurosci 20:681-689.

Mann JJ, Stanley M, Kaplan RD, Sweeney J, Neophytides A (1983) Central catecholamine
metabolism in vivo and the cognitive and motor deficits in Parkinson's disease. J Neurol
Neurosurg Psychiatry 46:905-910.

Maskri L, Zhu X, Fritzen S, Kuhn K, Ullmer C, Engels P, Andriske M, Stichel CC, Lubbert H (2004)
Influence of different promoters on the expression pattern of mutated human alpha-
synuclein in transgenic mice. Neurodegener Dis 1:255-265.

Masliah E, Rockenstein E, Veinbergs I, Mallory M, Hashimoto M, Takeda A, Sagara Y, Sisk A,
Mucke L (2000) Dopaminergic loss and inclusion body formation in alpha-synuclein
mice: implications for neurodegenerative disorders. Science 287:1265-1269.

Mavridis M, Degryse AD, Lategan AJ, Marien MR, Colpaert FC (1991) Effects of locus coeruleus
lesions on parkinsonian signs, striatal dopamine and substantia nigra cell loss after 1-
methyl-4-phenyl-1,2,3,6-tetrahydropyridine in monkeys: a possible role for the locus
coeruleus in the progression of Parkinson's disease. Neuroscience 41:507-523.

McCall JG, Al-Hasani R, Siuda ER, Hong DY, Norris AJ, Ford CP, Bruchas MR (2015) CRH
Engagement of the Locus Coeruleus Noradrenergic System Mediates Stress-Induced
Anxiety. Neuron 87:605-620.

McGeer PL, Itagaki S, Boyes BE, McGeer EG (1988) Reactive microglia are positive for HLA-DR in
the substantia nigra of Parkinson's and Alzheimer's disease brains. Neurology 38:1285-
1291.

Molinoff PB, Axelrod J (1971) Biochemistry of catecholamines. Annu Rev Biochem 40:465-500.

Murchison CF, Zhang XY, Zhang WP, Ouyang M, Lee A, Thomas SA (2004) A distinct role for
norepinephrine in memory retrieval. Cell 117:131-143.

Niu H, Shen L, Li T, Ren C, Ding S, Wang L, Zhang Z, Liu X, Zhang Q, Geng D, Wu X, Li H (2018)
Alpha-synuclein overexpression in the olfactory bulb initiates prodromal symptoms and
pathology of Parkinson's disease. Transl Neurodegener 7:25.

Norris JG, Benveniste EN (1993) Interleukin-6 production by astrocytes: induction by the
neurotransmitter norepinephrine. J Neuroimmunol 45:137-145.

Paukert M, Agarwal A, Cha J, Doze VA, Kang JU, Bergles DE (2014) Norepinephrine controls
astroglial responsiveness to local circuit activity. Neuron 82:1263-1270.

Pavese N, Rivero-Bosch M, Lewis SJ, Whone AL, Brooks DJ (2011) Progression of monoaminergic
dysfunction in Parkinson's disease: a longitudinal 18F-dopa PET study. Neuroimage
56:1463-1468.

Phillips RG, LeDoux JE (1992) Differential contribution of amygdala and hippocampus to cued
and contextual fear conditioning. Behav Neurosci 106:274-285.

Pifl C, Kish SJ, Hornykiewicz O (2012) Thalamic noradrenaline in Parkinson's disease: deficits
suggest role in motor and non-motor symptoms. Mov Disord 27:1618-1624.

Porter-Stransky KA, Centanni SW, Karne SL, Odil LM, Fekir S, Wong JC, Jerome C, Mitchell HA,
Escayg A, Pedersen NP, Winder DG, Mitrano DA, Weinshenker D (2019) Noradrenergic
Transmission at Alpha1-Adrenergic Receptors in the Ventral Periaqueductal Gray
Modulates Arousal. Biol Psychiatry 85:237-247.

Remy P, Doder M, Lees A, Turjanski N, Brooks D (2005) Depression in Parkinson's disease: loss
of dopamine and noradrenaline innervation in the limbic system. Brain 128:1314-1322.
Ritter S, Watts AG, Dinh TT, Sanchez-Watts G, Pedrow C (2003) Immunotoxin lesion of
hypothalamically projecting norepinephrine and epinephrine neurons differentially
affects circadian and stressor-stimulated corticosterone secretion. Endocrinology
144:1357-1367.
Rommelfanger KS, Weinshenker D (2007) Norepinephrine: The redheaded stepchild of
Parkinson's disease. Biochem Pharmacol 74:177-190.
Rommelfanger KS, Weinshenker D, Miller GW (2004) Reduced MPTP toxicity in noradrenaline
transporter knockout mice. J Neurochem 91:1116-1124.
Rommelfanger KS, Edwards GL, Freeman KG, Liles LC, Miller GW, Weinshenker D (2007)
Norepinephrine loss produces more profound motor deficits than MPTP treatment in
mice. Proc Natl Acad Sci U S A 104:13804-13809.
Rorabaugh JM, Chalermpalanupap T, Botz-Zapp CA, Fu VM, Lembeck NA, Cohen RM,
Weinshenker D (2017) Chemogenetic locus coeruleus activation restores reversal
learning in a rat model of Alzheimer's disease. Brain 140:3023-3038.
Ross GW, Petrovitch H, Abbott RD, Tanner CM, Popper J, Masaki K, Launer L, White LR (2008)
Association of olfactory dysfunction with risk for future Parkinson's disease. Ann Neurol
63:167-173.
Sawamoto K, Nakao N, Kobayashi K, Matsushita N, Takahashi H, Kakishita K, Yamamoto A,
Yoshizaki T, Terashima T, Murakami F, Itakura T, Okano H (2001) Visualization, direct
isolation, and transplantation of midbrain dopaminergic neurons. Proc Natl Acad Sci U S
A 98:6423-6428.
Schank JR, Ventura R, Puglisi-Allegra S, Alcaro A, Cole CD, Liles LC, Seeman P, Weinshenker D
(2006) Dopamine beta-hydroxylase knockout mice have alterations in dopamine
signaling and are hypersensitive to cocaine. Neuropsychopharmacology 31:2221-2230.
Schell H, Hasegawa T, Neumann M, Kahle PJ (2009) Nuclear and neuritic distribution of serine-
129 phosphorylated alpha-synuclein in transgenic mice. Neuroscience 160:796-804.
Singleton AB et al. (2003) alpha-Synuclein locus triplication causes Parkinson's disease. Science
302:841.
Song CH, Fan X, Exeter CJ, Hess EJ, Jinnah HA (2012) Functional analysis of dopaminergic
systems in a DYT1 knock-in mouse model of dystonia. Neurobiol Dis 48:66-78.
Song S, Jiang L, Oyarzabal EA, Wilson B, Li Z, Shih YI, Wang Q, Hong JS (2018) Loss of Brain
Norepinephrine Elicits Neuroinflammation-Mediated Oxidative Injury and Selective
Caudo-Rostral Neurodegeneration. Mol Neurobiol.
Sotiriou E, Vassilatis DK, Vila M, Stefanis L (2010) Selective noradrenergic vulnerability in alpha-
synuclein transgenic mice. Neurobiol Aging 31:2103-2114.
Spillantini MG, Schmidt ML, Lee VM, Trojanowski JQ, Jakes R, Goedert M (1997) Alpha-synuclein
in Lewy bodies. Nature 388:839-840.
Srinivasan J, Schmidt WJ (2003) Potentiation of parkinsonian symptoms by depletion of locus
coeruleus noradrenaline in 6-hydroxydopamine-induced partial degeneration of
substantia nigra in rats. Eur J Neurosci 17:2586-2592.
Szot P, Franklin A, Miguelez C, Wang Y, Vidaurrazaga I, Ugedo L, Sikkema C, Wilkinson CW,
Raskind MA (2016) Depressive-like behavior observed with a minimal loss of locus

coeruleus (LC) neurons following administration of 6-hydroxydopamine is associated
with electrophysiological changes and reversed with precursors of norepinephrine.
Neuropharmacology 101:76-86.
Tansey MG, Goldberg MS (2010) Neuroinflammation in Parkinson's disease: its role in neuronal
death and implications for therapeutic intervention. Neurobiol Dis 37:510-518.
Tong J, Hornykiewicz O, Kish SJ (2006) Inverse relationship between brain noradrenaline level
and dopamine loss in Parkinson disease: a possible neuroprotective role for
noradrenaline. Arch Neurol 63:1724-1728.
Valentino RJ, Van Bockstaele E (2008) Convergent regulation of locus coeruleus activity as an
adaptive response to stress. Eur J Pharmacol 583:194-203.
Vazey EM, Aston-Jones G (2014) Designer receptor manipulations reveal a role of the locus
coeruleus noradrenergic system in isoflurane general anesthesia. Proc Natl Acad Sci U S
A 111:3859-3864.
Wakamatsu M, Ishii A, Ukai Y, Sakagami J, Iwata S, Ono M, Matsumoto K, Nakamura A, Tada N,
Kobayashi K, Iwatsubo T, Yoshimoto M (2007) Accumulation of phosphorylated alpha-
synuclein in dopaminergic neurons of transgenic mice that express human alpha-
synuclein. J Neurosci Res 85:1819-1825.
Weinshenker D (2018) Long Road to Ruin: Noradrenergic Dysfunction in Neurodegenerative
Disease. Trends Neurosci 41:211-223.
Yao N, Wu Y, Zhou Y, Ju L, Liu Y, Ju R, Duan D, Xu Q (2015) Lesion of the locus coeruleus
aggravates dopaminergic neuron degeneration by modulating microglial function in
mouse models of Parkinsons disease. Brain Res 1625:255-274.
Zarow C, Lyness SA, Mortimer JA, Chui HC (2003) Neuronal loss is greater in the locus coeruleus
than nucleus basalis and substantia nigra in Alzheimer and Parkinson diseases. Arch
Neurol 60:337-341.
852
853

856
857
858 **Figure 1:** *hSNCA* cDNA open reading frame was targeted to the translational start
859 site of *DBH* by standard BAC recombineering methods

887
888 **Figure 2: Human *Snca* mRNA is expressed in DBH-hSNCA locus coeruleus**
889 **neurons.** Human *Snca* mRNA (hSnca; white) expression is detectable only in transgenic
890 (Tg: bottom row) LC neurons, where it co-localizes with mouse *Th* mRNA (mTh, green),
891 and mouse *Dbh* mRNA (mDbh; red) using RNA Scope Fluorescent Multiplex v2 assay.
892 Scale bar = 50µm.

921
922

923
924
Figure 3: Analysis of asyn expression in locus coeruleus of young DBH-hSNCA
mice. *A,* Immunofluorescent detection of human asyn (red) with a species-specific
antibody (Biolegend 807801) demonstrates co-localization with TH-expressing LC
neurons (green) in Tg mice (lower panel) but not in nTg mice (upper panel), or in ***C,*** TH-
expressing neurons (green) in the substantia nigra regardless of genotype. ***B,*** Human
asyn protein is expressed selectively in LC neurons of Tg mice by western blot. ***D,***
Immunoblot of LC protein with an antibody against asyn that detects the mouse and
human protein reveals a significant increase in total asyn protein expression in Tg LC
neurons as compared to that in LC of nTg littermate mice. Immunoblot data graphed as
arbitrary units (AU) normalized to total protein. All data are from 3-mo old nTg and Tg
mice. Scale bar = 50µm. Student's t-test ± SEM *$p<0.05$.

954
955

956
957

Figure 4: DBH-hSNCA mice display age-dependent behavioral abnormalities similar to the non-motor symptoms of PD. *A,* 3-mo and 14-mo Tg mice have increased sleep latency. *B,* Tg mice display anxiety-like behavior by marble-burying test at 14-mos and 24-mos and by *C,* latency to re-enter the center in the open field test at 14-mos. *D,* There is a significant effect of genotype at 14-mos with Tg mice exhibiting significantly more percent time freezing during fear training (center panel; bars represent tone presentation, bolt represents foot shock) and *E,* contextual learning**.** *F,* There was no effect of genotype in percent time freezing during cued test at any age. *G,* Ambulations do not differ by genotype at any age Students t-test of genotype ± SEM in A, B & C. Repeated measures ANOVA ± SEM in D, E, F, and G. *$p<0.05$, ***$p<0.001$.

969
970
971

972
973
974 **Figure 5: DBH-hSNCA LC neurons contain small asyn aggregates.** **A,** At 14-mos of
975 age, LC neurons in (TH; green) Tg mice contain significantly more oligomerized asyn
976 (red) in LC neurons as determined by proximity ligation assay (PLA). **B,** Representative
977 images. Scale bar = 15µm. Student's t-test ± SEM. *$p<0.05$.

986
987
Figure 6: Human asyn expression in LC neurons impacts striatal dopamine
metabolism in aged DBH-hSNCA mice. Catecholamine and catecholamine metabolite
tissue content from hippocampus and striatum was measured by HPLC. ***A,*** Hippocampal
NE is not affected by genotype. ***B,*** Striatal NE is reduced at 24-mos but does not reach
statistical significance. The NE metabolite MHPG to NE ratio does not differ at any age in
the ***C,*** hippocampus, or ***D,*** striatum. Dopamine content is not affected at any age in ***E,***
hippocampus, or ***F,*** striatum. ***G,*** The ratio of the DA metabolite DOPAC to DA is
unaffected in the hippocampus ***H,*** but is significantly reduced in the striatum of Tg mice
at 24-mos. Student's t-test of genotype for each age group ± SEM. *$p<0.05$.

997
998
999

1000
1001

1002
1003
1004
1005
1006 **Figure 7: Human asyn expression does not affect LC integrity.** NET-expressing LC
1007 cell bodies (red) percent immunoreactivity (IR) does not differ by genotype at 3-, 14-, or
1008 24-mos. Student's t-test of mean IR by genotype for each age group, *A,* graphed as fold
1009 change Tg from nTg mean ± SEM. *B,* Representative images. Scale bar = 50μm, *p<0.05.

1023
1024

1025
Figure 8: Tyrosine hydroxylase levels are elevated in young DBH-hSNCA Tg mice.
Tyrosine hydroxylase (TH: green) immunoreactivity is increased in LC neurons (NET: red)
at 3- and 14-mos of age. *A*, Representative immunofluorescent images. *B*, TH IR mean
normalized to NET IR mean and expressed as fold change Tg from nTg mean % IR ±
SEM. *C,* Western blot analysis of 3-mo old LC neurons confirms increased TH
abundance. Data graphed as arbitrary densitometric units (AU) was normalized to total
protein ± SEM. *D,* Representative western blot images of TH (top) and total protein stain
(bottom). Student's t-test of nTg and Tg at each age, n=5-10. Scale bar = 50µm. *p<0.05.
1034
1035
1036

1037

Fig 9. Human asyn in LC neurons affects expression of local inflammatory markers
There are fewer Iba1-positive cells (red) in the LC (TH: green) at 14-mos. **A**,
Representative immunofluorescent images. **B**, Microglial count in the LC graphed as fold
change Tg from nTg Iba1+ cell count ± SEM. Student's t-test of nTg and Tg for each age
group. Expression of astrocytic GFAP (red) is increased in the Tg LC (TH: green) at 24-
mos of age. **C**, Representative immunofluorescent images. **D**, Quantification of GFAP %
IR in the LC as percent ROI graphed as fold change Tg IR from nTg mean IR ± SEM.
Student's t-test of nTg and Tg for each age group. Scale bar = 50µm. *$p<0.05$.

1062
1063

1064 **Fig 10: Hippocampal astrocytic GFAP expression is increased at 14-mos**. At 14-
1065 mos, Tg mice have significantly more GFAP (red) expression than nTg in the CA1 region
1066 of the hippocampus. Representative immunofluorescent images of *A,* CA1, *C,* dentate
1067 gyrus, and *E,* CA3 regions. Quantification of GFAP IR as % area of ROI in *B,* CA1, *D,*
1068 dentate gyrus, and *F,* CA3. Student's t-test of nTg and Tg for each age group, graphed
1069 as fold change Tg from nTg mean ± SEM. Scale bar = 50µm. *$p<0.05$.

1070
1071
1072
1073

1074
1075
Fig 11: Fewer Iba1-expressing cells in hippocampal CA1 region in 14-mo old Tg
DBH-hSNCA mice. At 14-mos, Tg mice have fewer Iba1-expressing cells in hippocampal
region CA1. **A,** Representative immunofluorescent images of Iba1-expressing cells (red).
B, Quantification of Iba1-expressing cells in CA1 graphed as Tg fold change from nTg
mean ± SEM. Student's t-test by genotype for each age group. Nuclear stain in blue.
Scale bar = 150μm. *p<0.05.

Figure 12: Asyn expression in LC neurons results in age-dependent degeneration of hippocampal LC fibers. NET (red) IR is reduced in the dentate gyrus of Tg mice at 24-mos. Representative immunofluorescent images of **A**, CA1, **C**, dentate gyrus, and **E**, CA3 regions. Quantification of NET IR as percent area of ROI in **B**, CA1, **D**, dentate gyrus, and **F**, CA3. Student's t-test of nTg and Tg for each age group, graphed as fold change Tg from nTg mean ± SEM. Scale bar = 150μm. *p<0.05.

Figure 13: Asyn can directly affect neurotransmission. Asyn could enhance extracellular NE by several mechanisms. #1 – Increased expression of asyn can potentiate clustering of synaptic vesicles at the presynaptic membrane. #2 – Elevated asyn expression can reduce norepinephrine transporter (NET) expression at the plasma membrane. #3 – Asyn modulation of L-type voltage gated calcium channels can enhance neuronal excitability. #4 – Human asyn expression can enlarge the size of, and slow the closing of vesicular fusion pores, allowing more NE to spill into the extracellular space.

Table 1: Primary and secondary antibodies used in these studies

Immunofluorescence

Antigen	Host	Manufacturer	Cat. No.	Dilution
α-synuclein (Hu)	Mouse	Biolegend	807801	1:500
DAT	Rat	Millipore	MAB369	1:500
GFAP	Rabbit	Zymed	13-0300	1:500
Iba1	Rabbit	Wako	019-19741	1:500
NET	Mouse	Mab Technologies	NET05-2	1:1,000
Tyrosine hydroxylase	Rabbit	Millipore	AB152	1:1,000
Tyrosine hydroxylase	Chicken	Abcam	AB76442	1:1,000
VMAT2	Rabbit	*R.A. Cliburn et al, 2017*		1:500
Anti-chicken AlexaFluor488	Goat	Invitrogen	A11039	1:1,000
Anti-rabbit AlexaFluor488	Goat	Invitrogen	A11015	1:1,000
Anti-mouse AlexaFluor594	Goat	Invitrogen	A11020	1:1,000
Anti-rabbit AlexaFluor594	Goat	Invitrogen	A11012	1:1,000
Anti-rat AlexaFluor594	Goat	Invitrogen	A11007	1:1,000

Proximity ligation assay	Host	Manufacturer	Cat. No.	Dilution
α-synuclein (Hu)	Mouse	Abcam	AB80627	1:1000
α-synuclein- pSer129	Mouse	Biolegend	MMS-5091	1:1000
Syn-F1	Mouse	Biolegend	847802	1:1,000

Immunoblotting

Antigen	Host	Manufacturer	Cat. No.	Dilution
α-synuclein (Hu)	Mouse	Abcam	AB80627	1:1,000
α-synuclein (Ms + Hu)	Rabbit	Santa Cruz	SC-7011R	1:500
Tyrosine hydroxylase	Rabbit	Millipore	AB152	1:1,000
Anti-mouse HRP	Goat	Jackson ImmunoResearch	115-036-072	1:1,000
Anti-rabbit HRP	Goat	Jackson ImmunoResearch	111-036-047	1:1,000

1120
1121