A common pattern of influenza A virus single cell gene expression heterogeneity governs the innate antiviral response to infection

J. Cristobal Vera[1,2]*, Jiayi Sun[1]*, Yen Ting Lin[3], Jenny Drnevich[4], Ruian Ke[3], and Christopher B. Brooke[1,2]

[1] Department of Microbiology, University of Illinois at Urbana-Champaign, Urbana, IL.
[2] Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, IL.
[3] T-6, Theoretical Biology and Biophysics, Los Alamos National Laboratory, Los Alamos, NM.
[4] High-Performance Biological Computing at the Roy J. Carver Biotechnology Center, University of Illinois at Urbana-Champaign, Urbana, IL.

* These authors contributed equally

Corresponding author: Christopher Brooke (cbrooke@illinois.edu)

ABSTRACT

Viral infection outcomes are governed by the complex and dynamic interplay between the infecting virus population and the host response. It is increasingly clear that both viral and host cell populations are highly heterogeneous, but little is known about how this heterogeneity influences infection dynamics or viral pathogenicity. To dissect the interactions between influenza A virus (IAV) and host cell heterogeneity, we examined the combined host and viral transcriptomes of thousands of individual, single virion-infected cells. We observed complex, heterogeneous patterns of viral gene expression and the existence of multiple distinct host transcriptional responses to infection at the single cell level within populations of infected cells. Our analyses reveal that viral NS segment gene expression diverges from that of the rest of the viral genome within a subset of infected cells, and that this unique pattern of NS segment expression plays a dominant role in shaping the host cell response to infection. Finally, we show that seasonal human H1N1 and H3N2 strains differ significantly in patterns of host anti-viral gene transcriptional heterogeneity at the single cell level. Altogether, these data reveal a common pattern of viral gene expression heterogeneity across human IAV subtypes that serves as a major determinant of antiviral gene activation.

INTRODUCTION

RNA virus populations typically contain an enormous amount of sequence diversity due to an absence of virally encoded proofreading activity (1). In some cases, this genetic diversity can significantly influence infection outcomes (2–4)(4). In addition to genetic diversity, influenza A viruses (IAV) also exhibit substantial heterogeneity in the gene coding capabilities of individual virions. Most IAV virions are only capable of expressing variable, incomplete subsets of viral genes (5,6). The production and gene expression patterns of these semi-infectious particles (SIPs) can vary significantly between IAV strains (7). Altogether, the high degree of diversity within IAV populations means that patterns of viral gene expression can vary significantly between individual infected cells.

Numerous studies have leveraged recent advances in single cell analysis methods to assess the extent of cellular heterogeneity present during infection by a variety of viruses, including IAV (8–17). These studies connect to a larger body of work that has begun to explore single cell

heterogeneity within different host cell populations and tissues (18–20). To date, efforts to
characterize single cell heterogeneity during IAV infection have been hampered by limitations in
the number of cells analyzed or experimental design choices that complicated analysis. The
extent to which IAV population diversity influences patterns of single cell heterogeneity and the
ways in which these patterns shape broader infection dynamics and outcomes remain poorly
understood.
55
Here, we query the combined viral and host transcriptomes from thousands of individual cells,
each infected by a single virion. This high-resolution dissection of viral and host gene expression
patterns reveals that the transcriptional responses of individual infected cells can be highly
divergent, resulting from the interplay between underlying cellular and viral heterogeneity. Thus,
the host response to IAV infection consists of a heterogenous assemblage of highly variable
single cell responses. This approach also reveals critical differences in the interactions of H1N1
and H3N2 viruses with the host antiviral machinery at the single cell level and implicates variability
in NS segment expression as a major driver of the innate antiviral response to IAV infection.
Altogether, these results establish the interplay between viral and host cell heterogeneity as a
critical determinant of cellular infection outcomes.
66
RESULTS
Generation of viral and host transcriptional data from thousands of singly infected cells
To assess the effects of viral population heterogeneity on the host transcriptional response to
infection, we examined the combined viral and host transcriptional profiles from thousands of
single infected cells. To focus on the effects of viral heterogeneity in shaping the host response
to infection, we wanted to remove the variability that could arise from variation in cellular MOI
(21). To ensure that >99.99% of infected cells were each infected with a single virion, we infected
A549 cells with the 2009 H1N1 pandemic strain A/California/07/2009 (Cal07) at an MOI of 0.01,
and blocked secondary spread in the culture through the addition of NH4Cl (22). This resulted in
only a tiny fraction of cells being infected, so to obtain sufficient numbers of infected cells for our
analysis, we enriched infected cells by sorting based on surface expression of HA and/or M2 (**Fig.
1A**). This approach had the added benefit of allowing us to sort HA-M2- bystander cells from the
same culture to assess the paracrine effects of infection. In parallel, we sorted a group of mock-
infected cells as a control.
81
We used the 10X Genomics Chromium platform to generate oligo-dT-primed single cell RNAseq
libraries for mock, infected (HA+ and/or M2+), and bystander (HA-M2-) cells and sequenced them
on an Illumina Novaseq. We demultiplexed the reads and mapped them to a customized hybrid
reference containing both human and influenza sequences/annotation. Following quality control
filtering, we had high quality combined viral and host transcriptomes from 4662 single virion-
infected cells, as well as 3610 bystander cells, and 4597 mock cells. Clustering of these three
libraries by transcriptional similarity revealed that mock, bystander, and infected cells largely
clustered independently, as would be expected, though a subset of infected cells grouped closely
with the bystander cells (**Fig. 1B**).
91

Fig. 1. Generation of viral and host transcriptional data for thousands of singly infected cells. (A) Schematic depicting our strategy for generating single cell RNAseq libraries from thousands of cells infected at low MOI. In brief, we infect A549 cells with Cal07 at MOI=0.01 to ensure that infected cells are infected with a single virion. We then block secondary spread with NH4Cl treatment to make sure infection timing is uniform across all infected cells. Finally, we sort "infected" and "bystander" cells based on surface expression of HA and/or M2 and immediately generate single cell RNAseq libraries from these sorted cell populations using the 10X Chromium device. In parallel, mock cells are sorted and used as uninfected controls. **(B)** tSNE dimensionality reduction plot showing the extent of overlap of the indicated cell populations clustered based on transcriptional similarity.

92
93
Enormous single cell heterogeneity in viral gene expression patterns
We first asked whether we observed the same degree of heterogeneity in viral gene expression
between infected cells that has been reported previously (6,10,12). We calculated the fraction of
all transcripts within each cell that were viral in origin (**Fig. 2A**). Similar to prior studies, total viral
transcript levels ranged enormously between individual cells: from <1% up to ~90%. Notably, this
heterogeneity arose under conditions where both viral input (1 virion/cell) and infection timing
were equivalent across all cells.
101
This pattern of extreme cell-to-cell heterogeneity was also observed for individual viral transcripts,
again consistent with previous reports (**Fig. 2B**)(12). Expression distributions for individual viral
genes were highly over-dispersed compared with a Poisson distribution (**Fig 2C**). The tails of the
viral gene expression distributions were orders of magnitude longer than those of the top 10 most
abundant host transcripts (**Fig. 2D,E**), emphasizing that viral gene expression is substantially
noisier than host gene expression.
108

Fig. 2. Enormous heterogeneity in viral gene expression patterns. (A) Plot shows the distribution of Cal07-infected A549 cells, binned by the fraction of total cellular poly(A) RNA that is viral in origin. **(B)** Plots show the fraction of total poly(A) RNA per cell that maps to the indicated viral gene segment. Each dot represents a single cell, cells with no detectable reads mapping to the indicated segment were arbitrarily assigned a value of 0.000007 to show up on the log10 scale. **(C)** Distribution of normalized PB2-derived reads per cell (orange) compared with a Poisson distribution of equal mean (blue line) on a log-log scale. **(D)** The viral gene expression distributions normalized by the 70th quantile on a log-log scale. **(E)** Same analysis as in (D) but showing the top 10 most abundant host transcripts.

Significant heterogeneity in the host transcriptional response to infection

We next asked whether the observed variation in viral gene expression levels was associated with variation in host cell transcription. To test this, we first clustered all infected cells based on host transcription patterns. We identified multiple distinct transcriptional response groups to infection by Cal07, suggesting that there is not one standard transcriptional response to IAV infection and that a single cell type can simultaneously generate multiple distinct responses to the same virus population (**Fig. 3A,B**). It should be noted that these cluster definitions are not absolute and that substantial heterogeneity in the normalized expression levels of individual host genes existed within individual clusters.

We did not observe any clear associations between clustering pattern and total viral transcript levels (**Fig. 3C**); however, the clustering pattern did appear to be partially influenced by cell cycle status (**Fig. 3D**). Cells in G2M phase were primarily found within a distinct subset of clusters (1, 3, and 6) while cells in G1 and S phases exhibited no clear clustering patterns. Consistent with this, the expression levels of a number of host cell cycle-associated genes (*e.g.* CDK1, CENPF, MKI67, PTTG1, and CKS2) were significantly elevated in those same clusters (**Table S1**). While it is not surprising that cell cycle status would contribute to transcriptional heterogeneity during infection, these data highlight how little is known about how cell cycle status may influence the cellular response to infection.

Most of the observed heterogeneity could not be simply explained by cell cycle, however. The induction of numerous host genes known or likely to be involved in shaping IAV infection outcome varied significantly between clusters. One notable example is TRIM41, a protein previously shown

to inhibit IAV replication by marking viral NP for degradation (23). TRIM41 is clearly upregulated
in response to infection at the bulk level (**Fig. 3E**). At the single cell level, both the frequencies
and the overall expression levels of this transcript differed significantly between infected cell
clusters (**Fig. 3F,G**). We also observed substantial variation between clusters in upregulation of
the long non-coding RNA NEAT1, which is involved in inflammasome formation, regulation of
cytokine and chemokine expression, and nuclear paraspeckle formation (24–29). NEAT1
expression was upregulated during infection at the bulk level (**Fig. 3H**), but levels were
significantly higher in clusters 0 and 4 compared with other infected cells (adjusted p values =
$2.01*10^{-12}$ and $2.91*10^{-112}$ respectively) (**Fig. 3I,J**). Thousands of other host genes exhibited
similarly heterogeneous patterns of expression between individual infected cells, illustrating how
key drivers of infection outcomes may be primarily expressed by limited subsets of infected cells.
145

Fig. 3. Significant heterogeneity in the host transcriptional response to infection. (A) tSNE dimensionality reduction plot showing all Cal07-infected A549 cells clustered based on similarity of host transcriptional patterns. **(B)** Heat map showing differential expression of the top 10 characteristic host genes for each cluster from (A). Individual cells are each represented by a column, grouped by cluster, with individual rows representing relative expression of the top 10 specific host transcripts most significantly associated with each cluster. **(C)** Same tSNE plot of Cal07-infected cells shown in (A) with each cell colored by the proportion of total cellular poly(A) RNA that is viral in origin. **(D)** Same tSNE plot of Cal07-infected cells shown in (A) with each cell colored by predicted cell cycle stage, as determined by the scran package. **(E)** Comparison of normalized per cell counts of TRIM41 between mock and infected cells. **(F)** Fraction of cells in each infected cell cluster from (A) with detectable levels of TRIM41. **(G)** Comparison of normalized per cell counts of TRIM41 between clusters of infected cells shown in (A). **(H)** Comparison of normalized per cell counts of NEAT1 between mock and infected cells. **(F)** Fraction of cells in each infected cell cluster from (A) with detectable levels of NEAT1. **(G)** Comparison of normalized per cell counts of NEAT1 between clusters of infected cells shown in (A). **(K)** tSNE dimensionality reduction plot showing mock A549 cells clustered based on similarity of host transcriptional patterns. **(L)** Principle coordinate axes (PCoA) plot comparing the multivariate dispersions for mock (black) and Cal07-infected (red) A549 cells. The first two axes (PCoA 1 and PCoA 2) in the multivariate homogeneity of group dispersions analysis are used.

146
147

We also observed the existence of several distinct clusters within mock cells (**Fig. 3K**), leading
us to ask whether the heterogeneity we observed in infected cells was simply a reflection of
underlying intrinsic heterogeneity of the cell population prior to infection. In other words, does
infection actually increase the overall heterogeneity in host gene expression patterns beyond that
seen in mock cells? To quantify and compare heterogeneity in overall host transcription between
the two cell populations, we calculated the multivariate homogeneity of dispersions for mock and
infected cells using expression of all host genes (**Fig. 3L**)(30). We found that infection significantly
increased the overall single cell heterogeneity in host cell transcription patterns ($p<10^{-15}$ by
pairwise t test). Altogether, our data is consistent with a model in which the interaction between
viral population heterogeneity and pre-existing host cell heterogeneity gives rise to multiple
distinct transcriptional responses at the single cell level.

159
Many infected cells have undetectable levels of one or more viral transcripts
The vast majority of IAV virions fail to express one or more viral genes, resulting in the expression
of variable, incomplete subsets of viral gene products under low MOI conditions (5–7,31). We
next asked whether this variation in functional viral gene content within individual cells contributes
to the observed heterogeneity in host gene transcription. To assess the presence or absence of
each individual viral gene within infected cells while avoiding false positives due to RNA cross-
contamination, we took advantage of the fact that the vast majority of "bystander" cells in our
analysis were not infected but were collected from the same culture vessel as our infected cells.
We used these "true negative" cells to empirically determine the distribution of viral read counts
that could arise from mRNA cross-contamination during infection and drop-making (12). We then
used these values to generate cutoff thresholds that could be used to differentiate true positives
from false positives.
172

Fig. 4. Many infected cells have undetectable levels of one or more viral transcripts. (A) The fraction of all Cal07-infected cells that have detectable levels of transcripts derived from the indicated viral gene segments. **(B)** All Cal07-infected cells binned by the total number of detectable viral gene segments, with the actual numbers of cells in each group detailed above.

173
174
Using these cutoffs, we found that the fractions of infected cells that failed to express detectable
transcripts ranged from ~35% to nearly zero for the individual viral gene segments (**Fig. 4A**). It is
highly likely that our method of enriching for infected cells by sorting based on high level HA
and/or M2 expression significantly biased these results. Regardless, we still observe substantial
numbers of infected cells that fail to express detectable levels of individual viral transcripts, with
roughly half of all infected cells failing to express detectable levels of at least one viral gene (**Fig.
4B**). We were surprised to see so many cells that lacked detectable levels of the polymerase
transcripts because we expected that sorting infected cells based on surface protein expression
would bias cell collection against cells that could not synthesize new polymerase complexes. One
caveat with this approach is that we assessed transcript levels at 16 hpi and thus could have
missed transient early expression of viral genes.
186
**Dissecting the effects of individual viral genes on the host transcriptional response to
infection**

We next asked how the expression of individual viral genes influences the overall host response
to infection. We grouped all Cal07-infected cells into positive and negative populations based on
the expression of each individual viral gene segment and used two distinct differential single cell
gene expression analysis methods (MAST and NBID) to compare host transcript expression
between the two infected cell populations (32,33). For each viral gene, we generated a list of host
transcripts that were reported as significantly different between positive and negative cells by both
NBID and MAST (**Table S2-S9**). This approach allowed us to tease out the effects of individual
viral gene segments from the more general effects of infection.
197
We identified hundreds of host genes with expression levels that varied significantly based on the
expression status of individual viral genes (**Fig. 5A**). The one exception was the M segment, most
likely due to the lack of sufficient cells lacking M transcript for the analysis. Closer examination
revealed that many of these hits were found for multiple viral genes, suggesting that they may
correlate with overall viral gene expression levels or co-expression of multiple viral gene
segments. When we focused on the host genes that only exhibited differential expression in
association with the expression status of a single viral gene segment, we found that most were
associated with the PA, NP, or NS segments (**Fig. 5B, Table S10**).
206
We focused on the NS segment, as the NS-encoded NS1 protein plays a well-established, multi-
functional role in manipulating the host cell environment and the anti-viral response and could
thus serve as a positive control for our approach (34–37). We found that ~10% of infected cells
had background or undetectable levels of NS segment transcript (**Fig. 5C**). We observed nearly
450 host genes whose expression was significantly correlated with NS expression status,
however only 28 of these were uniquely influenced by NS. Notably, this list included SLFN5 (down
in NS-negative infected cells; FDR=$3.22*10^{-42}$), an interferon-stimulated gene (ISG) shown to
negatively regulate STAT1-dependent anti-viral gene transcription (38). We observed that both
the frequency and normalized per-cell expression levels of SLFN5 were significantly decreased
in infected cells that failed to express NS compared with those that did express NS (**Fig. 5D,E**).
Notably, expression of specific IFNs or other ISGs was not significantly affected by NS expression
status, echoing the previous finding that lack of NS expression does not deterministically result in
increased expression of these genes (12,13). These results both validate the utility of our
approach and identify the suppressor of antiviral gene transcription SLFN5 as a novel host target
of the NS gene segment.
222

Fig. 5. Dissection of the effects of individual viral genes on the host transcriptional response to infection. (A) The number of host transcripts for which expression levels significantly differ depending on whether the indicated viral gene segment is present or not, according to both MAST and NBID. **(B)** Same as in (A) but removing host genes that are differentially regulated by the expression status of more than one viral gene segment. **(C)** tSNE plot of all Cal07-infected A549 cells colored based on whether NS segment-derived transcripts are detected (Cyan) or not detected (Salmon). **(D)** Fractions of NS- and NS+ Cal07-infected A549 cells with detectable levels of SLFN5. **(E)** Distribution of SLFN5 normalized counts in cells shown in (D).

223
224
H1N1 and H3N2 viruses share a common pattern of single cell viral gene expression
heterogeneity
Since productively infected cells will by definition encode all viral genes, we asked how variation
in viral gene expression within infected cells that express all viral genes affects the host
transcriptional response. To more carefully dissect the relationships between the expression
patterns of individual viral genes, we examined the degree to which individual pairs of viral genes
were correlated with each other (**Fig. 6A, S1**). For Cal07, we found that most pairs of viral
transcripts were highly correlated. In particular all pairwise comparisons of the polymerase
components exhibited r^2 values of 0.796 or higher. In contrast, the NS segment was not generally
positively correlated with the other segments (**Fig. 6B**).
235
To determine whether this pattern was a unique feature of the Cal07 strain, we used the same
experimental approach detailed above to generate single cell RNAseq libraries for A549 cells
infected at low MOI with the human seasonal H3N2 strain A/Perth/16/2009 (Perth09). Analysis of
1197 Perth09-infected cells revealed a substantial amount of viral gene expression heterogeneity,
although the distributions differed somewhat from what we observed with Cal07 (**Fig. S2**). These
differences may be partially due to an artifact of the QC process by which we eliminated cell
doublets from our analysis (See Methods and **Fig. S4** for details). Regardless, Perth09-infected

cells exhibited the same pattern of discordance between NS and the rest of the genome segments
that we observed with Cal07 (**Fig. 6C, S3**).

To identify broader patterns of viral gene expression, we performed K means clustering based on
expression of the eight viral gene segments for all Cal07- and Perth09-infected cells that were
positive for all viral transcripts. Clustering patterns were highly similar for both viruses and
revealed the existence of three distinct infected cell populations that largely subdivided cells
based on both total viral gene expression levels and NS segment expression (**Fig. 6D**). Thus,
human H1N1 and H3N2 viruses share a common pattern of single cell viral gene expression in
which NS segment expression diverges from that of the rest of the viral genome in a subset of
infected cells.

Fig. 6. Antiviral gene transcription is governed by a common pattern of single cell viral gene expression heterogeneity. (A) Pairwise comparisons between normalized copy numbers of the indicated viral transcripts within Cal07-infected cells in which all viral gene segments are detected. Shown here are the comparisons of PB2 with PB1 (left) and NS (right), along with r² values of correlation. **(B)** Correlation values plotted as heatmap for all pairwise comparisons of viral transcripts within Cal07-infected cells that express all viral genes. **(C)** Same as in (B) for Perth09-infected cells. **(D)** All Cal07- (left) and Perth09- (right) infected A549 cells that are positive for all eight viral gene segments, plotted as a function of total viral gene expression versus NS segment expression. Each dot represents the viral transcriptome of a single cell, colored by cluster identity of K-means clustering based on viral gene expression. Cluster IDs are shown in the legend. **(E)** The fractions of Perth09-infected cells in the clusters shown in (D) with detectable levels of IFNL1. **(F)** Plots show normalized counts per cell for each of IFNL1 transcript from Perth09-infected cells in the clusters shown in (D). **(G, H)** Same as in (E,F) for IFNL2. **(I,J)** Same as in (E,F) for IFIT2. **(K,L)** Same as in (E,F) for ZC3HAV1.

255
256

Single cell variation in NS segment expression is a major driver of the IFN response to Perth09, but not Cal07 infection

To determine whether these distinct clusters of viral gene expression were associated with distinct
patterns of host transcription, we compared host transcript expression in each cluster to all other
infected cells in the analysis using both MAST and NBID. This approach identified dozens of host

genes differentially expressed in each cluster (**Table S11-16**). For Cal07, no IFN or ISG genes
were significantly differentially expressed between clusters, but for Perth09 there were clear
differences in the expression of numerous host antiviral defense genes between the three
clusters. The expression frequencies and/or levels for type III IFNs (IFNL1 and IFNL2) and
selected ISGs (IFIT2, ZC3HAV1) were significantly higher in cluster 2, the cluster with low NS
expression, compared with the other clusters (**Fig. 6E-H**). Thus, for Perth09 but not Cal07, single
cell heterogeneity in NS segment expression level is a major determinant of the activation of the
interferon response.
270
H1N1 and H3N2 strains can differ significantly in single cell patterns of IFN and ISG
transcription
Finally, we compared the activation of host innate anti-viral gene expression at the single cell level
between Cal07 and Perth09. Mirroring the previous observations, expression of type I and III IFNs
were only observed in a tiny fraction of Cal07-infected cells (**Fig. 7A**) (13,39). This observation
could partially be a function of the relatively late timepoint that we examined, as IFNβ expression
typically peaks earlier during infection (40). Expression of IFNα was also rare in bystander cells
(**Fig. 7B**). We also observed widespread expression of multiple ISGs within Cal07-infected cells
indicating the successful activation of host antiviral defenses within infected cells despite the low
frequency of IFN expression, as has been described previously (11,41). Similar to what we
observed with IFNα, ISG expression was minimal in bystander cells, suggesting that the inhibition
of IFN induction by Cal07 is sufficient to largely prevent paracrine ISG activation in this host
system (**Fig. 7C**).
284
During Perth09 infection, type III IFN transcription was roughly 20-fold more frequent compared
with Cal07, indicating a significant difference in the ability of the host to initiate an IFN response
to infection by these two strains (**Fig. 7D**). Consistent with this, some (but not all) ISGs were
expressed more frequently in bystander cells during Perth09 infection compared with Cal07,
suggesting that Perth09 is less able to prevent paracrine activation of bystander cells (**Fig. 7E**).
Altogether, these data reveal clear differences in single cell patterns of IFN and ISG transcription
between the human H1N1 and H3N2 strains tested.
292

Fig. 7. H1N1 and H3N2 strains differ significantly in single cell patterns of IFN and ISG transcription. (A) tSNE plots of all Cal07-infected A549 cells colored based on whether IFNB1 and IFNL1 are detected. Results are representative of all type III IFNs examined. **(B)** tSNE plot of all bystander A549 cells colored based on whether IFNA2 transcript is detected. Similar results are found for other IFN-α subtypes. **(C)** Histograms comparing distributions of per-cell normalized counts for the indicated host transcripts across the three libraries (mock, Bystander, and Infected) during Cal07 infection. Vertical lines indicate 95% cutoff thresholds based on mock libraries, with percentages of cells in each library above this threshold shown on right. **(D)** tSNE plot of all Perth09-infected A549 cells colored based on whether IFNL1 is detected. Results are representative of all type III IFNs examined. **(E)** Same as in (C) for Perth09 infection.

293
294
DISCUSSION
The host response to infection arises from the combined responses of many individual cells, both
infected and uninfected. To understand the factors that govern host responses at the tissue and
organismal levels, it is critical to define patterns of variation in single cell infection responses.
Here, we demonstrate that IAV infection gives rise to a heterogeneous collection of divergent
transcriptional responses at the single cell level. Notably, this heterogeneity occurred within a
single cell type, under conditions where per-cell viral input and infection timing were normalized.
Single cell responses *in vivo* may be significantly more variable. By examining patterns of viral
gene expression heterogeneity at the single cell level, we uncovered a surprising dynamic where
NS segment expression diverges from that of the other genome segments. This pattern was
shared between the human seasonal strains that we tested and played a major role in regulating
the host antiviral response to H3N2 infection. These data establish a clear role for viral

heterogeneity in modulating the host response to infection and highlight the power of single cell
approaches to reveal new determinants of viral infection outcomes.
309
Variation in gene expression patterns between individual infected cells appears to arise from a
combination of (1) genetic and genomic variation between individual virions, (2) underlying
heterogeneity within host cells, and (3) stochastic variation in infection processes. In line with
previous reports of significant genetic and phenotypic heterogeneity within A459 and other
transformed cell line populations, we observed substantial transcriptional heterogeneity between
uninfected A549 cells that could not be simply explained by cell cycle status (42–44). As a result,
virions that enter different cells establish infection within differing cellular environments that may
be more or less supportive and primed for differing transcriptional responses to infection. This
effect is likely enhanced by the stochastic variation inherent in different viral life cycle stages (45).
319
Layered on top of this intrinsic cellular heterogeneity and stochasticity is the genetic and genomic
heterogeneity characteristic of IAV populations (5). We and others have previously demonstrated
that well-characterized viral modulators of host cell function such as NS1 and PA-X are not
ubiquitously expressed during infection (6,12). Here, we show that this variability in viral gene
expression between individual infected cells has clear consequences for the host cell
transcriptional response. Interestingly, we found that two levels of viral gene expression
heterogeneity influence single cell infection outcomes: variation in the presence or absence of a
given viral gene and, for cells where a gene is expressed, variation in expression level. This effect
was most pronounced for the NS segment where variation in expression levels between cells
were associated with significant differences in the host transcriptional response to infection.
Altogether, it is clear that future efforts to understand the role of critical viral proteins such as NS1
in shaping infection outcome will have to account for heterogeneity in single cell expression
patterns.
333
The comparison of individual viral gene segment expression across thousands of infected cells
revealed surprisingly complex relationships between the NS segment and the rest of the genome.
Within a subset of infected cells, NS expression was markedly decreased compared to the other
viral genes. Importantly, this dynamic was not universal, as NS expression was well-correlated
with expression of the other gene segments in some clusters of infected cells. This suggests that
multiple distinct states of viral gene expression can arise during infection. Similar to what has
been shown during genome packaging, our data suggest that specific interactions between gene
segments play an important role in regulating overall viral gene expression patterns (46–48).
Future efforts will aim to understand the mechanism behind the unique expression pattern of the
NS segment.
344
The comparison of Cal07 and Perth09 revealed clear commonalities and differences in single cell
expression patterns between the two viruses that may have relevance for understanding
differences in the apparent pathogenicity of different human H1N1 and H3N2 viruses. The most
obvious difference between the two viruses was seen in the activation of IFN and ISG
transcription. In addition, while NS segment expression status significantly influenced the host
response to both viruses, only Perth09 showed a clear relationship between NS levels and the
activation of the host antiviral machinery. This finding is in line with previous work that showed
that the 2009 pandemic H1N1 NS1 protein was less effective at suppressing transcription of
human IFNs and ISGs compared with NS1 genes from other human IAV isolates (49). While the
NS segment of Cal07 appeared to play a less direct role in suppressing IFN and ISG expression,
overall expression frequencies for select IFNs and ISGs were actually lower during Cal07

infection, indicating that Cal07 is more effective at suppressing IFN activation in A549 cells.
Overall, our data make clear that there is still a lot that we do not understand about the interplay
between viral gene expression patterns, viral genotype, and the host antiviral response.
359
In this study, we took steps to eliminate two other sources of viral heterogeneity that are likely
common during natural IAV infection. The first is presence of defective interfering particles (DIPs),
which we minimized by generating our virus stocks under low MOI conditions. DIPs appear to be
common within IAV populations, even in humans, and can have complicated effects on both viral
and host gene expression (13,50–55). We would predict that the presence of DIPs would further
increase overall heterogeneity between individual cell responses to the virus. It is worth noting
that we could not reliably detect the presence of DIP-associated transcripts within our data due
to the unavoidably incomplete coverage of the viral gene segments generated by our short-read
sequencing approach. The second likely source of additional heterogeneity that we excluded from
this study is variability in the cellular MOI. We and others have demonstrated that cellular co-
infection can be common *in vivo* (7,56–58). This suggests that the number of viral genomes
entering individual cells is likely quite variable. We recently showed that this variability in cellular
MOI can have distinct phenotypic consequences, both for viral replication dynamics and for IFN
induction (21). Altogether, it appears likely that the heterogeneity that we describe here
underrepresents what would be observed *in vivo*.
375
Our results extend the previous results of other groups including our own in establishing the
enormous amount of heterogeneity in viral gene expression that occurs at the single cell level,
even under experimental conditions designed to minimizes sources of variability (6,10,12,45,59).
Phenotypes that are critical to viral fitness such as viral load dynamics, transmissibility, and
pathogenicity must emerge from the collective output of heterogeneous populations of infected
cells. This raises questions of how selection may act upon patterns of viral heterogeneity to alter
these emergent phenotypes and the extent to which these heterogeneity patterns are under viral
genetic control. These questions are especially pertinent for segmented viruses like IAV but are
likely relevant across diverse virus families.
385
Altogether, our results help establish the importance of considering the roles of viral and host cell
heterogeneity in influencing the pathogenesis of viral infections. Similar to the way that viral
populations are now viewed, our data clearly establishes that the host response to infection
should be seen as a heterogeneous assemblage of single cell responses that collectively give
rise to the bulk phenotypes that are generally measured. This creates the potential for complex
interactions between responding cell subsets and raises the question of how such a
heterogeneous system is effectively regulated. It also raises the possibility that viral and host
response dynamics may be disproportionately driven by small subsets of cells that are obscured
during bulk analysis. Dissection of these diverse constituents is likely to reveal new mechanisms
that govern the pathogenesis of influenza virus infection.
396
METHODS
Plasmids
The A/California/04/09 and A/Perth/16/2009 reverse genetics plasmids were generous gifts from
Drs. Jonathan Yewdell and Seema Lakdawala, respectively. Plasmids encoding
A/California/07/09 were generated by introducing A660G and A335G substitutions into HA and
NP respectively, to match the amino acid sequences of A/California/07/09 HA and NP (NCBI
accession numbers CY121680 and CY121683).
404

Cells

Madin-Darby canine kidney (MDCK) and human embryonic kidney HEK293T (293T) cells were maintained in Gibco's minimal essential medium with GlutaMax (Life Technologies). Human lung epithelial A549 cells were maintained in Gibco's F-12 medium (Life Technologies). MDCK and A549 cells were obtained from Dr. Jonathan Yewdell; 293T cells were obtained from Dr. Joanna Shisler. All media were supplemented with 8.3% fetal bovine serum (Seradigm). Cells were grown at 37°C and 5% CO_2.

Viruses

Recombinant A/California/07/09 (Cal07) and A/Perth/16/2009 (Perth09) viruses were rescued via the 8-plasmid reverse genetics approach. For the rescue of both viruses, sub-confluent 293T cells were co-transfected with 500 ng of the following plasmids: pDZ::PB2, pDZ::PB1, pDZ::PA, pDZ::HA, pDZ::NP, pDZ::NA, pDZ::M, and pDZ::NS, using JetPrime (Polyplus) according to the manufacturer's instructions. Plaque isolates derived from rescue supernatants were amplified into seed stocks in MDCK cells. Working stocks were generated by infecting MDCK cells at an MOI of 0.0001 $TCID_{50}$/cell with seed stock and collecting and clarifying supernatants at 48 hpi. All viral growth was carried out in MEM with 1 µg/ml trypsin treated with L-(tosylamido-2-phenyl) ethyl chloromethyl ketone (TPCK-treated trypsin; Worthington), 1 mM HEPES, and 100 µg/ml gentamicin. The titers of the virus stocks were determined via standard 50% tissue culture infectious dose ($TCID_{50}$) assay.

Viral infection and cell sorting for single cell RNAseq

Confluent A549 cell monolayers in 3 T-25 flasks were infected with Cal07 (or Perth09) at MOI of 0.01 $TCID_{50}$/cell for 1 h. At 1 hpi, monolayers were washed with phosphate-buffered saline (PBS) and incubated in serum-containing F-12 medium. At 3 hpi, the medium was changed to MEM with 50 mM HEPES and 20 mM NH_4Cl to block viral spread. At 16 hpi, monolayers were trypsinized and combined into single-cell suspension and washed with PBS. Cal07-infected cells were stained with Alexa Fluor 488-conjugated mouse anti-HA monoclonal antibody (mAb) EM4-CO4 (gift from Dr. Patrick Wilson) and Alexa Fluor 647-conjugated mouse anti-M2 mAb O19 (gift from Dr. Jonathan Yewdell). Perth09-infected cells were first stained with human anti-HA stem antibody FI6 (Gift from Dr. Adrian McDermott) and then stained with Alexa Fluor 488-conjugated donkey anti-human IgG (Jackson ImmunoResearch). After staining, cells were washed with PBS twice, and single live cells were sorted as "infected" or "bystander" populations based on the expression of HA and M2 on a BD FACS ARIA II sorter. Importantly, uninfected A549 cells from a separate flask were also trypsinized, stained, and sorted as "mock" population which served as a negative control.

Single cell RNAseq cDNA library generation

Sorted cell samples were counted and checked for viability on a BD20 cell counter (BIO-RAD) before they were diluted to equivalent concentrations with an intended capture of 4000 cells/sample. Each individual sample was used to generate individually barcoded cDNA libraries using the 10x Chromium Single Cell 3' platform (Pleasanton, CA) following the manufacturer's protocol. The Chromium instrument separates single cells into Gel Bead Emulsions (GEMs) that facilitate the addition of cell-specific barcodes to all cDNAs generated during oligo-dT-primed reverse transcription. The experiment with Cal07 used V2 reagent and the experiment with Perth09 used V3 reagent (all steps followed the manufacturer's protocol).

Illumina Library preparation and sequencing

Following ds-cDNA synthesis, individually barcoded libraries compatible with the Illumina
chemistry were constructed. The libraries were sequenced on an Illumina NovaSeq 6000 using
S4 flowcell for the experiment with Cal07 and S3 flowcell for the experiment with
Perth09. Basecalling and demultiplexing of raw data was done with the mkfastq command within
Cell Ranger 2.1.1 (10x Genomics).

Single cell RNAseq analysis
The three Cal07-associated 10x Chromium Single Cell v2 libraries (Infected, Bystander, and
Mock) were demultiplexed and reference mapped using Cell Ranger Count (version 2.2) for
alignment to a combined human+virus reference (human: hg38, version 1.2.0; Cal07: Genbank
Accessions: CY121680-CY121687), and then combined using Cell Ranger Aggr. The three
Perth09-associated 10x Chromium Single Cell v3 libraries (Infected, Bystander, and Mock) were
processed and combined using Cell Ranger (version 3.1) for alignment to a combined
human+virus reference (human: hg38, version 1.2.0; Perth09: Genbank Accessions:
KJ609203.1- KJ609210.1). The resulting raw count matrix for each virus set was imported into an
R pipeline using SimpleSingleCell (60), where it was filtered for empty droplets (61), low feature
cells (i.e. droplets removed if < 400 features/cell), low expressing features (i.e. droplets removed
if < 4cells/feature), and potential doublets (droplets removed if UMI counts of host transcripts > 2-
fold the median count number).

During the QC process for the Perth09 infection experiment, roughly half of the infected cells
registered as "doublets" based on total host mRNA content, despite our filtering out cell doublets
during flow sorting (**Fig. S4**). This appears to be a Perth09-specific effect, as we did not see this
during Cal07 infection. While it is possible that many of these are actually single cells in which
total RNA amounts are elevated due to infection, we removed these cells from our analysis.

The filtered matrix was scale normalized using the deconvolution method described and
implemented in the scran R package (60). Overall cellular infection status and individual virus
gene presence/absence was determined to be above background by comparison to expected
background counts generated from the subset of cells from the "Bystander" library containing
virus counts (assumed to be background molecules). For bystander cells with virus-mapping
reads, we calculated the 95th percentile values for total virus molecule counts ('Total Virus
Counts') as well as for each of the individual gene segments ('Virus Gene Counts') and used
these values to calculate the 'Expected Virus Background Count' for all cells. These values
represent the threshold below which we assume viral signal is due to background cross-
contamination rather than true viral gene expression. Cell cycle status was determined by running
the Cyclone tool in the scran R package (60). The filtered, normalized, and annotated matrix was
then imported into a Seurat pipeline for additional analysis and visualization (62), including
differential gene expression analysis using the MAST (33) and NBID tools (32), graph-based
clustering of the cells (63), and PCA/tSNE dimensional reduction visualization.

Differential Gene Analysis (DGE) gene lists were produced from the filtered, normalized count
matrices using R packages MAST and NBID (Finak et al., 2015; Chen et al., 2018). Individual
MAST and NBID results were intersected using a custom Perl script to obtain final lists. DGE lists
for missing virus genes were generated by first sub-setting the count matrices to contain only
'Infected' status cells and then using the individual virus gene status factors to test for cells with
'Present' versus 'Absent' status. DGE lists for individual cell clusters were generated by first sub-
setting for 'Infected' status cells and then by non-zero cluster ID status. Tests were then performed
for each cluster versus all other clusters.

502
503 All code used for single cell analysis, along with associated documentation, is available from:
504 https://github.com/BROOKELAB/SingleCell
505
Multivariate homogeneity of groups dispersions analysis
In the analyses, each host gene represents a variable/coordinate and thus each cell can be seen
as a point in the multivariate space. Multivariate homogeneity of groups dispersion analysis
(Anderson, 2006) was performed on log transformed host gene expressions of the two group of
cells, i.e. mock cells and infected cells, using the betadisper function in the R package
(http://www.R-project.org/). Distances between the points (i.e. cells) and their respective group
centroid in the principal coordinates were then used to test homogeneity of variances and
calculate the p-value. Note that similar analyses were performed on linear host gene expression
values and results remain the same, i.e. overall host gene expressions are significantly more
heterogenous than those in mock cells.
516
517 **K-means clustering**
518 The clusters are identified by applying standard k-means clustering analysis to partition the joint
519 8-viral gene segment expressions (on a log scale) into three distinct clusters (k=3).
520

ACKNOWLEDGEMENTS
We are grateful to Dr. Alvaro Hernandez and Ms. Chris Wright of the DNA Services Lab within
the Roy J. Carver Biotechnology Center for expert assistance in experimental planning and the
preparation and sequencing of single cell RNAseq libraries. We also would like to thank Dr. Chris
Fields of HPCBio, also within the Roy J. Carver Biotechnology Center, for assistance in
troubleshooting analysis. This work has been generously funded by NIAID (1R01AI139246-
01A1), the DARPA INTERCEPT program (DARPA INTERCEPT W911NF-17-2-0034 and R-
00676-19-0), and the Roy J. Carver Charitable Trust (17-4905).

REFERENCES
1. Andino R, Domingo E. Viral quasispecies. Virology. 2015 May;479–480:46–51.

2. Bordería AV, Isakov O, Moratorio G, Henningsson R, Agüera-González S, Organtini L, et
al. Group Selection and Contribution of Minority Variants during Virus Adaptation
Determines Virus Fitness and Phenotype. PLoS Pathog. 2015 May;11(5):e1004838.

3. Domingo-Calap P, Segredo-Otero E, Durán-Moreno M, Sanjuán R. Social evolution of
innate immunity evasion in a virus. Nat Microbiol. 2019 Jun;4(6):1006–13.

4. Xiao Y, Dolan PT, Goldstein EF, Li M, Farkov M, Brodsky L, et al. Poliovirus intrahost
evolution is required to overcome tissue-specific innate immune responses. Nat Commun.
2017 29;8(1):375.

5. Brooke CB. Population Diversity and Collective Interactions during Influenza Virus
Infection. J Virol. 2017 Nov 15;91(22).

6. Brooke CB, Ince WL, Wrammert J, Ahmed R, Wilson PC, Bennink JR, et al. Most influenza
a virions fail to express at least one essential viral protein. J Virol. 2013 Mar;87(6):3155–
62.

7.   Brooke CB, Ince WL, Wei J, Bennink JR, Yewdell JW. Influenza A virus nucleoprotein
selectively decreases neuraminidase gene-segment packaging while enhancing viral
fitness and transmissibility. Proc Natl Acad Sci U S A. 2014 Nov 25;111(47):16854–9.

8.   Drayman N, Patel P, Vistain L, Tay S. HSV-1 single cell analysis reveals anti-viral and
developmental programs activation in distinct sub-populations. eLife. 2019 May 15;8.

9.   Guo F, Li S, Caglar MU, Mao Z, Liu W, Woodman A, et al. Single-Cell Virology: On-Chip
Investigation of Viral Infection Dynamics. Cell Rep. 2017 Nov;21(6):1692–704.

10. Heldt FS, Kupke SY, Dorl S, Reichl U, Frensing T. Single-cell analysis and stochastic
modelling unveil large cell-to-cell variability in influenza A virus infection. Nat Commun.
2015 Nov 20;6:8938.

11. Ramos I, Smith G, Ruf-Zamojski F, Martinez-Romero C, Fribourg M, Carbajal EA, et al.
Innate immune response to influenza virus at single-cell resolution in human epithelial cells
revealed paracrine induction of interferon lambda 1. J Virol. 2019 Aug 2;

12. Russell AB, Trapnell C, Bloom JD. Extreme heterogeneity of influenza virus infection in
single cells. eLife [Internet]. 2018 Feb 16 [cited 2018 Mar 15];7. Available from:
https://elifesciences.org/articles/32303

13. Russell AB, Elshina E, Kowalsky JR, Te Velthuis AJW, Bloom JD. Single-cell virus
sequencing of influenza infections that trigger innate immunity. J Virol. 2019 May 8;

14. Schulte MB, Andino R. Single-cell analysis uncovers extensive biological noise in
poliovirus replication. J Virol. 2014 Jun;88(11):6205–12.

15. Steuerman Y, Cohen M, Peshes-Yaloz N, Valadarsky L, Cohn O, David E, et al. Dissection
of Influenza Infection In Vivo by Single-Cell RNA Sequencing. Cell Syst. 2018 Jun
27;6(6):679-691.e4.

16. Wang C, Forst CV, Chou T, Geber A, Wang M, Hamou W, et al. Cell-to-cell variation in
defective virus expression and effects on host responses during influenza virus infection.
bioRxiv [Internet]. 2018 Dec 10 [cited 2019 Jul 17]; Available from:
http://biorxiv.org/lookup/doi/10.1101/487785

17. Zanini F, Pu S-Y, Bekerman E, Einav S, Quake SR. Single-cell transcriptional dynamics of
flavivirus infection. eLife [Internet]. 2018 Feb 16 [cited 2018 Apr 6];7. Available from:
https://elifesciences.org/articles/32942

18. Lawson DA, Kessenbrock K, Davis RT, Pervolarakis N, Werb Z. Tumour heterogeneity and
metastasis at single-cell resolution. Nat Cell Biol. 2018 Dec;20(12):1349–60.

19. Papalexi E, Satija R. Single-cell RNA sequencing to explore immune cell heterogeneity.
Nat Rev Immunol. 2018 Jan;18(1):35–45.

20. Stuart T, Satija R. Integrative single-cell analysis. Nat Rev Genet. 2019 May;20(5):257–72.

21. Martin BE, Harris JD, Sun J, Koelle K, Brooke CB. Cellular co-infection increases the
speed and efficiency of influenza A virus production and shapes the interferon response to
infection. bioRxiv [Internet]. 2019 Aug 30 [cited 2019 Sep 3]; Available from:
http://biorxiv.org/lookup/doi/10.1101/752329

22. Martin K, Helenius A. Nuclear transport of influenza virus ribonucleoproteins: the viral
matrix protein (M1) promotes export and inhibits import. Cell. 1991 Oct 4;67(1):117–30.

23. Patil G, Zhao M, Song K, Hao W, Bouchereau D, Wang L, et al. TRIM41-Mediated
Ubiquitination of Nucleoprotein Limits Influenza A Virus Infection. Schultz-Cherry S, editor.
J Virol [Internet]. 2018 Jun 13 [cited 2019 Oct 7];92(16). Available from:
http://jvi.asm.org/lookup/doi/10.1128/JVI.00905-18

24. Beyleveld G, Chin DJ, Moreno Del Olmo E, Carter J, Najera I, Cillóniz C, et al. Nucleolar
Relocalization of RBM14 by Influenza A Virus NS1 Protein. Duprex WP, editor. mSphere
[Internet]. 2018 Nov 14 [cited 2019 Aug 1];3(6). Available from:
http://msphere.asm.org/lookup/doi/10.1128/mSphereDirect.00549-18

25. Imamura K, Imamachi N, Akizuki G, Kumakura M, Kawaguchi A, Nagata K, et al. Long
Noncoding RNA NEAT1-Dependent SFPQ Relocation from Promoter Region to
Paraspeckle Mediates IL8 Expression upon Immune Stimuli. Mol Cell. 2014
Feb;53(3):393–406.

26. Landeras-Bueno S, Ortín J. Regulation of influenza virus infection by long non-coding
RNAs. Virus Res. 2016 Jan;212:78–84.

27. Sasaki YTF, Ideue T, Sano M, Mituyama T, Hirose T. MENε/β noncoding RNAs are
essential for structural integrity of nuclear paraspeckles. Proc Natl Acad Sci. 2009 Feb
24;106(8):2525–30.

28. Zhang F, Wu L, Qian J, Qu B, Xia S, La T, et al. Identification of the long noncoding RNA
NEAT1 as a novel inflammatory regulator acting through MAPK pathway in human lupus. J
Autoimmun. 2016 Dec;75:96–104.

29. Zhang P, Cao L, Zhou R, Yang X, Wu M. The lncRNA Neat1 promotes activation of
inflammasomes in macrophages. Nat Commun [Internet]. 2019 Dec [cited 2019 Aug
1];10(1). Available from: http://www.nature.com/articles/s41467-019-09482-6

30. Anderson MJ. Distance-based tests for homogeneity of multivariate dispersions.
Biometrics. 2006 Mar;62(1):245–53.

31. Diefenbacher M, Sun J, Brooke CB. The parts are greater than the whole: the role of semi-
infectious particles in influenza A virus biology. Curr Opin Virol. 2018 Jul 24;33:42–6.

32. Chen W, Li Y, Easton J, Finkelstein D, Wu G, Chen X. UMI-count modeling and differential
expression analysis for single-cell RNA sequencing. Genome Biol. 2018 31;19(1):70.

33. Finak G, McDavid A, Yajima M, Deng J, Gersuk V, Shalek AK, et al. MAST: a flexible
statistical framework for assessing transcriptional changes and characterizing

heterogeneity in single-cell RNA sequencing data. Genome Biol [Internet]. 2015 Dec [cited
2019 Jul 15];16(1). Available from: http://genomebiology.com/2015/16/1/278

34.   Ayllon J, García-Sastre A. The NS1 protein: a multitasking virulence factor. Curr Top
Microbiol Immunol. 2015;386:73–107.

35.   Hale BG, Randall RE, Ortín J, Jackson D. The multifunctional NS1 protein of influenza A
viruses. J Gen Virol. 2008 Oct;89(Pt 10):2359–76.

36.   Krug RM. Functions of the influenza A virus NS1 protein in antiviral defense. Curr Opin
Virol. 2015 Jun;12:1–6.

37.   Marc D. Influenza virus non-structural protein NS1: interferon antagonism and beyond. J
Gen Virol. 2014 Dec 1;95(Pt_12):2594–611.

38.   Arslan AD, Sassano A, Saleiro D, Lisowski P, Kosciuczuk EM, Fischietti M, et al. Human
SLFN5 is a transcriptional co-repressor of STAT1-mediated interferon responses and
promotes the malignant phenotype in glioblastoma. Oncogene. 2017 Oct;36(43):6006–19.

39.   Killip MJ, Jackson D, Pérez-Cidoncha M, Fodor E, Randall RE. Single-cell studies of IFN-β
promoter activation by wild-type and NS1-defective influenza A viruses. J Gen Virol. 2017
Mar;98(3):357–63.

40.   Osterlund P, Strengell M, Sarin LP, Poranen MM, Fagerlund R, Melen K, et al. Incoming
Influenza A Virus Evades Early Host Recognition, while Influenza B Virus Induces
Interferon Expression Directly upon Entry. J Virol. 2012 Oct 15;86(20):11183–93.

41.   Schmid S, Mordstein M, Kochs G, García-Sastre A, tenOever BR. Transcription Factor
Redundancy Ensures Induction of the Antiviral State. J Biol Chem. 2010 Dec
31;285(53):42013–22.

42.   Ben-David U, Siranosian B, Ha G, Tang H, Oren Y, Hinohara K, et al. Genetic and
transcriptional evolution alters cancer cell line drug response. Nature. 2018
Aug;560(7718):325–30.

43.   Croce MV, Colussi AG, Price MR, Segal-Eiras A. Identification and characterization of
different subpopulations in a human lung adenocarcinoma cell line (A549). Pathol Oncol
Res POR. 1999;5(3):197–204.

44.   Watanabe N, Dickinson DA, Krzywanski DM, Iles KE, Zhang H, Venglarik CJ, et al. A549
subclones demonstrate heterogeneity in toxicological sensitivity and antioxidant profile. Am
J Physiol Lung Cell Mol Physiol. 2002 Oct;283(4):L726-736.

45.   Dou D, Hernández-Neuta I, Wang H, Östbye H, Qian X, Thiele S, et al. Analysis of IAV
Replication and Co-infection Dynamics by a Versatile RNA Viral Genome Labeling Method.
Cell Rep. 2017 Jul;20(1):251–63.

46.   Bolte H, Rosu ME, Hagelauer E, García-Sastre A, Schwemmle M. Packaging of the
Influenza Virus Genome Is Governed by a Plastic Network of RNA- and Nucleoprotein-
Mediated Interactions. J Virol. 2019 Feb 15;93(4).

47.   Giese S, Bolte H, Schwemmle M. The Feat of Packaging Eight Unique Genome
Segments. Viruses. 2016 17;8(6).

48.   Hutchinson EC, von Kirchbach JC, Gog JR, Digard P. Genome packaging in influenza A
virus. J Gen Virol. 2010 Feb;91(Pt 2):313–28.

49.   Hale BG, Steel J, Medina RA, Manicassamy B, Ye J, Hickman D, et al. Inefficient control of
host gene expression by the 2009 pandemic H1N1 influenza A virus NS1 protein. J Virol.
2010 Jul;84(14):6909–22.

50.   Alnaji FG, Holmes JR, Rendon G, Vera JC, Fields CJ, Martin BE, et al. Sequencing
Framework for the Sensitive Detection and Precise Mapping of Defective Interfering
Particle-Associated Deletions across Influenza A and B Viruses. Schultz-Cherry S, editor. J
Virol [Internet]. 2019 Mar 13 [cited 2019 Jul 26];93(11). Available from:
http://jvi.asm.org/lookup/doi/10.1128/JVI.00354-19

51.   Martin MA, Kaul D, Tan GS, Woods CW, Koelle K. The Dynamics of Influenza A H3N2
Defective Viral Genomes from a Human Challenge Study. bioRxiv [Internet]. 2019 Oct 22
[cited 2019 Oct 29]; Available from: http://biorxiv.org/lookup/doi/10.1101/814673

52.   Nayak DP, Chambers TM, Akkina RK. Defective-interfering (DI) RNAs of influenza viruses:
origin, structure, expression, and interference. Curr Top Microbiol Immunol. 1985;114:103–
51.

53.   Saira K, Lin X, DePasse JV, Halpin R, Twaddle A, Stockwell T, et al. Sequence analysis of
in vivo defective interfering-like RNA of influenza A H1N1 pandemic virus. J Virol. 2013
Jul;87(14):8064–74.

54.   Vasilijevic J, Zamarreño N, Oliveros JC, Rodriguez-Frandsen A, Gómez G, Rodriguez G,
et al. Reduced accumulation of defective viral genomes contributes to severe outcome in
influenza virus infected patients. Thomas PG, editor. PLOS Pathog. 2017 Oct
12;13(10):e1006650.

55.   Vignuzzi M, López CB. Defective viral genomes are key drivers of the virus-host
interaction. Nat Microbiol. 2019 Jun 3;

56.   Fukuyama S, Katsura H, Zhao D, Ozawa M, Ando T, Shoemaker JE, et al. Multi-spectral
fluorescent reporter influenza viruses (Color-flu) as powerful tools for in vivo studies. Nat
Commun. 2015 Mar 25;6:6600.

57.   Jacobs NT, Onuoha NO, Antia A, Steel J, Antia R, Lowen AC. Incomplete influenza A virus
genomes occur frequently but are readily complemented during localized viral spread. Nat
Commun. 2019 Aug 6;10(1):3526.

58.   Marshall N, Priyamvada L, Ende Z, Steel J, Lowen AC. Influenza Virus Reassortment
Occurs with High Frequency in the Absence of Segment Mismatch. Basler CF, editor.
PLoS Pathog. 2013 Jun 13;9(6):e1003421.

59.   Sjaastad LE, Fay EJ, Fiege JK, Macchietto MG, Stone IA, Markman MW, et al. Distinct
antiviral signatures revealed by the magnitude and round of influenza virus replication in
vivo. Proc Natl Acad Sci. 2018 Sep 18;115(38):9610–5.

60.   Lun ATL, McCarthy DJ, Marioni JC. A step-by-step workflow for low-level analysis of
single-cell RNA-seq data with Bioconductor. F1000Research. 2016;5:2122.

61.   Lun ATL, Riesenfeld S, Andrews T, Dao TP, Gomes T, participants in the 1st Human Cell
Atlas Jamboree, et al. EmptyDrops: distinguishing cells from empty droplets in droplet-
based single-cell RNA sequencing data. Genome Biol. 2019 22;20(1):63.

62.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, et al.
Comprehensive Integration of Single-Cell Data. Cell. 2019 Jun 13;177(7):1888-1902.e21.

63.   Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, et al. Highly Parallel
Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets. Cell. 2015
May 21;161(5):1202–14.

703
704 **FIGURE LEGENDS**
705
Fig. 1. Generation of viral and host transcriptional data for thousands of singly infected
cells. (A) Schematic depicting our strategy for generating single cell RNAseq libraries from
thousands of cells infected at low MOI. In brief, we infect A549 cells with Cal07 at MOI=0.01 to
ensure that infected cells are infected with a single virion. We then block secondary spread with
NH4Cl treatment to make sure infection timing is uniform across all infected cells. Finally, we sort
"infected" and "bystander" cells based on surface expression of HA and/or M2 and immediately
generate single cell RNAseq libraries from these sorted cell populations using the 10X Chromium
device. In parallel, mock cells are sorted and used as uninfected controls. **(B)** tSNE dimensionality
reduction plot showing the extent of overlap of the indicated cell populations clustered based on
transcriptional similarity.
716
Fig. 2. Enormous heterogeneity in viral gene expression patterns. (A) Plot shows the
distribution of Cal07-infected A549 cells, binned by the fraction of total cellular poly(A) RNA that
is viral in origin. **(B)** Plots show the fraction of total poly(A) RNA per cell that maps to the indicated
viral gene segment. Each dot represents a single cell, cells with no detectable reads mapping to
the indicated segment were arbitrarily assigned a value of 0.000007 to show up on the log10
scale. **(C)** Distribution of normalized PB2-derived reads per cell (orange) compared with a Poisson
distribution of equal mean (blue line) on a log-log scale. **(D)** The viral gene expression distributions
normalized by the 70th quantile on a log-log scale. **(E)** Same analysis as in (E) but showing the
top 10 most abundant host transcripts.
726
Fig. 3. Significant heterogeneity in the host transcriptional response to infection. (A) tSNE
dimensionality reduction plot showing all Cal07 infected A549 cells clustered based on similarity
of host transcription patterns. **(B)** Heat map showing differential expression of the top 10
characteristic host genes for each cluster from (A). Individual cells are each represented by a
column, grouped by cluster, with individual rows representing relative expression of the top 10
specific host transcripts most significantly associated with each cluster. **(C)** Same tSNE plot of
Cal07 infected cells shown in (A) with each cell colored by the proportion of total cellular poly(A)

RNA that is viral in origin. **(D)** Same tSNE plot of Cal07 infected cells shown in (A) with each cell
colored by predicted cell cycle stage, as determined by the scran package. **(E)** Comparison of
normalized per cell counts of TRIM41 between mock and infected cells. **(F)** Fraction of cells in
each infected cell cluster from (A) with detectable levels of TRIM41. **(G)** Comparison of normalized
per cell counts of TRIM41 between clusters of infected cells shown in (A). **(H)** Comparison of
normalized per cell counts of NEAT1 between mock and infected cells. **(F)** Fraction of cells in
each infected cell cluster from (A) with detectable levels of NEAT1. **(G)** Comparison of normalized
per cell counts of NEAT1 between clusters of infected cells shown in (A). **(K)** tSNE dimensionality
reduction plot showing mock A549 cells clustered based on similarity of host transcriptional
patterns. **(L)** Principle coordinate axes (PCoA) plot comparing the multivariate dispersions for
mock (black) and Cal07-infected (red) A549 cells. The first two axes (PCoA 1 and PCoA 2) in the
multivariate homogeneity of group dispersions analysis are used.
746
747 **Fig. 4. Many infected cells have undetectable levels of one or more viral transcripts. (A)**
748 The fraction of all Cal07-infected cells that had detectable transcripts derived from the indicated
749 gene segments. **(B)** All Cal07-infected cells binned by the total number of viral gene segments
750 that were detectable, with the actual numbers of cells in each group detailed above.
751
**Fig. 5. Dissection of the effects of individual viral genes on the host transcriptional
response to infection. (A)** The number of host transcripts for which expression levels
significantly differ depending on whether the indicated viral gene segment is present or not,
according to both MAST and NBID. **(B)** Same as in (A) but removing host genes that are
differentially regulated by the expression status of more than one viral gene segment. **(C)** tSNE
plot of all Cal07-infected A549 cells colored based on whether NS segment-derived transcripts
are detected (Cyan) or not detected (Salmon). **(D)** Fractions of NS- and NS+ Cal07-infected A549
cells with detectable levels of SLFN5. **(E)** Distribution of SLFN5 normalized counts in cells shown
in (D).
761
**Fig. 6. Antiviral gene transcription is governed by a common pattern of single cell viral
gene expression heterogeneity. (A)** Pairwise comparisons between normalized copy numbers
of the indicated viral transcripts within single cells for all Cal07 infected cells in which all viral gene
segments were detected. Shown here are the comparisons of PB2 with PB1 (left) and NS (right),
along with r^2 values of correlation. **(B)** Correlation values plotted for all pairwise comparisons of
Cal07 viral transcripts within infected cells that expressed all viral genes. Color indicates the r^2
correlation values. **(C)** Correlation values plotted for all pairwise comparisons of Perth09 viral
transcripts within infected cells that expressed all viral genes. Color indicates the r^2 correlation
values. **(D)** All Cal07 (Left) or Perth09 (right) infected A549 cells that were positive for all eight
viral gene segments, plotted as a function of both total viral gene expression and NS segment
expression. Each dot represents the viral transcriptome of a single cell, colored by cluster identity
by the K-means clustering based on viral gene expression within infected cells. Cluster IDs are
shown in the legend. Infected cells with undetectable levels of one or more viral gene segments
were not included in this analysis. **(E)** The fractions of Perth09-infected cells in the clusters shown
in (D) that had detectable levels of IFNL1 transcript. **(F)** Plots show normalized counts per cell for
each of IFNL1 transcript from Perth09-infected cells in the clusters shown in (D). **(G, H)** Same as
in (E,F) but for IFNL2 transcript. **(I,J)** Same as in (E,F) but for IFIT2 transcript. **(K,L)** Same as in
(E,F) but for ZC3HAV1 transcript.
780
**Fig. 7. H1N1 and H3N2 strains differ significantly in single cell patterns of IFN and ISG
transcription. (A)** tSNE plots of all Cal07-infected A549 cells colored based on whether the

indicated transcripts were detected. Results are representative of all type III IFNs examined **(B)**
tSNE plot of all bystander A549 cells colored based on whether IFNA2 transcript was detected.
Similar results were obtained for other IFNalpha subtypes. **(C)** Histograms comparing distributions
of per-cell normalized counts for the indicated host transcripts across the three libraries (mock,
bystander, and infected) during Cal07 infection. Vertical lines indicate 95% cutoff thresholds
based on mock libraries, with percentages of cells in each library above this threshold shown on
right. **(D)** tSNE plots of all Perth09-infected A549 cells colored based on whether IFNL1 was
detected. Results are representative of all type III IFNs examined **(E)** Same as (C) but for Perth09
infection.
792
793 **SUPPLEMENTAL FIGURES**
794
795 **Fig. S1. All pairwise Cal07 viral gene correlation plots.** Normalized per cell copy numbers for
796 the indicated Cal07 genes plotted against each other. Data only show infected cells that
797 are positive for all viral gene segments.
798
Fig. S2. Single cell heterogeneity in viral gene expression during low MOI Perth09
infection. (A) Plot shows the distribution of Perth09-infected A549 cells, binned by the fraction
of total cellular poly(A) RNA that is viral in origin. **(B)** Plots show the fraction of total poly(A) RNA
per cell that maps to the indicated viral gene segment. Each dot represents a single cell, cells
with no detectable reads mapping to the indicated segment were arbitrarily assigned a value of
0.000007 to show up on the log10 scale. **(C)** The fraction of all Perth09-infected cells that
have detectable levels of transcripts derived from the indicated viral gene segments. **(D)** All
Perth09-infected cells binned by the total number of detectable viral gene segments, with the
actual numbers of cells in each group detailed above.
808
809 **Fig. S3. All pairwise Perth09 viral gene correlation plots.** Normalized per cell copy numbers
810 for the indicated Perth09 genes plotted against each other. Data only show infected cells that
811 are positive for all viral gene segments.
812
Figure S4. Distributions of single cell host mRNA counts and doublet calling between
Cal07 and Perth09 infection. Single cell distributions of host UMI counts for mock, bystander,
and infected cell libraries during **(A)** Cal07 and **(B)** Perth09 infection. In each group, the vertical
line indicates a value equal to the median UMI count of host transcripts in combined mock,
bystander, and infected cells, multiplied by 2. All cells with host UMI counts greater than this value
are considered doublets and are excluded from subsequent analyses.
819
820 **SUPPLEMENTAL TABLES**
821
822 **Table S1: All host genes differentially expressed between Seurat clusters of Cal07 infected**
823 **cells**
824
825 **Table S2: Combined DGE list based on Cal07 PB2 expression status within infected cells**
826
827 **Table S3: Combined DGE list based on Cal07 PB1 expression status within infected cells**
828
829 **Table S4: Combined DGE list based on Cal07 PA expression status within infected cells**
830
831 **Table S5: Combined DGE list based on Cal07 HA expression status within infected cells**

832
Table S6: Combined DGE list based on Cal07 NP expression status within infected cells

833
834
Table S7: Combined DGE list based on Cal07 NA expression status within infected cells

835
836
Table S8: Combined DGE list based on Cal07 M expression status within infected cells

837
838
Table S9: Combined DGE list based on Cal07 NS expression status within infected cells

839
840
Table S10: Combined DGE list showing all host genes differentially expressed within infected cells based on expression status of individual viral gene segments

841
842
843
Table S11: Combined DGE list showing host genes differentially expressed in cluster 1 (from K means clustering of all Cal07 infected cells positive for all viral genes)

844
845
846
Table S12: Combined DGE list showing host genes differentially expressed in cluster 2 (from K means clustering of all Cal07 infected cells positive for all viral genes)

847
848
849
Table S13: Combined DGE list showing host genes differentially expressed in cluster 3 (from K means clustering of all Cal07 infected cells positive for all viral genes)

850
851
852
Table S14: Combined DGE list showing host genes differentially expressed in cluster 1 (from K means clustering of all Perth09 infected cells positive for all viral genes)

853
854
855
Table S15: Combined DGE list showing host genes differentially expressed in cluster 2 (from K means clustering of all Perth09 infected cells positive for all viral genes)

856
857
858
Table S16: Combined DGE list showing host genes differentially expressed in cluster 3 (from K means clustering of all Perth09 infected cells positive for all viral genes)

859
860
861