**Network Inference from Multi-omic Data Uncovers Dynamic Transcriptional**

**Regulation Modules in Pathogenic Fungus *Fusarium graminearum***

Li Guo[1, 2, #, §], Mengjie Ji[1, #], Kai Ye[1, §]

*[1] MOE Key Lab for Intelligent Networks & Network Security, Faculty of Electronics and*

*Information Engineering, Xi'an Jiaotong University, Xi'an, 710049, China*

*[2] The School of Life Science and Technology, Xi'an Jiaotong University, Xi'an, 710049,*

*China*

**Running title:** *Fusarium graminearum* module networks

[§] Corresponding author:

Li Guo

guo_li@xjtu.edu.cn

Kai Ye

kaiye@xjtu.edu.cn

# Equal contribution

## ABSTRACT

The filamentous fungus *Fusarium graminearum* causes devastating crop disease and produces harmful mycotoxins worldwide. Understanding the complex *F. graminearum* transcriptional regulatory networks (TRNs) is vital for effective disease management. Reconstructing *F. graminearum* dynamic TRNs, an NP-hard problem, remains unsolved using commonly adopted reductionist or co-expression based approaches. Multi-omic data such as fungal genomic, transcriptomic data and phenomic data are vital to but so far have been largely isolated and untapped for unraveling phenotype-specific TRNs. Here for the first time, we harnessed these resources to infer global TRNs for *F. graminearum* using a Bayesian network based algorithm, "module networks". The inferred TRNs contain 49 regulatory modules that show condition-specific gene regulation. Through a robust validation based on prior biological knowledge including functional annotations and TF binding site enrichment, our network prediction displayed high accuracy and concordance with existing knowledge, highlighted by its accurate capture of the well-known trichothecene gene cluster. In addition, we developed a new computational method to calculate the associations between modules and phenotypes, and discovered subnetworks responsible for fungal virulence, sexual reproduction and mycotoxin production. Finally, we found a clear compartmentalization of TRN modules in core and lineage-specific genomic regions in *F. graminearum*, reflecting the evolution of the TRNs in fungal speciation. This system-level reconstruction of filamentous fungal TRNs provides novel insights into the intricate networks of gene regulation that underlie key processes in *F. graminearum* pathobiology and offers promise for the development of improved disease control strategies.

**Keywords:** Fusarium head blight, transcription factor, phenomics, gene regulation, dynamic networks, systems biology

## INTRODUCTION

Agricultural plants worldwide commonly suffer from devastating diseases caused by

pathogenic fungi (Agrios, 2005), threatening food safety and human survival amid increasing

global climate change. Fusarium head blight (FHB) caused by *Fusarium graminearum* (*Fg*)

is a serious disease of cereal crops, reducing yield and polluting the grains with mycotoxins

such as deoxynivalenol (DON) and zearalenone (ZEA) (Leslie *et al.*, 2006). FHB

pathogenesis is tightly controlled by host and pathogen gene regulatory networks (GRNs).

For example, genes involved in *Fg* growth*,* infection and secondary metabolism are subject to

fine regulation (Ma *et al.*, 2013). Numerous studies have demonstrated that the expression of

*Fg* genes related to pathogenesis, such as those encoding effectors (Brown *et al.*, 2017) and

cell wall-degrading enzymes (Carapito *et al.*, 2008), is induced *in planta* but suppressed *in*

*vitro*. Similarly, host genes involved in defense and immune response are induced during

pathogen invasion (Pan *et al.*, 2018). Understanding GRNs is fundamental in solving medical

and agricultural problems (Emmert-Streib *et al.*, 2014) caused by microbial infections. GRNs

can inform disease control approaches by permitting the specific targeting of key pathogen

regulators, as reported in recent studies (Koch *et al.*, 2016). However, GRNs involved in

FHB and mycotoxin production remain poorly understood.

Genes and gene regulators such as signaling proteins and transcription factors (TFs) are

interconnected in GRNs. Many studies have attempted to dissect GRNs using a reductionist

approach by analyzing the gene expression profiles of *Fg* mutants (Lysoe *et al.*, 2011; Son *et*

*al.*, 2016; Kong *et al.*, 2018). Though conceptually valid, this approach is time-consuming

and unrealistic as a method of decoding the highly complex GRNs of eukaryotic cells.

Alternatively, protein interaction networks have been inferred using protein domain

homology. For instance, Zhao *et al.* constructed *Fg* protein-protein interaction (FPPI)

networks using protein domains that are conserved in *Fg* and *Saccharomyces cerevisiae* (*Sc*)

(Zhao *et al.*, 2009). Despite its usefulness in finding potentially interacting proteins, this

approach infers a network based solely on protein sequence features and therefore lacks

functional support. A more feasible approach is to use genome-wide expression data to

deduce regulatory networks. For example, Kim *et al.* predicted gene co-expression networks

involved in virulence of *F. verticillioides* using RNA-Seq data from the *FSR1* mutant (Kim *et*

*al.*, 2018). In addition, Liu *et al.* constructed a co-expression network based on gene

expression data and the FPPI database, identifying several hub pathogenicity genes and

subnetworks (Liu *et al.*, 2010). These approaches have indeed produced valuable insights into

*Fg* co-expression gene modules. However, co-expression does not necessarily indicate a true

regulatory relationship. Recently, Lysenko et al. used gene expression data combined with

data on protein interactions and sequence similarity to study the networks important for

virulence (Lysenko *et al.*, 2013). While integrating multiple sources of evidence is an

improvement, it still relies on co-expression evidence and does not prove actual regulatory

relationships. Furthermore, because the study focused on small gene sets that have an impact

on virulence, a systemic view of regulatory networks is lacking.

Regulatory relationships are typically inferred from large genome datasets using

computational methods built on mathematical models (Kim *et al.*, 2009). Boolean networks,

Bayesian networks, and Mutual Information have already proven to be powerful models for

inferring regulatory networks (Friedman *et al.*, 2000; Shmulevich *et al.*, 2002; Zhao *et al.*,

2016). Bayesian networks are probabilistic models that are ideal for studying regulatory

relationships using noisy data such as gene expression profiles (Friedman *et al.*, 2000).

Therefore, these models have been frequently adopted in various GRN-inference algorithms.

Previously, from a large collection of transcriptomic data, we reconstructed a global GRN for

*Fg* using the machine-learning method MinReg (Pe'er *et al.*, 2006) based on a Bayesian

network model, successfully predicting 120 top regulators for 13,300 *Fg* genes (Guo *et al.*,

2016). Despite the progress it represents, this first *Fg* GRN has obvious limitations. First, it

mainly focuses on master regulators that control general rather than fungal-specific biological

processes. Second, it is essentially static and offers little insight into how the networks adapt

to various changes in endogenous and environmental stimuli. Without such knowledge, it is

difficult to predict how gene regulation of diverse and specific biological processes operates

and to find the *bona fide* regulators in the system.

TFs regulate gene transcription via binding to the promoter regions of target genes.

Elucidation of transcription regulatory networks (TRNs) is a vital step in mapping global

GRNs. Here, for the first time, we reconstructed global TRNs for *Fg* by applying a module

network learning algorithm (Segal *et al.*, 2003) to a large collection of transcriptomic data

and integrating a phenomic database of *Fg* TFs that were reported previously (Son *et al.*,

2011). The integration of phenomics and transcriptomics data in this study allows us to

identify 49 module networks that are directly involved in the cellular processes that underlie

key phenotypes in *Fg*, yielding the novel and crucial knowledge that "regulator X regulates

target genes Y under condition Z". Validation of the networks demonstrates the high

accuracy of the inference. Association mining of the predicted module networks reveals links

between gene modules and fungal phenotypes. The condition-specific TRNs significantly

improve the resolution of the *Fg* transcriptional circuits controlling virulence, sexual

reproduction and mycotoxin production, laying a vital foundation for the development of

novel regimes to minimize FHB occurrence and mycotoxin contamination. The *Fg* module

network (FuNet) is available for public query and downloading (https://xjtu-funet-

source.github.io/FuNet/FuNet.html).

**MATERIALS AND METHODS**

**Fungal transcriptomic and TF phenomic data**

The *Fg* transcriptome data were downloaded from PLEXdb (www.plexdb.org) and from the

Filamentous Fungal Gene Expression database (http://bioinfo.townsend.yale.edu/) (Table

S1). The data and the normalization procedure have been described previously (Guo *et al.*,

2016). The phenome data for *Fg* TFs were obtained from literature (Son *et al.*, 2011) (Table

S2). The expression data and a candidate regulator list of 170 TFs showing phenotypic

changes in disruption mutants were used as the input data for module network inference.

**Module Network algorithm implementation**

Modularized TRNs were inferred using a probabilistic method called "module network"

(Segal *et. al* 2003). Using the input data, the algorithm determined both the partitioning of

genes to modules and the regulation program for each module in an iterative manner. For

each iteration, the procedure searched for a regulation program for each module and then

reassigned each gene to the module whose program best predicted its behavior. These two

steps were iterated until convergence was reached using the expectation maximization (EM)

algorithm, thereby returning the predicted regulatory modules containing a set of regulators

and target genes. Each module was represented as a decision tree that specified the conditions

under which target genes were regulated by a particular regulator and whether the regulation

was positive or negative (Segal *et al.*, 2003). The search process was repeated three times,

and the same modules and regulation programs were returned.

**Validation of module networks**

The predicted modules were first validated based on the consistency between regulator

phenotypes and target gene expression. Based on three major phenotypes of *Fg*, each module

was evaluated for consistency between the regulator phenotype and the experimental

conditions using the following three validation points: 1) sexual reproduction; 2) virulence;

and 3) mycotoxin production. To quantify the consistency of the evaluation, we developed a

scoring function (named $Score_{vp}$) for each validation point; this function was defined as

147                                     $$Score_{vp} = M_c / N_c$$

$N_c$ represents the total number of conditions included in this study, and $Mc$ represents the

number of conditions under which the regulator phenotype was matched with a

corresponding condition directly related to the phenotype.

The regulatory modules were also validated based on conservation of $Fg$ and $S.\ cerevisiae$ TF

binding site (TFBS). First, the 500-bp sequence upstream of the $Fg$ genes of each module

was extracted, and the $MEME$ algorithm (Bailey $et\ al.$, 2009) was used to search for

conserved sequence motifs. The top five enriched motifs (ranked by E-value) were

considered the candidate TFBS of each regulatory module. Each enriched TFBS was then

compared to the $YEASTRACT$ database of $S.\ cerevisiae$ using $Tomtom$ (Bailey $et\ al.$, 2009) to

find the conserved TFBS in budding yeast. The top conserved yeast motif for each $Fg$ TFBS

was then selected. With the existing knowledge of yeast TF-TFBS associations, the

conserved yeast motifs identified through $Tomtom$ identified corresponding TFs, which were

denoted as "motif-deduced TFs" (MTFs). Second, to examine how many of these TFBS are

potentially recognized by conserved TFs in $Fg$ and $S.\ cerevisiae$, a BLASTp search was

conducted in which the $Fg$ regulators in each module were searched against the $S.\ cerevisiae$

genome to find regulator orthologs (E-value < 1e-5); these were defined as "orthologous

TFs" (OTFs). The OTFs were overlapped with MTFs to find conserved TFs that also

potentially bind to conserved TFBS enriched in $Fg$ regulatory modules. Based on the two

separate analyses, a score was assigned to quantify the motif validation performance of each

module:

$$Score_{motif} = M_o \, / \, N_e$$

$M_o$ represents the number of motifs that showed conservation based on the above analysis,

and $N_e$ represents the total number of enriched motifs per module.

Lastly, we validated each module by finding consistency between the functional annotations

enriched in the module genes and the annotations associated with the enriched TFBS

(deduced from *YEASTRACT*). First, GO enrichment was performed on the target genes from

each module using MIPS Funcat (Ruepp *et al.*, 2004) with the *Fg* genome as a reference. We

then assigned functional annotations to each module by selecting the most highly enriched

GO terms for each module. The annotations of the conserved enriched TFBS from each

module were retrieved from *YEASTRACT* and compared to the GO terms enriched in the

module genes. Following this analysis, we scored the module annotations matched with motif

annotations as either 0 (no match) or 1 (match). We combined the scores from the above five

validation points to obtain a total score; based on this score, the module was categorized as a

high-confidence (>0.6), moderate-confidence (0.4~0.6) or low-confidence module (<0.4).

Modules with fewer than two validation points were not validated.

**Calculation of the module-phenotype association index**

We developed an in-house computational method to accurately quantify the association

between modules and phenotypes. We calculated a score called the association index (AI) for

each module-phenotype association using multiple variables. The first variable ($W_{ir}$) was the

weight of the regulators derived from the number of conditions affected by the regulators

specified by the regulation tree (Table S3). The number of experiments affected by each

regulator in each module was used to obtain $W_{ir}$ (the weight of each regulator in each

module; $i$ =2, 3, 4...49 for modules M02, M03, M04…M49 and $r$ =1, 2, 3...R for regulator 1,

regulator 2, regulator 3…regulator R) according to the ratio of the number of experiments

affected by the regulator. $N_{ir}$ indicates the number of conditions affected by the *r-th* regulator

in the *i-th* module. $W_{ir}$ (0~1) was calculated as follows:

$$W_{ir} = N_{ir} / \sum_{r=1}^{r=R} N_{ir} \tag{1}$$

We then computed AI for each module-phenotype combination. Using the variable $X_{irj}$,

which could take a value of 0, 1 or -1, we could represent the influence of any regulator on a

specific phenotype; the values 0, 1 and -1 indicated that the corresponding *r-th* regulator in

the *i-th* module has no effect on, enhances or reduces, respectively, the *j-th* phenotype (Table

S4). We calculated the AI ($P_{ij}$) by multiplying the influence of regulator *r* on a phenotype *j*,

denoted as $X_{irj}$, by the weight of the corresponding regulator $W_{ir}$ and finally summing the

product of all the regulators (*R*) in the module.

$$P_{ij} = \sum_{r=1}^{r=R} |X_{irj}| \bullet W_{ir} \tag{2}$$

**Association mining**

We created a correlation matrix of all modules based on phenotype associations. First, we

filtered out minor associations (AI < 0.3) to capture major module-phenotype associations.

The Pearson correlation coefficient (PCC) of each pair of modules was calculated and used to

create a PCC matrix using the association indexes of the modules across phenotypes.

Hierarchical clustering was used to find module clusters that are likely to contribute to similar

phenotypes. Each cluster was subjected to detailed downstream examination.

**Network compartmentalization analysis**

We identified 9,700 orthologous genes as core genes conserved among the three *Fusarium*

sister species *Fg, F. verticillioides* and *F. oxysporum* (Ma *et al.*, 2010). In total, 3,600 *Fg*

genes lacking orthologous sequences in the sister species were loosely defined as *Fg* lineage-

specific (LS) genes. We compared the observed ratio of LS and core genes in each predicted

regulatory module to the expected ratio for the *Fg* genome using two-tailed Fisher's exact test

to determine whether there is an enrichment of LS or core genes. A threshold p-value < 0.05

was applied to determine whether the module was enriched (either LS or core) or not (mix).

**Network visualization**

Cytoscape (version 3.6.1) (Shannon *et al.*, 2003) and Gephi (version 0.9.2) (Bastian *et al.*,

2009) were used for network visualizations. For building weight-based networks, modules

and regulators are presented as nodes, and the weight of the regulator ($W_{ir}$) in each module

was used as the connection value for the edges. For phenotype-module networks, association

indexes were used as the connection value of the edges.

**Network and code availability**

The source codes used in this study are available at https://github.com/XJTU-FuNet-

source/FuNet/tree/master/module-network_code-and-tool. The module networks of *Fg* are

available at https://xjtu-funet-source.github.io/FuNet/FuNet.html.

**RESULTS**

*1.   Fusarium graminearum* **module network inference**

To infer condition-specific TRNs in *Fg*, we applied the module network algorithm (Segal *et*

*al.*, 2003) to a public dataset of *Fg* transcriptomic profiles spanning 67 different experimental

conditions, including sexual reproduction and plant infection (Table S1). In addition, we used

a set of candidate regulators consisting of 170 TFs that were previously functionally

associated with key fungal phenotypes available in FgTFPD (*Fg* TF phenotype database)

(Fig. 1) (Son *et al.*, 2011). Combined expression data and phenotype-associated candidate

regulators were used as input data for the module network algorithm to reconstruct TRNs in

*Fg* (Fig. 1). Searching iteratively, the algorithm discovered 49 *Fg* gene regulatory modules,

48 of which had predicted regulators (Fig. 2a; Fig. S1; Table S5-S6).

Each of the 48 modules is a regulatory program composed of various regulators, target genes

and the expression profiles of target genes as a function of the expression level of the

regulators. The regulatory program is presented in a decision-tree structure that defines the

behavior of each regulator and the conditions under which the regulation takes place (Fig.

S1). Overall, we predicted 117 regulators for 48 modules in *Fg*. The average numbers of

target genes and regulators for each module are 268 and 7, respectively, with standard

deviations of 212.43 and 1.24, respectively (Fig. 2b and 2c). The regulator-module

association network (Fig. 2d) showed that 42 regulators were associated with only one

module and that 75 regulators were associated with two or more modules. Five regulators

including two ASPES proteins FGSG_04220 (13) and FGSG_10384 (7), a C2H protein

FGSG_07052 (7), an HMG protein FGSG_01366 (8) and a $Zn_2C_6$ DNA binding protein

FGSG_08626 (7) function as hub regulators associated with the greatest number of modules

(Fig. 2d). Unsurprisingly, these regulators are highly pleiotropic, especially FGSG_04220

and FGSG_10384, whose deletion mutants are defective in the majority of phenotypes

assayed (Table S2) (Son *et al.*, 2011).

**2. Validation using prior knowledge proves the high credibility of *Fg* module networks**

Following the network inference, we assessed its reliability based on its consistency with

prior knowledge. We scored each module by evaluating its performance based on multiple

pieces of evidence, including regulator phenotypes, experimental conditions, gene

annotations and cis-regulatory elements (Materials and Methods). A module was considered

high-confidence, moderate-confidence or low-confidence depending on the validation score

(Table S7). After discarding 16 modules for which there was little evidence, we identified 14

high-confidence, 13 moderate-confidence and 6 low-confidence modules. Overall, the high-

and moderate-confidence modules account for 81.8% of the evaluable modules (Table S7),

showing that our network inference has achieved a high degree of credibility. The following

are examples of validation results that indicate the high credibility of our predicted

modules.

*2.1. High concordance between regulator phenotypes and condition-specific gene regulation*

Transcriptional regulators and their target genes are usually involved in the same biological

processes. Based on this general premise, we validated all predicted modules by evaluating

the concordance between the phenotypes associated with the top regulators in each module

and our predicted condition-specific regulation using TF-phenotype associations and

expression data associated with the experimental conditions. Since our predicted regulatory

programs specify the regulators and the conditions under which the regulation occurs, the

inferred relationship is accurate if a regulator associated with a phenotype and its target genes

are both activated or suppressed under the experimental conditions that result in the

phenotype. For simplicity and clarity, we focused our validation on the three largest groups of

experimental conditions included in our data: sexual reproduction, plant infection and

secondary metabolism (Table S1); these groups correspond to the sexual reproduction,

virulence and mycotoxin (DON and ZEA) production phenotypes in FgTFPD,

respectively. We found that in 45 of 48 modules with predicted regulators (Table S6), the top

regulator has an effect on one or more of the phenotypes associated with sexual reproduction,

virulence (plant infection) or mycotoxin production (secondary metabolism). In 34 of the 45

modules, the top regulator activates or suppresses the target genes under experimental

conditions related to specific phenotypes (Table S7). Three specific examples, each

concerning a phenotype, are provided below.

Firstly, in 76% of the modules whose top regulators are associated with sexual reproduction,

regulation of the module genes by the top regulator was found under at least one sexual

reproduction condition; for over 50% of the modules, the regulation occurred under half of

the corresponding conditions (Table S7). For example, the top regulator of M30

(FGSG_06356) is essential for sexual reproduction. Our prediction showed that this TF and

M30 genes were highly expressed under all sexual reproduction conditions. Secondly, in 57%

of modules whose top regulators are associated with virulence, regulation of the module

genes by the top regulator was found under at least one plant infection condition, and in

nearly 30% of these modules, regulation occurred under half of the plant infection conditions

(Table S7). For example, the top regulator of M16 (FGSG_07928) is essential for virulence,

and our prediction showed that FGSG_07928 and the M16 genes were highly expressed

under 62.5% of plant infection conditions. Thirdly, in 70% of the modules whose top

regulators are associated with mycotoxin (DON or ZEA) production, regulation of module

genes by the top regulator was found under 50% or more of the conditions that lead to

mycotoxin induction (Table S7). For example, the top regulator for M46 (FGSG_03538) is

essential for DON production, and our prediction showed that FGSG_03538 and the M46

genes were highly expressed under all mycotoxin induction conditions. In summary, 24 of the

32 predicted top regulators (75%) for 34 of the 48 predicted modules (70%) showed high

concordance between the regulator phenotype and condition-specific gene regulation.

*2.2. Most predicted regulatory modules have functionally conserved TF binding sites*

TFs regulate genes via binding to upstream cis-regulatory gene regions. Co-expressed genes

(*e.g.*, genes in the same regulatory module) typically share TF binding sites (TFBS) that are

recognized by one or more TFs. Therefore, we validated the predicted *Fg* (*Fg*) network

modules by finding enriched TFBS in each module. Using the *MEME* algorithm, we first

identified the top five enriched motifs (E-value < 0.05) within 500 bp upstream of the coding

sequences for all *Fg* genes within a module (Table S7). Overall, 47 of 49 modules (96%)

have significantly enriched motifs, and 34 (70%) have at least three enriched motifs (E-value

<0.05). We then compared these significantly enriched motifs with the budding yeast *S.*

*cerevisiae* (*Sc*) TFBS database *YEASTRACT* using *Tomtom* to functionally annotate these *Fg*

motifs using conserved *Sc* motifs (Fig. 3; Table S8). To determine whether the enriched

motifs were consistent with the biological functions of each module, we identified the most

significantly enriched GO terms for each regulatory module (Fig. 3; Table S9) and compared

the GO terms with the functional annotations of the significantly enriched *Fg* motifs. We

found that the functional annotations of 27 of the 49 modules (55%) matched in the enriched

TFBS and GO enrichment (Fig. 3; Table S10). For example, we found a functional match

between M46 target genes and one of their enriched TFBS (*Yrm1p*) (Fig. 3); both are related

to detoxification and multidrug resistance. This is consistent with the fact that M46 was

highly associated with the mycotoxin production phenotype, as shown in later sections.

Secondly, to examine whether the functional conservation in TFBS was achieved through the

conservation of regulator genes, a BLASTp search was conducted in which predicted *Fg*

regulators were searched against the *Sc* genome to identify orthologous regulators (E-value <

1e-5). We then compared these yeast regulator orthologs with the *Sc* TFs derived from the *Sc*

TFBS homologous to the enriched *Fg* TFBS. By overlapping the regulators identified in the

two separate analyses, 10 different regulators regulating 36 modules (Table S11) were found,

suggesting that not only did the predicted *Fg* regulators in 70% of the modules have

conserved *Sc* homologs but also that conserved TFBS are likely associated with these fungal

TF homologs. For example, motif enrichment shows that modules M06, M24, and M38 were

enriched in a common TFBS (*Azf1p*) that is likely bound by YMR019W in *Sc*. One of the

predicted regulators of module FGSG_08028 was orthologous to YMR019W (E-value =

1.46e-7). Another example is M39; this module is enriched in TFBS (*Ace2p*), which is likely

bound by YLR131C in *Sc*. Interestingly, one of the predicted regulators of module

FGSG_01341 is orthologous to YLR131C (E-value = 3.34e-17). The *YEASTRACT* database

showed that this conserved TFBS and TF are involved in the biogenesis of cellular

components, and this was captured by the GO enrichment for *Fg* module genes.

*2.3. Predicted regulatory modules captured the best-known TRN model in Fg*

The best-understood model of a transcriptional regulation network in *Fg* is that of the

trichothecene biosynthesis gene cluster, known as the Tri-cluster. Previous gene functional

studies identified two TFs, *Tri6* (FGSG_03536) and *Tri10* (FGSG_03538), that reside within

the cluster and regulate the expression of the entire cluster (Seong *et al.*, 2009). Deletion of

*Tri6* or *Tri10* in *Fg* abolished fungal production of trichothecene mycotoxins such as DON

and its derivatives (Hohn *et al.*, 1999; Tag *et al.*, 2001). Remarkably, our network inference

yielded a 44-gene module (M46) capturing the majority (9 of 12) of the Tri-cluster genes and

accurately predicted *Tri6* and *Tri10* as the top two regulators of this module (Fig. 4a). To

validate this module, we overlapped the module gene list (Table S5) with previously

published lists of the genes that are down-regulated in *ΔTri6* and *ΔTri10* mutants (fold

change ≥ 2) (Seong *et al.*, 2009) as well as with lists of Tri6-binding genes identified by

ChIP-Seq (Nasmith *et al.*, 2011) and found that 34%, 22% and 11.3% of our module genes

were included in the *ΔTri6* and *ΔTri10* down-regulated genes and *Tri6* ChIP-seq targets,

respectively (Fig. 4b). Most of the overlapped genes were Tri-cluster genes. Furthermore, GO

enrichment showed that this module was enriched in terms such as "secondary metabolism",

"detoxification involving cytochrome P450", "sesquiterpenes metabolism", "isoprenoid

metabolism", and others (Fig. 4c), consistent with the idea that this module regulates both

primary and secondary metabolism as well as self-protection by *Tri10* and *Tri6* (Tag *et al.*,

2001). Notably, our prediction specified that the target genes were activated by *Tri10* and

*Tri6* during wheat and barley head infection, wheat coleoptile and crown rot infection, and

toxin-inducing conditions but suppressed during sexual reproduction. In addition, M46

captured the full eight-gene cluster (FGSG_08077 ~ FGSG_08084) for the biosynthesis of

the *Fusarium* mycotoxin butenolide, which is toxic to animals (Liu *et al.*, 2007). We

predicted that this gene cluster is regulated by *Tri6* and *Tri10*, providing further evidence that

the two TFs might be global regulators of secondary metabolism including trichothecene

production, as previously reported (Seong *et al.*, 2009; Nasmith *et al.*, 2011).

**3. Computational method revealing module-phenotype association networks**

To understand which modules are responsible for regulating key phenotypes in *Fg*, we

developed a computational framework and used it to calculate an association index (AI)

between each module and each phenotype (Fig. 5a; Table S12) using digitalized regulator

phenotypes (Table S4) weighted by regulator hierarchical positions on the decision tree

(Table S3) and the fraction of phenotype-matching conditions being regulated (Materials and

Methods). We found that the module-phenotype association was quite complex in that the

majority of modules were associated with multiple phenotypes. Under a threshold of AI >

0.3, we removed minor associations and clarified the primary associations between modules

and different phenotypes and built module-phenotype association networks (Fig. 5b). We

found that sexual reproduction (49), mycelial growth (33), mycotoxin production (31) and

virulence (30) were associated with the largest numbers of modules.

Regulatory modules typically work as groups to control cellular functions. Therefore, we

investigated the relationships among different modules with respect to their contribution to

phenotypes. We measured the correlation among all modules by calculating the pairwise

Pearson correlation coefficients (PCC) and then performed hierarchical clustering of the

modules based on their PCCs. The analysis identified five major clusters of gene modules;

each cluster showed a distinctive pattern of association with phenotypes (Fig. 5c). Cluster I

included four modules (M34, M39, M30 and M48) in which sexual reproduction and

mycotoxin production are major phenotypes as well as modules M34 and M39, which are

mildly linked to mycelial growth and/or virulence. Cluster II contained 12 modules (M23,

M44, M06, M40, M38, M33, M03, M15, M10, M18, M07 and M11) that are all linked to

mycelial growth, sexual reproduction, mycotoxin production and virulence. Cluster III was

composed of 11 modules (M13, M17, M37, M45, M41, M36, M46, M02, M12, M09, and

M35) that are associated with pigmentation, in addition to other phenotypes. Overall, the

phenotype associations of Clusters I to III were dominated by sexual reproduction and toxin

production, and M30, M48, M36, M46, M12 and M02 were exclusively associated with these

two phenotypes. Cluster IV contained eight modules (M28, M43, M24, M49, M22, M47,

M04, and M08) whose phenotype associations were dominated by sexual reproduction and

virulence and two modules that were exclusively associated with virulence. Cluster V

contained 13 modules (M29, M32, M19, M05, M25, M20, M27, M42, M16, M26, M31,

M14, and M21) associated with mycelial growth. However, except for the fact that M27 and

M42 were mostly associated with mycelial growth and sexual reproduction, no phenotype

associations were dominant.

**4. TRN subnetworks that control key fungal phenotypes**

From the association analysis results, we built subnetworks of *Fg* controlling virulence,

sexual reproduction and mycotoxin production (Fig. 6). In these subnetworks, the nodes

represent top regulators and modules, and the edges represent regulatory relationships

weighted by the degree of the regulator's influence on the module.

*4.1. Sexual reproduction TRNs*

Sexual reproduction is critical for *Fg*, which produces ascospores as the primary inocula in

the field. Based on the association index, we identified 10 modules (M30, M28, M24, M49,

M22, M47, M04, M08, M27 and M42) that are highly associated with sexual reproduction

and that have an association index of over 50% for all phenotype associations (Fig. 5c). The

top regulators of the 10 modules were all essential for normal development of perithecia, and

most act as positive regulators. For example, FGSG_04480 and FGSG_08890, the two top

regulators of M08, activated target genes during sexual development (Fig. S1-8).

Interestingly, FGSG_08890 is a mating-type locus gene (*MAT-1-3*) and therefore potentially

contributes to fungal mating by regulating sexual reproduction-related gene modules. In

contrast, the two TFs FGSG_06356 and FGSG_09308 are negative regulators of sexual

reproduction. For example, FGSG_06356, which is negatively associated with sexual

development (Son *et al.*, 2011), was predicted as a top regulator for M30 and M42.

Consistently, our regulation program for M30 predicted that FGSG_06356 acts as a

suppressor of M30 genes during sexual development (Fig. S1-30). Likewise, FGSG_09308

was predicted as a suppressor of M22 (Fig. S1-22), consistent with its previously reported

phenotype. In addition, some regulators that had connections with many modules acted as

hub regulators in sexual reproduction-associated networks, including FGSG_04220,

FGSG_00318, FGSG_01366, FGSG_13314, FGSG_07052, FGSG_08626 and FGSG_10384

(Fig. 6a). Unsurprisingly, mutants in which these regulators are disrupted display defective

sexual development (Table S2).

*4.2. Virulence TRNs*

*Fg* is a pathogen of wheat and barley. To understand the transcriptional regulatory circuits in

*Fg* infection, we built a network of gene modules and the top regulators associated with

virulence using an AI threshold of > 0.3 (Fig. 6b). In total, 30 modules were associated with

virulence (Fig. 5c); 5 of these (M06, M24, M49, M47 and M16) had the highest association

with virulence, which contributed 25% to 45% of their phenotype associations. Four of the 30

modules, FGSG_04220, FGSG_07052, FGSG_08028 and FGSG_10384, were hub regulators

of virulence. FGSG_08028, the top regulator for M06, M24 and M49, activated genes of the

three modules during wheat plant infection. The previous report that a disruption mutant of

FGSG_08028 loses pathogenicity but appears normal with respect to other phenotypes (Son

*et al.*, 2011) and our prediction that FGSG_08028 participates only in the virulence-

associated TRN (Fig. 6b) suggest that this TF might exclusively regulate virulence. Notably,

both positive and negative regulators were found in the virulence TRNs. For instance,

FGSG_13314 and FGSG_00318, which are essential for pathogenicity, were the top two

regulators positively regulating M47 genes under plant infection conditions (Fig. S1-47).

Mutants of these TFs display abnormal development of perithecia and ascospores, consistent

with their presence in the sexual reproduction TRN (Fig. 6a). On the other hand, the top two

regulators of M16, FSGG_07928 and FGSG_09019, negatively regulate M16 gene

expression during plant infection. However, disruption mutants of these TFs are

nonpathogenic to wheat. GO enrichment of the M16 genes suggested that the module is

highly enriched in anti-apoptosis processes. Loss of the function of its two main suppressors

might contribute to decreased fungal apoptosis during infection, which is generally

considered to be beneficial to pathogens during infection (Shlezinger *et al.*, 2011). We further

compared our predicted regulators with 70 existing genes reported in PHI-base whose

mutants showed reduced or lost virulence (Urban M et al., 2017) to identify overlapped

genes. Among these 70 virulence-associated genes reported in PHI-base, 63 were found in

our total predicted regulators,  44 of which were included in virulence TRNs modules (Table

S13), suggesting our network prediction has resulted in a virulence regulatory network that

captured the majority (63%) of the known *Fg* virulence-associated genes in public domain.

*4.3. Mycotoxin TRNs*

*Fg* produces toxic secondary metabolites, including DON and ZEA, that are harmful to

livestock and to humans who consume the mycotoxin-containing maize products. The

regulatory networks involved in fungal mycotoxin production are still poorly understood.

Here, we found 31 TRN modules that were associated with mycotoxin production under the

AI index threshold of 0.3 (Fig. 6c); six of these (M02, M12, M36, M46 and M48) were

highly associated with mycotoxins in 55% ~ 85% of all phenotypic associations (Fig. 5c).

The majority of the Tri-cluster was captured by another mycotoxin-associated module,

M46. Several regulators, including FGSG_04220, FGSG_05399, FGSG_00750,

FGSG_10517, FGSG_01173, FGSG_03538 and FGSG_10384, acted as hub regulators in

mycotoxin-associated modules. Not surprisingly, disruption mutants of these regulators show

abnormal production of DON and ZEA (Table S2). Two of them, FGSG_05399 and

FGSG_10157, are negative regulators of DON and ZEA, respectively. In addition, we

predicted FGSG_05399 as the top negative regulator of M02 in the mycotoxin production

and plant infection stages. Interestingly, GO enrichment suggested that the M02 genes are

highly enriched for isoprenoid metabolism, which generates the precursors of trichothecenes.

Although M02 did not include genes in the Tri-cluster, it contained a deacetylase gene,

FGSG_03544, that is located near the Tri-cluster and is co-expressed with Tri-cluster genes

(Sieber *et al.*, 2014).

**5. *Fg* module networks have core and lineage-specific (LS) compartments**

A key question in evolution is how newly acquired genes or chromosomes integrate with

existing or ancestral genomes in an organism and how the regulatory networks carried by two

distinct compartments, i.e., the ancestral and the novel genetic materials, are accommodated

to ensure the stability of the newly formed genome and its regulation during speciation.

Previous comparative genomic analysis suggested that the *Fg* genome can be roughly divided

into core and LS (Guo *et al.*, 2016). To determine whether our predicted *Fg* module networks

have a compartmentalized organization, we applied Fisher's exact test to the genes in each

module and identified modules significantly enriched for core and LS genomic regions. We

found that eight modules (M06, M09, M25, M27, M34, M38, M46 and M48) showed

significant enrichment of genes from LS regions, while 24 modules were enriched with genes

from core regions (Fig. 7). Phenotypically, these eight LS modules were primarily associated

with mycotoxin production, virulence and sexual reproduction (Fig. 6). Interestingly, among

the 117 predicted regulators, only ten were located in LS regions. However, we found that

four of them (FGSG_01327, FGSG_03536, FGSG_03538, and FGSG_08028) were

exclusively associated with LS modules and that six were associated with other modules.

Fisher's exact test showed these ten LS regulators showed a significant bias of regulator-

module association towards core or LS genomic regions (P-value < 0.05). FGSG_03536

(*Tri6*) and FGSG_03538 (*Tri10*) were the top two regulators of the LS module M46 (Fig. 4a),

and FGSG_08028 was the top regulator of the LS modules M06 and M38. These four LS

regulators, together with their associated LS modules, might be horizontally acquired and

may have contributed to the speciation of *Fg* via specifically regulating the fungal

biosynthesis of specific mycotoxins or plant infections. Lastly, a reciprocal regulation was

found for regulators and modules across the LS and core genomes (Table S14), suggesting

that there is communication between core and LS genomic regions and reflecting the

evolutionary signature of regulatory programs involving both vertically and horizontally

acquired gene regulatory elements.

**DISCUSSION**

GRNs are the heart of every cellular function. Given its high complexity, the complete

reconstruction of GRNs in multicellular organisms remains a daunting task. Traditional

reductionist approaches and gene co-expression analysis have failed to solve this challenge in

*Fg*. Here, we used a module network algorithm (Segal *et al.*, 2003) to reconstruct *Fg* TRNs.

The module network algorithm, which is a probabilistic method built on a Bayesian networks

model, is ideal for dealing with noisy data such as gene expression profiles. Importantly, its

strong performance in predicting dynamic and condition-specific gene regulation from large

transcriptome datasets has been demonstrated (Segal *et al.*, 2003). Furthermore, this TRN

inference used phenotype-associated TFs as candidate regulators, allowing us to pinpoint the

TRNs that directly mediate various fungal phenotypes. In contrast, previous GRNs were

comprehensive but coarse, focusing on the master regulatory networks that control generic

cellular functions. Lastly, we used phenomic data from TF knockout mutants as well as

transcriptome data to discover phenotypically specific regulatory networks.

Given such technical advantages, the TRNs inferred here present two major improvements

compared to previous GRNs. First, the TRNs predicted in this work improved the network

resolution compared to our previous GRNs which inferred 120 regulators, each regulating

329 genes on average (Guo et al., 2016). In comparison, each module in the new TRNs

contains an average of 268 genes and 7 regulators, significantly improving the network

resolution. Previous GRNs predicted 44 transcription factors as regulators, while the TRNs

yielded 117 TFs as regulators, suggesting the new TRNs found more regulators and therefore

increased the resolution of regulatory networks involving TFs. Second, the TRNs are

dynamic networks that specify the conditions under which a particular regulation occurs,

unlike previous static GRNs. For example, we found that the TRNs of this work and previous

GRNs shared eight regulators (Table S15). For these eight regulators, this TRN produced a

dynamic network model that specifies the conditions under which a particular regulation

occurs, while in previous static GRNs no such information existed. Third, we developed a

new computational method for establishing associations between key fungal phenotypes and

predicted network modules, while previous co-expression networks and GRNs lack any such

association. The module-phenotype associations suggest that cellular phenotypes are most

often controlled by complex networks of gene modules. Our association mining analysis also

revealed major groups of modules that contribute primarily to specific fungal phenotypes,

suggesting that biological processes in fungi are controlled by multiple intertwining gene

modules. Finding the links between regulatory modules and phenotypes enriches our

understanding of how fungal cells control their activities and, importantly, informs targeted

approaches for suppressing specific gene modules and key regulators for disease control.

The TRN inference in this study significantly improves our understanding of transcriptional

regulation in *Fg*. In previous functional studies, many TFs critical for fungal biology have

been characterized. However, little is known about what target genes they regulate, which

networks they use to regulate cellular functions, and, more importantly, when the regulation

occurs and how it changes when the environment is altered. We identified 49 TRN modules

in *Fg*. Each module represents a regulatory program in which a set of TFs regulates a number

of target genes, and the hierarchical organization of these TFs reflects their influence on the

module. Our biological validations suggested that our TRN inference achieved overall high

performance and that it successfully associated TFs and their target genes. Importantly, our

inference reveals the conditional specificity, the direction (positive vs. negative) and the

strength of gene regulation, factors that are not addressed in previous *Fg* network studies.

*Fg* is a pathogenic fungus that is capable of producing harmful mycotoxins. Sexual

reproduction plays a vital role in generating inocula for fungal infection. Therefore,

understanding the regulatory networks that control virulence, sexual reproduction and

mycotoxin production is a priority for research on *Fg* systems biology. For the first time, we

identified TRN subnetworks in filamentous fungi that control three *Fg* phenotypes. Unlike

studies that have focused on the contribution of a single gene or a set of co-expressed genes

to phenotypes, we reconstructed TRN subnetworks that are strongly associated with each

phenotype. The three subnetworks depicted the dynamic regulation of gene modules that are

highly associated with each of the phenotypes. This work provides the best knowledge to date

of the transcriptional regulation involved in these processes in *Fg* and lays an important

foundation for further systems biology studies in the fungus. We previously reported that

there appears to be a compartmentalization of *Fg* GRNs in core and LS genomic regions. The

TRNs inferred here confirmed such compartmentalization but also provided novel insights

into how the regulatory modules are organized into the core and LS compartments based on

our identification of 8 and 24 modules that are significantly enriched in target genes from the

LS and core genome regions, respectively. The top regulators in four of the eight LS-enriched

modules are located in the LS genome, suggesting that the regulatory circuits associated with

LS modules were very likely acquired with LS genome regions but that over time at least part

of these acquired circuits were integrated into the ancestral circuits carried by the fungus.

Overall, we reconstructed dynamic TRNs from a large scale of *Fg* transcriptomic data using a

module network algorithm. The overall accuracy of the network inference is high as

demonstrated by validation using prior knowledge including functional annotation, TF

binding site enrichment analyses and finally the deletion mutant transcriptome profiles in the

well-established Tri-cluster regulatory model. Despite the progress reported here, this

network remains an early step toward obtaining a complete understanding of authentic TRNs

and eventually achieving its predictive power. As high-throughput sequencing technologies

rapidly develop and costs continue to decrease, generation of additional large-scale genomic

data such as RNA-seq and ChIP-seq will be needed to improve the network model and realize

its predictive power in the future.

**ACKNOWLEDGMENTS**

This project was supported by the National Natural Science Foundation of China (31701739,

31671372) and National Key R&D Program of China (Grant Nos. 2018YFC0910400 and

2017YFC0907500). L.G. is also supported by a China Postdoctoral Foundation Grant

(2017M623188) and the Fundamental Research Fund of Xi'an Jiaotong University

(1191329155).

**AUTHOR CONTRIBUTIONS**

L.G. and K.Y. conceived and designed the project. L.G. and M.J. performed the network

inference and validations. M.J. conducted the module-phenotype association analysis. L.G.

and M.J. prepared the figures and tables. K.Y., L.G. and M.J. wrote the manuscript. L.G. and

M.J. contributed equally to this work. All authors read and approved the manuscript.

**REFERENCES**

**Agrios GN. (2005).** *Plant Pathology*. San Diego: Academic Press

**Bailey TL, Boden M, Buske FA, Frith M, Grant CE, Clementi L, Ren J, Li WW, Noble**

**WS. (2009).** MEME SUITE: tools for motif discovery and searching. *Nucleic Acids*

*Research* **37**(Web Server issue): W202-W208

**Bastian M, Heymann S, Jacomy M (2009)**. Gephi: an open source software for exploring

and manipulating networks.In E A, M H, T F, NS G, N N, BL T. *International AAAI*

*Conference on Web and Social Media*. Palo Alto, CA, USA: The AAAI Press. 361–

362.

**Brown NA, Evans J, Mead A, Hammond-Kosack KE. (2017).** A spatial temporal analysis

of the *Fusarium graminearum* transcriptome during symptomless and symptomatic

wheat infection. *Molecular Plant Pathology* **18**(9): 1295-1312

**Carapito R, Hatsch D, Vorwerk S, Petkovski E, Jeltsch J-M, Phalip V. (2008).** Gene

expression in *Fusarium graminearum* grown on plant cell wall. *Fungal Genetics and*

*Biology* **45**(5): 738-748

**Emmert-Streib F, Dehmer M, Haibe-Kains B. (2014).** Gene regulatory networks and their

applications: understanding biological and medical problems in terms of networks.

*Frontiers in Cell and Developmental Biology* **2**(38)

**Friedman N, Linial M, Nachman I, Pe'er D. (2000).** Using Bayesian Networks to analyze

expression data. *Journal of Computational Biology* **7**(3-4): 601-620

**Guo L, Zhao G, Xu J-R, Kistler HC, Gao L, Ma L-J. (2016).** Compartmentalized gene

regulatory network of the pathogenic fungus *Fusarium graminearum. New*

*Phytologist* **211**(2): 527-541

**Hohn TM, Krishna R, Proctor RH. (1999).** Characterization of a transcriptional activator

controlling trichothecene toxin biosynthesis. *Fungal Genetics and Biology* **26**(3): 224-

235

**Kim HD, Shay T, O'Shea EK, Regev A. (2009).** Transcriptional regulatory circuits:

predicting numbers from alphabets. *Science* **325**(5939): 429-432

**Kim M, Zhang H, Woloshuk C, Shim W, Yoon B. (2018).** Computational prediction of

pathogenic network modules in *Fusarium verticillioides*. *IEEE/ACM Transactions on*

*Computational Biology and Bioinformatics* **15**(2): 506-515

**Koch A, Biedenkopf D, Furch A, Weber L, Rossbach O, Abdellatef E, Linicus L,**

**Johannsmeier J, Jelonek L, Goesmann A, et al. (2016).** An RNAi-Based control of

*Fusarium graminearum* infections through spraying of long dsRNAs involves a plant

passage and is controlled by the fungal silencing machinery. *PLOS Pathogens* **12**(10):

e1005901

**Kong X, van Diepeningen AD, van der Lee TAJ, Waalwijk C, Xu J, Xu J, Zhang H,**

**Chen W, Feng J. (2018).** The *Fusarium graminearum* histone acetyltransferases are

important for morphogenesis, DON biosynthesis, and pathogenicity. *Frontiers in*

*Microbiology* **9**(654)

**Leslie JF, Summerell BA. (2006).** *Fusarium laboratory manual*: Blackwell Publishing

**Liu J-B, Wang Y-M, Peng S-Q, Han G, Dong Y-S, Yang H-Y, Yan C-H, Wang G-Q.**

**(2007).** Toxic effects of Fusarium mycotoxin butenolide on rat myocardium and

primary culture of cardiac myocytes. *Toxicon* **50**(3): 357-364

**Liu X, Tang W-H, Zhao X-M, Chen L. (2010).** A network approach to predict pathogenic

genes for *Fusarium graminearum*. *PLOS ONE* **5**(10): e13021

**Lysenko A, Urban M, Bennett L, Tsoka S, Janowska-Sejda E, Rawlings CJ, Hammond-**

**Kosack KE, Saqi M. (2013).** Network-based data integration for selecting candidate

virulence associated proteins in the cereal infecting fungus *Fusarium graminearum*.

*PLOS ONE* **8**(7): e67926

**Lysoe E, Pasquali M, Breakspear A, Kistler HC. (2011).** The transcription factor *FgStuAp*

influences spore development, pathogenicity, and secondary metabolism in *Fusarium*

*graminearum. Molecular plant-microbe interactions* **2011 v.24 no.1**(no. 1): pp. 54-50

**Ma LJ, Does HC, Borkovich KA, Coleman JJ, Daboussi MJ, Pietro A, Dufresne M,**

**Freitag M, Grabherr M, Henrissat B. (2010).** Comparative genomics reveals

mobile pathogenicity chromosomes in Fusarium. *Nature* **464**

**Ma LJ, Geiser DM, Proctor RH, Rooney AP, O'Donnell K, Trail F, Gardiner DM,**

**Manners JM, Kazan K. (2013).** Fusarium pathogenomics. *Annu Rev Microbiol* **67**:

399-416

**Martin Urban, Alayne Cuzick, Kim Rutherford, Alistair Irvine, Helder Pedro, Rashmi**

**Pant, Vidyendra Sadanadan, Lokanath Khamari, Santoshkumar Billal, Sagar**

**Mohanty, Kim E. Hammond-Kosack. (2017).** PHI-base: a new interface and further

additions for the multi-species pathogen–host interactions database. *Nucleic Acids*

*Research*, Volume 45, Issue D1, January 2017, Pages D604–D610

**Nasmith CG, Walkowiak S, Wang L, Leung WW, Gong Y, Johnston A, Harris LJ,**

**Guttman DS, Subramaniam R. (2011).** Tri6 is a global transcription regulator in the

phytopathogen *Fusarium graminearum. PLoS Pathog* **7**(9): e1002266

**Pan Y, Liu Z, Rocheleau H, Fauteux F, Wang Y, McCartney C, Ouellet T. (2018).**

Transcriptome dynamics associated with resistance and susceptibility against

fusarium head blight in four wheat genotypes. *BMC Genomics* **19**(1): 642-642

**Pe'er D, Tanay A, Regev A. (2006).** MinReg: a scalable algorithm for learning parsimonious

regulatory networks in yeast and mammals. *Journal of Machine Learning Research* **7**:

167-189

**Ruepp A, Zollner A, Maier D, Albermann K, Hani J, Mokrejs M, Tetko I, Güldener U,**

**Mannhaupt G, Münsterkötter M, et al. (2004).** The FunCat, a functional annotation

scheme for systematic classification of proteins from whole genomes. *Nucleic Acids*

*Research* **32**(18): 5539-5545

**Segal E, Shapira M, Regev A, Pe'er D, Botstein D, Koller D, Friedman N. (2003).**

Module networks: identifying regulatory modules and their condition-specific

regulators from gene expression data. *Nat Genet* **34**(2): 166-176

**Seong KY, Pasquali M, Zhou X, Song J, Hilburn K, McCormick S, Dong Y, Xu JR,**

**Kistler HC. (2009).** Global gene regulation by Fusarium transcription factors Tri6

and Tri10 reveals adaptations for toxin biosynthesis. *Mol Microbiol* **72**(2): 354-367

**Shannon P, Markiel A, Ozier O, Baliga NS, Wang JT, Ramage D, Amin N, Schwikowski**

**B, Ideker T. (2003).** Cytoscape: a software environment for integrated models of

biomolecular interaction networks. *Genome Research* **13**(11): 2498-2504

**Shlezinger N, Minz A, Gur Y, Hatam I, Dagdas YF, Talbot NJ, Sharon A. (2011).** Anti-

apoptotic machinery protects the necrotrophic fungus Botrytis cinerea from host-

induced apoptotic-like cell death during plant infection. *PLOS Pathogens* **7**(8):

e1002185-e1002185

**Shmulevich I, Dougherty ER, Kim S, Zhang W. (2002).** Probabilistic Boolean networks: a

rule-based uncertainty model for gene regulatory networks. *Bioinformatics* **18**(2):

261-274

**Sieber CMK, Lee W, Wong P, Münsterkötter M, Mewes H-W, Schmeitzl C, Varga E,**

**Berthiller F, Adam G, Güldener U. (2014).** The *Fusarium graminearum* genome

reveals more secondary metabolite gene clusters and hints of horizontal gene transfer.

*PLOS ONE* **9**(10): e110311-e110311

**Son H, Lim JY, Lee Y, Lee Y-W. (2016).** Utilization of a conidia-deficient mutant to study

sexual development in *Fusarium graminearum*. *PLOS ONE* **11**(5): e0155671

**Son H, Seo YS, Min K, Park AR, Lee J, Jin JM, Lin Y, Cao P, Hong SY, Kim EK, et al.**

**(2011).** A phenome-based functional analysis of transcription factors in the cereal

head blight fungus, *Fusarium graminearum. PLoS Pathog* **7**(10): e1002310

**Tag AG, Garifullina GF, Peplow AW, Ake C, Phillips TD, Hohn TM, Beremand MN.**

**(2001).** A novel regulatory gene, *Tri10*, controls trichothecene toxin production and

gene expression. *Applied and Environmental Microbiology* **67**(11): 5294

**Zhao J, Zhou Y, Zhang X, Chen L. (2016).** Part mutual information for quantifying direct

associations in networks. *Proceedings of the National Academy of Sciences* **113**(18):

5130

**Zhao XM, Zhang XW, Tang WH, Chen L. (2009).** FPPI: *Fusarium graminearum* protein-

protein interaction database. *J Proteome Res* **8**(10): 4714-4721

**FIGURE LEGENDS**

**Figure 1. System framework and main steps in constructing a modularized gene**

**regulatory network of *Fusarium graminearum***. Combined input consisting of gene

expression data and information on candidate regulators was provided to the module

networks algorithm (Segal *et al.* 2003). The inferred modules were then subjected to a robust

validation process involving functional annotations and transcription factor binding site

analysis. Then, the modules were analyzed for their phenotypic associations using an in-

house computational method. Finally, the genome compartmentalization of the regulatory

network was analyzed to provide evolutionary insights into the gene circuits.

**Figure 2. Overview of the module networks predicted for *Fusarium graminearum*.**

**(a).** Overview of inferred *F. graminearum* regulatory modules. The columns in the heatmap

represent *Fg* genes, and the rows represent the experimental conditions in the gene expression

data. Modules are delimited by vertical yellow lines. Red and blue represent gene activation

and suppression, respectively. **(b).** Distribution of the number of target genes per module. Y

axis represents the kernel density of target genes per module (X-axis). **(c).** Distribution of the

number of regulators per module. Y axis represents the kernel density of regulators per

module (X-axis). **(d).** An unweighted network of modules and regulators; the blue and red

nodes represent regulators and modules, respectively. For clarity of visualization, the prefixes

for regulator gene ID ("FGSG_") and module ("M") are omitted.

**Figure 3. Gene Ontology (GO) annotation and transcriptional factor (TF) binding**

**motifs enrichment in the predicted regulatory modules.** GO enrichment was conducted for

genes in each module, and major function associations were then assigned to each module

using the most enriched GO terms. Each module is annotated with 55 main functional

descriptions (P-value < 0.05). The *MEME* algorithm was used to find TF motif enrichment in

the 500-bp region upstream of the *F. graminearum* genes sequence for each module (E value

< 1e-5) followed by motif comparison with the budding yeast TF motif database using

*Tomtom* (Materials and Methods). The heatmap on the left shows the significantly enriched

GO terms per module; the color bar is scaled according to the P-value. The heatmap in the

middle shows the motif enrichment results sorted into a total of 56 motifs; the color bar is

scaled according to the E-value. The heatmap on the right shows the phenotype associations

of each module based on association index analysis.

**Figure 4. High accuracy of network inference highlighted by its capture of the**

**trichothecene biosynthetic gene cluster. (a).** Diagram of the trichothecene biosynthetic

gene cluster (*Tri*-cluster) in *F. graminearum*. **(b).** Regulation program of the predicted

module M46. The predicted regulators are arranged in a decision-tree structure that specifies

the conditions under which the regulation occurs; these are delineated by the yellow vertical

dotted lines. The heatmap underneath the tree shows the expression values of genes (rows)

under different experimental conditions (columns). The experimental conditions are grouped

into several major groups that are underlined; these are labeled at the bottom of the heatmap.

The black empty box highlights all the *Tri*-cluster genes (labeled on the left) predicted in this

module. **(c).** Venn diagram displaying the overlap of M46 genes with the previously

published down-regulated genes of *ΔTri6* and *ΔTri10* mutants as well as *Tri6*-binding genes

identified by ChIP-seq. **(d).** Dot plot showing the functional enrichment of M46 genes. The Y

axis represents the significantly enriched GO terms. The X axis represents the P value (-

Log10 transformed). The size of each dot is proportional to the gene ratio (Test/Reference),

which was calculated as the number of genes in M46 (Test) divided by the number of genes

in the *F. graminearum* genome (Reference) that are annotated with the GO term.

**Figure 5. Computational analysis of module-phenotype associations. (a).** Schematic

demonstration of the computational method used to mine the module-phenotype association.

The computational method calculates the association index based on the regulator's

phenotype, the regulator's hierarchical position on the regulation tree, and the proportion of

activated conditions in all experiments. **(b).** Module-phenotype association network based on

the association index. The outer circle nodes represent modules, and the inner nodes represent

phenotypes: M (mycelial growth), C (conidation), P (pigmentation), V (virulence), Toxin

(DON and ZEA production), S (sexual reproduction) and St (stress response). The edges

denote the associations between modules and phenotypes; the line sizes are proportional to

AI of the association. **(c).** Correlation analysis of all predicted modules using Pearson

correlation coefficients (PCC) based on the calculated association index. The modules are

clustered using hierarchical clustering, and a heatmap of the PCCs (the scale bar denotes the

PCC range) is then produced. The stacked bar graph adjacent to the heatmap summarizes the

phenotype association within each module based on the association index.

**Figure 6. *Fusarium graminearum* subnetworks involved in the regulation of key**

**phenotypes:** sexual reproduction **(a)**, virulence **(b)**, mycotoxin production **(c)**. In each

network, the edges represent the connections between the two nodes, which in turn represent

the contributions of different regulators to the module. The red line represents a positive

effect of the regulator on the corresponding phenotype, the green line represents a negative

effect, and the thickness and color depth of the connection indicates the strength of the effect.

**Figure 7. Compartmentalization of module networks into core and lineage-specific**

**genome regions of *Fusarium graminearum***. Each of 49 *F. graminearum* modules was

tested for enrichment of LS and core genes using two-tailed Fisher's exact test. The dot plot

summarizes the enrichment results; the X axis shows the module, the Y axis shows the test P-

value for each module (-log10 transformed), and the solid horizontal line represents the

threshold (P-value < 0.05). The color of the dot denotes the enriched compartment (green:

LS; red: core; blue: not enriched) for each module. The size of the dot is proportional to the

ratio of LS to core genes in each module.

**Supplementary Information**

**Figure S1.** Regulatory programs for all 49 regulatory modules in *F. graminearum*.

**Table S1.** Gene expression data sets used to predict module networks of *F. graminearum* in

this work.

**Table S2.** Phenotype summary of 170 transcription factors as candidate regulators extracted

from the originally reported by Son *et al.* 2011 in PLoS Pathogens.

**Table S3.** Weight of each regulator (R) in each module.

**Table S4.** Digital representation of various regulator phenotypes.

**Table S5.** Lists of target genes in each module.

**Table S6.** Summary of 49 regulatory modules, including information on the number of target

genes, the top regulators, and the top four enriched Gene Ontology terms (P value < 0.05).

**Table S7.** Summary of regulatory module validation results.

**Table S8.** Summary of transcription factor binding site (TFBS) enrichment and conservation

analysis for each module using MEME and *Tomtom*.

**Table S9.** Gene annotation enrichment of each module in 55 main functional categories (P

value < 0.05).

**Table S10.** Consensus functional terms associated with GO enrichment and enriched TF

binding sites for module target genes.

**Table S11.** Summary of *F. graminearum* and *S. cerevisiae* BLASTp results using predicted

top regulators. Only significant hits are listed (E-value < 1e-5).

**Table S12.** Phenotype association analysis for each module.

**Table S13.** Comparison of the virulence TRNs with virulence-associated genes in the PHI-

base database

**Table S14.** Regulatory networks compartmentalization analysis using Fisher's exact test. FS:

*F. graminearum*-specific; Core: Conserved genome. Mix: not enriched.

**Table S15.** Comparing the TRNs with previous *F. graminearum* GRNs（Guo *et al.*, 2016）

**Transcriptome Profiles**

*55 conditions x 13,300 genes*

**TF Phenome Data**

FgTFPD

**170 phenotype-associated TFs**

**INPUT DATA**

Top R1

TRUE

FALSE

Top R2

FALSE    TRUE

0          0          0

*Probablistic modeling*

**Module Networks Algorithm**

*50 iterations*

Gene Ontology

TFBS Analysis

FGSG_01173
FGSG_01350
FGSG_02083
FGSG_03538
FGSG_05399
FGSG_05604
FGSG_07928
FGSG_08626

5'    100    200    300    400    500    3'

*Regulation programs*

**Predicted TRN modules**

**Network Validation**

**Module-Phnotype Association Mining**

**Genome Compartment Analysis**

toxin
C
St
S
V
M
P

Core GRN

LS GRN

Core genome

LS genome

**(a)**

*F. graminearum* Network Modules (1 − 49)

Experiments

**(b)**

Distribution of target genes in each module

Density

Number of target genes

**(c)**

Distribution of regulators in each module

Density

Number of regulators

**(d)**

Regulators
Modules

GO annotations        Enriched TFBS (Yeast Homologs)        Phenotypes

(a) Sexual development TRN

(b) Virulence TRN

(c) Mycotoxin TRN

downregulate
module — regulator
upregulate

weak connection
module — regulator
strong connection