Charting the metabolic landscape of the facultative methylotroph *Bacillus*
methanolicus

Baudoin Delépine[a], Marina Gil López[b], Marc Carnicer[a*], Cláudia M. Vicente[a], Volker F.
Wendisch[b], Stéphanie Heux[a#]

[a] TBI, Université de Toulouse, CNRS, INRA, INSA, Toulouse, France.

[b] Genetics of Prokaryotes, Faculty of Biology & CeBiTec, Bielefeld University, Bielefeld,
Germany

Running Head: Metabolic states of the methylotroph *B. methanolicus*

Address correspondence to SH, heux@insa-toulouse.fr

Post: TBI - INSA de Toulouse, 135 avenue de Rangueil, 31077 Toulouse CEDEX 04; Phone
number: +33 (0)5 61 55 93 55

* Present address: IQS, Universitat Ramon Llull, Via Augusta 390, E-08017, Barcelona, Spain

ACKNOWLEDGEMENT

MetaToul (www.metatoul.fr), which is part of MetaboHub (ANR-11-INBS-0010, www.metabohub.fr) are gratefully acknowledged for their help in collecting, processing and interpreting 13-C NMR and MS data. We would like to thank Marcus Persicke, Maud Heuillet, Edern Cahoreau, Lindsay Periga, Pierre Millard, Gilles Vieira and Jean-Charles Portais for the insights they provided.

FUNDING

This work was supported by ERA-CoBioTech's project C1Pro (ANR-17-COBI-0003-05 and FNR-22023617).

ABSTRACT

Bacillus methanolicus MGA3 is a thermotolerant and relatively fast-growing methylotroph able to secrete large quantities of glutamate and lysine. These natural characteristics make *B. methanolicus* a good candidate to become a new industrial chassis organism, especially in a methanol-based economy. This has motivated a number of omics studies of *B. methanolicus* at the genome, transcript, protein and metabolic levels. Intriguingly, the only substrates known to support *B. methanolicus* growth as sole source of carbon and energy are methanol, mannitol, and to a lesser extent glucose and arabitol. We hypothesized that comparing methylotrophic and non-methylotrophic metabolic states at the flux level would yield new insights into MGA3 metabolism. ^{13}C metabolic flux analysis (^{13}C-MFA) is a powerful computational method to estimate carbon flows from substrate to biomass (i.e. the *in vivo* reaction rates of the central metabolic pathways) from experimental labeling data. In this study, we designed and performed a ^{13}C-MFA of the facultative methylotroph *B. methanolicus* MGA3 growing on methanol, mannitol and arabitol to compare the associated metabolic states. The results obtained validate previous findings on the methylotrophy of *B. methanolicus*, allowed us to characterize the assimilation pathway of one of the studied carbon sources, and provide a better overall understanding of this strain.

IMPORTANCE

Methanol is cheap, easy to transport and can be produced both from renewable and fossil resources without mobilizing arable lands. As such, it is regarded as a potential carbon source to transition toward a greener industrial chemistry. Metabolic engineering of bacteria and yeast able to efficiently consume methanol is expected to provide cell factories that will transform methanol into higher-value chemicals in the so-called methanol economy. Toward that goal, the study of

60 natural methylotrophs such as *B. methanolicus* is critical to understand the origin of their

61 efficient methylotrophy. This knowledge will then be leveraged to transform such natural strains

62 into new cell factories, or to design methylotrophic capability in other strains already used by the

63 industry.

64 **KEYWORDS**

ABBREVIATIONS

^{13}C-MFA: ^{13}C metabolic flux analysis; CID: carbon isotopologue distribution; FBA: Flux Balance Analysis; MS: mass spectrometry; NMR: nuclear magnetic resonance; PTS: phosphotransferase system

Pathways

SBPase: fructose bisphosphate aldolase/sedoheptulose bisphosphatase; TA: fructose bisphosphate aldolase/transaldolase; PPP: pentose phosphate pathway; RuMP: Ribulose Monophosphate Cycle; TCA: Krebs cycle

Reactions

akgdh: 2-oxoglutarate dehydrogenase and 2-oxoglutarate synthase; Araupt: arabitol uptake; detox: linear detoxification pathways; fum: fumarate reductase; glpx: sedoheptulose-bisphosphatase; gnd: phosphogluconate dehydrogenase; hps: 3-hexulose-6-phosphate synthase ; idh: isocitrate dehydrogenase; Manupt: mannitol uptake; mdh: methanol dehydrogenase; pdh: pyruvate dehydrogenase; pfk: 6-phosphofructokinase and fructose-bisphosphatase; pgi: glucose 6-phosphate isomerase ; pgk: glyceraldehyde 3-phosphate dehydrogenase, phosphoglycerate kinase, and phosphoglycerate mutase; phi: 6-phospho-3-hexuloisomerase ; pyk: pyruvate kinase; rpe: ribulose-phosphate 3-epimerase; rpi: ribose 5-phosphate isomerase; rpi: ribulose phosphate isomerase; ta: transaldolase; tkt1, tkt2: transketolases; zwf: glucose-6-phosphate dehydrogenase and 6-phosphogluconolactonase.

Metabolites

Ace: acetate; Aco: aconitate; AKG: 2-oxoglutarate; ala: alanine; Ara: arabitol; arg: arginine; asp: aspartate; Cit: citrate; DHAP: dihydroxyacetone phosphate; Ery4P: erythrose 4-phosphate; For:

formaldehyde; Fru6P: fructose 6-phosphate; FruBP: fructose 1,6-bisphosphate; Fum: fumarate; G3P: 3-phospho-D-glycerate; GAP: glyceraldehyde 3-phosphate; Glc6P: glucose 6-phosphate; glu: glutamate; gly: glycine; GlyOx: glyoxylate; Gnt6P: 6-phosphogluconate; Hex6P: hexulose 6-phosphate; his: histidine; ile: isoleucine; leu: leucine; lys: lysine; Mal: malate; Man: mannitol; MeOH: methanol; met: methionine; OAA: oxaloacetate; PEP: phosphoenolpyruvate; PGA: 2-phospho-D-glycerate; phe: phenylalanine; pro: proline; Pyr: pyruvate; Rib5P: ribose 5-phosphate; Ribu5P: ribulose 5-phosphate; Sed7P: sedoheptulose 7-phosphate; ser: serine; Suc: succinate; thr: threonine; tyr: tyrosine; val: valine; Xyl5P: xylulose 5-phosphate.

1. INTRODUCTION

^{13}C metabolic flux analysis (^{13}C-MFA) has emerged in the last decade as an outstanding experimental method to describe the metabolic states of microorganisms. It has successfully been used to identify new pathways (1), investigate responses to environmental changes (2), improve the titer of cell factories (3), screen strains based on their enzymatic capacity (4), and more generally to provide a better understanding of the metabolism of microorganisms (5) such as methylotrophs (6-8). Briefly, ^{13}C-MFA exploits a metabolic model and ^{13}C-isotope patterns measured from key metabolites to estimate reaction rates consistent with the observed labeling patterns (see (9, 10) for reviews). Specifically, cells are grown on a ^{13}C labeled carbon source, metabolites of the central metabolism or constituents of the biomass such as proteinogenic amino acids are sampled and quenched, and the labeling patterns of metabolites are monitored by mass spectrometry (MS) and/or nuclear magnetic resonance (NMR). With NMR carbon isotopomers are determined and hence positional information is provided whereas with MS, isotopologues are identified. A metabolic model is then used to fit these measurements to theoretical data that are simulated by optimizing reaction flux values from the metabolic model. Assuming mass balance, if the measurements are coherent and the topology of the metabolic model is correct, the experimental and simulated data will converge, yielding the estimated reaction fluxes. Finally, ^{13}C-MFA provides a flux map: a predicted snapshot of the metabolic fluxes through the organism of interest during the experiment, i.e. its metabolic state. ^{13}C-MFA flux maps are conceptually similar to those from flux balance analysis (FBA), a purely *in silico* method that, from a metabolic model (typically at the genome scale), computes the optimal reaction flux distribution to maximize an objective defined in terms of metabolite production, often designed to model cell growth. Importantly, ^{13}C-MFA flux maps are estimates based on experimental data, whereas FBA

flux maps are purely *in silico* predictions, which can be confirmed by gene deletion analysis or

[13]C-MFA for example.

Bacillus methanolicus MGA3 is a gram-positive bacterium that was first isolated in the 1990s

from freshwater marsh soil samples after an enrichment culture on methanol at 55 °C (11). Its

ability to grow quickly and to secrete large quantities of glutamate and lysine in methanol at high

temperature make it a good candidate for biotech applications. Methanol is indeed viewed as a

promising renewable feedstock because of its abundance and low price (12, 13), while high

temperature cultures are less prone to contamination and require less cooling when scaled up

(14). Furthermore, *B. methanolicus* is able to grow in seawater, which is also cheap and abundant

(15). Promising metabolic engineering studies have already established MGA3 as a cell factory

for the heterologous production of cadaverine (16) and GABA (17, 18). While the lack of genetic

tools must have impaired the development of new applications in the past (19), the establishment

of gene expression tools based on theta- and rolling-circle replicating plasmids have made *B.*

methanolicus amenable to the overproduction of amino acids and their derivatives (18), and there

is hope that recent breakthroughs from CRISPR interference (CRISPRi) will stimulate new

innovations (20).

As a facultative methylotroph, *B. methanolicus* MGA3 can also grow on non-methylotrophic

substrates such as ☐-mannitol, ☐-glucose and ☐-arabitol. The metabolic pathways involved in

the uptake of mannitol and glucose have been described (21), as has the organization of genes

involved in mannitol utilization (18). Both substrates enter the cells via a phosphotransferase

system (PTS), respectively as mannitol 1-phosphate and glucose 6-phosphate, and are further

converted to fructose 6-phosphate. Arabitol has recently been characterized as a fourth source of

carbon and energy for *B. methanolicus* (22). The operon responsible for arabitol assimilation was

identified as harboring a PTS system (AtlABC) and two putative arabitol phosphate

dehydrogenases (AtlD and AtlF), whose activities were demonstrated in crude extracts.

However, as the pathway was not completely characterized, it is unclear whether arabitol is

assimilated through arabitol 1-phosphate to xylulose 5-phosphate (Xyl5P), or to ribulose 5-

phosphate (Ribu5P) through arabitol 5-phosphate. It has been suggested that both routes operate

in parallel in *Enterococcus avium* and other gram-positive bacteria (23). A series of omics studies

comparing these carbon sources with methanol have contributed to a better understanding of

B. methanolicus metabolism at the genome (21, 24), transcriptome (21, 22, 25), proteome (26)

and metabolome levels (27, 28). However, a flux level description, which could validate previous

findings and provide new insights into the facultative methylotrophy of *B. methanolicus* and its

associated metabolic states, is still lacking.

In this study, we designed and performed [13]C-MFA of the facultative methylotroph *Bacillus*

methanolicus MGA3 growing on methanol, mannitol and arabitol. Methanol (CH_4O) and

mannitol ($C_6H_{14}O_6$) are the best known carbon sources for this strain (11, 14), with comparable

growth rates; while growth on arabitol ($C_5H_{12}O_5$) is significantly slower (22) and its assimilation

pathway has not yet been fully described. All three carbon sources are probably present in

MGA3's natural habitats, on plant leaves (29) or as plant degradation products (30). With their

wide availability and fast associated growth rate, methanol and mannitol are promising

feedstocks for industrial applications, while arabitol growth allows the facultative methylotrophy

of MGA3 to be studied with a less efficient C source and to finish characterizing its assimilation

pathway.

2. MATERIALS & METHODS

2.1 Strain

B. methanolicus wild-type MGA3 (ATCC 53907) strain was used for metabolic flux analyses. Strains used for cloning and expression are described in the section 2.5.1 and listed in Table 1.

2.2 Non-stationary ^{13}C fluxomics experiment

2.2.1 Culture conditions and parameters

For the carbon source methanol, two batch cultures were performed in 0.5 litre bioreactors (INFORS HT Multifors, The Netherlands) with a working volume of 0.40 litres coupled to a Dycor ProLine Process Mass Spectrometer (AMETEK Process Instruments, USA). The culture medium per litre was: 3.48 g Na$_2$HPO$_4$·12 H$_2$0, 0.606 g KH$_2$PO$_4$, 2.5 g NH$_4$Cl, 0.048 g yeast extract, 1 ml of 1 M MgSO$_4$ solution, 1 ml of trace salt solution, 1 ml of vitamins solution, 0.05 ml Antifoam 204 and 150 mM of methanol. The trace salt solution per litre was: 5.56 g FeSO$_4$·7 H$_2$O, 0.027 g CuCl$_2$·2 H$_2$O, 7.35 CaCl$_2$·2 H$_2$O, 0.040 g CoCl$_2$·6 H$_2$O, 9.90 g MnCl$_2$·4 H$_2$O, 0.288 g ZnSO$_4$·7 H$_2$O and 0.031 g H$_3$BO$_3$. The vitamin solution per litre was: 0.10 g □-biotin, 0.10 g thiamine·HCl, 0.10 g riboflavin, 0.10 g pyridoxine·HCl, 0.10 g pantothenate, 0.10 g nicotinamide, 0.02 g p-aminobenzoic acid, 0.01 g folic acid, 0.01 g vitamin B12 and 0.01 g lipoic acid. The pre-cultures were grown in two half litre shake flasks containing 150 ml of the pre-culture medium and inoculated with cryostock of *B. methanolicus* wild-type MGA3 cells. The cultures were grown overnight at 50 °C under shaking at 200 rpm, and used to inoculate the reactors. The aeration rate of 1 vvm was controlled by a mass flow meter (INFORS HT Multifors, The Netherlands) and pO2 were maintained above 25 % throughout all cultures.

188 Temperature, pH and stirring speed were maintained at 50 °C, pH 6.8 (with KOH 1 M) and 800

189 rpm, respectively. The N_2, O_2, Argon, CO_2, ^{13}C-CO_2 and methanol concentrations in the

190 bioreactors off-gas were measured on-line with the mass spectrometer.

191 To perform the pulse of tracer, 100 mM of ^{13}C-methanol (99 % ^{13}C; Euriso-Top, France) were

192 added to the cultures at an OD_{600} of 2.5. Growth curves are available in Supplementary Data 1.

2.2.2 Quantification of cells and supernatant NMR analysis

For determination of the dry weight of cells, a conversion factor of 0.389 g/l (dry weight) of cells

per OD_{600} unit was used. Supernatant samples were taken to analyse substrate methanol

consumption as well as by-product formation by subtracting 1 ml of culture and centrifuged it at

$13000 \times g$ for 60 s. Thereafter, supernatant was collected and stored until analysis at -20 °C.

Supernatant analysis was performed by ^1H 1D-NMR at 292 °K, using a 30° pulse and a

relaxation delay of 20 s, with an Avance 800 MHz spectrometer (Bruker, Germany). Deuterated

trimethylsilyl propionate (TSP-d4) was used as an internal standard for quantification.

2.2.3 Sampling and MS analysis of intra and extracellular pool sizes

When the cultures reached an OD_{600} of 2, metabolome samples were collected using the

optimized method described by (28). Briefly, total broth quenching with correction for

metabolites in the extracellular medium was performed in quadruplicates to assess the metabolite

pool sizes in the two cultivations performed. Metabolites pool sizes were quantified (Fig. S1) by

ion chromatography tandem mass spectrometry (IC-MS/MS) using cell extract of *Escherichia*

coli, cultivated on 99 % [$^{13}C_6$] glucose (Euroisotop, France), as internal standard (31). Liquid

anion exchange chromatography was performed as described previously (32).

2.2.4 Sampling and MS analysis of labeled metabolites

Label enrichments in the intracellular metabolites were followed after performing a pulse of 100 mM ^{13}C-methanol at an OD_{600} 2.5. Whole broth (internal + external pools; WB) and culture filtrate (external pools; CF) were sampled to indirectly track label incorporation in the intracellular metabolites. Specifically, 13 WB and 3 CF samples were collected in 3.5 min in each bioreactor. Exact sampling times can be seen in Supplementary Data 1. IC-MS/MS quantification was used to analyse the isotopologues of each metabolite as described by (32). The metabolites analysed were PEP, Rib5P+Ribu5P+Xyl5P, Sed7P, Gnt6P, Glc6P, FruBP, Fru6P, Gly3P, 13PG, G3P+PGA, Cit, Aco, Fum, Mal and Suc. After manual peak integration, the raw peak areas were corrected for the contribution of all naturally abundant isotopes using IsoCor software (33). Some cross-contamination was found in the isotopologues M+4 of Aco and M+2 and M+3 of Gnt6P that were subsequently removed from the analysis (Supplementary Data 1).

Additionally, the exact ratio between ^{13}C-methanol and ^{12}C-methanol after the pulse was measured by ^{1}H 1D-NMR as well as the evolution of $^{12}CO_2$ and $^{13}CO_2$ by the mass gas analyser.

2.2.5 Mass balance

Experimental data consistency of the measured rates was verified using standard data reconciliation procedures under the elemental mass balance constraints (34, 35). The biomass elemental composition used in the reconciliation procedure was taken from the closely related non-methylotrophic bacterium *Bacillus subtilis*, $CH_{1.646}N_{0.219}O_{0.410}S_{0.005}$ (35, 36). The ashes content were considered to be 6 % of the dry cell weight, average value obtained from different microorganisms (i.e. *Escherichia coli, Aspergillus niger, Penicicillium chrysogenum, Klebsiella aerogenes* (37)). After no proof of mismatch in the measurements, a better estimation of the physiological parameters were obtained as described by (34).

2.3 Stationary ^{13}C fluxomics experiments

2.3.1 Culture conditions

For the carbon sources mannitol and arabitol, the culture medium composition remained unchanged. The experiments were performed in 500 ml baffled shake flasks using 40 ml of media, and cells were grown at 50 °C and 200 rpm. Pre-cultures contained yeast extract and 15 mM unlabeled mannitol or arabitol. They were inoculated as stated previously and grown overnight. The pre-cultures were used to inoculate triplicate main cultures to an OD_{600} of 0.05, after centrifugation and re-suspension in media without yeast extract. The media to perform the ^{13}C MFA contained 15 mM [1-^{13}C] mannitol, 15 mM [5-^{13}C] arabitol or 15 mM of a mixture of 10 % [1-^{13}C] arabitol and 90 % [2-^{13}C] arabitol (99 % ^{13}C; Omicron Biochemicals, Inc., South Bend, IN, USA). An experimentally determined conversion factor of 0.22 g/litre (dry weight) of cells per OD_{600} unit was used. Growth curves are available in Supplementary Data 1.

2.3.2 Measurements of proteinogenic amino acids ^{13}C-isotopologues

Mannitol (resp. arabitol) cultures were sampled around 10 h (resp. 30 h) once they reached an OD_{600} of 1.3. The pellets obtained from the cellular extract were hydrolyzed 15 h at 110 °C with 500 µl HCL 6N. Samples were evaporated and washed twice with 500 µl of ultrapure water, evaporated to dryness, resuspended (625 µl, water), and diluted (1/1000, water) for the mass spectrometry analysis.

Amino acids were separated on a PFP column (150 × 2.1 mm i.d., particle size 5 µm; Supelco Bellefonte, PEN, USA). Solvent A was 0.1 % formic acid in H_2O and solvent B was 0.1 % formic acid in acetonitrile at a flow rate of 250 µL/min. The gradient was adapted from the method used

by (38). Solvent B was varied as follows: 0 min, 2 %; 2 min, 2 %; 10 min, 5 %; 16 min, 35 %; 20

min, 100 %; and 24 min, 100 %. The column was then equilibrated for 6 min at the initial

conditions before the next sample was analyzed. The volume of injection was 20 µL.

High-resolution experiments were performed with an Ultimate 3000 HPLC system (Dionex, CA,

USA) coupled to an LTQ Orbitrap Velos mass spectrometer (Thermo Fisher Scientific, Waltham,

MA, USA) equipped with a heated electrospray ionization probe. MS analyses were performed

in positive FTMS mode at a resolution of 60 000 (at 400 m/z) in full-scan mode, with the

following source parameters: the capillary temperature was 275 °C, the source heater

temperature, 250 °C, the sheath gas flow rate, 45 a.u. (arbitrary unit), the auxiliary gas flow rate,

20 a.u., the S-Lens RF level, 40 %, and the source voltage, 5 kV. Isotopic clusters were

determined by extracting the exact mass of all isotopologues, with a tolerance of 5 ppm.

Experimental carbon isotopologue distributions (CIDs) of alanine, glycine, valine, serine,

threonine, phenylalanine, aspartate, glutamate, histidine, isoleucine, leucine, lysine, arginine,

tyrosine, proline and methionine were obtained after correction of raw MS data for naturally

occurring isotopes other than carbon, using IsoCor (33).

Careful inspection of the CIDs revealed an overall excellent reproducibility between both the

technical and biological replicates (Supplementary Data 1). However, M+0 of valine and M+0-

M+1 of glycine had a higher variability which could be due to a signal closer to the noise level.

2.3.3 NMR measurements

Concentrations in supernatants were measured by [1]H 1D-NMR at 290 °K, using a 30° angle

pulse and a presaturation of water signal was applied during a relaxation delay of 8 s. TSP-d4

was used as internal standard for calibration and quantification.

The measurement of isotopomers and specific enrichments of targeted biomass components were

performed using the same samples used for proteinogenic amino acids ^{13}C-isotopologues MS

analysis, redried and suspended in 200 μl D_2O (0.1 % DCl). The positional isotopomer

distribution of alanine C2 and C3 was extracted from the analysis of ^{13}C–^{13}C couplings in 2D

^{1}H–^{13}C HSQC experiments (39). The carbon isotopic enrichments of alanine (C2 and C3) and

histidine (C2 and C5) were extracted from the analysis of ^{1}H–^{13}C couplings using 2D-zero

quantum filtered-TOCSY (ZQF-TOCSY) (40).

NMR spectra were collected on an Avance III 800 MHz spectrometer (Bruker, Germany),

equipped with a 5mm z-gradient QPCI cryoprobe. Every acquisition 1D, 2D and absolute

quantification were performed on ^{1}H-1D spectra using TopSpin 3.5 (Bruker, Germany).

2.4 Models and simulations

2.4.1 Metabolic Flux Analysis software

All simulations needed for Metabolic Flux Analysis (MFA) on methanol, mannitol and arabitol

were performed with *influx_si* software v4.4.4 (41), either in stationary or non-stationary mode.

influx has the advantage to allow for the integration of labeling data coming from different

experimental setups (MS, NMR) and to support several integration strategies (stationary, non-

stationary, parallel labeling).

All input and output files needed for reproducibility are available in an archive in Supplementary

Data 2. Importantly, this includes the models in FTBL and SBML file formats.

2.4.2 Metabolic models

We designed the models to cover central carbon metabolism and biomass needs of *B.*

methanolicus. *influx_si* uses a non-standard legacy file format (FTBL) to encode the metabolic

network and the associated atom-atom transitions. This format centralizes the metabolic network

with all biological measurements, i.e. metabolites pool sizes, fluxes, and carbon isotopologue

distribution. Consequently, we developed distinct model files for each carbon source. Models

share the same nomenclature and the same general topology between them, which is displayed in

Fig. 1 and detailed in Supplementary Data 1.

Assimilation pathways of methanol, mannitol and arabitol were included when relevant to

explain the incorporation of the tracer. The CO_2 pool was explicitly modeled within the system to

allow for re-incorporation of tracer via CO_2. Amino acids synthesis pathways were modeled as

part of biomass needs along with important precursors. Biomass equation was borrowed from

Bacillus subtilis genome-scale model (42). Reaction for phosphoenolpyruvate carboxykinase

(BMMGA3_RS13120) was not included in the final model as the associated expression level

was rather low in proteomics (26) and transcriptomics studies (21, 25).

Unless otherwise stated, each culture replicate was processed independently to estimate fluxes

for their respective carbon source condition.

For MFA on mannitol, we exploited carbon isotopologue distribution data of proteinogenic

alanine, glycine, valine, serine, threonine, phenylalanine, aspartate, glutamate, histidine,

isoleucine, leucine, lysine, arginine, tyrosine, proline and methionine. Acetate production was

modeled with an export flux from acetyl-CoA to which we associated the acetate flux measured

from supernatant data. Acetyl-CoA was further constrained using acetate positional labeling

measured by [1]H 1D-NMR spectroscopy from the supernatant.

For MFA on arabitol, we averaged the proteinogenic labeling measurements of the [5-[13]C]

arabitol experiment and exploited it as a parallel labeling dataset for each biological replicate of

the [1/2-[13]C] arabitol experiment. We analysed the same proteinogenic amino acids as those

mentioned above for mannitol. No acetate was observed in the supernatant and the associated

export reaction was consequently excluded from this model. Additionally, we exploited specific

labeling enrichment of histidine, alanine and ribulose-5-P, and positional isotopomer data of

alanine from labeling samples as described in section 2.3.3.

For MFA on methanol, the non-stationary nature of the experiment and the subsequent

importance of the pools on flux distributions forced us to explicit most of the reactions of the

central metabolism that were lumped for the stationary models. Measurements described above

were used to constrain intra and extracellular pools (section 2.2.3), and isotopologues profiles

(section 2.2.4) through *influx_si* optimization process. Only the measurement of the CO2 from

the exhaust gas was used but not it's labelling dynamic. The exchange fluxes of CO_2 and

methanol (feed and evaporation) were also exploited. No acetate production was observed.

Because some isotopologues of Gnt6P (M+2 & M+3) could not be quantified, these isotopic

measurements could not be included in the *influx_si* model, hence the flux through the oxidative

PPP could not be accurately resolved. To exploit these partial data and estimate the flux through

the oxidative PPP, we applied the ScalaFlux approach (43), which allows to quantify fluxes

through a given metabolic subnetwork of interest by modeling label propagation directly from

the metabolic precursor of this subnetwork. The flux calculations are thus purely based on

information from within the subnetwork of interest, and no additional knowledge about the

surrounding network is required. Briefly, we defined a metabolic subnetwork containing two

reactions: i) one reaction converting Glc6P into Gnt6P, and ii) one sink reaction operating at the

same rate to balance the 6PG pool, consistently with the metabolic steady-state assumption. This

metabolic model assumes direct formation of Gnt6P from Glc6P. Flux calculations were based

on the time-course profile of M+0 (relative to M+0,M+1,M+4,M+5,M+6), which showed the

highest range of variation during this experiment and was thus less affected by measurements

noise. Fraction M+0 of Glc6P was corrected for the presence of an extracellular unlabeled pool by normalizing its profile to the isotopic steady-state of Gnt6P. Experimental labelling dynamics of the precursor of this subnetwork (Glc6P) were used as label input, and the flux through the oxidative PPP was estimated by fitting the concentration and M+0 profile of Gnt6P. The goodness of fit was evaluated using a chi-square test, and the flux precision was estimated using the Monte-Carlo approach (with 200 iterations). Using such approach we obtain a flux through the oxidative PPP (i.e. zwf) of: 0.5909 +- 0.0578 and 0.6919 +- 0.0959 µmol/gDCW/s for the two replicates. Next we used those data to constraint the flux through zwf (as a "measured flux") through *influx_si* optimization process.

2.4.3. Quality checks

Experimental data were fitted to our models as per described above. For each culture replicate we performed a Monte Carlo sensitivity analysis (n=100) on the fit to assess its robustness to small variations around the fitted values. We also performed a chi-squared goodness-of-fit statistical test to ensure that simulated data for each biological replicate were significantly close to experimental data. All tests were significant with a significance level ($\square$) of 0.05 (Supplementary Data 2). For convenience, we provide figures of measured vs. simulated data points (Fig. S3).

Let us note here that we unfortunately were unable to estimate at a satisfactory precision the fluxes through malate dehydrogenase (BMMGA3_RS12590, 1.1.1.37), malic enzyme (*mae*, BMMGA3_RS12650, 1.1.1.38 or 1.1.1.40) and pyruvate carboxylase (*pyc*, BMMGA3_RS05255, 6.4.1.1). Those three reactions formed a cycle from pyruvate (Pyr) to oxaloacetate (OAA) and malate (Mal) in all our tested models. However, the Monte Carlo sensitivity analysis revealed that those fluxes were statistically undefined (not shown), meaning that virtually any value through the cycle would satisfy the constraints of the rest of the network.

370 In the absence of biological data to motivate any new constraint on the model, we preferred to

371 leave those reactions out of the analysis.

2.5 Analysis of arabitol phosphate dehydrogenases AtlD and AtlF

2.5.1 Strains and culture conditions

374 In this study, *Escherichia coli* DH5α (44) was used as the standard cloning host and recombinant

375 protein production was carried out with *E. coli* BL21(DE3) (45). A summary of the strains,

376 primers and plasmids constructed and used in this study can be found in Table 1. *E. coli* strains

377 were routinely cultivated at 37 °C and 180 rpm in Lysogeny Broth (LB) medium or on LB agar

378 plates supplemented with 100 µg ml^{-1} ampicillin and 0.5 mM IPTG when relevant.

2.5.1 Recombinant DNA work

Molecular cloning was performed as previously described (46) using primer sequences listed in

Table 1. Total DNA isolation from *B. methanolicus* was performed as described in (47). Inserts

were amplified by polymerase chain reactions (PCRs) with ALLin^{TM} HiFi DNA Polymerase

(HighQu, Kraichtal, Germany) and purified with the NucleoSpin® Gel and PCR Clean-up kit

(Macherey-Nagel, Düren, Germany). Plasmids were constructed from PCR-generated fragments

and pET16b vector cut with restriction enzymes using the isothermal DNA assembly method

(48). The GeneJET Plasmid Miniprep Kit (Thermo Fisher Scientific, Waltham, USA) was used

for plasmid isolation. For the transformation of chemically competent *E. coli* cells, the procedure

described by (49) was followed. Colony PCRs were performed using Taq polymerase (New

England Biolabs, Ipswich, England) with primers P192, P193, P194 and P195 (Table 1). All

cloned DNA fragments were verified by sequencing (Sequencing Core Facility, Bielefeld

University).

2.5.2 Overproduction and purification of AtlD and AtlF

Plasmids for protein production using *E. coli* BL21 (DE3) were constructed on the basis of pET16b (Novagen, Madison, WI, USA) and are presented in Table 1. The *atlD* and *atlF* genes were PCR-amplified from *B. methanolicus* MGA3 genomic DNA using the primers P192 and P193 or P194 and P195, respectively (Table 1). The resulting product was joined with *Bam*HI digested pET16b by applying the isothermal DNA assembly method (48), resulting in pET16b-*atlD* and pET16b-*atlF*. The pET16 vector allows for production of N-terminal His$_{10}$-tagged proteins. Protein production and purification was performed following the indications of (50), except for cell lysis which was performed by sonication (UP 200 S, Dr. Hielscher GmbH, Teltow, Germany) on ice at an amplitude of 55 % and a duty cycle of 0.5 for 8 min with a pause in between. Supernatants were subsequently filtered using a 0.2 µm filter and purified by nickel affinity chromatography with nickel-activated nitrilotriacetic acid-agarose (Ni-NTA) (Novagen, San Diego, CA, USA). His-tagged AltD and AtlF proteins eluted with 20 mM Tris, 300 mM NaCl, 5 % (vol/vol) glycerol and 50, 100, 200, or 400 mM imidazole were analysed by 12 % SDS-PAGE (51). Fractions showing the highest protein concentrations (with 100 and 200 mM or 100, 200 and 400 mM imidazole for AtlD and AtlF, respectively) were pooled and protein concentration was measured according to the Bradford method (52) using bovine serum albumin as reference. The purified protein was subsequently applied for enzymatic assays.

2.5.3 Arabitol phosphate dehydrogenase enzymatic assays

Determination of purified AtlD and AtlF activities in the reductive reaction using Xyl5P or Ribu5P as substrate were performed as previously described (23). The assay mixture contained 20 mM Tris-HCl buffer (pH 7.2), 1 mM DTT, 0.04 to 0.3 mM NADH or NADPH, 0.03 to 0.6 mM Xyl5P or 0.2 to 4 mM Ribu5P and 0.01 to 0.04 mg AtlD or 0.2 to 0.4 mg AtlF in a total

415   volume of 1 ml. The oxidation rate of NADH or NADPH was monitored at 340 nm and 30 ℃

416   for 3 min using a Shimadzu UV-1202 spectrophotometer (Shimadzu, Duisburg, Germany). In

417   order to confirm the presence of arabitol phosphate in the enzyme reactions after reduction of

418   Xyl5P and Ribu5P, samples were subjected to liquid chromatography-mass spectrometry (LC-

419   MS) analyses following the procedure described in (53).

420   2.6 Data availability

Gene locii mentioned throughout the text are from NCBI annotation of *B. methanolicus* MGA3

genome NZ_CP007739.1 (https://www.ncbi.nlm.nih.gov/nuccore/NZ_CP007739.1).

Supplementary Data 1 contains growth curves, processed MS and NMR data, a summary of the

reactions modelled and the absolute fluxes in mmol/gDCW/h.

Supplementary Data 2 contains raw models, input and output files for influx software and can be

downloaded from https://fairdomhub.org/data_files/3269?version=2.

3. RESULTS & DISCUSSION

3.1 *In vitro* assessment of arabitol assimilation

The operon responsible for arabitol assimilation in *B. methanolicus* consists of a PTS system (AtlABC) and two putative arabitol phosphate dehydrogenases (AtlD and AtlF) (22), which are chromosomally encoded and belong to the diverse superfamily of medium-chain dehydrogenases/reductases (MDRs). However, the physiological roles of AtlD and AtlF have not been described to date. Members of the MDR superfamily have high sequence conservation, but sequence-based prediction of their substrate scope is difficult since wide substrate specificity is common (54, 55) (see Supplementary Text for a discussion of their phylogeny). The substrate selectivity of AtlD and AtlF was therefore studied to identify whether arabitol is assimilated via arabitol 1-phosphate dehydrogenase to Xyl5P, and/or to Ribu5P via arabitol 5-phosphate dehydrogenase (23) (Fig. 2). The enzymes were purified as N-terminally His-tagged proteins from recombinant *E. coli* by nickel chelate chromatography. Arabitol phosphate oxidation could not be assayed because there are no commercial arabitol 1-phosphate or arabitol 5-phosphate standards. Therefore, the reverse reaction was tested, as described in material and methods, with either Xyl5P or Ribu5P as substrate for NAD(P)H dependent reduction. A number of potential sugars (□-ribose, □-fructose, □-xylose, □-mannose, □-arabinose, □-arabinose, □-sorbose, □-galactose, □-glucose, ribose 5-phosphate, glucose 6-phosphate, glucose 1-phosphate, fructose 6-phosphate and fructose 1-phosphate) and sugar alcohols (□-arabitol, □-mannitol, □-galactitol, □-sorbitol, □-arabitol, □-maltitol, □-xylitol, ribitol and meso-Erythritol) were also tested as substrates but no significant activity (i.e. > 0.05 U mg^{-1}) was detected. The kinetic parameters measured for AtlD and AtlF reduction using either Xyl5P or Ribu5P as substrate are summarized in Table 2. The K_M of 0.07 ± 0.03 mM for AtlD with Xyl5P as substrate and NADH as cofactor

was 2.5 times lower than the value obtained for AtlF (0.18 ± 0.05 mM). With Ribu5P as

substrate, the K_M for AtlD was 17 times lower (1.21 ± 0.42 mM) (Table 2), and for AtlF, no

activity was detected. The V_{max} for AtlD with Xyl5P as substrate and NADH as cofactor was 2.7

times higher than with Ribu5P as substrate (0.49 ± 0.06 U mg^{-1}) and 12 times higher than for

AtlF with Xyl5P (0.11 ± 0.01 U mg^{-1}) (Table 2). NADH was the preferred cofactor over NADPH

with a ten times lower K_M (0.01 ± 0.01 vs 0.11 ± 0.09 mM). The affinity and NADPH dependent

activity (0.24 ± 0.06 U mg^{-1}) could only be determined with AtlD and Xyl5P, as no significant

activity was detected for any of the other reactions. LC-MS analyses were performed to confirm

the formation of arabitol phosphate in the enzyme reactions catalyzed by AtlD. Although arabitol

1-phosphate and arabitol 5-phosphate could not be distinguished, arabitol phosphate was clearly

produced with both Xyl5P and Ribu5P as substrates (Fig. S2).

Overall, these data suggest that AtlD has a major role in arabitol catabolism *in vitro* and that

AtlD and AltF both prefer Xyl5P. This suggests that arabitol is mainly assimilated through the

arabitol 1-phosphate pathway. However, assimilation via arabitol 5-phosphate cannot be

excluded since AtlD can also use Ribu5P, albeit with reduced efficiency as shown both by the

kinetic parameters (Table 2) and the significant residual Ribu5P detected in the enzyme reactions

(Fig. S2). Moreover, *in vitro* enzymatic analyses of purified arabitol phosphate dehydrogenase

from *E. avium* (23) and *Bacillus halodurans* (56) showed that both could convert arabitol 1-

phosphate and arabitol 5-phosphate into both Xyl5P and Ribu5P. By assessing metabolic

operations *in vivo*, [13]C-MFA experiments should identify which assimilation pathway is actually

used *in vivo*.

3.2 ^{13}C-MFA experimental design

Experimental design is a key step to define the isotopic composition of the label input and the isotopic data to be measured, thereby improving both the number of fluxes that can be estimated from a set of isotopic data and the precision of the flux values. To address this point, we used a dedicated program, IsoDesign (57), using as input metabolic models of each carbon source and several labels as described in the material and methods section. For arabitol and mannitol, good flux precision could be obtained with mass spectrometry labelling data from proteinogenic amino acids. This is advantageous for two reasons. First, the experimental setup is simpler than for intracellular metabolites because no quenching is required and the cellular pellet can just be collected by centrifugation. Second, the labelling can be measured by NMR and MS, providing crucial positional information to distinguish between the two pathways (i.e. via arabitol 5-phosphate or via arabitol 1-phosphate dehydrogenase). Among the different label inputs tested, 100% [5-^{13}C] arabitol appeared ideal to identify whether arabitol is converted into arabitol 1-phosphate only or into arabitol 5-phosphate only. In addition, a 9:1 mix of [1-^{13}C] and [2-^{13}C] arabitol was used to see if both pathways are active. For mannitol, the best label input was 100 % [1-^{13}C]. Finally, since methanol is a C1 compound, a stationary ^{13}C-MFA approach is not suitable since the amino acids become fully labeled in the isotopic steady-state. Non-stationary ^{13}C fluxomics should be used instead to follow the incorporation of the tracer after a pulse of labeled substrate; however significantly more ^{13}C data are required for this approach (58, 59) (Supplementary Data 1).

All labeling samples were collected in the exponential growth phase at metabolic steady state and analyzed by MS or 1D ^{1}H NMR. The labelling profile of the analyzed metabolites, as measured by their CIDs, were inspected manually and then used with additional NMR and

494 physiological data to fit a model of *B. methanolicus*'s central metabolism (Supplementary Data

495 1).

3.3. Measurement of physiological parameters

Assessment of physiological parameters is a prerequisite for flux calculation. Here, *B.*

methanolicus was grown in batch on three different carbon sources. For each cultivation, growth

rate and consumption and production rates were determined. Results are given in Table 3. No

significant differences between the physiological parameters obtained for the culture replicates

were observed. On methanol (batch cultures at 50 °C), around 35 % of the carbon source was

directly evaporated, and biomass and carbon dioxide were the only products formed in detectable

amounts. The maximal growth rate obtained here with methanol (0.46 h^{-1}) is slightly higher than

reported in a previous proteomic study of *B. methanolicus* MGA3 (0.40 h^{-1}, (14)) or for the

growth of the related strain *B. methanolicus* PB1 (0.32 h^{-1}, (15)). Biomass yields did not differ

significantly from 0.5 g/g indicating that approximately half of the consumed methanol went to

biomass and the other half was oxidized to CO_2. The growth rates with mannitol and arabitol are

consistent with published values (22)).

B. methanolicus MGA3 is known to overproduce glutamate at up to 50 g/l (21) on methanol

under magnesium or methanol limitation (60, 61). When grown on arabitol or methanol, no

metabolite accumulated in the supernatant at concentrations above the NMR detection limit

(approximately 100 µM). There was therefore little or no metabolite secretion by MGA3 under

our chosen methanol and arabitol growth conditions, which is consistent with previous studies

and the growth conditions studied here (27, 28). However, acetate was produced at up to 2

mmol/gDCW/h in mannitol cultures (yield, 0.3 mol/mol). To the best of our knowledge, acetate

production has never previously been reported for *B. methanolicus*. Based on genomics data

(21), we hypothesized that acetate is synthesized from acetyl-CoA in a classical two-step process

involving phosphate acetyltransferase (EC:2.3.1.8, BMMGA3_RS15725) and acetate kinase

(EC:2.7.2.1, BMMGA3_RS12735). Acetate has moreover been discussed at length in the context

of overflow metabolism, a special metabolic state in which fermentation pathways are used even

though further oxidation (respiration) would be more ATP-efficient (62-65).

Overall, we show the reproducible growth characteristics of our cultures, and show an

unexpected production of acetate that may have an impact for industrial applications, as overflow

metabolism leads to carbon and energy waste.

3.4 *In vivo* characterization of arabitol assimilation

The *in vitro* enzymatic analyses of the two dehydrogenases, AltD and AltF, suggest that both

assimilation pathways (i.e. via arabitol 1-phosphate and via arabitol 5-phosphate) may operate *in*

vivo. To confirm whether either or both arabitol assimilation pathways are operative in *B.*

methanolicus, as suggested for *E. avium* (23) and *Bacillus halodurans* (56), we carried out a ^{13}C-

MFA specifically designed to discriminate between the two pathways (see section 3.1, and Fig

2). Interestingly, while the possibility to assimilate arabitol through Xyl5P or Ribu5P or both was

a free parameter, the optimal solution found during the fitting process exclusively used the route

through Xyl5P (Fig. 1C). This indicates that the (low) activity observed *in vitro* through Ribu5P

was not present at a detectable level in our cultures, and that the PTS system imports arabitol as

arabitol 1-phosphate. The kinetic parameters obtained for AtlD and AtlF are in line with the flux

data, i.e. entry of arabitol into the pentose phosphate pathway (PPP) via PTS-mediated uptake

and phosphorylation to arabitol 1-phosphate followed by oxidation to Xyl5P with AtlD as the

major dehydrogenase (Fig. 2 and Table 2).

Overall, these data demonstrate for the first time how arabitol is assimilated in *B. methanolicus*

and rule out the hypothesis of additional catabolism through arabitol 5-phosphate and Ribu5P

derived from our enzymatic analyses (Table 2) and previous reports in *E. avium* (23) and *B.*

halodurans (56).

543   3.5. *In vivo* operation of the pentose phosphate pathway with the

544   different carbon sources

B. methanolicus assimilates methanol through the ribulose monophosphate (RuMP) cycle that

condenses formaldehyde (For) and ribulose 5-P (Ribu5P) into hexulose 6-P (Hex6P) (66). The

regenerative part of the RuMP cycle that maintains a pool of Ribu5P overlaps with the non-

oxidative pentose phosphate pathway (PPP). Strong flux through the non-oxidative part of the

PPP is therefore expected on methanol and indeed, ribulose-phosphate 3-epimerase's relative

flux accounted for 62 % of methanol assimilation (*rpe*, Fig. 1A). Interestingly, mannitol and

arabitol are closely connected to the PPP since mannitol is converted to fructose 6-P (F6P) (just

like methanol), whereas arabitol is converted to Xyl5P as discussed above. On mannitol and

arabitol (Fig. 1B), PPP utilization was lower compare to methanol. However on arabitol, the

fluxes associated with the PPP accounted for a larger fraction of the assimilated carbon flow than

on mannitol (the estimated *rpe* flux was 15% and 28 % of respectively mannitol and arabitol

assimilation; Fig 1B-C). This indicates that the non-oxidative PPP is more important for carbon

assimilation on arabitol than on mannitol, in spite of similar expression levels (22). Simulations

carried without transaldolase activity indicated that its activity was essential to fit the labeling

data given the network topology used (notably *glpx* been irreversible) on arabitol. A possibility

for future studies would be to take advantage of adaptive laboratory driven evolution, or

overexpression of key enzymes such as transaldolase, to investigate if the PPP could be adjusted

to increase growth rates when arabitol is the sole source of carbon and energy.

The RuMP cycle has several variants, which differ in their efficiency (67). Genes for two of

these have been identified in MGA3, namely, the fructose bisphosphate aldolase/sedoheptulose

bisphosphatase (SBPase) cycle (53, 68) and the fructose bisphosphate aldolase/transaldolase

(TA) cycle(69), which as their names suggest favor the regeneration of Ribu5P through

sedoheptulose bisphosphatase and transaldolase, respectively. It is generally accepted that MGA3

uses the SBPase variant (66). The main evidence for this is the presence of a copy of a

characteristic gene of the SBPase variant ($glpX^P$) on the pBM19 plasmid, whereas there is only

one transaldolase gene (ta^C) in the chromosome. Proteomic (26) and transcriptomic (21) studies

have also associated $glpX^P$ with a significant increase in expression in methanol compared with

mannitol, whereas the expression associated with ta^C remained constant. According to the ^{13}C-

MFA, both $glpx$ (associated to $glpX^P$) and ta (associated to ta^C) carried a comparable flux, thus

both variant may be active (Fig. 3). This arrangement may serve as a fail-safe to guarantee the

replenishment of Ribu5P. Alternatively, since transaldolase activity is essential to fit the isotopic

data for growth on arabitol, an advantage of the TA cycle may be that it increases the flexibility

of the PPP and allows the regeneration of important precursors such as Ribu5P from different

carbon sources.

The labeling data suggest that the oxidative part of the PPP was similar on methanol and on

arabitol   and almost five times higher on mannitol (the estimated relative glucose 6-phosphate

isomerase (pgi) flux was 31, 12 and 9 %, respectively; Fig. 1). This follows the conclusions of

previous studies showing that the oxidative part of the PPP is slightly upregulated on mannitol

compare to methanol (14) but suggested that this route should be used on methanol to provide

NADPH   while detoxifying formaldehyde (14, 27) via the so-called cyclic dissimilatory RuMP

pathway (21, 70). The utilization of the oxidative part of the PPP on methanol is of particular

interest because, intuitively, one imagines that decarboxylation should be avoided when growth

occurs on a C1 to avoid wasting carbon. This intuition proved to be correct in Bennett et al.'s

(71) study of an *E. coli* synthetic methylotroph in which they knocked-out *pgi* to increase the

regeneration of Ribu5P through the non-oxidative part of the PPP. However, we cannot exclude

the possibility that the oxidative part of the PPP serves as a backup formaldehyde dissimilation

pathway when the linear dissimilation pathways become saturated at high methanol

concentrations. *B. methanolicus* is indeed quite sensitive to variations in methanol concentration

(72) and we can assume that this critical biological function is tightly controlled.

As expected, the ^{13}C-MFA shows that the PPP is critical on methanol as it overlaps with the

RuMP responsible for methanol assimilation. The SBPase variant of the RuMP is active,

however we could not rule out a parallel operation of the TA variant. We suggest that a parallel

operation on mixed carbon sources may benefit *B. methanolicus* to replenish important

precursor's pools.

3.6 *In vivo* operation of the TCA cycle with the different carbon

sources

B. methanolicus has a full gene set for a functional tricarboxylic acid cycle (TCA) and glyoxylate

shunt (14, 21). This seems to contrast with some methylotrophs, including some that use the

RuMP pathway, that do not need a complete TCA to fulfill their energy requirements (73).

Results (Fig. 1A) indicated that the TCA was used much less than the RuMP pathway on

methanol, with relative fluxes 3% up to isocitrate (ICit) and almost no flux afterwards (Fig. 1A).

This small flux is needed to support the synthesis of biomass precursors. On mannitol and

arabitol in contrast, the TCA was used intensively, with relative flux of respectively 57 % and

110% for isocitrate dehydrogenase (*idh*) (Fig.1B-C). These results are in agreement with

previous measurements of the actual usage of the TCA during methylotrophic growth. At the

transcript and protein levels, it has been suggested that the TCA should be more active on

mannitol than on methanol (21, 26). Activity assays in crude cell extracts also showed very low

2-oxoglutarate dehydrogenase (*akgdh*) activity (60). Finally, isotopic labeling experiments have

shown slow isotopic enrichment of key TCA metabolites (citrate, 2-oxoglutarate, fumarate) on

methanol (27).

Although *B. methanolicus* also has glyoxylate shunt genes (BMMGA3_RS01750, 2.3.3.9; and

BMMGA3_RS01755, 4.1.3.1), the reported expression levels suggest that this pathway does not

carry a high flux under mannitol or under methanol growth conditions (21, 26). In agreement

with these findings, the glyoxylate shunt fluxes estimated in this study were negligible under all

the tested conditions (Fig. 1).

Overall, as reported before, the TCA is mainly active in non-methylotrophic conditions and the

glyoxylate shunt was inactive in our conditions.

3.7 Analysis of cofactor usage with the various carbon sources

To assess how *B. methanolicus* balances its redox and energy needs, we used the estimated fluxes

from the ^{13}C-MFA to infer production and consumption rates of ATP, NADH and NADPH, as

displayed in Fig. 4. ^{13}C-MFA is constrained by carbon balance and the distribution of the isotopic

tracer, but unlike flux balance analysis it is typically not constrained by cofactor balance. We

used the estimated fluxes from the MFA and the expected stoichiometry of the associated

reactions to assess the absolute rates of cofactor production and consumption. In the absence of

specific measurements, we use the biomass requirements of *B. subtilis* (8, 74). Additionally,

since the samples were collected in metabolic pseudo steady-state, the production and

consumption rates of each cofactor were assumed to be balanced; this allowed us to identify rates

that would have been impossible to estimate otherwise such as the proportion of NAD (and ATP)

formed from the respiratory chain (with a P/O=1.5) and other production/consumption rates not

accounted for by our other assumptions.

Despite their different growth rates, which influence their cofactor requirements for growth

(Table 3), the cells grown on mannitol and arabitol had a similar absolute usage of NADH,

NADPH and ATP (Fig. 3). ATP and NADH came mostly from the same sources. However, the

proportion of NADPH generated in the oxidative part of the PPP was three times higher in

mannitol growth conditions than in arabitol (3.5 against 1.0 mmol/gDCW/h) which is consistent

with the higher fluxes observe in the oxidative part of the PPP on mannitol. This difference was

compensated by a higher flux in the TCA for arabitol (4.7 against 3.5 mmol/gDCW/h) in excess

of the estimated NADPH requirements for growth and even contributing to the production of

~3.4 mmol/gDCW/h NADPH whose consumption remains unaccounted for. Discrepancies

between estimates of the cofactor requirements for biomass formation and those derived from

measured isotopic data are typical for this kind of analysis (75, 76), and can be attributed to an

underestimation of cofactor requirements, notably for non-essential processes such as cell

motility (for ATP).

Comparing methylotrophic and non-methylotrophic growth, it is not surprising that NADH was

produced at a higher rate on methanol (45.6 mmol/gDCW/h, versus 30.4 and 25.7

mmol/gDCW/h on mannitol and arabitol respectively), because of the conversion of methanol

into formaldehyde by methanol dehydrogenase (77). This suggests that at the same growth rate

(i.e. methanol vs mannitol), O_2 consumption should be higher on methanol to provide the

additional NAD^+ required. The NADPH total production was estimated to be largely higher on

methanol (9.2 mmol/gDCW/h) compare to mannitol and arabitol (Fig. 3). It was mainly provided

by the oxidative part of the PPP (which would be part of the cyclic dissemination pathway; 4.6

mmol/gDCW/h) as on mannitol but also by the linear formaldehyde detoxification pathways,

while usage of the TCA was negligible. These estimates suggest that linear detoxification

pathways play an important role in the generation of NADPH on methanol, in addition to their

established protective function against formaldehyde.

Overal, non-methylotrophic growth on mannitol and arabitol share the same features despite

their associated different growth rates (with the notable exception of the contribution of the

oxidative PPP for NADPH production). On methanol, however, we observe more differences in

the origin of cofactors production which clearly highlight a different metabolic state.

Importantly, the linear detoxification pathways share with the PPP an important role in cofactor

regeneration.

3.8. Conclusion

In summary, this MFA of *B. methanolicus* MGA3 provides three snapshots of its metabolic states

for growth on methanol, mannitol or arabitol. Isotopic data are consistent with prior knowledge

of MGA3 methylotrophy, showing greater flux in the RuMP cycle than in the TCA. The ^{13}C-

MFA provided new insights related to the utilization of cyclic RuMP versus linear dissimilation

pathways, and between the RuMP cycle variants; and finally, the characterization of the arabitol

assimilation pathway was completed using enzymatic data. In future studies, these validated flux

maps will be used as references for constraint based modelling to validate genome-scale model

predictions. Overall, the information provided in this work and previous omics studies on *B.*

methanolicus metabolism can be used to improve design strategies for new strains (i.e. by multi-

omics analysis). For instance, the comparison between methanol and arabitol might help

identifying key control steps in the metabolic network that allows shifting from a cyclic (i.e.

methanol with the Rump cycle) to a linear (i.e. arabitol with the PPP) mode of operation. The

experimental path outlined here likely leads to *B. methanolicus* becoming a viable alternative to

existing cell factories.

References

1. Peyraud R, Kiefer P, Christen P, Massou S, Portais J-C, Vorholt JA. 2009. Demonstration of the ethylmalonyl-CoA pathway by using 13C metabolomics. Proceedings of the National Academy of Sciences of the United States of America 106:4846-4851.
2. Fendt S-M, Oliveira AP, Christen S, Picotti P, Dechant RC, Sauer U. 2010. Unraveling condition-dependent networks of transcription factors that control metabolic pathway activity in yeast. Molecular Systems Biology 6:432.
3. Becker J, Zelder O, Häfner S, Schröder H, Wittmann C. 2011. From zero to hero--design-based systems metabolic engineering of Corynebacterium glutamicum for L-lysine production. Metabolic Engineering 13:159-168.
4. Heux S, Poinot J, Massou S, Sokol S, Portais J-C. 2014. A novel platform for automated high-throughput fluxome profiling of metabolic variants. Metabolic Engineering 25:8-19.
5. Klingner A, Bartsch A, Dogs M, Wagner-Döbler I, Jahn D, Simon M, Brinkhoff T, Becker J, Wittmann C. 2015. Large-Scale 13C flux profiling reveals conservation of the Entner-Doudoroff pathway as a glycolytic strategy among marine bacteria that use glucose. Applied and Environmental Microbiology 81:2408-2422.
6. Van Dien SJ, Strovas T, Lidstrom ME. 2003. Quantification of central metabolic fluxes in the facultative methylotroph methylobacterium extorquens AM1 using 13C-label tracing and mass spectrometry. Biotechnology and Bioengineering 84:45-55.
7. Fu Y, He L, Reeve J, Beck DAC, Lidstrom ME. 2019. Core Metabolism Shifts during Growth on Methanol versus Methane in the Methanotroph Methylomicrobium buryatense 5GB1. mBio 10.
8. Peyraud R, Schneider K, Kiefer P, Massou S, Vorholt JA, Portais J-C. 2011. Genome-scale reconstruction and system level investigation of the metabolic network of Methylobacterium extorquens AM1. BMC systems biology 5:189.
9. Niedenführ S, Wiechert W, Nöh K. 2015. How to measure metabolic fluxes: a taxonomic guide for (13)C fluxomics. Current Opinion in Biotechnology 34:82-90.
10. Heux S, Bergès C, Millard P, Portais J-C, Létisse F. 2017. Recent advances in high-throughput 13C-fluxomics. Current Opinion in Biotechnology 43:104-109.
11. Schendel FJ, Bremmon CE, Flickinger MC, Guettler M, Hanson RS. 1990. L-lysine production at 50 degrees C by mutants of a newly isolated and characterized methylotrophic Bacillus sp. Applied and Environmental Microbiology 56:963-970.
12. Pfeifenschneider J, Brautaset T, Wendisch VF. 2017. Methanol as carbon substrate in the bio-economy: Metabolic engineering of aerobic methylotrophic bacteria for production of value-added chemicals. Biofuels, Bioproducts and Biorefining 11:719-731.
13. Iaquaniello G, Centi G, Salladini A, Palo E, Perathoner S. 2018. Waste to Chemicals for a Circular Economy. Chemistry (Weinheim an Der Bergstrasse, Germany) 24:11831-11839.
14. Müller JEN, Heggeset TMB, Wendisch VF, Vorholt JA, Brautaset T. 2015. Methylotrophy in the thermophilic Bacillus methanolicus, basic insights and application for commodity production from methanol. Applied Microbiology and Biotechnology 99:535-551.
15. Komives CF, Cheung LY-Y, Pluschkell SB, Flickinger MC. 2005. Growth of Bacillus methanolicus in seawater-based media. Journal of Industrial Microbiology & Biotechnology 32:61-66.

16. Naerdal I, Pfeifenschneider J, Brautaset T, Wendisch VF. 2015. Methanol-based
cadaverine production by genetically engineered Bacillus methanolicus strains. Microbial
Biotechnology 8:342-350.
17. Irla M, Nærdal I, Brautaset T, Wendisch VF. 2017. Methanol-based γ-aminobutyric acid
(GABA) production by genetically engineered Bacillus methanolicus strains. Industrial
Crops and Products 106:12-20.
18. Irla M, Heggeset TMB, Nærdal I, Paul L, Haugen T, Le SB, Brautaset T, Wendisch VF.
2016. Genome-Based Genetic Tool Development for Bacillus methanolicus: Theta- and
Rolling Circle-Replicating Plasmids for Inducible Gene Expression and Application to
Methanol-Based Cadaverine Production. Frontiers in Microbiology 7:1481.
19. Drejer EB, Hakvåg S, Irla M, Brautaset T. 2018. Genetic Tools and Techniques for
Recombinant Expression in Thermophilic Bacillaceae. Microorganisms 6.
20. Schultenkämper K, Brito LF, López MG, Brautaset T, Wendisch VF. 2019. Establishment
and application of CRISPR interference to affect sporulation, hydrogen peroxide
detoxification, and mannitol catabolism in the methylotrophic thermophile Bacillus
methanolicus. Applied Microbiology and Biotechnology 103:5879-5889.
21. Heggeset TMB, Krog A, Balzer S, Wentzel A, Ellingsen TE, Brautaset T. 2012. Genome
sequence of thermotolerant Bacillus methanolicus: features and regulation related to
methylotrophy and production of L-lysine and L-glutamate from methanol. Applied and
Environmental Microbiology 78:5170-5181.
22. López MG, Irla M, Brito LF, Wendisch VF. 2019. Characterization of D-Arabitol as
Newly Discovered Carbon Source of Bacillus methanolicus. Frontiers in Microbiology
10:1725.
23. Povelainen M, Eneyskaya EV, Kulminskaya AA, Ivanen DR, Kalkkinen N, Neustroev
KN, Miasnikov AN. 2003. Biochemical and genetic characterization of a novel enzyme
of pentitol metabolism: D-arabitol-phosphate dehydrogenase. The Biochemical Journal
371:191-197.
24. Irla M, Neshat A, Winkler A, Albersmeier A, Heggeset TMB, Brautaset T, Kalinowski J,
Wendisch VF, Rückert C. 2014. Complete genome sequence of Bacillus methanolicus
MGA3, a thermotolerant amino acid producing methylotroph. Journal of Biotechnology
188:110-111.
25. Irla M, Neshat A, Brautaset T, Rückert C, Kalinowski J, Wendisch VF. 2015.
Transcriptome analysis of thermophilic methylotrophic Bacillus methanolicus MGA3
using RNA-sequencing provides detailed insights into its previously uncharted
transcriptional landscape. BMC genomics 16:73.
26. Müller JEN, Litsanov B, Bortfeld-Miller M, Trachsel C, Grossmann J, Brautaset T,
Vorholt JA. 2014. Proteomic analysis of the thermophilic methylotroph Bacillus
methanolicus MGA3. Proteomics 14:725-737.
27. Müller JEN, Meyer F, Litsanov B, Kiefer P, Vorholt JA. 2015. Core pathways operating
during methylotrophy of Bacillus methanolicus MGA3 and induction of a bacillithiol-
dependent detoxification pathway upon formaldehyde stress. Molecular Microbiology
98:1089-1100.
28. Carnicer M, Vieira G, Brautaset T, Portais J-C, Heux S. 2016. Quantitative metabolomics
of the thermophilic methylotroph Bacillus methanolicus. Microbial Cell Factories 15:92.
29. Izhaki I, Fridman S, Gerchman Y, Halpern M. 2013. Variability of bacterial community
composition on leaves between and within plant species. Current Microbiology 66:227-
235.
30. Fall R, Benson AA. 1996. Leaf methanol — the simplest natural product from plants.
Trends in Plant Science 1:296-301.

31.   Mashego MR, Wu L, Van Dam JC, Ras C, Vinke JL, Van Winden WA, Van Gulik WM,
Heijnen JJ. 2004. MIRACLE: mass isotopomer ratio analysis of U-13C-labeled extracts.
A new method for accurate quantification of changes in concentrations of intracellular
metabolites. Biotechnology and Bioengineering 85:620-628.

32.   Kiefer P, Nicolas C, Letisse F, Portais J-C. 2007. Determination of carbon labeling
distribution of intracellular metabolites from single fragment ions by ion chromatography
tandem mass spectrometry. Analytical Biochemistry 360:182-188.

33.   Millard P, Delépine B, Guionnet M, Heuillet M, Bellvert F, Létisse F. 2019. IsoCor:
isotope correction for high-resolution MS labeling experiments. Bioinformatics (Oxford,
England) 35:4484-4487.

34.   van der Heijden RT, Romein B, Heijnen JJ, Hellinga C, Luyben KC. 1994. Linear
constrain relations in biochemical reaction systems III. Sequential application of data
reconciliation for sensitive detection of systematic errors. Biotechnology and
Bioengineering 44:781-791.

35.   Dauner M, Sauer U. 2001. Stoichiometric growth model for riboflavin-producing
Bacillus subtilis. Biotechnology and Bioengineering 76:132-143.

36.   Dauner M, Bailey JE, Sauer U. 2001. Metabolic flux analysis with a comprehensive
isotopomer model in Bacillus subtilis. Biotechnology and Bioengineering 76:144-156.

37.   Stephanopoulos G, Locher G, Duff MJ, Kamimura R, Stephanopoulos G. 1997.
Fermentation database mining by pattern recognition. Biotechnology and Bioengineering
53:443-452.

38.   Boudah S, Olivier M-F, Aros-Calt S, Oliveira L, Fenaille F, Tabet J-C, Junot C. 2014.
Annotation of the human serum metabolome by coupling three liquid chromatography
methods to high-resolution mass spectrometry. Journal of Chromatography B, Analytical
Technologies in the Biomedical and Life Sciences 966:34-47.

39.   Massou S, Nicolas C, Letisse F, Portais J-C. 2007. NMR-based fluxomics: quantitative
2D NMR methods for isotopomers analysis. Phytochemistry 68:2330-2340.

40.   Massou S, Nicolas C, Letisse F, Portais J-C. 2007. Application of 2D-TOCSY NMR to
the measurement of specific(13C-enrichments in complex mixtures of 13C-labeled
metabolites. Metabolic Engineering 9:252-257.

41.   Sokol S, Millard P, Portais J-C. 2012. influx_s: increasing numerical stability and
precision for metabolic flux analysis in isotope labelling experiments. Bioinformatics
(Oxford, England) 28:687-693.

42.   Sauer U, Hatzimanikatis V, Hohmann HP, Manneberg M, van Loon AP, Bailey JE. 1996.
Physiology and metabolic fluxes of wild-type and riboflavin-producing Bacillus subtilis.
Applied and Environmental Microbiology 62:3687-3696.

43.   Millard P, Schmitt U, Kiefer P, Vorholt JA, Heux S, Portais J-C. 2019. ScalaFlux: a
scalable approach to quantify fluxes in metabolic subnetworks. bioRxiv
doi:10.1101/735308:735308.

44.   Hanahan D. 1983. Studies on transformation of Escherichia coli with plasmids. Journal of
Molecular Biology 166:557-580.

45.   Studier FW, Moffatt BA. 1986. Use of bacteriophage T7 RNA polymerase to direct
selective high-level expression of cloned genes. Journal of Molecular Biology 189:113-
130.

46.   Green MR, Sambrook J, Sambrook J. 2012. Molecular cloning: a laboratory manual, 4th
ed ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.

47.   Eikmanns BJ, Thum-Schmitz N, Eggeling L, Lüdtke KU, Sahm H. 1994. Nucleotide
sequence, expression and transcriptional analysis of the Corynebacterium glutamicum
gltA gene encoding citrate synthase. Microbiology (Reading, England) 140 (Pt 8):1817-

1828.

48.   Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO. 2009.
Enzymatic assembly of DNA molecules up to several hundred kilobases. Nature Methods
6:343-345.

49.   Mandel M, Higa A. 1970. Calcium-dependent bacteriophage DNA infection. Journal of
Molecular Biology 53:159-162.

50.   Lindner SN, Vidaurre D, Willbold S, Schoberth SM, Wendisch VF. 2007. NCgl2620
encodes a class II polyphosphate kinase in Corynebacterium glutamicum. Applied and
Environmental Microbiology 73:5026-5033.

51.   Laemmli UK. 1970. Cleavage of structural proteins during the assembly of the head of
bacteriophage T4. Nature 227:680-685.

52.   Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram
quantities of protein utilizing the principle of protein-dye binding. Analytical
Biochemistry 72:248-254.

53.   Stolzenberger J, Lindner SN, Persicke M, Brautaset T, Wendisch VF. 2013.
Characterization of fructose 1,6-bisphosphatase and sedoheptulose 1,7-bisphosphatase
from the facultative ribulose monophosphate cycle methylotroph Bacillus methanolicus.
Journal of Bacteriology 195:5112-5122.

54.   Nordling E, Jörnvall H, Persson B. 2002. Medium-chain dehydrogenases/reductases
(MDR). Family characterizations including genome comparisons and active site
modeling. European Journal of Biochemistry 269:4267-4276.

55.   Persson B, Hedlund J, Jörnvall H. 2008. Medium- and short-chain
dehydrogenase/reductase gene and protein families : the MDR superfamily. Cellular and
molecular life sciences: CMLS 65:3879-3894.

56.   Soroka NV, Kulminskaya AA, Eneyskaya EV, Shabalin KA, Uffimtcev AV, Povelainen
M, Miasnikov AN, Neustroev KN. 2005. Synthesis of arabinitol 1-phosphate and its use
for characterization of arabinitol-phosphate dehydrogenase. Carbohydrate Research
340:539-546.

57.   Millard P, Sokol S, Letisse F, Portais J-C. 2014. IsoDesign: a software for optimizing the
design of 13C-metabolic flux analysis experiments. Biotechnology and Bioengineering
111:202-208.

58.   Wiechert W, Nöh K. 2005. From stationary to instationary metabolic flux analysis.
Advances in Biochemical Engineering/Biotechnology 92:145-172.

59.   Nöh K, Grönke K, Luo B, Takors R, Oldiges M, Wiechert W. 2007. Metabolic flux
analysis at ultra short time scale: isotopically non-stationary 13C labeling experiments.
Journal of Biotechnology 129:249-267.

60.   Brautaset T, Williams MD, Dillingham RD, Kaufmann C, Bennaars A, Crabbe E,
Flickinger MC. 2003. Role of the Bacillus methanolicus citrate synthase II gene, citY, in
regulating the secretion of glutamate in L-lysine-secreting mutants. Applied and
Environmental Microbiology 69:3986-3995.

61.   Shiio I, Sano K, Nakamori S. 1972/12/26/ 1972. Method of producing l-lysine by
fermentation patent US3707441A.

62.   Szenk M, Dill KA, de Graff AMR. 2017. Why Do Fast-Growing Bacteria Enter Overflow
Metabolism? Testing the Membrane Real Estate Hypothesis. Cell Systems 5:95-104.

63.   Bernal V, Castaño-Cerezo S, Cánovas M. 2016. Acetate metabolism regulation in
Escherichia coli: carbon overflow, pathogenicity, and beyond. Applied Microbiology and
Biotechnology 100:8985-9001.

64.   Enjalbert B, Millard P, Dinclaux M, Portais J-C, Létisse F. 2017. Acetate fluxes in
Escherichia coli are determined by the thermodynamic control of the Pta-AckA pathway.

Scientific Reports 7:42135.

65.   Polen T, Rittmann D, Wendisch VF, Sahm H. 2003. DNA microarray analyses of the
877   long-term adaptive response of Escherichia coli to acetate and propionate. Applied and
878   Environmental Microbiology 69:1759-1774.

879   66.   Jakobsen ØM, Benichou A, Flickinger MC, Valla S, Ellingsen TE, Brautaset T. 2006.
880   Upregulated transcription of plasmid and chromosomal ribulose monophosphate pathway
881   genes is critical for methanol assimilation rate and methanol tolerance in the
882   methylotrophic bacterium Bacillus methanolicus. Journal of Bacteriology 188:3063-
883   3072.

884   67.   Anthony C. 1991. Assimilation of carbon by methylotrophs. Biotechnology (Reading,
885   Mass) 18:79-109.

886   68.   Stolzenberger J, Lindner SN, Wendisch VF. 2013. The methylotrophic Bacillus
887   methanolicus MGA3 possesses two distinct fructose 1,6-bisphosphate aldolases.
888   Microbiology (Reading, England) 159:1770-1781.

889   69.   Pfeifenschneider J, Markert B, Stolzenberger J, Brautaset T, Wendisch VF. 2020.
890   Transaldolase in Bacillus methanolicus: biochemical characterization and biological role
891   in ribulose monophosphate cycle. BMC Microbiology 20:63.

892   70.   Chistoserdova LV, Chistoserdov AY, Schklyar NL, Baev MV, Tsygankov YD. 1991.
893   Oxidative and assimilative enzyme activities in continuous cultures of the obligate
894   methylotroph Methylobacillus flagellatum. Antonie Van Leeuwenhoek 60:101-107.

895   71.   Bennett RK, Gonzalez JE, Whitaker WB, Antoniewicz MR, Papoutsakis ET. 2018.
896   Expression of heterologous non-oxidative pentose phosphate pathway from Bacillus
897   methanolicus and phosphoglucose isomerase deletion improves methanol assimilation
898   and metabolite production by a synthetic Escherichia coli methylotroph. Metabolic
899   Engineering 45:75-85.

900   72.   Bozdag A, Komives C, Flickinger MC. 2015. Growth of Bacillus methanolicus in 2 M
901   methanol at 50 °C: the effect of high methanol concentration on gene regulation of
902   enzymes involved in formaldehyde detoxification by the ribulose monophosphate
903   pathway. Journal of Industrial Microbiology & Biotechnology 42:1027-1038.

904   73.   Chistoserdova L, Kalyuzhnaya MG, Lidstrom ME. 2009. The expanding world of
905   methylotrophic metabolism. Annual Review of Microbiology 63:477-499.

906   74.   Oh Y-K, Palsson BO, Park SM, Schilling CH, Mahadevan R. 2007. Genome-scale
907   reconstruction of metabolic network in Bacillus subtilis based on high-throughput
908   phenotyping and gene essentiality data. The Journal of Biological Chemistry 282:28791-
909   28799.

910   75.   Gonzalez JE, Long CP, Antoniewicz MR. 2017. Comprehensive analysis of glucose and
911   xylose metabolism in Escherichia coli under aerobic and anaerobic conditions by 13C
912   metabolic flux analysis. Metabolic Engineering 39:9-18.

913   76.   Revelles O, Millard P, Nougayrède J-P, Dobrindt U, Oswald E, Létisse F, Portais J-C.
914   2013. The carbon storage regulator (Csr) system exerts a nutrient-specific control over
915   central metabolism in Escherichia coli strain Nissle 1917. PloS One 8:e66386.

916   77.   Arfman N, Hektor HJ, Bystrykh LV, Govorukhina NI, Dijkhuizen L, Frank J. 1997.
917   Properties of an NAD(H)-containing methanol dehydrogenase and its activator protein
918   from Bacillus methanolicus. Eur J Biochem 244:426-33.

Figures Caption

Fig. 1: Flux map of *B. methanolicus* central metabolism relative to substrate intake for methanol (A), mannitol (B) and arabitol (C). The thickness of the reaction lines represent the flux proportions relative to substrate incorporation into the metabolism: hps (A), Manupt (B), Araupt (C). Flux proportions were averaged among the biological replicates ($n_{methanol}=2$, $n_{arabitol} = n_{mannitol} = 3$). The direction of the reaction arrows represent the estimated net flux directionality. Large grey arrows represent a flux directed toward amino acid synthesis and biomass requirements for growth. Pathways discussed in the text are represented by a background patch of color. Metabolites are represented by circles and have a solid bottom half if they are duplicated on the same panel, as per recommended by the SBGN standard. Metabolites subjected to experimental measurement of ^{13}C labelling are marked by a yellow box. Some intermediate metabolites are not needed for ^{13}C-MFA and are consequently lumped in mannitol and arabitol models (B, C); the same metabolites can be explicitly modelled in methanol model (A) since non-stationary ^{13}C-MFA requires all intermediates pools (e.g.: G3P, Cit, Aco). Reactions toward amino acid pools and biomass are not represented.

Fig. 2: Alternative assimilation pathways of D-arabitol with an example of the measured and expected labeling of Rib5P. Arabitol entry point into the metabolism is expected to be Ribu5P or Xyl5P, depending on the substrate specificity of the PTS system and the arabitol phosphate dehydrogenase. Taking [5-^{13}C]arabitol as an example, we show that the labeling of downstream metabolites can be used to identify which pathway is operating *in vivo* (Ribu5P, orange; Xyl5P, blue). We exemplify our approach with one data point: the NMR specific enrichment of Rib5P and the associated estimates from the best fit of different scenarios in which

PTS1 or PTS5 are allowed to carry flux in the model (barplot). Unlike the others, the model in

which only PTS5 was allowed to carry flux did not fit the data (red crossmark, ☐). Circles are

carbon atoms: solid for ^{13}C (●), empty for ^{12}C (○), dotted when irrelevant.

Fig. 3: Estimated absolute rates of production and consumption of NADPH,

NADH/FADH2, and ATP. Rates (mmol/gDCW/h) are calculated as the sum of the estimated

flux of the reactions modeled in the MFA that are expected to produce (positive value) or

consume (negative value) those cofactors. Values are averaged over the biological replicates and

the error bars are the associated standard deviation ($n_{methanol} = 2$, $n_{arabitol} = n_{mannitol} = 3$). The

growth requirements ("biomass", red) are computed from measures on *B. subtilis* (mmol/gDCW)

and the growth rates we observed. Production and consumption rates should be balanced in our

conditions, so we mark putative production/consumption rates as "unknown" to complete the

balance when needed. The full production of NADH and FADH2 is assumed to be consumed in

the respiratory chain ("oxidative phosphorylation") and to produce ATP (with P/O$_{NADH}$ = 1.5,

P/O$_{FADH2}$ = 1). We modeled the PTS systems of mannitol and arabitol as consumers of one

equivalent ATP (in "uptake"). NADPH production: glucose 6-phosphate dehydrogenase (*zwf*),

phosphogluconate   dehydrogenase   (*gnd*),   isocitrate   dehydrogenase   (*idh*),   and

methylenetetrahydrofolate dehydrogenase from the linear detoxification pathway (detox). NADH

production: glyceraldehyde-3-phosphate dehydrogenase (*pgk*), pyruvate dehydrogenase (*pdh*), 2-

oxoglutarate dehydrogenase (*akgdh*), malate dehydrogenase (assimilated to *fum*), methanol,

mannitol and arabitol dehydrogenase (*mdh*, manupt, araupt), and formate dehydrogenase from

the linear detoxification pathway (detox). FADH2 production: succinate dehydrogenase (*fum*).

ATP consumption: 6-phosphofructokinase (*pfk*), PTS systems of mannitol and arabitol (manupt,

araupt). ATP production: phosphoglycerate kinase (*pgk*), pyruvate kinase (*pyk*), succinyl-CoA

synthetase (assimilated to *akgdh*), acetate kinase (*out_Ac*), and formate-tetrahydrofolate ligase

from the linear detoxification pathway (detox).

 # Tables

 Table 1. Strains, primers and plasmids used in this study.

Strain, plasmid or primer	Relevant characteristics	Reference
Strains		
E. coli DH5α	F⁻ *thi*-1 *endA1 hsdR17*(r⁻ m⁻) *supE44* Δ*lacU169* (φ80*lacZ*ΔM15) *recA1 gyrA96 relA1*	(43)
E. coli BL21 (DE3)	F⁻ *ompT hsdSB*(r$_B$⁻ m$_B$⁻) *gal dcm* (DE3)	(44)
B. methanolicus MGA3	Wild type strain (ATCC 53907)	(11)
Plasmids		
pET16b	Amp^R; overproduction of decahistidine-tagged proteins in *E. coli* (pBR322 oriVE.c., *PT7*, *lacI*)	Novagen
pET16b-*atlD*	pET16b derivative for the production of *B. methanolicus* His$_{10}$-tagged AtlD from *E. coli* BL21 (DE3)	This study

Primers	Sequence [5' → 3']	Characteristics
pET16b-*atlF*	pET16b derivative for the production of *B. methanolicus* His$_{10}$-tagged AtlF from *E. coli* BL21 (DE3)	This study

Primers	Sequence [5' → 3']	Characteristics
P192	agcttcctttcgggctttgttagcagccgTTAATCAATAGGTGTCAACAATAC	Amplification of *atlD* for pET16b-*atlD*
P193	gccatatcgaaggtcgtcatatgctcgagATGAAAGCTTTAGTCAAAAAAG	
P194	agcttcctttcgggctttgttagcagccgTTATGATTTTTCTGGATGGAAG	Amplification of *atlF* for pET16b-*atlF*
P195	gccatatcgaaggtcgtcatatgctcgagATGAAAGCATTAAAGCTGTATG	

973   Amp^R: ampicillin resistance; overlapping regions are shown in lower case.

974　**Table 2.** Kinetic data of purified AtlD and AtlF

Condition[a]: protein, substrate, cofactor[b]	Substrate K_M (mM)	V_{max} (U mg^{-1})[c]	V_{max} / K_M
AtlD, Xyl5P, NADH	0.07 ± 0.03	1.33 ± 0.23	19
AtlD, Ribu5P, NADH	1.21 ± 0.42	0.49 ± 0.06	0.4
AtlF, Xyl5P, NADH	0.18 ± 0.05	0.11 ± 0.01	0.6

[a] The following reactions were also analysed, but no significant activity (i.e. < 0.05 U mg^{-1}) could be detected: AtlD,
Ribu5P, NADPH; AtlF, Xyl5P, NADPH; AtlF, Ribu5P, NADH; AtlF, Ribu5P, NADPH.
[b] Cofactor K_M values were analysed using 0.2 mM Xyl5P and were 0.01 ± 0.01 and 0.11 ± 0.09 for NADH and
NADPH, respectively.
[c] The V_{max} for AtlD, Xyl5P, NADPH was calculated using 0.2 mM of substrate and 0.3 mM of cofactor, and resulted
in 0.24 ± 0.06 U mg^{-1}.

Table 3: Physiological parameters of *B. methanolicus* cultures used for Metabolic Flux
Analysis. Growth rate and biomass quantities were deduced from OD$_{600}$ measurements in the
exponential growth phase. Carbon source evolution rates were measured from supernatant
samples and analyzed by NMR. Labelled CO_2 was monitored by MS in the bioreactor's outgoing
gas flow. All cultures were aerobic. n.a.: not measured; n.d.: not detected. The uncertainties
shown are standard errors between biological replicates.

Method	Methanol (CH3OH) 13C INST-MFA bioreactor (n=2)	Mannitol (C6H14O6) [1-13C] MFA flask (n=3)	Arabitol (C5H12O5) Mix. [1-13C] & [2-13C] MFA, flask (n=3)	Arabitol (C5H12O5) [5-13C] MFA flask (n=3)
Growth rate [h-1]	0.46 ± 0.002	0.36 ± 0.02	0.15 ± 0.004	0.1411 ± 0.02
Biomass yield [gDCW/gSubstrate]	0.55 ± 0.01	0.34 ± 0.08	0.19 ± 0.02	0.23 ± 0.02
C source [mmolC/gDCW/h]	21.9 ± 2.2	36 ± 7.3	21.4 ± 1.2	24.5 ± 0.75
CO2$_{out}$ [mmol/gDCW/h]	7.6 ± 0.04	n.a.	n.a.	n.a.
Acetate$_{out}$ [mmol/gDCW/h]	n.d.	1.75 ± 0.32	n.d.	n.d.

990
991
992
993

Fig. 1: Flux map of *B. methanolicus* central metabolism relative to substrate intake for methanol (A), mannitol (B) and arabitol (C). The thickness of the reaction lines represent the flux proportions relative to substrate incorporation into the metabolism: hps (A), Manupt (B), Araupt (C). Flux proportions were averaged among the biological replicates ($n_{methanol}=2$, $n_{arabitol} = n_{mannitol} = 3$). The direction of the reaction arrows represent the estimated net flux directionality. Large grey arrows represent a flux directed toward amino acid synthesis and biomass requirements for growth. Pathways discussed in the text are represented by a

background patch of color. Metabolites are represented by circles and have a solid bottom half if they are duplicated on the same panel, as per recommended by the SBGN standard. Metabolites subjected to experimental measurement of ^{13}C labeling are marked by a yellow box. Some intermediate metabolites are not needed for ^{13}C-MFA and are consequently lumped in mannitol and arabitol models (B, C); the same metabolites can be explicitly modeled in methanol model (A) since non-stationary ^{13}C-MFA requires all intermediates pools (e.g.: G3P, Cit, Aco). Reactions toward amino acid pools and biomass are not represented.

Fig. 2: Alternative assimilation pathways of D-arabitol with an example of the measured and expected labeling of Rib5P. Arabitol entry point into the metabolism is expected to be Ribu5P or Xyl5P, depending on the substrate specificity of the PTS system and the arabitol phosphate dehydrogenase. Taking [5-13C]arabitol as an example, we show that the labeling of downstream metabolites can be used to identify which pathway is operating *in vivo* (Ribu5P, orange; Xyl5P, blue). We exemplify our approach with one data point: the NMR specific enrichment of Rib5P and the associated estimates from the best fit of different scenario in which PTS1 or PTS5 are allowed to carry flux in the model (barplot). Unlike the others, the model in which only PTS5 was allowed to carry flux did not fit the data (red cross-mark, ☐). Circles are carbon atoms: solid for ^{13}C (●), empty for ^{12}C (○), dotted when irrelevant.

Fig. 3: Estimated absolute rates of production and consumption of NADPH, NADH/FADH2, and ATP. Rates (mmol/gDCW/h) are calculated as the sum of the estimated flux of the reactions modeled in the MFA that are expected to produce (positive value) or consume (negative value) those cofactors. Values are averaged over the biological replicates and the error bars are the associated standard deviation ($n_{methanol}$=2, $n_{arabitol}$ = $n_{mannitol}$ = 3). The growth requirements ("biomass", red) are computed from measures on *B. subtilis* (mmol/gDCW) and the growth rates we observed. Production and consumption rates should be balanced in our conditions, so we mark putative production/consumption rates as "unknown" to complete the balance when needed. The full production of NADH and FADH2 is assumed to be consumed in the respiratory chain ("oxidative phophorylation") and to produce ATP (with P/O$_{NADH}$=1.5, P/O$_{FADH2}$=1). We modeled the PTS systems of mannitol and arabitol as consumers of one equivalent ATP (in "uptake"). NADPH production: glucose-6-phosphate dehydrogenase (*zwf*), phosphogluconate dehydrogenase (*gnd*), isocitrate dehydrogenase (*idh*), and methylenetetrahydrofolate dehydrogenase from the linear detoxification pathway (detox). NADH production: glyceraldehyde-3-phosphate dehydrogenase (*pgk*), pyruvate dehydrogenase (*pdh*), 2-oxoglutarate dehydrogenase (*akgdh*), malate dehydrogenase (assimilated to *fum*), methanol mannitol and arabitol dehydrogenase (*mdh*, manupt, araupt), and formate dehydrogenase from the linear detoxification pathway (detox). FADH2 production: succinate dehydrogenase (*fum*). ATP consumption: 6-phosphofructokinase (*pfk*), PTS systems of mannitol and arabitol (manupt, araupt). ATP production: phosphoglycerate kinase (*pgk*), pyruvate kinase (*pyk*), succinyl-CoA synthetase (assimilated to *akgdh*), acetate kinase (*out_Ac*), and formate-tetrahydrofolate ligase from the linear detoxification pathway (detox).