Maternal- and Somatic-type snoRNA Expression and Processing in Zebrafish Development

Johanna F. B. Pagano[1+], Mauro D. Locati[1+], Wim A. Ensink[1], Marina van Olst[1], Selina van Leeuwen[1], Wim C. de Leeuw[1], Ulrike Nehrdich[2], Herman P. Spaink[2], Han Rauwerda[1], Martijs J. Jonker[1], Rob J. Dekker[1] and Timo M. Breit[1,*]

* To whom correspondence should be addressed. Tel: +31 20 5257058; Fax: +31 20 5257762; Email: t.m.breit@uva.nl

+ First two authors contributed equally to this publication

[1] RNA Biology & Applied Bioinformatics research group, Swammerdam Institute for Life Sciences, Faculty of Science, University of Amsterdam, Amsterdam 1090 GE, the Netherlands

[2] Department of Molecular Cell Biology, Institute of Biology, Leiden University, Gorlaeus Laboratories - Cell Observatorium, Leiden 2333 CE, the Netherlands

Maternal- and somatic-type snoRNAs in zebrafish

U3 U8 snoRNA, maternal expression, RNA processing, embryogenesis

ABSTRACT

Small nucleolar RNAs (snoRNAs) are non-coding RNAs that play an important role in the

complex maturation process of ribosomal RNAs (rRNAs). SnoRNAs are categorized in classes,

with each class member having several variants present in a genome. Similar to our finding

of specific rRNA expression types in zebrafish embryogenesis, we discovered preferential

maternal- and somatic-expression for snoRNAs. Most snoRNAs and their variants have

higher expression levels in somatic tissues than in eggs, yet we identified three snoRNAs;

U3, U8 and snoZ30 of which specific variants show maternal- or somatic-type expression.

For U3 and U8 we also found small-derived snoRNAs that lack their 5' rRNA recognition part

and are essentially Domain II hairpin structures (U-DII). These U-DII snoRNAs from variants

showed similar preferential expression, in which maternal-type variants are prominently

expressed in eggs and subsequently replaced by a somatic-type variants during

embryogenesis. This differential expression is related to the organization in tandem repeats

(maternal type) or solitary (somatic-type) genes of the involved U snoRNA loci. The

collective data showed convincingly that the preferential expression of snoRNAs is achieved

by transcription regulation, as well as through RNA processing. Finally, we observed small-

RNAs derived from internal transcribed spacers (ITSs) of a U3 snoRNA loci that via

complementarity binding, may be involved in the biosynthesis of U3-DII snoRNAs.

Altogether, the here described maternal- and somatic-type snoRNAs are the latest addition

to the developing story about the dual ribosome system in zebrafish development.

INTRODUCTION

Small nucleolar RNAs (snoRNAs) are a class of non-coding RNA molecules of variable length

(the majority being 60-200 nucleotides long), found in archaea and eukaryotes (1). SnoRNAs

are thought to mainly be involved in post-transcriptional modifications and maturation of

ribosomal RNAs (rRNAs) (2,3). However, recently additional functions have been ascribed to

specific snoRNAs, from regulation of mRNA editing and splicing (4) to post-transcriptional

gene silencing (5,6). SnoRNAs do not possess any intrinsic catalytic or modification activity,

but act both as a scaffold for partner proteins, forming small nucleolar ribonucleoproteins

(snoRNPs) and as guide for target specificity (7). Based on base-pairing interactions with

their target RNA, snoRNAs can thus direct the associated catalytic protein subunits to

accurately modify a specific RNA site (8).

In general, eukaryotic genomes can contain up to 200+ unique snoRNA genes (9). Based on

the presence of conserved sequence motifs, the majority of snoRNAs are classified into two

distinct classes: box C/D snoRNAs, which guide 2'-O-methylation of ribose, and box H/ACA

snoRNAs, which are involved in the isomerization of specific uridine residues to

pseudouridine (1). C/D snoRNAs are defined by the presence of two conserved motifs, the C

box (UGAUGA) and the D box (CUGA), found near the 5'- and 3'-end, respectively (10). In

the folded C/D snoRNA molecule, these two motifs are in close proximity of each other by

means of a hairpin structure and serve as a binding site for interacting proteins (11–13). In

addition, many C/D box snoRNAs may have less-well-conserved copies of the C and D motifs

(called C' and D'), which are also involved in the interaction with specific proteins (14–16). A

conserved region of 7-20 nucleotides upstream of the D (and the D', if present) box interacts

via base-complementarity with the rRNA, pinpointing the to-be-methylated RNA base,

which is usually the 5[th] nucleotide from the D or D' box (17–20). Additional base interactions

with the target rRNA can stimulate methylation by up to five-fold (16). H/ACA snoRNAs

typically feature a secondary structure consisting of two hairpins linked by a hinge region

that contains the H-box (ANANNA), followed by a short tail with the ACA-box (ACA) (21). The

hairpin regions contain internal bulges known as pseudouridylation pockets in which a

conserved region of 6-20 nucleotides is complementary to the target, but leaves a uridine

residue unpaired, marking it for enzymatic modification (22,23). However, more and more

snoRNAs are discovered that don't follow this classification. For example, there are snoRNAs

that have both C/D box and H/ACA box (24,25), or C/D snoRNAs that are not involved in 2'-

O-methylation, like snoRNA U3 and U8, which instead are essential in the processing of 18S

and 28S, 5.8S rRNAs, respectively (26–28).

Most vertebrate snoRNA genes are located in introns of genes that usually encode for

proteins related to ribosome biogenesis and protein synthesis (29) (Figure 1A). After

transcription of these genes by RNA polymerase II, the formation of functional snoRNAs

requires the processing of intronic RNA sequences, which are released during pre-mRNA

splicing (2). In vertebrates, only a few snoRNAs genes have an independent promoter and

are directly transcribed by RNA polymerase II or III (30,31). snoRNA genes are present either

as solitary snoRNA genes, or as clusters of multiple coding units; such clusters can consist of

the same or different snoRNA genes (9) (Figure 1A).

An increasing number of recently identified snoRNAs show differential expression among

different cell types and tissues, suggesting a role in distinct physiological processes (6). For

example, several studies discovered that certain snoRNAs are expressed mainly in the brain

where they might affect organ function and/or development (32,33).

Our group has recently described the presence of two distinct types of rRNAs expressed

during zebrafish development: a maternal-type, exclusively accumulated during oogenesis,

and replaced throughout the embryogenesis by a somatic-type, which is the only one rRNA

type present in adult somatic tissue (34,35). Since maternal- and somatic-type rRNAs show

ample sequence differences combined with the fact that snoRNAs interact with rRNA in a

sequence specific way, we hypothesized that there also might exist specific maternal- and

somatic-type snoRNAs. For this, we investigated the expression of the snoRNAome

throughout zebrafish development from egg to adult, by small-RNA-seq and discovered that

indeed complete and partial transcripts of snoRNA variants exist that show either maternal-

or somatic-preferential expression. Moreover, we determined that this developmentally

regulated expression is likely regulated both at the level of transcription, as well as RNA

processing. Together, with our recent finding that also a maternal-type spliceosome with

specific snRNA variants exists for early embryogenesis (36) everything points to the

existence of a comprehensive dual translation system in zebrafish embryogenesis.

RESULTS AND DISCUSSION

Cataloguing the zebrafish snoRNAs

In our previous studies (34,35), we identified two distinct types of rRNAs in zebrafish

development: maternal and somatic. The maternal-type rRNAs make up virtually all the

rRNA present in oocytes and are gradually, yet completely, replaced by the somatic-type

rRNAs during embryogenesis. There are, for each rRNA species, 5S, 5.8S, 18S and 28S,

significant sequence differences between the two types of rRNA, indicative of a substantial

functional difference. Given that snoRNAs are intricately involved in the complex

maturation process of rRNA via sequence specific interactions, we investigated whether

there are specific snoRNAs being co-expressed with maternal- and somatic-type rRNAs.

As the zebrafish genome is quite well annotated, we started by making an inventory of the

known genomic snoRNA sequences. In general, snoRNAs are categorized in two main

families based on the presence of conserved sequence motifs: the C/D box and H/ACA box

(37) and they can be found in the databases snOPY (38) and Ensemble 89 (39)

(Supplemental Table ST1 and Supplemental Figure SF1). Collectively, 257 snoRNA loci were

identified on the 25 zebrafish chromosomes. However, since some of these loci contained

identical sequences, 250 zebrafish snoRNA sequences were found to be unique

(Supplemental File SF1). In contrast, several snoRNAs appear to be present as different

variants, which we labeled with a unique identifier (Supplemental Table ST1). Most snoRNA

loci are present in introns (*intronic, 81%*) and rely for their expression on the transcription of

the associated genes (Figure 1A and 1C). Yet, a few snoRNA loci, mainly belonging to the

C/D box snoRNAs, are expressed as independent transcriptional units, consisting of one (*4%*)

or several (*15%*) snoRNA genes (Figure 1A and 1C, Table 1 and Supplemental Figure SF1).

Most transcriptionally self-regulated snoRNAs have been named U3 and U8 (40). The

snoRNA loci appear to be non-randomly distributed over the chromosomes, but no

apparent co-location with known rRNA sequences was observed (Supplemental Table ST1).

Figure 1. Genomic organization and RNA molecules from zebrafish snoRNA genes.
A: Different configurations of genomics organization of snoRNA genes illustrated by
exemplifying variants (upper row) and the different forms of RNA transcripts (middle rows).
For the U snoRNA genes also derived transcripts are indicated (lower row): snoRNA-DII
(green) are transcripts that exist just of the Domain-II part of the U snoRNA gene; sitsRNA
(yellow) are snoRNA-ITS-derived-small-RNAs that originate from ITS regions in clusters of
the tandemly repeated U snoRNA genes. The dotted lines indicate possible RNA processing
routes for snoRNA-DII transcripts.
B: The various snoRNA boxes are indicated in a schematic representation of the U3 snoRNA
(left panel). The possible interaction of a chromosome 14 U3 sitsRNA with a complete U3
snoRNA is indicated (middle panel), which may play a role in processing the latter at the
position of indicated by the red arrow to produce a U3-DII transcript. A schematic
representation of the U3-DII transcript found in this study (right panel).
C: Several characteristics of zebrafish snoRNA genes organized by their H/ACA or CD boxes.
[1]Distribution plot of lengths of snoRNA genes (Supplemental Figure SF1);
[2]snoU2_19 and SNORD94_a are present twice in the genome;
[3]U3_l and U3_k were found two and five times in the genome, respectively.
126

Developmental-stage-specific expression of two snoRNA types

After the identification and annotation of all snoRNAs in the zebrafish genome (GRCz10), we

investigated whether, similar to rRNAs, maternal- and somatic-type snoRNAs exist. For this,

we analyzed the differential expression between egg and adult zebrafish of the 204

expressed snoRNAs found by small-RNA-seq. The distribution of snoRNAs based on their

differential expression was clearly bimodal, with no snoRNA-variant being equally expressed

in egg and adult zebrafish (Figure 2A). Hence, all these snoRNAs displayed preferential

expression, as either maternal- (n = 18) or somatic-type (n = 186) snoRNA (Figure 2), albeit

less absolute as previously observed with maternal- and somatic-type rRNAs (Supplemental

Table ST2). The maternal-type snoRNAs turned out to be variants of just three snoRNAs: U3,

U8 and snoZ30, each of which also has one somatic-type variant (Figure 2B). Remarkably,

for the U3 and U8 snoRNAs, the variants in tandem repeats showed a maternal-preferential

expression profile, whereas the solitary variants were somatic-preferential (Figure 2B). For

both U3 and U8 in tandem repeats, there were some variants that showed no expression

(Supplemental Figure SF2). For the snoZ30 snoRNAs, the maternal- and somatic-type

variants are also organized in a different way on the genome with the maternal-type being a

solitary snoRNA, whereas the somatic-type is located intronically (Figure 2B). Thus, there is

a clear association between the genome organization of the snoRNA variants and their

expression during development.

There are ample sequence differences between the maternal and somatic snoRNA types

(Figure 3A). We used these sequence differences to confirm the developmental-stage

preferential expression of maternal- and somatic-type snoRNAs, by qRT-PCR-analysis for

several selected maternal- and somatic-type U3, U8 and snoZ30 snoRNAs in egg and adult

tissue. The qRT-PCR results are in in line with the small-RNA-seq results (Figure 3C).

To further characterize the shift from maternal-type snoRNA expression to somatic-type

expression, we determined the expression of the selected maternal- and somatic snoRNAs

during twelve stages of zebrafish embryogenesis. Similar to rRNAs, the predominant

maternal-type snoRNAs in eggs are, during embryogenesis, gradually replaced by somatic-

type snoRNAs (Figure 3B), extending the zebrafish dual ribosome system with snoRNAs.

156
Figure 2. Identifying maternal and somatic types snoRNAs.
A: Distribution plot of differential snoRNA expression between zebrafish egg and adult-male tissue.
Maternal-type snoRNA (blue), 18 snoRNA variants with a positive fold change, i.e. with a higher
expression in egg than in adult-male tissue; somatic-type snoRNA (red), 186 snoRNA variants with a
negative fold change.
B: Overview of maternal- and somatic-type snoRNA variants of U3, U8 and snoZ30 genes.
[1] Names of the associated variants within each of the indicated snoRNA locus.
[2] Genomic organization: intronic, solitary or in clusters of tandemly repeated (tand rep) genes.
Chromosome number (chrom) plus the number (n) of variant genes in a snoRNA locus.
[3] Normalized NGS-read counts of the snoRNA variants (cf. Supplemental Table ST2) in the same
cluster of tandemly repeated genes were added up.

168
Figure 3. Expression of maternal- and somatic-type snoRNAs in zebrafish development.
A: Sequence alignment of relevant maternal-type (M) and somatic-type (S) snoRNA variants;
identical nucleotides are indicated as dots, while gaps as dashes (For sequence alignment of all U
snoRNA cf. Supplemental Figure SF3). The qRT-PCR primers (panel C, Supplemental Table ST4) are
indicated with half arrows specific for the maternal-type (blue) or specific somatic-type (red). The
grey boxed sequences in the U3 and U8 snoRNAs indicate the absent regions for the observed U
snoRNA-DII transcripts (Supplemental Figures SF2 and SF3).
B: Relative expression of maternal-type (blue) and somatic-type (red) snoRNA variants indicated by
comparative percentage of U3-DII, U8-DII, and snoZ30 NGS-reads, respectively. Prot-mouth:
protruding-mouth; Adult FT: adult female-tail; Adult MWB: adult-male whole-body.
C: Relative expression of maternal-type (blue) and somatic-type (red) snoRNA variants indicated by
comparative percentage qRT-PCR of U3-DII, U8-DII, and snoZ30 NGS, respectively for qRT-PCR
analyses on zebrafish eggs and adult male-tail (Adult MT) tissue using RT-PCR primers as indicated in
panel A (Supplemental Table ST3).
D: Relative expression of maternal-type (blue) and somatic-type (red) U3 and U8 snoRNA variants
indicated by comparative percentages of RT-PCR-qSeq on zebrafish eggs and adult male-tail (A MT)
tissue using RT-PCR primers and subsequent quantitative sequencing (Supplemental Table ST4).

Structural characteristics of maternal- and somatic-types snoRNAs

A significant characteristic of the observed U3 and U8 snoRNA sequences was their length. Whereas all snoZ30 NGS-reads mapped to the whole genes (98 nt) (Supplemental Table ST1), almost all U3 NGS-reads (~130 nt.) represented just the 3' part (Domain II (DII), Figures 1A and B), including the C/D box, of the ~215 nt U3 genes (Supplemental Figure SF3). Hence, these U3 NGS-reads are missing the U3 5' part, including the A and A' boxes, which guides the snoRNA in interactions with target RNAs. Consequently, the length of these U3-DII snoRNAs is quite similar to all other H/ACA box snoRNAs (Supplemental Figure SF1). Similarly, almost all U8 NGS-reads (~86 nt) represented just the 3' part (Domain II, Figures 1A and 4), with C/D box, of the ~131 nt U8 genes, again missing the guiding 5' part. Whether complete U transcripts are truly absent or merely go undetected is addressed in the next section. The length of the U8-DII snoRNAs is quite similar to most C/D box snoRNAs (Table 1 and Supplemental Figures SF1 and SF3). U3-DII and U8-DII snoRNAs are reported before (13,41) and are likely a result of the removal of the mono-methylated (m^7G) 5' cap by the decapping complex, which is regulated by PNRC1 (42) although we cannot completely rule out that they also could originate from an alternative transcriptional start site (Figure 1A). The U-DII snoRNAs are not merely degradation products of the complete U3 and U8 snoRNAs, as can be seen by their gradual disappearance during the embryogenesis (Figure 2B) and by the strictly defined read coverage (Supplemental Figure SF3). Thus, beside an earlier reported dominant-negative effect on ribosome biogenesis (10) , we speculate that these partial U-DII snoRNAs may exert other specific functions. Since their guiding parts are missing (1,43), such function would be different from that of the traditional U snoRNA.

As a starting point for finding biological relevance of the observed maternal- and somatic-

expression profiles for different snoRNAs, we compared the sequences of the maternal-type

to the somatic-type variants of the U3, U8, and snoZ30 snoRNAs. There are quite some

sequence differences (8%, 16%, and 36%, respectively), which may be indicative for

functional diversity (Figure 3A and Supplemental Figure SF4). One obvious difference could

relate to the usual target RNAs for each snoRNA.

Several potential snoRNA-rRNA interactions could be pinpointed by complementary

sequences, in line with those reported in literature (1). However, many are in the 5' guiding

part of the involved snoRNAs, which is absent in our reads (Supplemental Figure SF5).

Conversely, many (known) interactions are found in the ETS regions of the rRNA genes

(Supplemental Figure SF5). Yet we observed that the involved maternal-type U-DII snoRNAs

are present throughout embryogenesis, much like the maternal-type rRNAs. This, combined

with the assumption that maternal-type rRNAs are only transcribed and processed during

oogenesis and somatic-type rRNAs during embryogenesis and adulthood, it seems unlikely

that maternal-type snoRNAs have a role in the processing of maternal-type rRNAs. Hence,

the observed U-DII snoRNA likely have other functions, much like snoZ30 that binds to

snRNA U6 (44). Given these constraints we were unable to identify any promising

interactions between the many possible interactions of U-DII snoRNAs and rRNAs

(Supplemental Figure SF5). It should be noted that such interactions are hard to find given

the sometimes seemingly feeble reverse complement base pairing in these situations.

Preferential expression of complete snoRNAs

The fact that virtually only U-DII snoRNA sequences were found by small-RNAseq, did not

match with the current knowledge of complete U3 and U8 snoRNAs, as well as a qRT-PCR

analysis which showed their abundant presence in the tested zebrafish samples (result not

shown). Given that we sequenced nearly all zebrafish tissues lead to the conclusion that,

with the NGS platform (Ion Proton) employed in this study, we are not able to sequence

complete U3 and U8 snoRNAs. This is also the case for many 5S rRNAs plus almost all tRNAs

and is likely caused by the cumulative effect of 5'/3' modifications, strong secondary

structures, and modified nucleotides.

To still determine whether the complete U snoRNAs also display preferential maternal or

somatic expression, we developed a PCR-based strategy that focused on the missing 5' parts

of the U snoRNA genes. To overcome the high similarity between the U3 variants, as well as

between the U8 variants, this approach starts with a RT-PCR, using as much as possible

generic PCR primers (Supplemental Table ST4), after which sequencing of the PCR products

reveals the distribution of variants (Supplemental Table ST4). This RT-PCR-qSeq approach

revealed that the complete U3 and U8 snoRNAs display the same expression patterns as the

shorter U3DII and U8DII (Figure 3D and Supplemental Table ST4).

Possible involvement of ITS sequences in U3-DII processing

While investigating the maternal-type U3 snoRNA tandem repeats, we observed additional

small RNAs that originate from the ITS regions between the snoRNA genes in the U3

genomic regions. These small RNAs either have their own promoter, or the U snoRNA genes

in tandem repeats generate one transcript which is later on processed into individual

mature U snoRNAs. As these small RNAs are derived from the U-ITS region, we named them

snoRNA-ITS-small-RNAs (sitsRNA). For instance, for the U3 loci on chromosome 14, we

detected essentially five different sitsRNAs (25 nt to 30 nt) that came from a highly

conserved region of 265 nt, just 379 nt upstream of each U3 variant sequence. We noticed

that there are several sitsRNAs from the various U snoRNA in tandem repeats that show

complementarity with U snoRNA sequences, in particular, one 26 nt sitsRNAs from the U3

loci on chromosome 14 showed two regions (each 7 nt long) that are reverse complement

to U3 snoRNA a sequence in the 5' part and one in the D-II region (Figure 1B). This raises the

intriguing possibility that this particular sitsRNAs may somehow be involved in the

processing from full length U3 snoRNA to U3-DII snoRNA as the locations of interaction

between the sitsRNAs and the U3 snoRNA span the cutting site (Figure 1B). Although similar

sites were found in the other U3 and U8 clusters, we were unable to find another sitsRNAs

that would interact like this. Still we feel there are enough indications that warrant further

investigation of the possible role of U-ITS sequences in snoRNA processing.

CONCLUSION

In this study, similar to our previously report on developmental-specific expression of

rRNAs, we observed a specific subset of snoRNAs; U3, U8 and snoZ30 of which variants

show distinct expression profiles during early zebrafish embryogenesis. All other snoRNAs

are about eight times higher preferentially expressed in non-embryonic developmental

stages, which may be a logical consequence of the fact that the rRNAs are already processed

in an egg, thus only requiring snoRNAs for other tasks than rRNA maturation during early

embryogenesis.

We discovered next to the complete U snoRNAs, also U3-DII and U8-DII partial snoRNAs,

which miss their 5' rRNA recognizing part and essentially consist of just the Domain-II

hairpin structures. These U-DII partial snoRNAs variants also showed maternal- or somatic

preferential expression, which correlated nicely with their genomic organization in tandem

repeats and solitary, respectively. The complete versus partial U snoRNAs show the same

preferential expression.

Though, at least part of the detected differential gene expression is likely caused by

associated promotors of the involved snoRNA genes, we were unable to find any sequences

in the 200 bp upstream promoter region that could discriminate the maternal- from the

somatic-type snoRNA genes (results not shown).

While the function of the U-DII snoRNAs is still unclear, given that the DII part of complete U

snoRNAs is known to bind to several proteins (45,46), the intact hairpin in a U-DII snoRNA

hints at a role as ribonucleoprotein. In any case, since they seem to be strictly regulated, U-

DII snoRNAs probably have a significant role in the zebrafish embryogenesis.

How they come about is another fascinating puzzle. We observed small RNAs, which

originate from the ITS regions of U3 snoRNA loci, that show convincing complementarity

with U3 sequences. This raises the possibility that they somehow may be involved in the

biosynthesis of the U3-DII snoRNAs. In any case, there is a notable analogy between the

relatively-small RNA-processing snoRNAs that are located in introns of genes usually

involved in ribosome biogenesis and small RNAs, which are located in the ITS of snoRNAs

they possibly support processing.

MATERIAL AND METHODS

Biological materials, RNA-isolation and small-RNA-seq

Adult zebrafish (strain ABTL) were handled in compliance with local animal welfare

regulations and maintained according to standard protocols (http://zfin.org). The breeding

of adult fish was approved by the local animal welfare committee (DEC) of the University of

Leiden, the Netherlands. All protocols adhered to the international guidelines specified by

the EU Animal Protection Directive 86/609/EEC.

For this study we used samples of two pools of unfertilized eggs (oocyte clutches) and two

male-adult zebrafish tails. The harvesting of the biological materials and RNA-isolation have

been described previously in (34) and (35).

Source data

Next-generation data previously generated in our group (34) and available through the

BioProject database with accession number PRJNA347637 has been used in this study with

respect to i) Three pools of unfertilized eggs (oocytes); ii) one embryo at each of the 12

developmental stages: 64 cells (2 hours post-fertilization (hpf)); high stage (3.3 hpf); 30%

epiboly stage (4.7 hpf); 70% epiboly stage (7 hpf); 90% epiboly stage (9 hpf); 4-somite stage

(11.3 hpf); 12-somite stage (15 hpf); 22-somite stage (20 hpf); prim-5 stage (24 hpf); prim-16

(31 hpf); long-pec stage (48 hpf); protruding-mouth stage (72 hpf); and iii) one whole-body

male-adult zebrafish sample.

qRT-PCR analysis

Forward and reverse PCR primers were designed for the maternal-type snoRNA genes U3_k,

U8_d and snoZ30_b, and for the somatic-type snoRNA genes U3_a, U8_e and snoZ30_a

(Supplemental Table ST3). Reverse transcription was done in two independent reactions

primed with the combined reverse PCR primers of either the three maternal-type variants

or the three somatic-type variants. Both reactions were performed on zebrafish egg pool

and whole-body adult male total RNA, in a total of four reactions. SuperScript IV Reverse

Transcriptase (Thermo Fisher Scientific) was used according to the manufacturer's

instructions. Quantitative real-time PCR (qRT-PCR) was performed on 10-fold dilutions of

the cDNA using a QuantStudio 3 Real-Time PCR System (Thermo Fisher Scientific).

RT-PCR-qSeq analysis

Forward and reverse PCR primers were designed for all U3 and U8 variants, in such a way

that: 1) as much as possible of the 5'-end of the full-length variants is included in the final

amplicon, and 2) generic primers were selected that bind to the maternal-type, as well as

the somatic-type variants (Supplemental Table ST4). cDNA was prepared as described above

and used in regular PCR reactions for each of the variants independently. Amplification was

performed using the Q5 High-Fidelity DNA Polymerase (New England Biolabs). The resulting

amplicons were purified using the QIAquick PCR Purification Kit (Qiagen) and their size was

verified on a 2200 TapeStation System (Agilent). Barcoded sequencing libraries were

prepared using a modified version of the Ion Xpress Plus Fragment Library Kit (Thermo

Fisher Scientific). Massive-parallel sequencing was performed on an Ion Proton System

(Thermo Fisher Scientific) using an Ion PI Chip Kit v3.

Bioinformatics analyses

Known snoRNA sequences. Known snoRNA sequences of *D. rerio* were downloaded from

Ensemble 91 (39) and from snOPY (38) in May 2017. A union was made of these two set of

sequences and sequence annotations. Characters were added to the snoRNA names to

uniquely distinguish the multiple variants of the same snoRNA and a FASTA file containing

only the unique snoRNA sequences was created.

Zebrafish snoRNA similarity tree. A hierarchical clustering was used to compare the snoRNA

sequences. The dendrogram was made by using the *hclust* and *stringDist* functions from the

R version 3.2.1 package 'stats' and 'Biostrings' respectively (47)

Mapping NGS-reads. At both 5' and 3' end of each snoRNA sequence, 5 Ns were added to

facilitate the alignments in the NGS-read mapping. NGS-reads longer than 20 nt from all

experiments, were mapped against the unique snoRNA sequences (Supplemental File

SFile1) using Bowtie2 (48) with the following settings: *-np* to 0, *- - score-min* to L, -1, -0.3 for

zebrafish in order to limit the maximal amount of mismatches to 5%. SAMtools v1.2 (49)

was used to convert the alignment to the BAM file format and to retrieve the mapped NGS-

read counts per snoRNA sequence. NGS-reads that were smaller than 50% of the length of a

snoRNA sequence were discarded.

Analysis of NGS-read mapping results. The NGS-read counts were scaled using total raw 5.8S

RNA NGS-read counts for each sample (35). The average of the three egg and the three

adult-male technical replicates was taken. A cutoff of at least 100 NGS-read count for egg

and adult-male combined was used to determine whether a snoRNA sequence was present

and to available for the analysis of differential gene expression, which was calculated in a

log2 scale of the egg over adult-male NGS-read counts.

Analysis of the RT-PCR-qSeq reads. A subsequence defined by the primers was selected for

U3, U8 snoRNA (*Supplemental Table 4*). These subsequences were then used to search

exact matching reads in the FASTQ files generated by the RT-PCR-qSeq experiments. The

number of exact matches is reported as read count for each variant of U3 and U8 for all

samples. The read counts belonging to the same tissues (clutch and adult male) for each

variant were added up. The percentage of each variant was calculated based on total tissue

reads.

snoRNA-rRNA interactions. BLASTn (50) was used to detect possible U3 and U8 snoRNA-

rRNA base-pairing interactions starting from known interactions in human. The word-size

parameter of BLASTn was set to 7 and of the resulting alignments the forward-reverse were

selected with at least 10 matching base-pairs. For snoRNA-ITS interaction analysis, this was

set to 7 matching base pairs.

ACKNOWLEDGMENTS

We are grateful for the valuable input by (anonymous) reviewers, which allowed us to

improve our manuscript. We acknowledge the support of The Netherlands Organization for

Scientific Research (NWO) grant number 834.12.003

REFERENCES

1. Watkins NJ, Bohnsack MT. The box C/D and H/ACA snoRNPs: Key players in the modification, processing and the dynamic folding of ribosomal RNA. Wiley Interdiscip Rev RNA. 2012;3(3):397–414.

2. Kiss T. Small nucleolar RNA-guided post-transcriptional modification of cellular RNAs. EMBO J. 2001;20(14):3617–22.

3. Jorjani H, Kehr S, Jedlinski DJ, Gumienny R, Hertel J, Stadler PF, et al. An updated human snoRNAome. Nucleic Acids Res. 2016;44(11):5068–82.

4. Falaleeva M, Pages A, Matuszek Z, Hidmi S, Agranat-Tamir L, Korotkov K, et al. Dual function of C/D box small nucleolar RNAs in rRNA modification and alternative pre-mRNA splicing. Proc Natl Acad Sci [Internet]. 2016;113(12):E1625–34. Available from: http://www.pnas.org/lookup/doi/10.1073/pnas.1519292113

5. Scott MS, Ono M. From snoRNA to miRNA: Dual function regulatory non-coding RNAs. Biochimie [Internet]. 2011;93(11):1987–92. Available from: http://dx.doi.org/10.1016/j.biochi.2011.05.026

6. Brameier M, Herwig A, Reinhardt R, Walter L, Gruber J. Human box C/D snoRNAs with miRNA like functions: Expanding the range of regulatory RNAs. Nucleic Acids Res. 2011;39(2):675–86.

7. Ellis JC, Brown DD, Brown JW. The small nucleolar ribonucleoprotein (snoRNP) database. Rna. 2010;16(4):664–6.

8. Sloan KE, Warda AS, Sharma S, Entian KD, Lafontaine DLJ, Bohnsack MT. Tuning the ribosome: The influence of rRNA modification on eukaryotic ribosome biogenesis and function. RNA Biol [Internet]. 2017;14(9):1138–52. Available from: https://doi.org/10.1080/15476286.2016.1259781

9. Matera AG, Terns RM, Terns MP. Non-coding RNAs: lessons from the small nuclear and small nucleolar RNAs. Nat Rev Mol Cell Biol [Internet]. 2007 Mar [cited 2014 Jul 11];8(3):209–20. Available from: http://www.ncbi.nlm.nih.gov/pubmed/17318225

10. Samarsky DA, Fournier MJ, Singer RH, Bertrand E. The snoRNA box C/D motif directs nucleolar targeting and also couples snoRNA synthesis and localization. EMBO J. 1998;17(13):3747–57.

11. Watkins NJ. Cbf5p, a potential pseudouridine synthase, and Nhp2p, a putative RNA-binding protein, are present together with Gar1p in all H BOX/ACA-motif snoRNPs and constitute a common bipartite structure. Rna. 1998;4(12):1549–68.

12. Watkins NJ, Ségault V, Charpentier B, Nottrott S, Fabrizio P, Bachi A, et al. A common

core RNP structure shared between the small nucleoar box C/D RNPs and the
spliceosomal U4 snRNP. Cell. 2000;103(3):457–66.

13.   Newman DR, Kuhn JF, Shanab GM, Maxwell ES. Box C/D snoRNA-associated proteins:
Two pairs of evolutionarily ancient proteins and possible links to replication and
transcription. Rna. 2000;6(6):861–79.

14.   Decatur WA, Fournier MJ. RNA-guided Nucleotide Modification of Ribosomal and
Other RNAs. J Biol Chem. 2003;278(2):695–8.

15.   Filipowicz W, Pogačić V. Biogenesis of small nucleolar ribonucleoproteins. Curr Opin
Cell Biol. 2002;14(3):319–27.

16.   Van Nues RW, Granneman S, Kudla G, Sloan KE, Chicken M, Tollervey D, et al. Box C/D
snoRNP catalysed methylation is aided by additional pre-rRNA base-pairing. EMBO J
[Internet]. 2011;30(12):2420–30. Available from:
http://dx.doi.org/10.1038/emboj.2011.148

17.   Nicoloso M, Qu LH, Michot B, Bachellerie JP. Intron-encoded, antisense small
nucleolar RNAs: The characterization of nine novel species points to their direct role
as guides for the 2'-O-ribose methylation of rRNAs. J Mol Biol. 1996;260(2):178–95.

18.   Kiss-László Z, Bachellerie J-P, Caizergues-Ferrer M, Kiss T, Henry Y. Site-Specific Ribose
Methylation of Preribosomal RNA: A Novel Function for Small Nucleolar RNAs. Cell.
1996;85(7):1077–88.

19.   Cavaille J, Nicoloso M, Bachellerie J. Targeted ribose methylation of RNA in vivo
directed by tailored antisense RNA guides. 1996;383(October):732–5.

20.   Chen CL, Perasso R, Qu LH, Amar L. Exploration of Pairing Constraints Identifies a 9
Base-pair Core within Box C/D snoRNA-rRNA Duplexes. J Mol Biol. 2007;369(3):771–
83.

21.   Ganot P, Caizergues-Ferrer M, Kiss T. The family of box ACA small nucleolar RNAs is
defined by an evolutionarily conserved secondary structure and ubiquitous sequence
elements essential for RNA accumulation. Genes Dev. 1997;

22.   Ganot P, Bortolin ML, Kiss T. Site-specific pseudouridine formation in preribosomal
RNA is guided by small nucleolar RNAs. Cell. 1997;89(5):799–809.

23.   Kehr S, Bartschat S, Tafer H, Stadler PF, Hertel J. Matching of soulmates: Coevolution
of snoRNAs and their targets. Mol Biol Evol. 2014;31(2):455–67.

24.   Jády BE, Kiss T. A small nucleolar guide RNA functions both in 2'-O-ribose methylation
and pseudouridylation of the U5 spliceosomal RNA. EMBO J. 2001;20(3):541–51.

25.   Marz M, Gruber AR, Höner Zu Siederdissen C, Amman F, Badelt S, Bartschat S, et al.
Animal snoRNAs and scaRNAs with exceptional structures. RNA Biol. 2011;8(6).

26.   Peculis BA, Steitz JA. Disruption of U8 nucleolar snRNA inhibits 5.8S and 28S rRNA
processing in the Xenopus oocyte. Cell. 1993;73(6):1233–45.

27.   Hughes JMX. Functional Base-pairing Interaction Between Highly Conserved Elements
of U3 Small Nucleolar RNA and the Small Ribosomal Subunit RNA. J Mol Biol
[Internet]. 1996;259(4):645–54. Available from:
https://doi.org/10.1006/jmbi.1996.0346

28.   Peculis BA, DeGregorio S, McDowell K. The U8 snoRNA gene family: Identification and
characterization of distinct, functional U8 genes in Xenopus. Gene. 2001;274(1–
2):83–92.

29.   Lui L, Lowe T. Small nucleolar RNAs and RNA-guided post-transcriptional
modification. Essey Biochem. 2013;(54):53–77.

30.   Dieci G, Preti M, Montanini B. Eukaryotic snoRNAs: A paradigm for gene expression

flexibility. Genomics [Internet]. 2009;94(2):83–8. Available from:
http://dx.doi.org/10.1016/j.ygeno.2009.05.002

31. Bratkovič T, Rogelj B. Biology and applications of small nucleolar RNAs. Vol. 68,
Cellular and Molecular Life Sciences. 2011. p. 3843–51.

32. Bachellerie JP, Cavaillé J, Hüttenhofer A. The expanding snoRNA world. Vol. 84,
Biochimie. 2002. p. 775–90.

33. Bratkovič T, Rogelj B. The many faces of small nucleolar RNAs. Biochim Biophys Acta -
Gene Regul Mech. 2014;1839(6):438–43.

34. Locati MD, Pagano JFB, Ensink WA, van Olst M, van Leeuwen S, Nehrdich U, et al.
Linking maternal and somatic 5S rRNA types with different sequence-specific non-LTR
retrotransposons. RNA [Internet]. 2017;23(4):446–56. Available from:
http://rnajournal.cshlp.org/lookup/doi/10.1261/rna.059642.116

35. Locati MD, Pagano JFB, Girard G, Ensink WA, van Olst M, van Leeuwen S, et al.
Expression of distinct maternal and somatic 5.8S, 18S, and 28S rRNA types during
zebrafish development. RNA [Internet]. 2017;23(8):1188–99. Available from:
http://rnajournal.cshlp.org/lookup/doi/10.1261/rna.061515.117

36. Pagano JFB, Dekker RJ, Ensink WA, van Olst M, Bos A, van Leeuwen S, et al. An
alternative spliceosome defined by distinct snRNAs in early zebrafish embryogenesis.
Not Publ.

37. Dupuis-Sandoval F, Poirier M, Scott MS. The emerging landscape of small nucleolar
RNAs in cell biology. Wiley Interdiscip Rev RNA. 2015;6(4):381–97.

38. Yoshihama M, Nakao A, Kenmochi N. snOPY: a small nucleolar RNA orthological gene
database. BMC Res Notes. 2013;6(1):426.

39. Zerbino DR, Achuthan P, Akanni W, Amode MR, Barrell D, Bhai J, et al. Ensembl 2018.
Nucleic Acids Res. 2018;46(D1):D754–61.

40. Olson MOJ. The Nucleolus. 1st ed. Olson MOJ, editor. Springer US; 2004. 364 p.

41. Watkins NJ, Leverette RD, Xia L, Andrews MT, Stuart Maxwell E. Elements essential
for processing intronic U14 snoRNA are located at the termini of the mature snoRNA
sequence and include conserved nucleotide boxes C and D. RNA. 1996;

42. Gaviraghi M, Vivori C, Pareja Sanchez Y, Invernizzi F, Cattaneo A, Santoliquido BM, et
al. Tumor suppressor PNRC 1 blocks r RNA maturation by recruiting the decapping
complex to the nucleolus . EMBO J. 2018;

43. Dutca LM, Gallagher JEG, Baserga SJ. The initial U3 snoRNA:pre-rRNA base pairing
interaction required for pre-18S rRNA folding revealed by in vivo chemical probing.
Nucleic Acids Res. 2011;39(12):5164–80.

44. Qu LH, Zhou H, Chen YQ, Du YP. The Schizosaccharomyces pombe mgU6-47 gene is
required for 2 '-O-methylation of U6 snRNA at A41. Nucleic Acids Res.
2002;30(4):894–902.

45. Watkins NJ, Lemm I, Ingelfinger D, Schneider C, Hoßbach M, Urlaub H, et al. Assembly
and maturation of the U3 snoRNP in the nucleoplasm in a large dynamic multiprotein
complex. Mol Cell. 2004;16(5):789–98.

46. Granneman S, Kudla G, Petfalski E, Tollervey D. Identification of protein binding sites
on U3 snoRNA and pre-rRNA by UV cross-linking and high-throughput analysis of
cDNAs. Proc Natl Acad Sci [Internet]. 2009;106(24):9613–8. Available from:
http://www.pnas.org/cgi/doi/10.1073/pnas.0901997106

47. R Development Core Team. R: A Language and Environment for Statistical Computing.
R Found Stat Comput Vienna Austria [Internet]. 2016;0:{ISBN} 3-900051-07-0.

Available from: http://www.r-project.org/

48.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012 Apr;9(4):357–9.

49.   Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009 Aug;25(16):2078–9.

50.   Madden T. Chapter 16 : The BLAST Sequence Analysis Tool. NCBI Handbook[internet].
2002;1–15.

SUPPLEMENTAL FIGURES

519   SF1.pdf: Zebrafish snoRNA characteristics.

520   SF2.pdf: Tandem repeat DII snoRNA expression distribution.

521   SF3.pdf: Maternal- and somatic-type snoRNA read coverage.

522   SF4.pdf: Sequence alignments of snoRNA variants.

523   SF5.pdf: Possible snoRNA-rRNA interactions

SUPPLEMENTAL TABLES AND FILES

526   SFile1.fa : Zebrafish snoRNA sequences

527   ST1.xlsx: Genome annotation of zebrafish snoRNAs

528   ST2.xlsx: Read counts from rna-seq experiment mapping to zebrafish snoRNAs

529   ST3.xlsx: qRT-PCR primers

530   ST4.xlsx: RT-PCR-qSeq experiments set-up and results.

A

U snoRNA

solitary

tandem repeats (TR)

DNA

pre-snoRNA

snoRNA

snoRNA-DII
sitsRNA

snoRNA

intronic

host gene

pre-mRNA

snoRNA

B

sitsRNA

A
A'
B
C
C
D
hinge

U3

U3-DII

C

Characteristics of zebrafish snoRNA genes

snoRNA		Avg. length[1]	Genomic organization			Total
Box	Subset		Intronic	Solitary	TR	
H/ACA	all	141	89	3	0	**92**
C/D	rest	91	119	3	0	**122[2]**
C/D	U3	215	0	3	35	**38[3]**
C/D	U8	131	0	1	4	5
			81%	*4%*	*15%*	*257*

Figure1

Figure3

A

B

Zebrafish snoRNA genes of which specific variants display maternal-type expression during development

snoRNA		Genomic organization (chrom, n)[2]	Normalized read counts[3]		Fold change range Egg / Adult male (log2)	Expression type
Name	Variant[1]		Egg	Adult male		
U3	a	solitary (c1, 1)	1,211	8,370	-2.8	somatic
	b-h	tand rep (c1, 7)	1,452	4	5.4 to 9.0	maternal
	k-t	tand rep (c14, 15)	22,297	7	8.1 to 12.0	maternal
	u-ag	tand rep (c20, 13)	1,318	3	7.0 to 8.8	maternal
U8	a-d	tand rep (c10, 4)	3,947	6	8.5 to 9.5	maternal
	e	solitary (c10, 1)	2,223	24,519	-3.5	somatic
snoZ30	a	intronic (3, 1)	450	8,382	-4.2	somatic
	b	solitary (c5, 1)	10,152	522	4.3	maternal
other		solitary intronic	*variable*	*variable*	-9.1 to -0.4	somatic

Figure2