In silico design of a T-cell epitope vaccine candidate for parasitic helminth infection

Ayat Zawawi[1, #a], Ruth Forman[1], Hannah Smith[1], Iris Mair[1], Murtala Jibril[1], Munirah Albaqshi[1], Andrew Brass[2], Jeremy P. Derrick[1,*], and Kathryn J. Else[1,*]

[1]Lydia Becker Institute of Immunology and Inflammation, School of Biological Sciences, Faculty of Biology, Medicine, and Health, University of Manchester, Manchester Academic Health Science Centre, Manchester M13 9PT, United Kingdom,

[2]Faculty of Biology, Medicine and Health, Division of Informatics, Imaging and Data Sciences, The University of Manchester, Manchester, UK.

[#a]Current address: Faculty of Applied Medical Sciences, King Abdul Aziz University, Jeddah, Saudi Arabia.

*Corresponding author

Email: Kathryn.j.else@manchester.ac.uk or Jeremy.derrick@manchester.ac.uk

Abstract

Trichuris trichiura **is a parasite that infects 500 million people worldwide, leading to colitis, growth retardation and Trichuris dysentery syndrome. There are no licensed vaccines available to prevent *Trichuris* infection and current treatments are of limited efficacy. *Trichuris* infections are linked to poverty, reducing children's educational performance and the economic productivity of adults. We employed a systematic, multi-stage process to identify a candidate vaccine against trichuriasis based on the incorporation of selected T cell epitopes into virus-like particles. We conducted a systematic review to identify the most appropriate *in silico* prediction tools to predict histocompatibility complex class II (MHC-II) molecule T-cell epitopes. These tools were used to identify candidate MHC-II epitopes from predicted ORFs in the *Trichuris* genome, selected using inclusion and exclusion criteria. Selected epitopes were incorporated into Hepatitis B core antigen virus-like particles (VLPs). A combined VLP vaccine containing four *Trichuris* MHC-II T-cell epitopes stimulated dendritic cells and macrophages to produce pro-inflammatory and anti-inflammatory cytokines. The VLPs were internalized and co-localized in the antigen presenting cell lysosomes. Upon challenge infection, mice vaccinated with the VLPs+T-cell epitopes showed a significantly reduced worm burden, and mounted *Trichuris*-specific IgM and IgG2c antibody responses. The protection of mice by VLPs+T-cell epitopes was characterised by the production of mesenteric lymph node (MLN)-derived Th2 cytokines and goblet cell hyperplasia. Collectively our data establishes that a combination of *in silico* genome-based CD4+ T cell epitope prediction, combined with VLP delivery, offers a promising pipeline for the development of an effective, safe and affordable helminth vaccine.**

Author Summary

The soil transmitted helminth *Trichuris trichiura* is a major parasite in developing countries; development of a comprehensive vaccine has been elusive. Here we used a systematic approach based on *in silico* identification of MHC-II T cell epitopes from genome sequences, their incorporation into a virus-like particle (VLP), characterization of the assemblies and testing in an *in vivo* murine infection model. Animals vaccinated with a preparation of four different VLP-antigen fusions showed significant reductions in intestinal worm burdens and associated antibody responses consistent with protection. The results suggest that a pipeline based on *in silico* prediction of potent MHC-II T cell epitopes, followed by incorporation into VLPs, could be a strategy which enables rapid translation into a vaccine against *Trichuris trichiura*.

Introduction

Trichuriasis, caused by the whipworm *Trichuris trichiura*, is one of the most widespread soil-transmitted helminths (STH) in the world (1). Global mass drug administration (MDA) programmes are being implemented, but cure rates are low, repeated treatments are costly and may prevent the development of acquired immunity. Further, the existence of drug-resistant parasites is a constant concern (2-4). Using the mouse model of human trichuriasis, *Trichuris muris* excretory/secretory (ES) products (5), ES fractions (6), extracellular vesicles (EVs) (7), and, more recently, *T. muris* whey acidic protein (8) in the context of the adjuvant alum, have shown considerable potential in a number of pre-clinical protection trials. Despite these successes, developing a vaccine based on native antigens is associated with many manufacturing challenges, including cost, time consumption, difficulties in purifying large quantities of worm antigens and control over differences between batches (9, 10). The advent of the genome era has provided alternative strategies for vaccine development (11). For example, the reverse vaccinology (RV) approach combines genome information with immunological and bioinformatics tools to overcome some of the limitations of conventional methods of screening vaccine candidates (12-14).

These observations, combined with the economic challenges of producing a low-cost vaccine, prompted us to examine the potential of using virus-like particles (VLPs) as a scaffold for the presentation of predicted *Trichuris* MHC-II epitopes. The Hepatitis B core protein (HBc) has been widely used as a VLP: it forms stable self-assemblies, which can accommodate T cell epitopes, is cheap to produce and is safe for human use (15, 16). Structural analysis has shown that each HBc monomer forms helical structures, which protrude as spikes from the capsid (17). Antigens, in the form of short T cell epitopes or whole globular domains, can be inserted at the tip of each spike. Designing vaccines based on 'multi-stage' antigens which are expressed at different stages of infection have shown promising results against several complex pathogens, such as *M. tuberculosis* and *Plasmodium* (18, 19). Further, CD4+ Th2

cells play essential roles in the development of protective immunity against *Trichuris* spp (20).The principal objective of this study was therefore to develop a novel MHC II T-cell epitope-based vaccine predicted from multi-stage *Trichuris* proteins, which will induce Th2 protective immunity. To achieve this aim, first, a systematic review was performed to select the optimal MHC class II *in silico* prediction tools. Second, potential *Trichuris* MHC-II T-cell epitope vaccine candidates obtained from the *Trichuris* genome were identified using the selected *in silico* prediction tool. Third, these epitopes were produced in a commercially viable manner by fusing epitopes in the hepatitis B core antigen (HBc-Ag) virus-like particle (VLP) vaccine delivery system. These VLPs+T-cell epitope vaccine candidates were then tested *in vitro* for their ability to activate antigen presenting cells (APCs). Finally, *in vivo* experiments were conducted to test the protective capacity of VLPs expressing different *Trichuris* T-cell epitopes *in vivo* using *T. muris* infection of mice. Collectively the results of this research represent the first significant progress towards identifying a novel, epitope-based vaccine for trichuriasis.

Results

The IEDB and NetMHC-II 2.2 tools exhibited similarly high levels of sensitivity for predicting epitopes with strong affinities. Based on a list of search terms (Supplementary Table S1) used on Google and other websites (Supplementary Table S2), 88 servers that predict T-cell epitopes based on MHC class I and II binding were identified (Supplementary Table S3). Of these 88, only 48 tools which could predict MHC-II epitopes were identified. Since our primary focus was MHC-II T-cell epitope prediction, only tools with that functionality were further evaluated using the inclusion criteria (Fig 1A). Five tools met the inclusion criteria: IEDB, SYPEITHI, NetMHC-II 2.2, Rankpep and ProPred. These tools were then scanned for the ability to predict MHC-II T-cell epitopes for two mouse alleles, I-Ab and I-Ad. The ProPred tool was excluded from further analysis because it only predicted HLA-DR

binding sites. The four epitope prediction tools that met all the selection criteria were

subsequently evaluated using an epitope training set to calculate the sensitivity with which

they could predict MHC II T-cell epitopes. Using the epitope training set (Supplementary Table

S4), it was observed that IEDB and NetMHC-II 2.2 tools had high levels of sensitivity

(~78.00%) for predicting MHC-II T-cell binding epitopes with high affinities, while Rankpep and

SYFPETHI exhibited low sensitivity (10.61% and 8.33%, respectively). Collectively, the data

indicate that the best prediction tools across all MHC-II T-cell prediction servers considered in

this study are IEDB and NetMHC-II 2.2.

Data acquisition and identification of potential vaccine component proteins. The

stichosome, which forms the majority of the whipworm anterior region, is thought to release

ES products which trigger the host immune response and induce host immunity after infection

(7, 21). Research conducted by Dixon et al., (22) and Else et al., (23), showed that antibody

recognition of high molecular weight proteins in both *T. muris* adult and larval ES correlated

with resistance to *T. muris* infection. To increase the chances of the identification of antigens

which induce a strong immune response, secreted and surface-exposed proteins associated

with the anterior region of L2, L3, male and female adult worms were selected for this study

(24).

Predicted ORFs from the 85-Mb genome (~11,004 protein-encoding genes) of *T. muris* and

the 73-Mb genome (~9650 protein-coding genes) of *T. trichiura* (24) were scanned for

potential vaccine candidates; 637 proteins were selected. From this subset, only proteins that

possess signal peptides and did not exhibit transmembrane domains were included, reducing

the total to a sample of 156 proteins. Full-length sequences of the *Trichuris* proteins were

obtained from the Universal Protein Resource (UniProt) database in FASTA format

http://www.uniprot.org/.

Elimination of closely homologous mouse and human proteins. To eliminate

potential autoimmune reactions when tested in mice and humans, the 156-protein subset was

checked to determine homology with human and mouse proteins using the basic local

alignment search tool (BLAST). All proteins with any degree of homology with humans or mice were excluded, leaving 60 candidate proteins. Of these, only upregulated proteins in the anterior region of L2, L3 and adult worms were selected based on their transcript expression level. This criterion was based on high-throughput transcriptome data generated from the RNA of *T. muris* and Gene Ontology (GO) term enrichment analyses, a transcriptional upregulation in a particular protein refers to a ≥10 Log2 (normalised read count) transcript expression level (24). Implementing this criterion, 27 proteins were selected for MHC-II T-cell epitope prediction (Fig 1B).

Prediction of *Trichuris* MHC-II T-cell binding epitopes. All 27 selected proteins were screened to predict T-cell MHC class II epitopes using the IEDB (consensus method) prediction tool (25). The analysis was carried out to predict the binding affinity to the MHC class II allele I-Ab mouse strain and the 11 most prevalent human class II HLA allele supertypes (25-28). The consensus method was used to select only those peptides with a low median percentile rank according to three different prediction methods to reduce the chance of failure during prediction. To cover common global alleles, peptides were selected based on their ability to bind to at least three different human alleles.

Conservation and allergens. To assess how well the predicted MHC-II T-cell peptides were conserved within the *T. trichiura* genome, the IEDB conservancy analysis tool was used. This tool calculates the degree of conservancy (i.e. similarity) of a peptide within a specified protein sequence (29). Only peptides that were >70% conserved with at least one homologous *T. trichiura* protein were selected for further analysis. Of the 219 MHC-II T-cell peptides, only 33 met these criteria. The 33 MHC-II peptides were then assessed for the prediction of IgE epitopes and allergenic potential using the AllerTOP v.2.0 server http://www.ddg-pharmfac.net/AllerTOP/ (30). The final set of 10 *Trichuris* MHC-II T-cell epitopes containing 33 overlapping peptides were predicted to have no allergenic potential. The 10 epitopes were further triaged to a final four epitopes based on solubility once expressed on HBc-Ag VLPs. A

flow diagram summarising the approach used is shown in Fig 1B, with the final 4 *Trichuris*

MHC II T-cell epitopes detailed in Fig 1C.

Purification and assembly of VLPs expressing *Trichuris* T-cell epitopes. Four

HBc-Ag fusion proteins were designed, incorporating each predicted T-cell epitope into the

major immunodominant region. The construct also included a Strep tag at the C-terminus, for

affinity purification (Supplementary Figure S1). A second round of purification using size

exclusion chromatography was performed to produce a homologous population of assembled

VLPs. HBc-Ag preparations expressing *Trichuris* MHC-II T-cell epitopes were pure by SDS

PAGE (Supplementry Figure S1 B-E). The endotoxin levels in all purified VLPs used in this

study were <0.2 endotoxin units (EU) as assessed using the ELISA-based endotoxin detection

assay (Data not shown).

VLPs expressing *Trichuris* T-cell epitopes induced the production of

proinflammatory cytokines *in vitro* and were internalised and co-localized in

APCs. BMDCs stimulated with different VLPs, irrespective of whether they include T-cell

epitopes, produced high levels of IL-6 (Fig 2A) and TNF-α (B) at levels equivalent to LPS and

ES-stimulated BMDCs. Similiarly, all VLPs activated BMDMs, inducing the secretion of high

levels of both TNF-α (Fig. 2E), and IL-6 (F). To visualise VLP internalisation by APCs, BMDCs

and BMDMs were stimulated with or without fluorescein-conjugated VLPs. Images of

individual cells taken by merging the brightfield (BF) and FITC (green) channels demonstrated

that the fluorescein-conjugated VLPs were internalised by both APCs (Fig 2C & G). To confirm

that the VLPs were co-localized within the APC lysosome and not the cell surface, BMDCs

and BMDMs were stained with the lysosome-specific LysoTracker dye. Cells were then

subsequently stimulated with or without fluorescein-conjugated VLPs. Merged images of the

brightfield (BF), FITC (green), and Lysotracker (red) channels revealed that the fluorescein-

conjugated VLPs were accumulated in the lysosome compartment of the BMDCs (Fig 2D) and

BMDMs (Fig 2H).

Immunization of mice with VLPs expressing *Trichuris* T-cell epitopes induced a

significant reduction in worm burden following challenge infection. The protective

capacity of the 4 T cell epitopes was assessed in the *T. muris* – mouse accelerated expulsion

model (31) (Fig 3A). Mice vaccinated with pre-mixed four VLPs+T-cell epitopes showed a

statistically significant reduction in worm burden by day 14 p.i. compared to the native VLP

(HBc-Ag; (P<0.01) and PBS (P<0.05) immunised groups (Fig 3B). In comparison, ES/Alum

immunised mice harboured no parasites (Fig 3C).

Immunization of VLPs expressing *Trichuris* T-cell epitopes induced humoral

immunity following challenge infection. To evaluate *T. muris* - specific serum antibody

responses induced by vaccination with VLPs+T-cell epitopes, parasite-specific IgM, IgG1, and

IgG2c serum antibody levels were determined at d14 p.i. Following vaccination and infection,

VLPs+T-cell epitopes and ES/Alum vaccinated mice had statistically significant higher levels

of parasite-specific IgM (Fig 3D) and IgG2c (F) compared to the control PBS and native VLP

(HBc-Ag) injected mice. However, high levels of parasite-specific IgG1 were only detected in

the serum of mice vaccinated with ES/Alum following *Trichuris* infection (Fig 3E). There was

no or very low levels of parasite-specific IgM, IgG1, and IgG2c detected in the serum of native

VLP (HBc-Ag) and PBS/alum injected mice at day 14 p.i., as shown in Fig. (3D-F). Similarly,

statistically significant higher levels of VLPs+T-cell epitopes-specific IgM (Fig 3G), IgG1 (H)

and IgG2c (I) were produced following *Trichuris* infection of mice vaccinated with VLPs+T-cell

epitopes, compared to mice given native VLP (HBc-Ag) or PBS. Notably, mice vaccinated with

ES/Alum also produced high levels of VLPs+T-cell epitopes-specific IgM following *Trichuris*

infection, as shown in Fig 3 (G).

Immunisation of mice with VLPs expressing *Trichuris* T-cell epitopes induces a

mixed Th1/Th2 immune response following challenge infection. To analyse the

cellular immune responses at the primary site of adaptive immune cell activation following *T.*

muris infection (32), MLN cells of mice vaccinated with VLPs+T-cell epitopes were re-

stimulated *in vitro* with *Trichuris* ES. Supernatants were assayed for Th2 cytokines (IL-4, IL-5,

IL-9 and IL-13), Th1/Th17 cytokines (IL-2, IFN-γ and IL-17), proinflammatory cytokines (IL-6

and TNF-α), and the anti-inflammatory cytokine (IL-10) production by CBA (Fig 4 and

Supplementary Figure S2). Levels reached significance versus control vaccinated mice for IL-

5 (Fig 4A; ES vaccinated mice) and IFN-γ (Fig 4D; VLP+T cell epitope vaccinated mice) with

non-significant increases in IL-9 and IL-13 (Fig 4B, 4C). MLN cells from all vaccinated mouse

groups also produced IL-4, IL-6, IL-10, IL-17 and TNF upon their re-stimulation *in vitro*

(Supplementary Figure S2 A-E). Collectively these data support the *in vivo* immunogenicity of

the novel VLP+ T cell epitope vaccine, evidencing the presence of a mixed Th1/Th2 immune

response and are in keeping with the worm expulsion and antibody data. Proximal colon

goblet cells were quantified in mice vaccinated with ES/Alum, PBS, VLP (HBc-Ag) and

VLPs+T-cell epitopes day 14 post *T. muris* infection. (Fig 4E). Interestingly, VLPs+T-cell

epitopes and ES/Alum vaccinated mice exhibited a significantly elevated goblet cell

hyperplasia (P<0.05) compared to mice injected with PBS or native VLP (HBc-Ag) (Fig 4F).

272   Discussion

Trichuris trichiura is one of the most common human STH parasites and remains a major

health concern for humans worldwide (1). A number of pre-clinical vaccines against trichuriasis

have been reported, containing whole *Trichuris* antigens or fractions (6, 7, 22). However,

developing a vaccine for *Trichuris* based on native antigens has several limitations (10). An

alternative strategy embraces the use of informatics to predict and assess MHC-II T cell

epitopes using criteria to maximize their *in vivo* protective potential. The fast-growing

application of VLP-based vaccines against several parasites, including *Plasmodium* spp (33,

34), *Leishmania infantum* (35), *Toxoplasma gondii* (36), *Trichinella spiralis* (37), and

Clonorchis sinensis (38) prompted our interest in developing a VLP-based vaccine against

trichuriasis.

Identification of novel *Trichuris* MHC-II T-cell epitopes as promising vaccine candidates. Initiation of an antigen-specific immune response requires presentation of antigenic peptides to CD4+ T-cells in the context of MHC Class II molecules (39). Selection of appropriate MHC Class II binding peptides is therefore a critical first step in developing an epitope-based vaccine. There are more than 80 computer-based prediction tools for identifying peptides that bind to MHC class I and II molecules, but not all are equivalent; some epitope prediction tools may fail to predict all significant epitopes (40). The performance of *in silico* prediction tools is affected by several factors. For example, MHC-II CD4+ T-cell epitope prediction tools have much lower accuracy than MHC-I tools because the MHC-I binding groove is closed, while the MHC-II groove is open at both ends (40, 41). Also, the use of a limited training dataset to evaluate the tools may affect performance (25, 42, 43). Further, the low performance of MHC class II prediction tools may not only be due to poor algorithm performance; the genetic diversity of human populations presents additional challenges (44). Thus, choosing the best bioinformatics tool to predict MHC class II T-cell epitopes is critical when designing epitope-based vaccines (45, 46). This study presents a systematic review of existing MHC-II restricted T-cell epitope prediction tools and evaluated four tools in order to establish the most appropriate bioinformatics tools currently available for predicting *Trichuris* MHC-II T-cell epitopes.

The IEDB and NetMHC-II 2.2 tools achieved similarly high levels of sensitivity of predicting binding epitopes with high affinities, while Rankpep and SYFPETHI exhibited low sensitivity. Each tool applies different methods of prediction; the IEDB tool uses a quantitative consensus method that combines the strengths of various methods (25); the NetMHC-II 2.2, another quantitative tool, uses an NN-align algorithm and weight matrix (28); and both SYFPEITHI (47) and Rankpep (48) are qualitative tools that use motif PSSMs. All the tools were 'user-friendly' but SYFPETHI was limited in its coverage of mouse and human MHC-II alleles. The IEDB tool has features that the other three do not, including a browser to input protein sequence formats

in a National Centre for Biotechnology Information (NCBI) database, seven different prediction

methods and an easy method of downloading the prediction output into an Excel spreadsheet.

Several studies have compared the performance of different MHC class II peptide binding

prediction tools (49-51), but the comparison presented in this study is different for two main

reasons. First, the MHC-II prediction tools were selected in a systematic way using

inclusion/exclusion criteria. Second, a unique dataset consisting of a new 'test' dataset which

has not been used to build or evaluate IEDB was used. For example, Zhao and Sher (2018)

evaluated the MHC-II prediction tools hosted on the IEDB analysis resource server using

newly available, untested data of both synthetic and naturally processed epitopes. Among the

18 predictors that were benchmarked, NetMHC-II outperformed all other tools, including

NetMHCIIpan and the consensus method for both MHC class I and class II predictions (52).

Furthermore, Andreatta et al. (2018) created an automated platform to benchmark six

commonly used MHC class II peptide binding prediction tools using 59 datasets that were

newly entered into the IEDB database and had not yet been made public to prevent biased

assessment of the available prediction tools. Their evaluation suggested that NetMHCIIpan is

currently the most accurate tool, followed by NN-align and the IEDB (consensus) tool (53).

Despite differences in the datasets used for comparison, these studies agree with the

comparison study conducted in our study, which found that the IEDB (consensus) and

NETMHC-II 2.2 (ANN) tools are among the best MHC-II prediction tools. However,

NetMHCIIpan could not be included in this study because it did not meet the inclusion criteria.

Collectively, we recommend the use of IEDB and NETMHC-II 2.2 prediction tools in any MHC

class II epitope prediction study to reduce the experimental cost of identifying epitopes.

However, the output of these tools needs to be carefully evaluated *in vitro* and *in vivo* before

they are used to bring an epitope-based vaccine to trial.

Since the advent of immunoinformatics tools for prediction of antigenic epitopes and protein

analysis, several VLP-based vaccines have been engineered to carry foreign antigens and

have proven to be highly immunogenic (54). To our knowledge, the data presented in this

study is the first to identify novel *Trichuris* MHC-II T-cell epitopes as potential vaccine antigen

candidates. The framework used here to identify potential epitope vaccine candidates within

the *T. muris* and *T. trichiura* genome could be used in the future to identify potential vaccine

candidates for other parasite species. The final set of *Trichuris* MHC-II T-cell epitope vaccine

candidates were derived from chitin-binding domain-containing proteins and chymotrypsin-like

serine proteases (Fig 1C).

Chitin-binding domains genes are highly expressed in different life stages in many nematodes

including the parasites *T. trichiura*, *Ascaris lumbricoides* and *Ancylostoma ceylanicum* and the

free-living nematode *Caenorhabditis elegans*, (55-58). In particular, these proteins are thought

to be associated with eggshell formation and early development at the single-cell stage (59,

60). Furthermore, given that, in addition to *T. muris*, more than 40 other helminth species

express high levels of chymotrypsin-like serine proteases, these proteins may also be

promising vaccine candidates for other helminths (24, 61). Chymotrypsin-like serine proteases

are thought to play central roles in either the invasion process or modulation of the host

immune response to enhance parasite survival (62-65). In addition, numerous publications,

using pre-clinical models of parasitic infection, have noted that protective immunity can be

consistently achieved using helminth protease molecules (66). For example, vaccinating

BALB/c mice with the whole recombinant serine protease of *T. spiralis* prior to challenge

infection led to a reduction in worm burden and induced a mixed Th1/Th2 immune responses

(67-69). Furthermore, Shears et al. (6) showed that vaccinating mice with the *T. muris* ES

fraction containing serine proteases induced high parasite-specific antibody responses.

Remarkably all the proteins selected in this study have been identified within the most

immunogenic fractions of *T. muris* ES following vaccination of mice (6).

Identification of a novel VLP-based vaccine against trichuriasis. All VLP

recombinant proteins stimulated a non-specific inflammatory response characterized by the

secretion of high levels of proinflammatory cytokines. Furthermore, by identification of

intracellular co-localization with lysosomes, VLPs were shown to be taken up by both BMDC

and BMDMs. These results are consistent with those of Serradell et al. (70) and Wahl-Jensen

et al. (71) who examined the activation of APCs in response to VLPs. For example, Serradell

et al. developed a (VLP-HA/VSP-G) vaccine composed of an influenza virus expressing

influenza virus hemagglutinin (HA) antigen, co-expressed with the extracellular region of

Giardia lamblia variant-specific surface proteins (VSPs) as an adjuvant. The recombinant

protein (VLP-HA/VSP-G) induced the secretion of high levels of TNF-α, IL-10, and IL-6

cytokines *in vitro*. They also showed that oral vaccination with the recombinant protein

protected mice from influenza infection and generated protective humoral and cellular

immunity. These results suggest that the VLPs are well placed to act as delivery systems to

drive immune responses. Further, the data raises the exciting prospect of modifying the VLPs

using adhesion molecules and/or cytokines co-displayed on the VLP surface, in order to target

the epitopes to specific APC subsets, thus enhancing the activation of antigen-specific T- cells

(72).

Remarkably, upon challenge with *T. muris* infection, mice vaccinated with 50 µg of pre-mixed

VLP+T-cell epitopes (HBc-CBD$_{1243-1259}$, HBc-CBD$_{241-257}$, HBc-CLSP$_{143-158}$, and HBc-CLSP$_{398-}$

$_{416}$), in the absence of any additional adjuvant, showed a significantly reduced worm burden.

Parasite-specific IgM and IgG2c were detected in the sera of mice vaccinated with the

VLPs+T-cell epitopes. Levels were equivalent to the control ES in alum-vaccinated mice, and

significantly higher than control vaccinated and infected mice. These results suggest that

these VLPs+T-cell epitopes are antigenic and can boost antibody response sufficiently to

recognize specific small peptides in the *T. muris* ES. Importantly, analysis of the serum from

immunised mice showed that vaccination with the VLPs+T-cell epitopes elicited high levels of

IgM, IgG1 and IgG2c to the VLP+T cell epitope recombinant protein pool (HBc-CBD$_{1243-1259}$,

HBc-CBD$_{241-257}$, HBc-CLSP$_{143-158}$, and HBc-CLSP$_{398-416}$) with limited recognition of the native

VLP (HBc-Ag) protein. In keeping with these data, the malaria VLP-based vaccine (Malarivax)

(73-75) composed of HBc-Ag expressing *P. falciparum* T cell and B-cell epitopes identified

from the circumsporozoite protein developed long-lasting immunity, eliciting a CD4+ T cell

immune response and is currently undergoing a clinical trial (15, 76, 77). The results of these

studies support some critical insights into VLP-based vaccines, including confirming that an

HBc virus-like particle, in particular, is an excellent delivery system for developing potential

vaccine candidates for parasites.

A proportion of mice vaccinated with VLP+T-cell epitopes produced detectable levels of MLN-

derived Th2 cytokines IL-5, IL-9, and IL-13 in response to re-stimulation *in vitro* with *Trichuris*

ES. The Th1 cytokine IFN$-\gamma$ was also significantly elevated above levels detected in control

mice. These data indicate that vaccine-induced protective immunity is characterized by a

mixed Th1/Th2 immune response, as has been previously reported (21, 78). Similarly, Gu et

al (78), reported that the protective immunity to *Trichinella spiralis* infection induced by

vaccination with CD4+ T cell epitopes was associated with both Th1 and Th2 cytokines.

Whilst the mechanism by which vaccination protects mice from *T. muris* infection remains

unclear, this study reveals that vaccination of mice with VLPs+T-cell epitopes or ES/Alum

promoted a marked goblet cell hyperplasia (79). Goblet cells, and the mucins they produce,

have been implicated in Th2-mediated defence in mice resistant to a primary *T. muris* infection

(80, 81). However, it remains to be determined whether the goblet cell hyperplasia seen here

in vaccinated mice is simply a Th2 correlate or is functionally important in the protection

observed. In summary, the current study describes the development and efficacy of a novel

epitope-based vaccine against trichuriasis. VLPs expressing different *Trichuris* MHC-II T-cell

epitopes, predicted from chitin-binding domain containing proteins and chymotrypsin-like

serine proteases, were shown to promote protective immunity *in vivo*. Collectively, given the

right combination of immunoinformatics and immunogenicity screening tools, epitope-based

vaccines will undoubtedly limit the cost and effort associated with bringing a *Trichuris* vaccine

to trial.

Methods

Search strategy. A protocol was designed to identify the bioinformatics tools that can predict

MHC-II T-cell epitopes in accordance with the well-defined Preferred Reporting Items for

Systematic Reviews and Meta-Analyses (PRISMA) (82). The search was limited to the English

language and used the search terms shown in (Supplementary Table S1). A list of the search

terms was first used in December 2015 on Google and other websites (Supplementary Table

S2) to screen for MHC class I and II *in silico* prediction tools. The number of citations (>200),

number of publications (>200), online availability, last update and community were considered

to determine whether the tools would be selected for further analysis. The number of

publications and citations for each tool were obtained from Google Scholar. In the case of

duplicate citations, the highest number of citations was used. A flow diagram of the systematic

review screening process for MHC-II T-cell bioinformatics prediction tools is shown in Fig. 1A.

Construction of an epitope training set. To evaluate the performance of the tools, a

literature search for MHC-II CD4+ T-cell peptide binding datasets was performed using the

search terms 'universal T-cell epitopes' and 'MHC class II T-cell epitopes' on Google Scholar

in December 2015. These datasets included epitopes with publicly available sequences that

the literature had experimentally validated to be immunogenic in mice. Because training sets

of peptides were used for the development of epitope prediction tools, care was taken to

exclude the training datasets used for tool development (83-85).

In order to find the optimal set of epitopes, two different sets of epitopes were analysed. The

first set was composed of publicly available series of peptide sequences from 15 different

proteins that can bind to MHC class II molecules. The second set included one untested

protein of *T. muris*, which served as a control to evaluate the performance of the tools.

Collectively, the training dataset was composed of 145 epitopes from 16 different proteins

(Supplementary Table S4).

Prediction of *Trichuris* MHC-II T-cell epitopes. Full-length sequences of proteins containing

T-cell epitopes were obtained from the Universal Protein Resource (UniProt) database

http://www.uniprot.org in FASTA format. The Immune Epitope Database (IEDB), NetMHC-II

2.2, Rankpep and SYFPEITHI tools were used to predict T-cell epitopes for mouse strains

with I-Ab and I-Ad mouse alleles. Each tool applied a different prediction method and scoring

system to generate the predictions output. For instance, the IEDB tool uses the consensus

method for prediction, which combines the NN-align, SMM-align, combinatorial library and

Sturniolo methods (25), while the NetMHC 2.2 tool uses ANNs (28). The median percentile

rank (%) of the three prediction methods was used to generate the rank for the consensus

method. A small numbered percentile rank (%) indicates that a peptide has a high binding

affinity to MHC class II alleles (25).The output of the IEDB and NetMHC 2.2 tools (the binding

affinities) were expressed as half-maximal inhibitory concentration IC50 (nM) values. Epitopes

that bind with an affinity of <50 nM are considered to have high affinity, those that bind with

<500 nM have an intermediate affinity and those that bind with <5000 nM have a low affinity

(28). All epitopes with high or intermediate affinity are considered 'true binders', while epitopes

with low affinity are considered 'nonbinders'. No known T-cell epitope has an IC50 value of

>5000 nM (25). The scoring system of the SYFPEITHI prediction tool depends on whether

peptide amino acids are frequently occur in anchor positions. Optimal anchor residues are

given the value 15 and scores of -1 or -3 points are given to amino acids that have a negative

effect on epitopes' binding ability at a certain sequence position. Epitopes that bind strongly

are among the top 2% of all peptides predicted in 80% of all predictions results (47). The

Rankpep tool uses position-specific scoring matrices (PSSMs) to predict MHC-II T-cell

epitopes (48). A high peptide score percentage indicates that the epitope is likely to bind to

the set of aligned peptides that bind to a given MHC-II molecule (48). The peptide lengths in

all the resulting sets were based on the complex of 15 mer core region peptides of MHC class

II molecules.

Evaluation and statistical analysis. Using the training set of epitopes, the performance of

the four MHC-II epitope prediction tools, selected through our inclusion/exclusion criteria, were

assessed as weak, intermediate or high binders. The prediction results were classified into

two categories, true positive (TP) and false negative (FN), based on the threshold values (86).

In addition, the evaluation assessed sensitivity (TP/ [TP+FN]). Nonparametric Spearman

correlation and Bland-Altman analyses were performed to show the relationship and

agreement between the scores derived from the NetMHC-II 2.2 and IEDB tools. The level of

significance was set at $p < 0.05$ for the correlation test.

Mice and Parasites. 6–8 weeks male C57BL/6 mice (Envigo) were fed autoclaved food and water and were maintained under specific pathogen-free conditions. Parasite maintenance, ES collection from adult *T. muris* worms and the method used for infection and evaluation of worm burden were carried out as described previously (87).

Ethics Statement. All animal experiments were approved by the University of Manchester Animal Welfare and Ethical Review Board and performed under the regulation of the Home Office Scientific Procedures Act (1986), and the Home Office approved licence 70/8127.

Plasmid construction. The coding sequence for native VLP (HBc-Ag) was engineered with BamHI + EcoRI sites to allow insertion of peptide antigen sequences into the major immunodominant region (MIR). The entire HBc-Ag coding sequence was inserted into the pET17b expression vector between the NdeI (CATATG) and XhoI (CTCGAG) restriction sites. The insertion of each MHC-II T-cell epitope into the MIR was obtained by annealing the relevant oligonucleotide primers (Supplementary Table S5) with BamHI + EcoRI restriction sites and ligation into BamHI/EcoRI cut HBc-Ag in the pET-17b vector. Constructs containing MHC-II T-cell epitopes were confirmed by DNA sequencing.

Production and purification of VLP recombinant proteins. The recombinant plasmids were transformed into ClearColi-BL21 (DE3) electrocompetent cells (Lucigen) by heat-shock transformation. The bacterial culture was grown in LB media containing 100 μg/mL ampicillin and incubated at 37°C for 14 hours in a shaker incubator. The transformed cells at a starting optical density (0.1 OD600) were inoculated into LB liquid media supplemented with 100 μg/mL ampicillin for 3-4 hours until the optical density at 595 nm (OD600) reached 0.6. Isopropyl-b-D-thiogalactopyranoside (IPTG) was added to the culture to a final concentration of 0.4 mM, and the culture was grown by continuous shaking for 12-16 hours at 16°C. The cells were then harvested by centrifugation, resuspended and sonicated in Strep-Tag washing buffer (100 mM Tris-HCl, 150 mM NaCl, 1 mM EDTA) containg 1 tablet of cOmplete TM EDTA- free protease inhibitor cocktail (Sigma-Aldrich) per 50 mls of the resuspended cells along with 5 μg/ml DNase I (Sigma-Aldrich). The cell suspension was then disrupted by ultra-sonication on ice using Banddelinuw 3200 (Sonoplus) amplitude (Am) in 35% for 5 sec on and

10 sec off for 5-8 minutes. The supernatant containing soluble recombinant protein was

harvested following centrifugation at 18,900 x g for 40 mins at 4°C using Sorvall RC Plus with

the Fiber-Lite F21 8x50y rotor. Finally, the soluble supernatant was filtered through 0.22 µm

pore size filters. After filtration, Strep(II)-tag proteins were purified using affinity column

chromatography using a StrepTrap™ column prepacked with StrepTactin Sepharose by

following the manufacturer's protocol (GE Healthcare). The VLP recombinant proteins were

further purified by size exclusion chromotogrophy (Superose 6, 10/300 GL;GE Healthcare).

The level of endotoxin in all the purified VLPs was measured with an ELISA-based endotoxin

detection assay (Hyglos) following the manufacturer's protocol.

SDS-PAGE. The VLP recombinant protein samples were subjected to 10% SDS-PAGE and

subsequently characterized by TEM as described previously [20]. Briefly, the VLP

recombinant proteins were separated on 10% polyacrylamide gels and stained with instant

blue protein stain (Expedeon).

***In vitro* stimulation of BMDCs and BMDMs.** BMDCs were harvested from mice into

RPMI 1640© (Sigma-Aldrich) supplemented with 5% heat-inactivated FCS (Hyclone, Logon,

UT), 100 µg /mL penicillin and 100 µg/mL streptomycin (Sigma-Aldrich), 2 mM L-Glutamine

(Sigma-Aldrich), 50 µM β-mercaptoethanol (Gibco) and 20 ng/ml granulocyte-macrophage

colony-stimulating factor (GM-CSF) (eBioscience). BMDMs were harvested from mice into

modified Eagle's Medium (DMEM) – high glucose (Sigma-Aldrich) supplemented with 5%

heat-inactivated FCS (Hyclone, Logon, UT), 100 µg/mL penicillin and 100 µg/mL streptomycin

(Sigma-Aldrich), Eagle's minimum essential medium (MEM) non-essential amino acid solution

(Sigma-Aldrich), and 0.1 mg/ml macrophage colony-stimulating factor (M-CSF) (eBioscience).

BMDMs and BMDCs on day 8 were collected at $1X10^6$/ml and incubated with 10 µg/ml VLP

recombinant protein for overnight at 37°C. ES at a final concentration 50 µg/ml and

lipopolysaccharide (LPS) at final concentration 0.1 µg/ml were used as positive controls, and

untreated cells were used as a negative control. After 24 hours, the supernatants were

harvested separately and stored at -20°C until analysed by LEGENDplex for cytokine content

according to the manufacturer's instructions.

Fluorescein-conjugated VLP internalization and localization in APCs. BMDCs and BMMs on day 8 were collected at $1X10^6$/ml and incubated with 10 µg/ml with fluorescein-conjugated VLP separately (HBc-CBD$_{1243-1259}$, HBc-CBD$_{241-257}$, HBc-CLSP$_{143-158}$ and HBc-CLSP$_{398-416}$) for overnight at 37°C in 5% CO_2. Unstimulated BMDCs and BMMs were used as negative controls. Next day, cells were incubated with 50 mM LysoTracker dye obtained from (Invitrogen) for 45 min prior to harvesting to visualise the lysosomal localisation of fluorescein-conjugated VLPs by ImageStreamX cytometry. A Brightfield-1 filter was employed to image dendritic cells and macrophages, a fluorescein isothiocyanate (FITC-488 nm) filter to image fluorescein-conjugated VLPs, and an (APC-592 nm) filter to image lysosomes. The data were analysed using IDEAS software version 6.2.187.0. The degree of co-localization was measured by the Bright Detail Similarity (BDS-R3) on a cell-by-cell basis. A Bright Detail Similarity value of 1.0 indicates a high degree of similarity between two images in the same spatial location (correlated) and a value around 0 has no significant similarity (uncorrelated).

Immunization schedule and challenge infection Mice were divided randomly into 4 groups: mice were inoculated s.c with overnight ES antigens emulsified with an equal volume of Alum- an aluminum salt adjuvant (88) obtained from Thermo Scientific to achieve a vaccine dose of 100 µg ES in 100 µl Alum as a positive control. The negative group was inoculated s.c with 200 µl of sterile PBS. Alternatively, mice were vaccinated with 200 µl of 50 µg VLP (HBc-Ag). The vaccinated mouse group was administrated with 200 µl of 50 µg pre-mixed VLPs+T-cell epitopes (HBc-CBD$_{1243-1259,}$ HBc-CBD$_{241-257,}$ HBc-CLSP$_{143-158}$ and HBc-CLSP$_{398-416}$). All experimental groups were vaccinated in the scruff of the neck at (day -20) and were boosted with the same dose at (day -10) in the same site. At day 10 post vaccination, mice were challenged orally with a high dose of *T. muris* (200 eggs) per mouse. Mice were sacrificed 14 days after challenge infection by an overdose of CO_2.

Ag-specific antibody detection in the serum. Blood samples were collected immediately from mice by cardiac puncture and left at room temperature to clot. Parasite-specific and VLP recombinant protein-specific antibodies (IgM, IgG1 and IgG2c) were determined in sera by enzyme-linked immunosorbent assay (ELISA) as previously described (89). Briefly, 96-well

plates were coated with 50 µg/well of the overnight *T. muris* E/S antigen at 5 µg/ml or with 50

µg/well of the purified VLP recombinant protein in 0.5 M carbonate-bicarbonate buffer for

overnight at 4°C. The plates were washed and then blocked with 3% BSA (Sigma-Aldrich) in

PBS Tween-20 (PBS-T20) (0.05% Tween 20, Sigma-Aldrich) for 1 h at 37 °C. After washing,

diluted sera were added and incubated for 1 h at 37 °C. Antibody responses were detected

using the Biotinylated rat anti-mouse IgM, IgG1 and IgG2a/c (BD Biosciences). After washing,

streptavidin peroxidase (Sigma-Aldrich) was added to the plates and incubated for 1 hour at

room temperature. The TMB ELISA substrate (3, 3', 5, 5'-tetramethylbenzidine- Thermo) was

used to develop color and stopped with 0.003% (H_2SO_4). The optical density at 450 nm with

a reference of 570 nm was measured with a Dynex MRX11 plate reader (DYNEX

Technologies) at 450 nm with a reference of 570 nm.

***In vitro* stimulation of MLN.** MLNs were harvested from mice at autopsy and dissociated in

RPMI-1640 medium (Sigma-Aldrich) supplemented with 10% foetal calf serum (FCS), 1% L-

Glutamine, and 1% penicillin/streptomycin (all Invitrogen Ltd., Paisley, UK). 5×10^6/ml MLN

cells were re-stimulated *in vitro* in 96-well plates with 50 µg/ml *T. muris* E/S antigens and with

10 µg/ml pre-mixed VLPs+T-cell epitopes proteins (HBc-CBD$_{1243-1259}$, HBc-CBD$_{241-257}$, HBc-

CLSP$_{143-158}$, and HBc-CLSP$_{398-416}$) at 37 °C for 48 hours. Cytokine concentration was quantified

in the supernatants by cytometric bead array (CBA) according to the manufacturer's

instructions from the Mouse/Rat Soluble Protein Flex Set System kit (BD Biosciences

Pharmingen, Oxford, UK). CBA data was analysed on a MACSQuant analyser with

MACSQuantify™ Software and the CBA analysis software package (BD Biosciences).

Histology. At autopsy, colonic tissue samples were removed and fixed at room temperature

for overnight in 10% neutral buffered formalin, prior to storage in 70% EtOh and processing

and embedding in paraffin wax. 5 µm thick serial sections were cut on a Microrn HM325

microtome (Microm International, Germany), de-waxed in citroclear and rehydrated prior to

periodic acid–Schiff staining. Stained slides of proximal colon sections were scanned using

3D Histech Pannoramic 250 flash slide scanner. Photographs of the sections were taken at

100X magnification using panoramic viewer version 1.15.4 software. The number of goblet

cells was determined as the total number of PAS-positive cells in 60 randomly selected crypts

in three fields of view from each section.

Statistics Statistical analyses were performed using Graph Pad Prism, version 7.00 software.

In all tests, P≤0.05 were considered statistically significant and were determined using the

Kruskal–Wallis non-parametric ANOVA for comparing multiple groups.

Acknowledgments

593   We thank the University of Manchester Histology, Bio-imaging and Flow Cytometry Core

594   Facility for facilitating the image analyses.

595
References
597

1.   Alexander TY, Blackburn BG. Soil-Transmitted Helminths: *Ascaris*, *Trichuris*, and *Hookworm*
Infections. Water and Sanitation-Related Diseases and the Environment: In the Age of Climate Change.
2019:95.
2.   WHO WhO. Soil-transmitted helminthiasis: Deworming campaign improves child health,
school attendance in Rwanda Geneva: WHO Library. 2015.
3.   Farrell SH, Coffeng LE, Truscott JE, Werkman M, Toor J, de Vlas SJ, et al. Investigating the
Effectiveness of Current and Modified World Health Organization Guidelines for the Control of Soil-
Transmitted Helminth Infections. Clinical Infectious Diseases. 2018;66(suppl_4):S253-S9.
4.   Speich B, Ali SM, Ame SM, Bogoch II, Alles R, Huwyler J, et al. Efficacy and safety of albendazole
plus ivermectin, albendazole plus mebendazole, albendazole plus oxantel pamoate, and mebendazole
alone against *Trichuris trichiura* and concomitant soil-transmitted helminth infections: a four-arm,
randomised controlled trial. The Lancet infectious diseases. 2015;15(3):277-84.
5.   Jenkins SN, Wakelin D. The source and nature of some functional antigens of *Trichuris muris*.
Parasitology. 1977;74(2):153-61.
6.   Shears RK, Bancroft AJ, Sharpe C, Grencis RK, Thornton DJ. Vaccination Against Whipworm:
Identification of Potential Immunogenic Proteins in *Trichuris muris* Excretory/Secretory Material.
Scientific reports. 2018;8(1):4508.
7.   Shears R, Bancroft A, Hughes G, Grencis R, Thornton D. Extracellular vesicles induce protective
immunity against *Trichuris muris*. Parasite immunology. 2018:e12536.
8.   Briggs N, Wei J, Versteeg L, Zhan B, Keegan B, Damania A, et al. *Trichuris muris* whey acidic
protein induces type 2 protective immunity against whipworm. PLoS pathogens.
2018;14(8):e1007273.
9.   Geldhof P, De Maere V, Vercruysse J, Claerebout E. Recombinant expression systems: the
obstacle to helminth vaccines? Trends in parasitology. 2007;23(11):527-32.
10.   Hewitson JP, Maizels RM. Vaccination against helminth parasite infections. Expert review of
vaccines. 2014;13(4):473-87.
11.   Del Tordello E, Rappuoli R, Delany I. Reverse Vaccinology: Exploiting Genomes for Vaccine
Design. Human Vaccines: Emerging Technologies in Design and Development. 2017:65-86.
12.   Capecchi B, Serruto D, Adu-Bobie J, Rappuoli R, Pizza M. The genome revolution in vaccine
research. Current issues in molecular biology. 2004;6:17-28.
13.   Moise L, Cousens L, Fueyo J, De Groot AS. Harnessing the power of genomics and
immunoinformatics to produce improved vaccines. Expert opinion on drug discovery. 2011;6(1):9-15.

14.   Arnon   R,   Ben-Yedidia   T.   Old   and   new   vaccine   approaches.   International
immunopharmacology. 2003;3(8):1195-204.
15.   Gregson AL, Oliveira G, Othoro C, Calvo-Calle JM, Thorton GB, Nardin E, et al. Phase I trial of
an alhydrogel adjuvanted hepatitis B core virus-like particle containing epitopes of *Plasmodium*
falciparum circumsporozoite protein. PloS one. 2008;3(2):e1556.
16.   Roose K, Baets SD, Schepens B, Saelens X. Hepatitis B core–based virus–like particles to
present heterologous epitopes. Expert review of vaccines. 2013;12(2):183-98.
17.   Wynne S, Crowther R, Leslie A. The crystal structure of the human hepatitis B virus capsid.
Molecular cell. 1999;3(6):771-80.
18.   Mosavat A, Soleimanpour S, Farsiani H, Sadeghian H, Ghazvini K, Sankian M, et al. Fused
Mycobacterium tuberculosis multi-stage immunogens with an Fc-delivery system as a promising
approach for the development of a tuberculosis vaccine. Infection, Genetics and Evolution.
2016;39:163-72.
19.   Cabrera-Mora M, Fonseca JA, Singh B, Zhao C, Makarova N, Dmitriev I, et al. A recombinant
chimeric Ad5/3 vector expressing a multistage *plasmodium* antigen induces protective immunity in
mice   using   heterologous   prime-boost   immunization   regimens.   The   Journal   of   Immunology.
2016:1501926.
20.   Bouchery T, Kyle R, Ronchese F, Le Gros G. The Differentiation of CD4(+) T-Helper Cell Subsets
in the Context of Helminth Parasite Infection. Front Immunol. 2014;5:487.
21.   Dixon H, Little MC, Else KJ. Characterisation of the protective immune response following
subcutaneous vaccination of susceptible mice against *Trichuris muris*. International journal for
parasitology. 2010;40(6):683-93.
22.   Dixon H, Johnston C, Else K. Antigen selection for future anti-Trichuris vaccines: a comparison
of cytokine and antibody responses to larval and adult antigen in a primary infection. Parasite
immunology. 2008;30(9):454-61.
23.   Else K, Wakelin D, Wassom D, Hauda K. MHC-restricted antibody responses to *Trichuris muris*
excretory/secretory (E/S) antigen. Parasite immunology. 1990;12(4-5):509-27.
24.   Foth BJ, Tsai IJ, Reid AJ, Bancroft AJ, Nichol S, Tracey A, et al. Whipworm genome and dual-
species transcriptome analyses provide molecular insights into an intimate host-parasite interaction.
Nature genetics. 2014;46(7):693.
25.   Wang P, Sidney J, Dow C, Mothe B, Sette A, Peters B. A systematic assessment of MHC class II
peptide   binding   predictions   and   evaluation   of   a   consensus   approach.   PLoS   Comput   Biol.
2008;4(4):e1000048.
26.   Wang P, Sidney J, Kim Y, Sette A, Lund O, Nielsen M, et al. Peptide binding predictions for HLA
DR, DP and DQ molecules. BMC bioinformatics. 2010;11:568.
27.   Nielsen M, Justesen S, Lund O, Lundegaard C, Buus S. NetMHCIIpan-2.0-Improved pan-specific
HLA-DR predictions using a novel concurrent alignment and weight optimization training procedure.
Immunome Res. 2010;6(9):7580-6.
28.   Nielsen M, Lund O. NN-align. An artificial neural network-based alignment algorithm for MHC
class II peptide binding prediction. BMC bioinformatics. 2009;10(1):1.
29.   Bui HH, Sidney J, Li W, Fusseder N, Sette A. Development of an epitope conservancy analysis
tool   to   facilitate   the   design   of   epitope-based   diagnostics   and   vaccines.   BMC   Bioinformatics.
2007;8:361.
30.   Dimitrov I, Bangov I, Flower DR, Doytchinova I. AllerTOP v. 2—a server for in silico prediction
of allergens. Journal of molecular modeling. 2014;20(6):1-6.
31.   Bancroft AJ, Else KJ, Humphreys NE, Grencis RK. The effect of challenge and trickle *Trichuris*
muris infections on the polarisation of the immune response. International journal for parasitology.
2001;31(14):1627-37.
32.   King IL, Mohrs K, Meli AP, Downey J, Lanthier P, Tzelepis F, et al. Intestinal helminth infection
impacts the systemic distribution and function of the naive lymphocyte pool. Mucosal immunology.
2017;10(5):1160.

33.   Nardin EH, Calvo-Calle JM, Oliveira GA, Nussenzweig RS, Schneider M, Tiercy J-M, et al. A
totally synthetic polyoxime malaria vaccine containing *Plasmodium falciparum* B cell and universal T
cell epitopes elicits immune responses in volunteers of diverse HLA types. The Journal of Immunology.
2001;166(1):481-9.
34.   Salman AM, Montoya-Díaz E, West H, Lall A, Atcheson E, Lopez-Camacho C, et al. Rational
development of a protective *P. vivax* vaccine evaluated with transgenic rodent parasite challenge
models. Scientific reports. 2017;7:46482.
35.   Moura APV, Santos LC, Brito CRN, Valencia E, Junqueira C, Filho AA, et al. Virus-like particle
display of the α-Gal carbohydrate for vaccination against *Leishmania infection*. ACS central science.
2017;3(9):1026-31.
36.   Guo J, Zhou A, Sun X, Sha W, Ai K, Pan G, et al. Immunogenicity of a Virus-Like-Particle Vaccine
Containing Multiple Antigenic Epitopes of *Toxoplasma gondii* Against Acute and Chronic
Toxoplasmosis in Mice. Front Immunol. 2019;10:592.
37.   Lee S-H, Kim S-S, Lee D-H, Kim A-R, Quan F-S. Evaluation of protective efficacy induced by
virus-like particles containing a *Trichinella spiralis* excretory-secretory (ES) protein in mice. Parasites
& vectors. 2016;9(1):384.
38.   Lee D-H, Kim A-R, Lee S-H, Quan F-S. Virus-like particles vaccine containing *Clonorchis sinensis*
tegumental protein induces partial protection against *Clonorchis sinensis* infection. Parasites &
vectors. 2017;10(1):626.
39.   Dhanda SK, Usmani SS, Agrawal P, Nagpal G, Gautam A, Raghava GPS. Novel *in silico* tools for
designing peptide-based subunit vaccines and immunotherapeutics. Brief Bioinform. 2017;18(3):467-
78.
40.   Yang X, Yu X. An introduction to epitope prediction methods and software. Reviews in medical
virology. 2009;19(2):77-96.
41.   Godkin AJ, Smith KJ, Willis A, Tejada-Simon MV, Zhang J, Elliott T, et al. Naturally processed
HLA class II peptides reveal highly conserved immunogenic flanking region sequence preferences that
reflect antigen processing rather than peptide-MHC interactions. The Journal of Immunology.
2001;166(11):6720-7.
42.   Lin HH ZG, Tongchusak S, Reinherz EL, Brusic Evaluation of MHC-II peptide binding prediction
servers: applications for vaccine research. BMC Bioinformatics. 2008;9:Suppl 12.
43.   Hegde NR, Gauthami S, Kumar HMS, Bayry J. The use of databases, data mining and
immunoinformatics in vaccinology: where are we? Expert Opinion on Drug Discovery. 2018;13(2):117-
30.
44.   Paul S, Arlehamn CSL, Scriba TJ, Dillon MB, Oseroff C, Hinz D, et al. Development and validation
of a broad scheme for prediction of HLA class II restricted T cell epitopes. Journal of immunological
methods. 2015;422:28-34.
45.   Schumacher FR, Delamarre L, Jhunjhunwala S, Modrusan Z, Phung QT, Elias JE, et al. Building
proteomic tool boxes to monitor MHC class I and class II peptides. Proteomics. 2017;17(1-2).
46.   Pappalardo F, Flower D, Russo G, Pennisi M, Motta S. Computational modelling approaches to
vaccinology. Pharmacol Res. 2015;92:40-5.
47.   Rammensee H-G, Bachmann J, Emmerich NPN, Bachor OA, Stevanović S. SYFPEITHI: database
for MHC ligands and peptide motifs. Immunogenetics. 1999;50(3-4):213-9.
48.   Reche PA, Glutting J-P, Reinherz EL. Prediction of MHC class I binding peptides using profile
motifs. Human immunology. 2002;63(9):701-9.
49.   Gowthaman U, Agrewala JN. In silico tools for predicting peptides binding to HLA-class II
molecules: more confusion than conclusion. Journal of proteome research. 2007;7(01):154-63.
50.   Borrás-Cuesta F, Golvano J-J, Garcıa-Granero M, Sarobe P, Riezu-Boj J-I, Huarte E, et al. Specific
and general HLA-DR binding motifs: comparison of algorithms. Human immunology. 2000;61(3):266-
78.
51.   Lin HH, Ray S, Tongchusak S, Reinherz EL, Brusic V. Evaluation of MHC-II peptide binding
prediction servers: applications for vaccine research. BMC bioinformatics. 2008;9((Suppl 12)):10.

52. Zhao W, Sher X. Systematically benchmarking peptide-MHC binding predictors: From
synthetic to naturally processed epitopes. PLoS Comput Biol. 2018;14(11):e1006457.
53. Andreatta M, Trolle T, Yan Z, Greenbaum JA, Peters B, Nielsen M. An automated benchmarking
platform for MHC class II binding prediction methods. Bioinformatics. 2018;34(9):1522-8.
54. Dobano C, Sanz H, Sorgho H, Dosoo D, Mpina M, Ubillos I, et al. Concentration and avidity of
antibodies to different circumsporozoite epitopes correlate with RTS,S/AS01E malaria vaccine
efficacy. Nat Commun. 2019;10(1):2174.
55. Veronico P, Gray L, Jones J, Bazzicalupo P, Arbucci S, Cortese M, et al. Nematode chitin
synthases: gene structure, expression and function in *Caenorhabditis elegans* and the plant parasitic
nematode *Meloidogyne artiellia*. Molecular Genetics and Genomics. 2001;266(1):28-34.
56. Wei J, Damania A, Gao X, Liu Z, Mejia R, Mitreva M, et al. The hookworm *Ancylostoma
ceylanicum* intestinal transcriptome provides a platform for selecting drug and vaccine candidates.
Parasites & vectors. 2016;9(1):518.
57. Foor WE. Ultrastructural aspects of oocyte development and shell formation in *Ascaris
lumbricoides*. The Journal of parasitology. 1967:1245-61.
58. Santos LN, Silva ES, Santos AS, De Sá PH, Ramos RT, Silva A, et al. De novo assembly and
characterization of the *Trichuris trichiura* adult worm transcriptome using Ion Torrent sequencing.
Acta tropica. 2016;159:132-41.
59. Johnston WL, Krizus A, Dennis JW. The eggshell is required for meiotic fidelity, polar-body
extrusion and polarization of the *C. elegans* embryo. BMC biology. 2006;4(1):35.
60. Johnston WL, Krizus A, Dennis JW. Eggshell chitin and chitin-interacting proteins prevent
polyspermy in *C. elegans*. Current Biology. 2010;20(21):1932-7.
61. Cuesta-Astroz Y, de Oliveira FS, Nahum LA, Oliveira G. Helminth secretomes reflect different
lifestyles and parasitized hosts. International journal for parasitology. 2017;47(9):529-44.
62. Balasubramanian N, Toubarro D, Simoes N. Biochemical study and in vitro insect immune
suppression by a trypsin-like secreted protease from the nematode Steinernema carpocapsae.
Parasite immunology. 2010;32(3):165-75.
63. Santos LN, Gallo MB, Silva ES, Figueiredo CAV, Cooper PJ, Barreto ML, et al. A proteomic
approach to identify proteins from *Trichuris trichiura* extract with immunomodulatory effects. Parasite
immunology. 2013;35(5-6):188-93.
64. Yang Y, jun Wen Y, Cai YN, Vallée I, Boireau P, Liu MY, et al. Serine proteases of parasitic
helminths. The Korean journal of parasitology. 2015;53(1):1.
65. Toubarro D, Lucena-Robles M, Nascimento G, Santos R, Montiel R, Veríssimo P, et al. Serine
protease-mediated host invasion by the parasitic nematode Steinernema carpocapsae. Journal of
biological chemistry. 2010:jbc. M110. 129346.
66. Dalton JP, Brindley PJ, Knox DP, Brady CP, Hotez PJ, Donnelly S, et al. Helminth vaccines: from
mining genomic information for vaccine targets to systems used for protein expression. International
journal for parasitology. 2003;33(5-6):621-40.
67. Wang B, Wang ZQ, Jin J, Ren HJ, Liu LN, Cui J. Cloning, expression and characterization of a
Trichinella spiralis serine protease gene encoding a 35.5 kDa protein. Experimental parasitology.
2013;134(2):148-54.
68. Feng S, Wu X, Wang X, Bai X, Shi H, Tang B, et al. Vaccination of mice with an antigenic serine
protease-like protein elicits a protective immune response against *Trichinella spiralis* infection. The
Journal of parasitology. 2013;99(3):426-32.
69. Xu J, Bai X, Wang LB, Shi HN, Van Der Giessen JW, Boireau P, et al. Immune responses in mice
vaccinated with a DNA vaccine expressing serine protease-like protein from the new-born larval stage
of *Trichinella spiralis*. Parasitology. 2017;144(6):712-9.
70. Serradell MC, Rupil LL, Martino RA, Prucca CG, Carranza PG, Saura A, et al. Efficient oral
vaccination by bioengineering virus-like particles with protozoan surface proteins. Nature
Communications. 2019;10(1):361.

782 71. Wahl-Jensen V, Kurz SK, Hazelton PR, Schnittler H-J, Ströher U, Burton DR, et al. Role of Ebola
783 virus secreted glycoproteins and virus-like particles in activation of human macrophages. Journal of
784 virology. 2005;79(4):2413-9.

785 72. Lahoud MH, FatmaKitsoulis, SusieSan Wan, SooVremec, DavidLee, Chin-NienPhipson,
786 BelindaShi, WeiSmyth, Gordon K, Lew AM. Targeting antigen to mouse dendritic cells via Clec9A
787 induces potent CD4 T cell responses biased toward a follicular helper phenotype. The Journal
788 of Immunology. 2011(2):842-50.

789 73. Birkett A, Thornton B, Milich D, Oliveira G, Siddique A, Nussenzweig R, et al. Hepatitis B virus
790 core antigen particles containing minimal T and B cell epitopes of *Plasmodium falciparum* CS protein
791 elicit high levels of malaria specific immune responses in mice and non-human primates. Am J Trop
792 Med Hyg. 2001;65:258.

793 74. Birkett A, Lyons K, Schmidt A, Boyd D, Oliveira G, Siddique A, et al. A modified hepatitis B virus
794 core particle containing multiple epitopes of the *Plasmodium falciparum* circumsporozoite protein
795 provides a highly immunogenic malaria vaccine in preclinical analyses in rodent and primate hosts.
796 Infection and immunity. 2002;70(12):6860-70.

797 75. Milich DR, Hughes J, Jones J, Sällberg M, Phillips TR. Conversion of poorly immunogenic
798 malaria repeat sequences into a highly immunogenic vaccine candidate. Vaccine. 2001;20(5-6):771-
799 88.

800 76. Nardin EH, Oliveira GA, Calvo-Calle JM, Wetzel K, Maier C, Birkett AJ, et al. Phase I testing of a
801 malaria vaccine composed of hepatitis B virus core particles expressing *Plasmodium falciparum*
802 circumsporozoite epitopes. Infection and immunity. 2004;72(11):6519-27.

803 77. Oliveira GA, Wetzel K, Calvo-Calle JM, Nussenzweig R, Schmidt A, Birkett A, et al. Safety and
804 enhanced immunogenicity of a hepatitis B core particle *Plasmodium falciparum* malaria vaccine
805 formulated in adjuvant Montanide ISA 720 in a phase I trial. Infection and immunity. 2005;73(6):3587-
806 97.

807 78. Gu Y, Huang J, Wang X, Wang L, Yang J, Zhan B, et al. Identification and characterization of
808 CD4+ T cell epitopes present in Trichinella spirals paramyosin. Veterinary parasitology. 2016;231:59-
809 62.

810 79. Datta R, Hedeler C, Paton N, Brass A, Else K. Identification of novel genes in intestinal tissue
811 that are regulated after infection with an intestinal nematode parasite. Infection and immunity.
812 2005;73(7):4025-33.

813 80. Hasnain SZ, Wang H, Ghia JE, Haq N, Deng Y, Velcich A, et al. Mucin gene deficiency in mice
814 impairs host resistance to an enteric parasitic infection. Gastroenterology. 2010;138(5):1763-71. e5.

815 81. Hasnain SZ, McGuckin MA, Grencis RK, Thornton DJJPNTD. Serine protease(s) secreted by the
816 nematode Trichuris muris degrade the mucus barrier. 2012;6.

817 82. Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic
818 reviews and meta-analyses: the PRISMA statement. International journal of surgery. 2010;8(5):336-
819 41.

820 83. Yasser E-M, Dobbs D, Honavar V. On evaluating MHC-II binding peptide prediction methods.
821 PloS one. 2008;3(9):e3268.

822 84. Murugan N, Dai Y. Prediction of MHC class II binding peptides based on an iterative learning
823 model. Immunome research. 2005;1(1):6.

824 85. Nielsen M, Lundegaard C, Worning P, Hvid CS, Lamberth K, Buus S, et al. Improved prediction
825 of MHC class I and class II epitopes using a novel Gibbs sampling approach. Bioinformatics.
826 2004;20(9):1388-97.

827 86. Altman DG, Bland JM. Diagnostic tests. 1: Sensitivity and specificity. BMJ: British Medical
828 Journal. 1994;308(6943):1552.

829 87. Wakelin DJP. Acquired immunity to *Trichuris muris* in the albino laboratory mouse.
830 1967;57(3):515-24.

831 88. HogenEsch H. Mechanism of immunopotentiation and safety of aluminum adjuvants.
832 Frontiers in immunology. 2013;3:406.

833 89.   Else K, Entwistle G, Grencis R. Correlations between worm burden and markers of Th1 and
834 Th2 cell subset induction in an inbred strain of mouse infected with *Trichuris muris*. Parasite
835 immunology. 1993;15(10):595-600.
836
837

Supporting Information Legends

S1 Fig. The HBc-antigen fusion recombinant protein purification and assembly. Left:
Chromatogram of the VLP recombinant proteins HBc-Ag (A), HBc-CBD$_{1243-1259}$ (B), HBc-CBD$_{241-}$
$_{257}$ (C), HBc-CLSP$_{143-158}$, (D), and HBc-CLSP$_{398-416}$ (E) purified using SEC. Separation was carried
out on a Superose 6, 10/300 GL (GE Healthcare), at a flow rate of 0.2 ml/min with 100 mM Tris-HCl,
150 mM NaCl, 1 mM EDTAatpH 8. VLP recombinant proteins were eluted in 0.5 ml fraction and a
sample of SEC the elution peak was visualised by SDS-PAGE. Vertical red lines indicate the elution
volume of mass standards, from left to right: 2MDa, 670kDa. Right: Purification visualised by SDS-
PAGE (10% gel) analysis stained with Coomassie blue. M: protein marker (Precision Plus protein
standard, unstained, Bio-Rad); Lane 1: VLP recombinant protein after Strep-Tag affinity purification
using the StrepTrap HP, and Lane 2: VLP recombinant protein after SEC purification. The arrow
represents VLP recombinant protein monomer with an expected molecular mass of ~22 kDa.
850
S2 Fig. Cytokine productions by mesenteric lymph node cells and quantification of goblet cell
numbers in mice vaccinated with VLPs+T-cell epitopes and control mice. (A) MLNs from mice
vaccinated with 50 µg of pre-mixed of VLPs+T-cell epitopes epitopes (HBc-CBD$_{1243-1259}$, HBc-CBD$_{241-}$
$_{257}$, HBc-CLSP$_{143-158}$ and HBc-CLSP$_{398-416}$), 50 µg of VLP (HBc-Ag), 50 µg ES/Alum, or PBS on day -20
and boosted on day -10 were stimulated at $5x10^6$/ml with 50 µg/ml *T. muris* ES. After 48 hours of
stimulation, cell culture supernatants were harvested and assayed by cytometric bead array for IL-4
(A), IL-6 (B), IL-10 (C), IL-17 (D), and TNF (E) production. Statistical analyses were carried out using
the Kruskal-Wallis test (multiple comparisons). Significant differences between groups are represented
by (*P≤0.05) with a line. Results are shown as mean ± SEM. n= 11 mice per group. The results
presented are from two separated experiment pooled together.

S1 Table. List of Keywords used to screen for MHC I and II *in silico* prediction tools.

S2 Table. List of MHC class I and II and HLA binders peptides *in silico* prediction tools.
S3 Table. MHC-II T-cell epitopes dataset used to evaluate the MHC-II *in silico* prediction tools.
S4 Table. List of MHC-II T-cell epitope sequences and primers used to amplify the target.
865
866 **Figure Legends**
867
Fig 1. (A) PRISMA flow diagram of the systematic review screening process for
restricted MHC-II T-cell bioinformatics prediction tools. A list of search terms was used
in Google website and other website sources to screen for MHC class I and II *in silico*
prediction tools. Aspects relating to the number of citations (>200), number of publications,
online availability, last update and community were used as inclusion/exclusion criteria to
select the top MHC-II *in silico* prediction tools. n= number of key words, websites sources and
in silico prediction tools. (B) The flow diagram of reverse vaccinology approach to identify
potential vaccine candidates (MHC-II T-cell epitopes) from the *T. muris* genome. The number
on the arrow represents the number of proteins or epitopes selected for the next step. (C) List
of 4 *Trichuris* MHC-II T-cell epitopes which have potential as vaccine candidates.
878
879
Fig 2. (A) Inflammatory cytokines production by mouse bone marrow-derived DCs
(BMDCs) and mouse bone marrow-derived macrophages (BMDMs) (B) in response to
VLPs. BMDCs and BMDMs at $1X10^6$/ml were stimulated *in vitro* with 10 µg/ml VLPs (HBc-Ag,
HBc-H$_{112-128}$, HBc-CBD$_{1243-1259}$, HBc-CBD$_{241-257}$, HBc-CLSP$_{143-158}$ and HBc-CLSP$_{398-416}$) and
with 50 µg/ml ES and 0.1 µg/ml LPS as positive controls. Unstimulated BMDCs and BMDMs
served as negative controls. Supernatants were harvested after 24 hours for IL-6 and TNF α

cytokine analyses measured by CBA. The bars represent mean ± SEM. Statistical analyses
were carried out using the Kruskal-Wallis test (multiple comparisons). Significant differences
between groups are represented by (*P≤0.05) with a line. Chart bars represent BMDCs and
BMDMs grown from three individual mice from one representative experiment of two separate
experiments. (C) Representative images of fluorescein-conjugated VLPs internalisation in the
BMDCs and BMDMs (G). BMDCs and BMDMs at $1X10^6$/ml were incubated with 10 µg/ml
fluorescein-conjugated VLP (HBc-Ag and HBc-CLSP$_{398-416}$) for 24 hours. As a negative
control, unstimulated BMDCs and BMDMs were examined. Cell internalisation was
determined by Amnis ImageStreamX cytometer compared to unstimulated BMDCs and
BMDMs. Images shown, from left to right, show individual Brightfield images (BF) in the white
channel, fluorescent-labelled stimulus (FITC) in the green channel and the combination of both
BF/FITC merged channels. The internalisation mean absolute deviation (MAD) is included
above its images. The positive MAD value represents internalisation, and negative values
represent poor internalisation. (D) Representative images of fluorescein-conjugated VLPs co-
localization in the BMDCs and BMDMs (H). BMDCs and BMDMs at $1X10^6$/ml were stained
with Lysotracker to visualise the cellular lysosome compartment and subsequently stimulated
with 10 µg/ml fluorescein-conjugated VLP (HBc-Ag, and HBc-CLSP$_{398-416}$) for 24 hours. As a
negative control, unstimulated BMDCs and BMDMs were examined. Intracellular co-
localization was determined by Amnis ImageStreamX cytometer. Images shown, from left to
right, show individual Brightfield images (BF) in the white channel, fluorescent-labelled
stimulus (FITC) in the green channel, stained lysosome (Lysotracker) in the red channel and
the combination of both FITC/ Lysotracker merged channels. The Similarity bright detail score
(SBDS) from the IDEAS quantitative co-localization analysis is included above its image.
SBDS values around 1 represent co-localization, and 0 values represent poor co-localization.
Scale bars represent 10 µm.
911
912
Fig 3. Experimental schedule, worm burden, parasite-specific and VLP-specific
antibodies levels of mice vaccinated and challenged with a *T. muris* infection. 6-week
old male C57BL/6 mice were vaccinated subcutaneously with 50 µg of pre-mixed of VLPs+T-
cell epitopes (HBc-CBD$_{1243-1259}$, HBc-CBD$_{241-257}$, HBc-CLSP$_{143-158}$, and HBc-CLSP$_{398-416}$), 50 µg
of VLP (HBc-Ag), or PBS on day -20 and boosted on day -10. Mice were infected by oral
gavage with approximately 150 infective *T. muris* eggs on day 0 and sacrificed at d14 post-
infection. (B) Comparison of worm burden at day 14 post-infection. (C) Worm burden in mice
vaccinated with 50 µg of ES. n= 11 mice per group. The results presented are from two
separated experiments pooled together. Day 14 p.i. sera were titrated against *T. muris* ES
antigens to assess parasite-specific IgM (D) IgG1 (E), and IgG2c (F) levels in VLPs+T-cell
epitopes, VLP (HBc-Ag), PBS and in ES vaccinated mice by ELISA (reading at 405 nm). Day
14 p.i. sera were titrated against premixed VLPs+ T- cell epitopes antigens to assess VLPs+
T- cell epitopes-specific IgM (G) IgG1 (H), and IgG2c (I) levels in VLPs+T-cell epitopes, VLP
(HBc-Ag), PBS and in ES vaccinated mice. Statistical analyses were carried out using Kruskal-
Wallis test (multiple comparisons). Significant differences between groups are represented by
(*P≤0.05, **P≤0.01, ***P≤0.001, ****P≤0.0001) with a line. Results are shown as mean ± SEM.
n= 6 mice per group. This experiment was repeated two times, and the ELISA results shown
here are representative of the two experiments.
931
Fig 4. Cytokine productions by mesenteric lymph node cells and quantification of
goblet cell numbers in mice vaccinated with VLPs+T-cell epitopes and control mice. (A)
MLNs from mice vaccinated with 50 µg of pre-mixed of VLPs+T-cell epitopes epitopes (HBc-
CBD$_{1243-1259}$, HBc-CBD$_{241-257}$, HBc-CLSP$_{143-158}$ and HBc-CLSP$_{398-416}$), 50 µg of VLP (HBc-Ag),
50 µg ES/Alum, or PBS on day -20 and boosted on day -10 were stimulated at $5x10^6$/ml with
50 µg/ml *T. muris* ES. After 48 hours of stimulation, cell culture supernatants were harvested
and assayed by cytometric bead array for IL-5 (A), IL-9 (B), IL-13 (C), IFN-□ (D) production.
(E) Representative photographs of gut sections stained with Periodic acid–Schiff staining
(PAS). The slides were scanned, and the goblet cells were quantified using Panoramic Viewer

software. All photographs were taken with 20X magnification. (F) Goblet cells were quantified
by counting the number of alcian blue-stained cells per crypt unit in three fields of view from
each section and are shown as mean cell numbers per 20 crypt units (cu) ±SEM. Statistical
analyses were carried out using the Kruskal-Wallis test (multiple comparisons). Significant
differences between groups are represented by (*$P \leq 0.05$, ****$P \leq 0.0001$) with a line. Results
are shown as mean ± SEM. n= 11 mice per group. The results presented are from two
separated experiment pooled together.
948

Figure 1

Figure 2

Figure 3

Figure 4