***Pseudomonas aeruginosa* is capable of natural transformation in biofilms**

Laura M. Nolan[1,2], Lynne Turnbull[1], Marilyn Katrib[1], Sarah R. Osvath[1], Davide Losa[1], James J.

Lazenby[3] and Cynthia B. Whitchurch[1,3,4]*

Author affiliations:

[1] The ithree institute, University of Technology Sydney, Ultimo, New South Wales, 2007, Australia.

[2] National Heart and Lung Institute, Imperial College London, London, SW3 6LR, UK.

[3] Microbes in the Food Chain Programme, Quadram Institute Bioscience, Norwich Research Park,

Norwich, NR4 7UQ, UK

[4] School of Biological Sciences, University of East Anglia, Norwich, NR4 7TJ, UK

* Corresponding author:

Email: Cynthia.Whitchurch@quadram.ac.uk

Current Addresses:

CBW: Microbes in the Food Chain Programme, Quadram Institute Bioscience, Norwich Research

Park, Norwich, NR4 7UQ, UK

LMN: National Heart and Lung Institute, Imperial College London, London, SW3 6LR, UK.

DL: Department of Cellular, Computational and Integrative Biology (CIBIO), University of Trento,

Trento, TN 38123, Italy

Abbreviations:

extracellular DNA; eDNA, type IV pili; T4P

Abstract

Natural transformation is a mechanism that enables competent bacteria to acquire naked, exogenous

DNA from the environment. It is a key process that facilitates the dissemination of antibiotic

resistance and virulence determinants throughout bacterial populations. *Pseudomonas aeruginosa* is

an opportunistic Gram-negative pathogen that produces large quantities of extracellular DNA

(eDNA) that is required for biofilm formation. *P. aeruginosa* has a remarkable level of genome

plasticity and diversity that suggests a high degree of horizontal gene transfer and recombination but

is thought to be incapable of natural transformation. Here we show that *P. aeruginosa* possesses

homologs of all proteins known to be involved in natural transformation in other bacterial species.

We found that *P. aeruginosa* in biofilms is competent for natural transformation of both genomic and

plasmid DNA. Furthermore, we demonstrate that type IV pili (T4P) facilitate but are not absolutely

essential for natural transformation in *P. aeruginosa*.

Introduction

The continued increase in antimicrobial resistance (AMR) levels is considered to be a significant global threat[1]. Horizontal gene transfer (HGT) is a key source of bacterial genome variation and evolution and is largely responsible for the acquisition of antibiotic resistance genes by bacterial pathogens[2]. Bacteria can acquire and heritably incorporate new genetic information via three HGT mechanisms: conjugation, transduction and natural transformation. Conjugation is a cell-contact dependent mechanism that transfers DNA directly from the cytoplasm of one bacterial cell into another. Transduction involves encapsidation of DNA into a bacteriophage which then injects the DNA into the recipient cell. The third HGT mechanism is natural transformation which involves the import of naked DNA from the environment through a specialised DNA transport apparatus[3,4].

Extracellular DNA (eDNA) is present in significant quantities in both clinical and environmental settings, and provides a vast reservoir of genetic material that can be sampled by bacteria that are competent for natural transformation[5]. In many naturally competent bacterial species Type IV pili (T4P) are required for natural transformation[4]. While the exact role of T4P in natural transformation is unclear, the generally accepted model is that DNA binds to the pilus structure, which retracts and pulls the DNA to the cell surface. It is unclear whether or not DNA is translocated across the outer membrane through the PilQ secretin pore. The incoming DNA can then be accessed by the ComEA DNA translocation machinery in the periplasm, which mediates DNA uptake possibly by a ratchet mechanism[4,6]. In Gram-positive bacterial species that do not produce T4P, natural transformation involves a number of proteins with homology to T4P proteins which are thought to form a pseudopilus structure that spans the cell wall and is coupled to the DNA translocation complex at the cytoplasmic membrane[4,7]. Once exogenous DNA has been taken up by the cell it can be stably incorporated into the genome via recombination or transposition, or be maintained as an independent replicon if plasmid DNA is taken up by an appropriate host[7]. Natural plasmid transformation has also been

described in *Escherichia coli*[8]. This appears to be a distinct mechanism which does not require

homologs of the T4P/competence pseudopilus or DNA translocation machinery[9].

Pseudomonas aeruginosa is a highly antibiotic resistant Gram-negative bacterium which is a part of

the 'ESKAPE' group of pathogens that pose a serious health risk worldwide. *P. aeruginosa* readily

acquires antibiotic resistance determinants, and demonstrates a high degree of genomic diversity and

malleability similar to that seen in naturally transformable bacteria[10,11]. Despite this, *P. aeruginosa*

has long been thought to be incapable of natural transformation[12]. *P. aeruginosa* is a model organism

for studying T4P[13]. Interestingly, *P. aeruginosa* produces copious quantities of eDNA under

conditions that promote T4P production such as in static broth cultures[14], biofilms[15] and during

twitching motility-mediated biofilm expansion[16,17]. We therefore hypothesised that *P. aeruginosa*

may be competent for natural transformation under conditions that promote both T4P expression and

eDNA production. Here we show that some clinical and laboratory strains of *P. aeruginosa* are in

fact capable of natural transformation under these conditions.

Methods

Strains, plasmids and growth conditions

P. aeruginosa strains used in this study were PAO1 (ATCC 15692), PAK[18], PA14[19], PA103[20],

PAO1$_{GFP}$ which contains mini-Tn7-Gm^R-P$_{A1/04/03}$-*egfp* encoding *gfp* and *aac1* (Gm^R) inserted

downstream of *glmS*[21], PAO1$_{CTX}$ which contains miniCTX2 encoding *tet* (Tc^R) inserted into the *attB*

site of the chromosome[22] and T4P mutants PAK*pilA*::*TcR*[23], and Tn5-B21 mutants of *pilQ*[24], *pilT*[25],

pilV[26]*, fimV*[27]. The *P. aeruginosa* CF sputum clinical isolates were obtained from David Armstrong

at Monash Medical Centre (Melbourne, Australia), and the otitis externa *P. aeruginosa* clinical

isolates were obtained from Di Olden at Gribbles Pathology Melbourne (Australia). The pUCPSK

plasmid used is a non-conjugative *E. coli-P. aeruginosa* shuttle vector encoding *bla* which confers

carbenicillin resistance (Carb[R]) in *P. aeruginosa*[28]. *E. coli* Dh5α (*recA, endA1, gyrA96, hsdR17, thi-*

1, supE44, relA1, φ80, dlacZΔM15) was used as a host strain for pUCPSK and was miniprepped from

P. aeruginosa and *E. coli* strains using a Qiagen miniprep kit according to manufacturer's

instructions.

P. aeruginosa was cultured on lysogeny broth (LB) solidified with agar at 1.5% (w/v) for routine

maintenance and at 1.5% or 1% (w/v) for colony biofilm assays and grown in cation-adjusted Mueller

Hinton Broth (CAMHB) at 37°C for all static broth and flow biofilm assays. Antibiotics were used

at the following concentrations (w/v) as required: ampicillin 50 µg/ml for *E. coli* and carbenicillin

250 µg/ml, gentamicin 100 µg/ml and tetracycline 100 µg/ml for *P. aeruginosa*.

Bioinformatics and data and statistical analyses

Homologs of proteins involved in natural transformation were identified in *P. aeruginosa* PAO1

using BLASTp[29]. The Pseudomonas.com resource[30] and the PAK genome[31] were used to identify *P.*

aeruginosa orthologs.

Data was graphed and analysed using Graph Pad Prism version 8.0. The number of replicates and any

statistical tests are described in Figure legends.

Colony biofilm assay

Overnight cultures of *P. aeruginosa* were grown in 2 ml CAMHB at 37°C, shaking at 250 rpm. A 10

µL plastic loop was used to generate a 1 cm patch of the overnight culture on a dry 1% LBA plate.

This was then incubated overnight at 37°C. The next day 10 µL of pUCPSK plasmid DNA (at a final

concentration of 5 µg) was spotted onto the established colony biofilm and allowed to dry into the

cells. The plate was then incubated with the agar downwards at 37°C for the indicated time. After

incubation the colony biofilm was harvested from the plate into 1 mL LB, vortexed to resuspend and

then incubated at 37°C for 30 min to fully resuspend the cells. The cell suspension was then spread

plated between two 150 mm LBA plates with appropriate antibiotic selection and incubated for 24 hr

at 37°C.

Static broth assay

Overnight cultures of *P. aeruginosa* were grown in 2 ml CAMHB at 37°C, shaking at 250 rpm. 40μL

of overnight culture was subcultured into 2 ml fresh CAMHB with DNA added at the indicated

concentration. The media, cells and DNA were then mixed and incubated at 37°C statically for 24 hr.

Note for the shaking broth assay the same setup was used however the culture was incubated with

shaking at 250 rpm. In both cases after incubation the cell suspension was then spread plated between

two 150 mm LBA plates with appropriate antibiotic selection and incubated for 24 hr at 37°C.

To image the aggregates present in static broth cultures *P. aeruginosa* was grown as described for the

static broth assay in a glass-bottomed Ibidi® μ-Dish and visualised using DeltaVision Elite inverted

research microscope with a × 100 1.4 numerical aperture UPlanFLN objective, InsightSSI

illumination, SoftWorX acquisition software, fitted with a WeatherStation environmental chamber

(Applied Precision, GE Healthcare, Issaquah, WA, USA) and a scientific CMOS 15-bit camera

(pco.edge, PCO AG, Kelheim, Germany).

Isolation of DNA for use in continuous flow biofilm assays

Chromosomal DNA (gDNA) was purified from PAO1Tn7::*gfp-aac1* cells using the Epicentre®

Masterpure DNA purification kit. Extracellular DNA (eDNA) was purified from a confluent lawn of

PAO1Tn7::*gfp-aac1* cultured overnight on MacConkey agar containing 5% (v/v) glycerol. Bacteria

were suspended in sterile phosphate buffered saline (PBS), centrifuged and the supernatant filtered

through 0.2 μm PES membrane. eDNA present in the supernatant was further purified by removal of

proteins and ethanol precipitation as reported previously[32]. Sterility of all DNA samples was

confirmed prior to use by plating onto LB agar.

Continuous flow biofilm assays

10 cm lengths of Tygon laboratory tubing (2mm ID) were inoculated with 1/100 dilution of an

overnight culture of *P. aeruginosa* in CAMHB and allowed to attach for 2 hours under static

conditions after which continuous flow was commenced at a rate of 80 µL/min at room temperature.

Influent media was CAMHB containing either no added DNA or DNA added at a final concentration

of 1 µg/mL for pUCPSK plasmid DNA, or 0.5 mg/mL for gDNA or eDNA. At harvest, the attached

biofilm was removed by sonication and biofilm-associated bacteria collected by centrifugation.

Transformants were selected by plating onto LB agar with appropriate antibiotic selection and

incubated for 24 hr at 37°C. To visualise natural transformation by PAO1 continuous flow biofilms,

an IBIDI® µ-slide I (with flow kit) was inoculated and cultured as described for the Tygon tubing

biofilms. Biofilms were imaged using an Olympus IX71 inverted research microscope with a × 100

1.4 numerical aperture UPlanFLN objective, FViewII monochromatic camera and AnalySIS

Research acquisition software (Olympus Australia, Notting Hill, VIC, Australia) fitted with an

environmental chamber (Solent Scientific, Segensworth, UK) for fluorescent imaging.

Confirmation of natural transformation events from continuous flow biofilms

The presence of mini-Tn7-Gm^R-$P_{A1/04/03}$-*egfp* at the chromosomal attTn7 site was confirmed by PCR

using primers Tn7$_{-up}$ (5'CGTATTCTTCGTCGGCGTGAC3') and Tn7$_{-down}$

(5'CGAAGCCGCCGACAAGGG3'). Expression of GFP was confirmed by epifluorescence

microscopy on an Olympus IX71. The presence of mini-CTX2 at the chromosomal *attB* site was

confirmed by PCR using primers P$_{ser-up}$ (5'CGAGTGGTTTAAGGCAACGGTCTTGA3') and P$_{ser-}$

$_{down}$ (5'AGTTCGGCCTGGTGGAACAACTCG 3')[22]. To confirm the presence of pUCPSK in *P.*

aeruginosa, plasmid DNA was extracted from *P. aeruginosa*, transformed into *E. coli*, extracted and

confirmed by sequencing with M13-FUP (5'TGTAAAACGACGGCCAGT3').

Results

Bioinformatic analyses of the sequenced *P. aeruginosa* strains PAO1, PA14 and PAK show that each

of these strains encode homologs of all genes known to be involved in natural transformation in other

bacterial species (Table 1). To determine if *P. aeruginosa* might be capable of natural transformation

in biofilms, we established biofilms with a 1:1 mixture of PAO1$_{GFP}$ (Gm^R) and PAO1 [pUCPSK]

(Carb^R) under continuous flow conditions. Biofilm effluent was collected each day for 4 days and

bacteria tested for their ability to grow on LB agar plates containing both gentamicin and

carbenicillin. An average of 50-100 Gm^R/Carb^R colonies (resistant to both gentamicin and

carbenicillin) were obtained from the effluent of mixed PAO1$_{GFP}$ and PAO1 [pUCPSK] biofilms on

day 1 and confluent lawns of Gm^R/Carb^R colonies obtained from day 2 onwards. The presence of

mini-Tn7-Gm^R-P$_{A1/04/03}$-*egfp* at the chromosomal *attTn7* site in these colonies was confirmed by PCR

amplification. To confirm that these colonies also possessed pUCPSK, plasmid DNA was extracted,

transformed into *E. coli* and confirmed by sequencing. Neither the PAO1$_{GFP}$ or PAO1 [pUCPSK]

strains used to establish these mixed biofilms, or effluent from control single strain biofilms were

able to grow on the dual antibiotic selection plates. As PAO1 lacks a prophage capable of transduction

and pUCPSK is a non-conjugative plasmid, these results suggest that HGT of plasmid DNA and/or

chromosomal DNA might occur via natural transformation in *P. aeruginosa* biofilms.

To determine if HGT of chromosomal DNA did indeed occur via natural transformation, and the

above result was not just due to HGT of plasmid DNA, we established biofilms with a 1:1 mixture of

PAO1$_{GFP}$ (Gm^R) and PAO1$_{CTX}$ (Tc^R) under continuous flow conditions. Biofilm effluent was

collected each day for 8 days and bacteria tested for their ability to grow on LB agar plates containing

both gentamicin and tetracycline. Whilst no Gm^R/Tc^R colonies were obtained from days 1-4, from

days 5-8 an average of 3 Gm^R/Tc^R colonies that were resistant to both antibiotics and expressed GFP

were recovered per day. The presence of both mini-Tn7-Gm^R-$P_{A1/04/03}$-*egfp* and mini-CTX2 in these

colonies was confirmed by PCR. Importantly, neither the $PAO1_{GFP}$ or $PAO1_{CTX}$ strains used to

inoculate the mixed biofilms or effluent from control single-strain biofilms were able to grow on the

dual antibiotic selection plates. As neither conjugation or transduction is likely to account for these

HGT events in PAO1 biofilms, these results suggest that *P. aeruginosa* is able to acquire and

incorporate chromosomal DNA and plasmids encoding antibiotic resistance genes via natural

transformation in biofilms.

To confirm that *P. aeruginosa* is indeed capable of natural transformation and to rule out any

possibility of HGT through transduction or conjugation, we performed a series of experiments to

follow the uptake of purified, sterile exogenous DNA. We firstly examined plasmid DNA uptake

under continuous flow biofilm conditions. PAO1 flow biofilms were cultured in the presence or

absence of purified pUCPSK plasmid DNA (at a final concentration of 1 μg/mL) and the amount of

natural transformation within the biofilm biomass and effluent assessed at days 3, 4 and 5. $Carb^R$

colonies were obtained from both the biofilm biomass and biofilm effluent (Fig. 4a). No $Carb^R$

colonies were obtained in the no DNA control.

We also examined if natural transformation by uptake of exogenous chromosomal DNA occurs in

biofilms cultured under continuous flow. *P. aeruginosa* PAO1 flow biofilms were cultured in the

presence and absence of sterile gDNA or eDNA obtained from $PAO1_{GFP}$ (Gm^R) (at a final

concentration of 0.5 mg/mL) in the media influent. After 5 days the number of Gm^R colonies

recovered from the biofilm biomass were counted. This revealed extremely variable rates of natural

transformation of gDNA by cells within the biofilm biomass across multiple experiments (Fig. 1b).

This is not unexpected as the rate is likely to be dependent upon the time at which the natural

transformation event occurred. If this event occurred early in the assay, we would expect many

transformants recovered due to proliferation of the transformed cells. However, if transformation

occurred later we would expect far fewer transformants as these did not have as long to proliferate.

For the eDNA experiments, while some Gm^R transformants were obtained (Fig. 1c), the rate of natural

transformation was overall much lower than for gDNA (Fig. 1b). This may be due to the integrity of

the DNA as we observed via agarose gel electrophoresis that the eDNA used in these experiments

was quite degraded compared with the gDNA, presumably through the action of nucleases present in

the extracellular milieu. No Gm^R colonies were obtained for continuous flow biofilms in the absence

of gDNA or eDNA indicating that the gentamicin resistant cells recovered from these assays was due

to the presence of the exogenous chromosomal DNA. To further rule out the possibility of

spontaneous resistance, the presence of the mini-Tn7-Gm^R-$P_{A1/04/03}$-*egfp* at the chromosomal attTn7

site in the biofilm-derived Gm^R colonies was confirmed by PCR. The presence of the *gfp* gene in the

Gm^R colonies was also confirmed by visualisation of GFP expression using epifluorescence

microscopy (Fig. 2b). No GFP expression was observed in the PAO1 inoculum strain (Fig. 2a). As it

was not possible to directly visualise biofilms cultured in Tygon tubing, PAO1 continuous flow

biofilms were cultured in transparent flow cells over 5 days in the presence and absence of gDNA

obtained from PAO1$_{GFP}$. Epifluorescence microscopy revealed microcolonies of GFP-expressing

bacteria within the biofilm (Fig. 2c-f). No GFP expression was observed in the no DNA control

biofilms.

To further investigate natural transformation by *P. aeruginosa* we set out to determine if this process

with also occurring in colony biofilms on agar plates. We first grew wildtype strains PAK and PAO1

on LB agar overnight to form a colony biofilm on the surface of the agar. We then added 5 µg of

sterile DNA (or the equivalent volume of sterile water) of the plasmid pUCPSK, onto the surface of

the colony biofilm. After 2 hr incubation at 37°C the colony was resuspended and cells plated onto

media containing carbenicillin to select for transformants that had acquired the plasmid. Only colony patches that had been exposed to plasmid DNA yielded Carb^R colonies (Fig. 3a), whereas colony patches exposed to sterile water yielded none. The transformation efficiency for PAK and PAO1 was 24.7 ± 10.1 and 5.8 ± 1.9 transformants/µg plasmid DNA, respectively. To confirm that the carbenicillin resistant colonies had acquired pUCPSK, plasmid DNA was extracted, re-transformed into *E. coli* and confirmed by sequencing. These observations indicate that a proportion of cells within colony biofilms of *P. aeruginosa* are competent for natural transformation and are able to take up and maintain plasmid DNA.

Given that both *P. aeruginosa* strains PAK and PAO1 appeared to be naturally competent, we wanted to determine if this was also the case for two other commonly utilised lab strains (PA14 and PA103) and clinical isolates. All *P. aeruginosa* strains were first confirmed to be carbenicillin sensitive prior to use. Thirteen *P. aeruginosa* otitis externa and twelve cystic fibrosis (CF) lung sputum isolates were assayed for the ability to uptake pUCPSK plasmid DNA in a colony biofilm. Of these, 7/12 otitis externa and 6/10 CF isolates were able to uptake exogenous plasmid DNA (Fig. 3b). No Carb^R colonies were obtained in the no plasmid DNA controls for each strain. Interestingly, transformation ability varied amongst both clinical and lab strains. Of the lab strains, PA14 was the least capable of natural transformation with PAK the most efficient. These results demonstrate that many lab and clinical isolate strains of *P. aeruginosa* are capable of natural transformation within colony biofilms, albeit with different efficiencies.

P. aeruginosa also expresses T4P when cultured in static nutrient broth[13]. Under these conditions *P. aeruginosa* forms biofilms and suspended microcolony aggregates that contain eDNA[14,33]. To determine if natural transformation also occurred in static broth cultures, 10 µg of sterile pUCPSK plasmid DNA (or the equivalent volume of sterile water) was added to a subculture of *P. aeruginosa*

wildtype PAK or PAO1 and incubated statically at 37°C for 24 hrs. Cells were then recovered and

plated onto media containing carbenicillin to select for transformants. Carb^R colonies were obtained

for both PAO1 and PAK under these conditions, whereas no Carb^R colonies were identified in the

water control. As was observed with colony biofilm transformations (Fig. 3a), PAK was more

efficient for natural transformation of pUCPSK than PAO1 in static broth cultures, with efficiencies

of 24.6 ± 5.5 and 1.0 ± 0.3 transformants/μg plasmid DNA, respectively (Fig. 4a).

To determine if transformation efficiency was dependent on the amount of DNA added, we performed

static broth transformation assays with increasing amounts of plasmid DNA. Whilst we observed an

increase in the number of transformants with increasing amounts of sterile plasmid DNA added (Fig.

S1), the transformation efficiency (transformants/μg DNA) was relatively unchanged over the range

of DNA quantities used for transformation (0-30 μg) (Fig. 4b).

We were also interested in determining whether *P. aeruginosa* was also able to uptake exogenous

chromosomal DNA and integrate this into the chromosome in static broth cultures. To examine this,

chromosomal DNA from PAO1$_{GFP}$ (Gm^R) was purified from either a sterile whole cell lysate (gDNA)

or the total sterile eDNA from confluent agar plate culture and 15 μg added to static broth cultures of

PAK or PAO1 for 24 hr. Cells were recovered and cultured on agar containing gentamicin to select

for transformants. These assays revealed that natural transformation of gDNA occurred at a low

frequency for both PAK and PAO1 in static broth cultures (Fig. S2). No natural transformation with

sterile eDNA was observed in static broth cultures for either PAK or PAO1 (Fig. S2). As mentioned

above for eDNA in continuous flow biofilms, this may be due to the degraded nature of the eDNA

compared to the gDNA as visualised by agarose gel electrophoresis. No gentamicin resistant colonies

were obtained in the no DNA controls.

P. aeruginosa produces more T4P when cultured in static broth cultures than under shaking

conditions[13]. We investigated the effects on natural transformation efficiency of culturing under static

or shaking conditions and found that although some natural transformation was still observed under

shaking conditions, more transformants were obtained with static culture conditions (Fig. 5a),

consistent with a role of T4P in natural transformation. To directly examine the role of T4P in natural

transformation of *P. aeruginosa,* we added 10 μg sterile pUCPSK plasmid DNA to static broths of

PAK mutants defective in the production of the pilin subunit (*pilA*), in T4P assembly (*pilV, pilQ,*

fimV) and in T4P retraction (*pilT*). Interestingly, all T4P mutants were capable of some natural

transformation of pUCPSK, however a significant reduction in transformation efficiency compared

to wildtype PAK was observed (Fig. 5b). No Carb^R colonies were identified in the no DNA controls.

There was no apparent difference in the transformation efficiency of mutants which either didn't have

any surface-assembled T4P (*pilA, pilV, pilQ, fimV*) or were unable to retract T4P (*pilT*) (Fig. 5b).

Given that wildtype *P. aeruginosa* forms aggregates in a static broth[14,33] we considered the possibility

that natural transformation was occurring within these aggregates in our static broth transformation

assay. To determine if the reduction in transformation efficiency by T4P mutants (Fig. 5b) could be

accounted for by the inability to form aggregates, we used microscopy to image wildtype PAK and

T4P mutants under static broth conditions. This revealed that mutants unable to make surface-

assembled T4P (*pilA, pilV, pilQ, fimV*) were also unable to form aggregates (Fig. 5c). A *pilT* mutant,

which has a T4P retraction defect, was however still able to form aggregates (Fig. 5c). While the total

number of aggregates was less than that observed for wildtype (Fig. 5c), this decrease does not

account for the observed decrease in natural transformation efficiency (Fig. 5b). This suggests that it

is the lack of functional T4P effects natural transformation and not other factors associated with

aggregate formation.

313 Overall these data suggest that in *P. aeruginosa*, T4P facilitate transport of DNA to the cell surface

314 but are not essential for natural transformation, at least of plasmid DNA. Furthermore, these

315 observations indicate that during natural transformation in *P. aeruginosa* the PilQ secretin pore is not

316 absolutely required for translocation of plasmid DNA across the outer membrane.

318 **Discussion**

319 Here we have demonstrated, in contrast to current dogma, that *P. aeruginosa* is capable of natural

320 transformation of both plasmid and chromosomal DNA under conditions that promote the expression

321 of T4P and eDNA production, such as in static broth cultures and biofilms. We found that whilst T4P

322 appear to be involved in facilitating plasmid DNA uptake, T4P are not absolutely required for natural

323 transformation of plasmid DNA in *P. aeruginosa.* Furthermore, our data suggests that the PilQ

324 secretin pore is not absolutely required for translocation of plasmid DNA across the outer membrane

325 in this organism.

327 It is thought that separate uptake mechanisms for plasmid or chromosomal DNA are present in *E.*

328 *coli*[8]. While natural transformation of chromosomal DNA and integration into the genome has not

329 been demonstrated in *E. coli*, T4P/competence pseudopilus and DNA translocation machinery

330 homologs have been shown to be responsible for uptake of chromosomal DNA for use as a nutrient

331 source[34]. Natural plasmid transformation in *E. coli* does not utilise the competence pseudopilus

332 uptake system although the precise mechanism for this process has not been determined[9]. At this

333 stage we cannot rule out the possibility that there are two mechanisms in *P. aeruginosa* involved in

334 the natural transformation of plasmid DNA or chromosomal DNA.

336 The finding that *P. aeruginosa* is capable of natural transformation is a paradigm shift in our

337 understanding of how this pathogen acquires genetic diversity. Indeed, recombination has been

identified as a major means of genetic diversity in *P. aeruginosa* cystic fibrosis (CF) lung isolates although the source of DNA and the mechanism of HGT was not determined[35]. Natural transformation may be an important mechanism for the acquisition of antibiotic resistance and virulence genes in this ESKAPE pathogen and a significant contributor to the rapid increase in the number of multidrug resistant *P. aeruginosa* strains that are an emerging problem worldwide.

Acknowledgements

We thank Jacob Bertrand, Elizabeth Tran, Lisa MacAskill, Dervilla McGowan, Heather Smith and Kate Rainzcuk for technical assistance and helpful discussions. L.M.N was supported by an Imperial College Research Fellowship. D.L. was supported by the Swiss National Science Foundation (SNSF, grant n. P2GEP3_161769). C.B.W. was supported by a National Health and Medical Research Council of Australia (NHMRC) Career Development Award and a Senior Research Fellowship (571905).

Author information

Contributions

L.M.N, L.T, M.K, S.R.O, D.L, J.J.L. and C.B.W performed experiments. L.M.N, L.T and C.B.W analyzed data. C.B.W. provided project administration and funding. L.M.N. and C.B.W. wrote the manuscript.

Ethics declarations

The authors declare no conflict of interest.

References

1. O'Neill, J. *Tackling drug-resistant infections globally: Final report and recommendations*.

(UK Government and Wellcome Trust, 2016).

2. Pallen, M. J. & Wren, B. W. Bacterial pathogenomics. *Nature* **449,** 835–842 (2007).

3. Davison, J. Genetic exchange between bacteria in the environment. *Plasmid* **42,** 73–91 (1999).

4. Dubnau, D. & Blokesch, M. Mechanisms of DNA uptake by naturally competent bacteria.

Annu. Rev. Genet. (2019). doi:10.1146/annurev-genet-112618-043641

5. Ibáñez de Aldecoa, A. L., Zafra, O. & González-Pastor, J. E. Mechanisms and regulation of

extracellular DNA release and its biological roles in microbial communities. *Front. Microbiol.*

8, 1390 (2017).

6. Ellison, C. K. *et al.* Retraction of DNA-bound type IV competence pili initiates DNA uptake

during natural transformation in Vibrio cholerae. *Nat. Microbiol.* **3,** 773–780 (2018).

7. Chen, I. & Dubnau, D. DNA uptake during bacterial transformation. *Nat. Rev. Microbiol.* **2,**

241–249 (2004).

8. Hasegawa, H., Suzuki, E. & Maeda, S. Horizontal Plasmid Transfer by Transformation in

Escherichia coli: Environmental Factors and Possible Mechanisms. *Front. Microbiol.* **9,** 2365

(2018).

9. Sun, D. *et al.* Transforming DNA uptake gene orthologs do not mediate spontaneous plasmid

transformation in Escherichia coli. *J. Bacteriol.* **191,** 713–719 (2009).

10. Kung, V. L., Ozer, E. A. & Hauser, A. R. The accessory genome of Pseudomonas aeruginosa.

Microbiol. Mol. Biol. Rev. **74,** 621–641 (2010).

11. Shen, K. *et al.* Extensive genomic plasticity in Pseudomonas aeruginosa revealed by

identification and distribution studies of novel genes among clinical isolates. *Infect. Immun.* **74,**

5272–5283 (2006).

12. Carlson, C. A., Pierson, L. S., Rosen, J. J. & Ingraham, J. L. Pseudomonas stutzeri and related

species undergo natural transformation. *J. Bacteriol.* **153,** 93–99 (1983).

13. Whitchurch, C. B. in *Pseudomonas Volume 4 Molecular Biology of Emerging Issues* (ed. J. L.

Ramos and R. C. Levesque) Springer USA: 139-188 (2006).

14. Allesen-Holm, M. *et al.* A characterization of DNA release in *Pseudomonas aeruginosa*

cultures and biofilms. *Mol. Microbiol.* **59,** 1114–1128 (2006).

15. Whitchurch, C. B., Tolker-Nielsen, T., Ragas, P. C. & Mattick, J. S. Extracellular DNA

required for bacterial biofilm formation. *Science* **295,** 1487 (2002).

16. Gloag, E. S. *et al.* Self-organization of bacterial biofilms is facilitated by extracellular DNA.

Proc. Natl. Acad. Sci. USA **110,** 11541–11546 (2013).

17. Turnbull, L. *et al.* Explosive cell lysis as a mechanism for the biogenesis of bacterial

membrane vesicles and biofilms. *Nat. Commun.* **7,** 11220 (2016).

18. Takeya, K. & Amako, K. A rod-shaped Pseudomonas phage. *Virology* **28,** 163–165 (1966).

19. Rahme, L. G. *et al.* Common virulence factors for bacterial pathogenicity in plants and

animals. *Science* **268,** 1899–1902 (1995).

20. Liu, P. V. Exotoxins of Pseudomonas aeruginosa. I. Factors that influence the production of

exotoxin A. *J. Infect. Dis.* **128,** 506–513 (1973).

21. Klausen, M. *et al.* Biofilm formation by Pseudomonas aeruginosa wild type, flagella and type

IV pili mutants. *Mol. Microbiol.* **48,** 1511–1524 (2003).

22. Hoang, T. T., Kutchma, A. J., Becher, A. & Schweizer, H. P. Integration-proficient plasmids

for *Pseudomonas aeruginosa*: site-specific integration and use for engineering of reporter and

expression strains. *Plasmid* **43,** 59–72 (2000).

23. Watson, A. A., Mattick, J. S. & Alm, R. A. Functional expression of heterologous type 4

fimbriae in Pseudomonas aeruginosa. *Gene* **175,** 143–150 (1996).

24. Martin, P. R., Hobbs, M., Free, P. D., Jeske, Y. & Mattick, J. S. Characterization of pilQ, a

new gene required for the biogenesis of type 4 fimbriae in Pseudomonas aeruginosa. *Mol.*

Microbiol. **9,** 857–868 (1993).

25. Whitchurch, C. B., Hobbs, M., Livingston, S. P., Krishnapillai, V. & Mattick, J. S.

Characterisation of a Pseudomonas aeruginosa twitching motility gene and evidence for a specialised protein export system widespread in eubacteria. *Gene* **101,** 33–44 (1991).

26. Alm, R. A. & Mattick, J. S. Identification of a gene, pilV, required for type 4 fimbrial biogenesis in Pseudomonas aeruginosa, whose product possesses a pre-pilin-like leader sequence. *Mol. Microbiol.* **16,** 485–496 (1995).

27. Semmler, A. B., Whitchurch, C. B., Leech, A. J. & Mattick, J. S. Identification of a novel gene,. *Microbiology (Reading, Engl.)* **146 (Pt 6),** 1321–32 (2000).

28. Watson, A. A., Alm, R. A. & Mattick, J. S. Construction of improved vectors for protein production in *Pseudomonas aeruginosa. Gene* **172,** 163–164 (1996).

29. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment search tool. *J. Mol. Biol.* **215,** 403–410 (1990).

30. Winsor, G. L. *et al.* Enhanced annotations and features for comparing thousands of Pseudomonas genomes in the Pseudomonas genome database. *Nucleic Acids Res.* **44,** D646-53 (2016).

31. Cain, A. K. *et al.* Complete Genome Sequence of Pseudomonas aeruginosa Reference Strain PAK. *Microbiol. Resour. Announc.* **8,** (2019).

32. Miller, S. A., Dykes, D. D. & Polesky, H. F. A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res.* **16,** 1215 (1988).

33. Schleheck, D. *et al.* Pseudomonas aeruginosa PAO1 preferentially grows as aggregates in liquid batch cultures and disperses upon starvation. *PLoS One* **4,** e5513 (2009).

34. Palchevskiy, V. & Finkel, S. E. Escherichia coli competence gene homologs are essential for competitive fitness and the use of DNA as a nutrient. *J. Bacteriol.* **188,** 3902–3910 (2006).

35. Dettman, J. R., Rodrigue, N. & Kassen, R. Genome-wide patterns of recombination in the opportunistic human pathogen Pseudomonas aeruginosa. *Genome Biol. Evol.* **7,** 18–34 (2014).

Tables and Figures

Table 1. Homologs of proteins involved in natural transformation in a range of bacteria.

Competence Protein	*Bacillus subtilis*	*Streptococcus pneumoniae*	*Haemophilus influenzae*	*Thermus thermophilus*	*Pseudomonas stutzeri*	*Neisseria gonorrhoeae*	*Pseudomonas aeruginosa**
T4P/Competence Pseudopilus							
Traffic NTPase(s)	ComGA	ComGA	PilB	PilF	PilT, PilU	PilF, PilT	PilB, PilT, PilU
Polytopic membrane protein	ComGB	ComGB	PilC	PilC	PilC	PilG	PilC
Pilins or pseudopilins	ComGC, -GD, -GE, -GG	CglC, CglD	PilA	PilA1, -A2, -A3, -A4	PilA1	PilE, ComP	PilA, -V, -W, -X, -E, FimT, FimU
Prepilin peptidase	ComC	CilC	PilD	PilD		PilD	PilD
Secretin/pilot	na	na	ComE	PilQ		PilQ/PilP	PilQ/PilP
DNA translocation machinery							
DNA receptor	ComEA	ComEA		ComEA		ComE	PA3140
Membrane channel	ComEC	ComEC	Rec-2	ComEC	ComA	ComA	PA2984
ATP-binding protein	ComFA	ComFA			ExbB		PA2983
Other							
			DprA (Smf)				PA0021
			TfoX (Sxy)				PA4703
			CRP				Vfr
			CyaA				CyaA, CyaB
			ComM				PA5290
			ComF				PA0489

*Using *P. aeruginosa* PAO1 gene nomenclature

Figure 1. *P. aeruginosa* is capable of natural transformation in continuous flow biofilms.

Carbenicillin or gentamicin resistant transformants obtained from continuous flow biofilms of

PAO1 in Tygon tubing. For (a) pUCPSK was added to the media influent (at a final concentration

of 1 μg/mL) with either the biofilm biomass or effluent harvested and plated on carbenicillin

selective agar plates on the indicated day. For (b-c) sterile gDNA or eDNA obtained from PAO1GFP

(GmR) (at a final concentration of 0.5 mg/mL) was added to the media influent and the biofilm

biomass harvested and plated on gentamicin selective agar plates after 5 days. The mean of each set

of technical triplicates was calculated to give an n≥3 which is presented as mean ± S.E.M.

Figure 2. *P. aeruginosa* is capable of natural transformation of exogenous chromosomal DNA.

The inoculum PAO1 strain did not express GFP (a). Sterile PAO1$_{GFP}$ gDNA (at a final

concentration of 0.5 mg/mL) was added to the media influent of continuous flow biofilms of PAO1

in Tygon tubing (b) or flow cells (c-f) and incubated for 5 days. Gentamicin resistant (Gm^R)

colonies obtained from Tygon tubing biofilms were resuspended in PBS and visualized by

epifluorescence microscopy which showed all cells from Gm^R colonies expressed GFP (B). The

biofilm biomass from PAO1 flow cells was visualized by phase contrast (c, e) or epifluorescence

microscopy (d, f) which showed the presence of representative biofilm microcolonies expressing

GFP. Scale bar 100 μm.

Figure 3. Lab and clinical strains of *P. aeruginosa* are capable of natural transformation of

plasmid DNA within colony biofilms. pUCPSK (5 μg) DNA was applied to colony biofilms of (a)

P. aeruginosa PAK or PAO1, or (b) *P. aeruginosa* lab or clinical strains and incubated for 2 hr.

Cells were then harvested and the number of carbenicillin resistant transformants determined by

spread plating on selective media. For (a) the mean of each set of technical triplicates was

calculated to give an n≥4 which is presented as mean ± S.E.M. For (b) the values presented are

from n=1. For the CF sputum isolates the designation M refers to mucoid phenotype, NM is non-

mucoid.

Figure 4. *P. aeruginosa* is capable of natural transformation of plasmid DNA in static broth cultures. Carbenicillin resistant transformants obtained from static broth cultures incubated for 24 hr with (a) 10 μg pUCPSK or (b) 0-30 μg pUCPSK. The mean of each set of technical triplicates was calculated to give an n≥3 which is presented as mean ± S.E.M. For (a) ** P <.005; Mann-Whitney U-test compared to PAK.

Figure 5. T4P are not absolutely required for natural transformation of plasmid DNA. Static (a-c) or shaking (a) broth cultures were incubated for 24 hr with 10 μg pUCPSK and the number of (a-b) carbenicillin resistant transformants or (c) aggregates observed determined. For (a-b) the mean of each set of technical triplicates was calculated to give an n≥3 which is presented as mean ± S.E.M. For (a) $P > .05$; Mann-Whitney U-test. For (b) ** $P <.005$; Mann-Whitney U-test compared to PAK. For (c) the total number of aggregates from 60 random fields imaged on 3 separate days (20 fields/day) is presented.

Supplementary Figures

Supplementary Figure 1. Transformant numbers increase with plasmid DNA concentration

in static broth cultures. Static broth cultures of PAK or PAO1 with 10 μg pUCPSK added were

incubated and the numbers of carbenicillin resistant transformants determined after 24 hr. The mean

of each set of technical triplicates was calculated to give an n≥3 which is presented as mean ±

S.E.M.

Supplementary Figure 2. Natural transformation of gDNA and eDNA is inefficient in static broth cultures of *P. aeruginosa*. Static broth cultures of PAK or PAO1 with 15 μg gDNA or eDNA were incubated and the numbers of gentamicin resistant transformants determined after 24 hr. The mean of each set of technical triplicates was calculated to give an n≥3 which is presented as mean ± S.E.M.