Distinct life histories impact dikaryotic genome evolution in the rust fungus *Puccinia striiformis* causing stripe rust in wheat

Authors:

Benjamin Schwessinger[6,#,$], Yan-Jun Chen[1,#], Richard Tien[2], Josef Korbinian Vogt[3], Jana Sperschneider[4], Ramawatar Nagar[6], Mark McMullan[7], Thomas Sicheritz-Ponten[1], Chris K. Sørensen[5], Mogens Støvring Hovmøller[5], John P. Rathjen[$], and Annemarie Fejer Justesen[5,$]

[1]University of Copenhagen, Faculty of Health and Medical Sciences, The GLOBE Institute, Øster Farimagsgade 5, DK-1353 Copenhagen K, Denmark
[2]School of Dentistry, The University of Western Australia, Nedlands WA 6009, Australia
[3]University of Copenhagen, Novo Nordisk Foundation Center for Basic Metabolic Research, Maersk Tower, Panum, Blegdamsvej 3B, DK-2200, Copenhagen N, Denmark
[4]Biological Data Science Institute, The Australian National University, Acton ACT 2601, Australia.
[5]Department of Agroecology, Faculty of Science and Technology, Aarhus University, Forsøgsvej 1, DK-4200 Slagelse, Denmark
[6]Research School of Biology, The Australian National University, Acton ACT 2601, Australia.
[7]Earlham Institute, Norwich Research Park, Norwich, NR4 7UZ, UK.

# These authors contributed equally to this work.
$Author for Correspondence:
Benjamin Schwessinger, Research School of Biology, The Australian National University, Acton, ACT 2601, Australia, +61 2 6125 7794, benjamins.schwessinger@anu.edu.au
Annemarie Fejer Justesen, Department of Agroecology, Aarhus University, Forsøgsvej 1, DK-4200 Slagelse, Denmark, +45 87158135, annemariefejer.justesen@agro.au.dk
John P. Rathjen, Research School of Biology, The Australian National University, Acton, ACT 2601, Australia, +61 2 6125

Abstract

Stripe rust of wheat, caused by the obligate biotrophic fungus *Puccinia striiformis* f. sp. *tritici*, is a major threat to wheat production world-wide with an estimated yearly loss of US $1 billion. *P. striiformis* may undergo a complex life cycle with several spore types and an alternate host for completion of the sexual cycle. It infects the gramineous host (e.g., wheat) during its asexual stage via repeated cycles of infection by dikaryotic urediniospores. The recent advances in long-read sequencing technologies and tailored-assembly algorithms enabled us to disentangle the two haploid genomes of *Pst*. This provides us with haplotype-specific information at a whole-genome level. Exploiting this novel information, we perform whole genome comparative

genomics of two *P. striiformis* f. sp. *tritici* isolates with contrasting life histories. We compare one
isolate of the old European lineage (PstS0), which has been asexual for over 50 years, and a
Warrior isolate (PstS7 lineage) from a novel incursion into Europe in 2011 from a sexual
population in the Himalayan region. This comparison provides evidence that long-term asexual
evolution leads to genome expansion, accumulation of transposable elements, and increased
heterozygosity at the single nucleotide, structural and allele levels. At the whole genome level,
candidate effectors are not compartmentalized and do not exhibit reduced levels of synteny. Yet
we were able to identify two subsets of candidate effector populations. About 70% of candidate
effectors are invariant between the two isolates while 30% are hypervariable. The latter might
be involved in host adaptation on wheat and explain the different phenotypes of the two isolates.
Overall this detailed comparative analysis of two haplotype-aware assemblies of *P. striiformis* f.
sp*. tritici* are the first steps in understanding the evolution of dikaryotic rust fungi at a whole
genome level.
54
55 **Keywords:** Basidiomycetes, dikaryon, haplotype, plant pathogen, asexual reproduction, sexual
56 reproduction, wheat yellow rust, wheat stripe rust
57
58
59 **Introduction**
60
Fungi have complex life cycles involving clonal as well as sexual reproduction. Sex in fungi is
often associated with dispersal, dormancy and survival under adverse conditions such as the
absence of hosts (Anikster 1986). Sexual reproduction creates genotypic diversity and enables
adaptation to new environments by combining beneficial alleles and eliminating harmful genetic
traits (Seudre et al. 2018; Heitman 2015). The advantage of asexual reproduction, on the other
hand, is the rapid multiplication and spread of the most well-adapted individuals. In some
species, sexual reproduction is so rare or cryptic that the sexual cycle historically was thought to
be absent and only recently has been discovered (Zeyl 2009; Taylor et al. 2015). Studies of the
genomic architecture of fungal pathogens with different reproduction strategies can help us to
understand the genetic and genomic processes behind pathogen adaptation. This knowledge is
important for predicting the response of pathogens to challenges such as climate change,
drugs, fungicides, host resistance or immunity, and will strengthen the development of methods
for combating fungal pathogens (Gladieux et al. 2014).
74

Rust fungi, the *Pucciniales*, are a large group of fungal plant pathogens, some of which impact
severely on agricultural and forest production (Lorrain et al. 2019). They depend on living host
cells and have complex life cycles, which may comprise many spore types over two
phylogenetically distinct host plants for completion of the sexual life cycle. *Puccinia striiformis*
causes yellow (stripe) rust on cereals and grasses and has historically been restricted to
temperate-cool and wet climates (Rapilly 1979). Since 2000, new aggressive strains adapted to
higher temperatures have caused the expansion of the disease into warmer regions, and it is
currently one of the most prevalent and damaging diseases on common wheat (*Triticum*
aestivum L.) worldwide (Beddow et al. 2015; Wellings 2011; Savary et al. 2019; Singh et al.
2016). During the asexual life cycle, which recurs several times during the wheat growing
season, *P. striiformis* produces dikaryotic urediniospores which reinfect wheat (Schwessinger
2017). These spores are wind borne and may spread over long distances to introduce exotic
strains into new areas (Hovmøller et al. 2016; Ali, Gladieux, Leconte, et al. 2014). Historically,
the sexual cycle of *P. striiformis* has been a great mystery and many years of search for the
alternate host was fruitless until it was shown that barberry (*Berberis* spp.) can serve as the
alternate host for completion of the sexual cycle (Jin et al. 2010). The significance of *Berberis*
spp. as an alternate host for *P. striiformis* in nature is still unknown, but high levels of genetic
diversity in *P. striiformis* populations in China and Pakistan indicates that sexual reproduction is
taking place under natural conditions, and the near-Himalayan region is the likely the centre of
genetic diversity for *P. striiformis* (Mboup et al. 2012; Duan et al. 2010; Ali et al. 2016; Ali,
Gladieux, Rahman, et al. 2014). In contrast, reproduction of *P. striiformis* in Europe, America
and Australia is clonal, and adaptation to host resistance happens through stepwise mutations
(Chen et al. 2014; Hovmøller & Justesen 2007; Steele et al. 2001; Markell & Milus 2008). These
mutations lead to the loss of recognition of *P. striiformis* effectors which previously were
recognized by specific plant resistance (*R*) gene products and *P. striiformis* is then termed
virulent to the specific *R* gene. In the absence of recognition, e.g. in a wheat plant that lacks the
corresponding *R* gene, effectors suppress plant immune responses and facilitate the infection
process (Ellis et al. 2014).
103
Only few clonal lineages of *P. striiformis* are responsible for the yellow rust epidemics
worldwide, and stepwise mutation and clonal reproduction has been the major driving force for
the evolution of new virulent races that can overcome widely deployed resistance genes (Walter
et al. 2016; Hovmøller et al. 2008; Ali et al. 2017). Combination of virulence alleles by sexual or
somatic recombination may also generate strains with new virulence profiles, yet the frequency

of these mechanisms in nature is not yet known. Long-term asexual isolates though produce
fewer teliospores, which are obligatory for completing the sexual life cycle and this indicates that
these long-term clonal strains may have a reduced ability to undergo sexual reproduction (Ali et
al. 2010).

In 2011, a new *P. striiformis* lineage, PstS7, known generally as Warrior was detected in high
frequency on both wheat and triticale (*x Triticosecale*) in many European countries (Hubbard et
al. 2015; Hovmøller et al. 2016). It was able to cause disease on cultivated host varieties that
had previously been resistant (Sørensen et al. 2014). Moreover, it produced high amounts of
telia and was able to complete the sexual cycle on barberry under experimental conditions
(Rodriguez-Algaba et al. 2014). Warrior is genetically very different from the clonal lineages
present in Europe before 2011 (e.g., PstS0) and similar to isolates originating from the centre of
diversity in the near-Himalayan region (Hovmøller et al. 2016; Hubbard et al. 2015; Ali,
Gladieux, Leconte, et al. 2014). Determining the mechanisms behind the generation of new
virulent strains of *P. striiformis* such as Warrior is important. However, sequencing technologies
available to date have provided only limited understanding of genetic recombination, comprising
somatic, parasexual and sexual recombination, as a driving force for generating new strains of
the pathogen.

Until recently, most draft genome assemblies of *P. striiformis* were based largely on short-read
sequencing data (Cantu et al. 2011, 2013; Zheng et al. 2013). However, short-read sequencing
data is not well suited to the dikaryotic rusts due to the high proportion of repetitive sequences
(>50%), high heterozygosity, leading to the difficulty in phasing and high level of fragmentation
of the two haploid genomes. Hence, short-read assemblies typically lack contiguity and
individual haplotype information. Single molecule sequencing technologies such as PacBio and
Nanopore produce far longer reads, which in combination with novel assembly algorithms have
provided more accurate (Xia, Wang, Yin, Cornejo, Scot Hulbert, et al. 2018) and complete
genomes with effective haplotype phasing (Schwessinger et al. 2018). Here we provide a
genome assembly of a Warrior isolate sampled in Denmark (Pst-DK0911) using long-read
sequencing and a diploid-aware genome assembler (Schwessinger et al. 2018; Chin et al.
2016). We previously produced a highly phased and contiguous assembly of the presumed
asexual Australian isolate Pst-104E belonging to the old European clonal lineage PstS0
(Schwessinger et al. 2018; Wellings 2007). This new genome assembly allows whole genome
comparisons between two isolates representing contrasting lifestyles of *P. striiformis.* It provides

detailed insights into the adaptive divergence and evolution of *P. striiformis* and rust fungi in
general. Specifically, we explore question around the impact of sexual and asexual reproduction
on structural genomic evolution, the generation of genetic diversity, and host adaptation.
146
147
148   **Methods and materials**
149

***Puccinia striiformis* f. sp. *tritici* pathotype, growth conditions and spore amplification**
The Pst-DK0911 isolate was purified from a single lesion sampled from the commercial wheat
cv. Holeby in Denmark in 2011. The isolate was race phenotyped according to Hovmøller et al.
2017 (Hovmøller et al. 2017) and genotyped as described previously (Hovmøller et al. 2016;
Rodriguez-Algaba et al. 2014). Prior to DNA extraction, the isolate was multiplied on susceptible
seedlings of wheat cv. Morocco, which were inoculated using the method described (Sørensen
et al. 2016). The inoculated seedlings were misted with water and incubated at 10°C in the dark
for 24 h under high relative humidity in an incubation room. After incubation the inoculated
plants were transferred to quarantine spore-proof greenhouse cabinets with a temperature
regime of 17°C day and 12°C night, and a light regime of 16 h photoperiod of natural light and
supplemental sodium light (100 µmol s^{-1} m^{-2}), and 8 h dark. The plants were covered with
cellophane bags 5-7 days after infection (dai) to avoid cross-contamination before sporulation.
The urediniospores were harvested at 15–20 dai and freeze-dried before DNA extraction.
163

DNA extraction and genome sequencing
We extracted DNA from freeze-dried urediniospores as described previously (Schwessinger &
Rathjen 2017) with few modifications. A volume of 875 µl lysis buffer was used per 60 mg
spores in a 2 ml eppendorf tube. Library preparation and PacBio sequencing were performed at
the Earlham Institute (Norwich, UK). One DNA library was sequenced on a PacBio RSII
instrument using P6-C4 chemistry and 16 SMRT cells were sequenced in total. Moreover, a
DNA sample from the same isolate were sequenced using Illumina short read technology.
TruSeq library was sequenced on a HiSeq 2000 instrument as a 100 bases paired-end library at
the Danish National High-Throughput DNA Sequencing Centre (Copenhagen, Denmark).
173

RNAseq
Pst-DK0911 was point inoculated on leaves of the susceptible cv. Avocet S with a pipette using
5 µl spore suspension in 3M^{TM} Novec^{TM} 7100 (2 mg spores per ml) as described previously

(Sørensen et al. 2016) without fixation of leaves. Three replicates were prepared and leaf
segments of 10-12 mm of the 2nd leaves of each replicate were harvested at 7 and 9 dai for
transcriptome sequencing. We isolated total RNAs with the RNeasy Plant Mini kit (Qiagen) from
100 mg of infected leaf tissues including a DNaseI (Qiagen) treatment according to the
manufacturer's instructions. Transcriptome libraries were generated using the TruSeq Stranded
Total RNA with Ribo-Zero plant kit (Illumina) according to the standard protocol to remove
cytoplasmic, mitochondrial and chloroplast ribosomal RNA. Next-generation sequencing (NGS)
for 6 multiplexed samples was performed using an Illumina HiSeq2000 sequencer at BGI
(Copenhagen, Denmark).
186

Genome Assembly
For genome assembly we used FALCON and FALCON -Unzip (Chin et al. 2016) github tag
1.7.4 with the parameters described in Supplemental Data 1 and 2. We checked the resulting
contigs for eukaryotic contamination by blastn searches against the NCBI nucleotide reference
database (downloaded 04052016) (Altschul et al. 1990). None of the contigs had predominant
non-eukaryotic sequences as best blast hits at a given position. We performed two manual
curation steps. In the first step we reasoned that some of the primary contigs without
corresponding haplotigs might actually represent haplotigs that could not be assigned to their
respective primary contigs in the assembly graph because of too large a difference between the
two haplotypes. We aligned all primary contigs without haplotigs to primary contigs with
haplotigs using MUMmer version 3 (Kurtz et al. 2004). We screened the best alignments of
each primary contig without haplotig for percentage alignment, length of alignment, and if they
aligned to regions in the primary contigs that previously had not been covered by a haplotig
alignment. Using this approach, we re-assigned five primary contigs without haplotigs (~6 Mb)
to haplotigs.
202

Gene model annotation
We annotated genes on primary contigs and haplotigs independently. We combined RNAseq
guided *ab initio* prediction using CodingQuarry v2.0 (Testa et al. 2015) and braker v1.9 (Hoff et
al. 2016) with *de novo* transcriptome assembly approaches using Trinity v2.2.0 (Grabherr et al.
2011) and pasa v2.0.1 (Haas et al. 2008). Gene models were unified using evidencemodeler
v1.1.1 (Haas et al. 2008) using the weights given in Supplemental Data 3.
We mapped the trimmed RNAseq reads described in this study (see above) and previously
(Dobon et al. 2016; Schwessinger et al. 2018) against the complete assembly (primary contigs

and haplotigs) using hisat2 v2.1.0 (--max-intronlen 10000 --min-intronlen 20 --dta-cufflinks) (Kim

et al. 2015). For *ab initio* predictions we reconstructed transcripts using stringtie v1.3.3b (-f 0.2)

(Pertea et al. 2015). We ran CodingQuarry (-d) in the pathogen mode using SignalP4 (Petersen

et al. 2011) for secretome prediction on the soft-masked genome using RepeatMasker v4.0.5 (-

xsmall -s -GC 43). Similarly, we used the stringtie reconstructed transcripts as training set for

the *ab initio* prediction pipeline braker 1 v1.9 (Hoff et al. 2016) and used the non-repeat masked

genome as reference.

We used Trinity v2.2.0 to obtain *P*. *striiformis* f. sp*. tritici* transcripts both in the *de novo* mode

and the genome guided mode. All RNAseq samples contained host and pathogen RNA as they

were prepared from infected wheat tissue. We first mapped all reads to the genome using hisat2

(see above). We extracted mapped RNAseq reads using picard tools SamToFastq. Only those

reads that mapped against *P. striiformis* f. sp*. tritici* contigs were used in the *de novo* pipeline of

Trinity (--seqType fq). For genome guided assembly we used bam files generated with hisat2 as

starting point for Trinity (--jacard_clip, --genome_gudied_max_intron 10000). We used the pasa

pipeline v2.0.2 to align both sets of Trinity transcripts against *P. striiformis* f. sp*. tritici* contigs

with blat and gmap.

The different gene models were combined using evidencemodeler v.1.1.1 to get the initial gene

sets for primary contigs and haplotigs. These were filtered for homology with proteins encoded

in transposable elements. We used blastp to search for homology in the Repbase v21.07

peptides database with an e-value cut-off of $1*e^{-10}$. In addition, we used transposonPSI (Bao et

al. 2015). We used the outer union of both approaches to remove genes coding for proteins

associated with transposable elements from our list of gene models.

Repeat annotation and analysis

Repeat regions were annotated as described previously (xx) (Schwessinger et al. 2018)using

the REPET pipeline v2.5 (Quesneville et al. 2005; Flutre et al. 2011; Maumus & Quesneville

2014) for repeat annotation in combination with Repbase v21.05 (Bao et al. 2015). For de novo

identification, we predicted repeat regions in primary contigs of Pst_104E_v13 and DK0911_v04

independently using TEdenovo. We combined the resulting de novo repeat libraries with the

script 'RemoveRedundancyBasedOnCI.py' within the REPET pipeline using the configurations

provided in Supplemental Data 4. We named the resulting repeat library

104E_p_DK0911_p_classif. We annotated the primary contigs and haplotigs of both assemblies

separately using three repeat libraries (repbase2005_aaSeq, repbase2005_ntSeq, and

104E_p_DK0911_p_classif). We generated superfamily classifications as described previously

(Schwessinger et al. 2018). See jupyter notebooks
DK0911_TE_filtering_and_summary_p_contigs_fix.ipynb,
DK0911_TE_filtering_and_summary_h_contigs_fix.ipynb,
Pst_104E_v14_TE_filtering_and_summary_p_contigs_fix.ipynb, and
Pst_104E_v13_TE_filtering_and_summary_h_contigs_fix.ipynb in the github repo for full
analysis details.
We analyzed the relative identity distribution of TEs as proxy of TE age as previously described
(Maumus & Quesneville 2014). The average identity and genome coverage per TE family is
provided by the REPET pipeline. We used these tables as input and plotted the relative
coverage frequency of TE families within average identity size bins of one (Supplemental Figure
2A). We plotted the cumulative difference in TE genome coverage between Pst-104E and Pst-
DK0911 against the average TE family identity normalizing the current observed TE coverage
difference to one. See jupyter notebook DK0911_TE_variation_analysis.ipynb for details.
258

Protein annotation
We performed protein annotation as described previously including homology to swissprot
proteins (uniref90, downloaded 22/09/2016) (Bairoch & Apweiler 2000), to carbohydrate-active
enzymes (dbCAN, downloaded 22/09/2016) (Yin et al. 2012), to peptidases (MEROPS v10.0)
(Rawlings et al. 2016) (see jupyter notebooks in the protein_annotation folder on github). We
complemented this analysis by interproscan v5.21-60 (-iprlookup -goterms -pa) (Jones et al.
2014), eggnog-mapper v0.99.2 (-m diamond and –d euk) (Huerta-Cepas et al. 2016), SignalP 3
(Bendtsen et al. 2004), and EffectorP v2.0 (Sperschneider et al. 2015, 2018).
267

268 **BUSCO analysis**
269 We used BUSCO3 to identify core conserved genes and to assess genome completeness
270 (Simão et al. 2015). In all cases we ran BUSCO3 in the protein mode using the basidiomycota
271 reference database downloaded 01/09/2016 (-l basidiomycota_odb9 -m protein).
272

Inter-haplotype variation analysis
We performed the inter-haplotype variation analysis similar as described previously
(Schwessinger et al. 2018). Trimmed Illumina reads derived from DK0911, Pst_104E, and
PST78 were mapped against the genomes (DK0911 and Pst_104E) using BWA-MEM v0.7.15-
r1142-dirty using the standard parameters (Li 2013). We called SNPs in bulk using freebays
(Garrison & Marth 2012) and following hard filtering by vcffilter with DP > 10 and QUAL > 20

(2017b). Variants for DK0911, Pst_104E, and PST78 resequencing data were parsed out and
summarized with real time genomic vcfstats v3.8.4 (2017a).

We performed large-scale variation analysis as described previously using nucmer and
assemblytics (Kurtz et al. 2004; Nattestad & Schatz 2016). See jupyter notebooks
DK_0911_nucmer_assemblytics_mapping.ipynb, DK_0911_assemblytics_analysis.ipynb, and
DK_0911_Pst104E_assemblytics_plot.ipynb for details.

Within genome allele variation analysis
We used proteinortho v5.16 in synteny mode with default parameters (-synteny) to identify
alleles between the primary assembly and haplotigs (Lechner et al. 2011). We restricted our
analysis to high quality allele pairs that are located on linked primary contigs and haplotigs. We
assessed the variation of high-quality allele pairs by calculating the Levenshtein distance on the
codon-based CDS alignments. The pairwise protein alignments were generated with muscle
v3.8.31 (Edgar 2004), and converted into codon-based CDS alignments using PAL2NAL v14
(Suyama et al. 2006).   Analysis details can be found in jupyter notebook
DK0911_vs_Pst104E_gene_pair_analysis.ipynb.

Coverage analysis and identification of unphased regions in primary contigs
We identified unphased regions in Pst-DK0911 as reported previously for Pst-104E
(Schwessinger et al. 2018). We performed detailed read depth coverage analysis to obtain a
genome level insight into the relative quantity of fully phased, homozygous collapsed, and
hemizygous regions in the two *P. striiformis* f. sp*. tritici* genomes. We determined the
sequencing depth in 1-kb sliding windows with 200-base intervals. Read depth coverage was
calculated by mapping short reads derived from each *P. striiformis* f. sp*. tritici* isolate against its
own genome using BWA-MEM v0.7.15-r1142-dirty and the standard parameters (Li 2013). We
mapped reads against primary contigs only (p) and both primary contigs and haplotigs (ph). We
normalized sequencing depth to one at the main "haploid" sequencing depth peak on primary
contigs when mapping against primary contigs and haplotigs (Figure 2 B). In a perfect fully
phased assembly this is the main sequencing depth peak, because of the absence of any
homozygous collapsed regions. We plotted the normalized sequencing depth for primary
contigs (both p and ph mapping), haplotigs (ph mapping), regions in primary contigs that have
corresponding phased haplotigs and regions in primary contigs without assigned haplotigs. For
details   on   the   generation   of   required   bed   files   see   jupyter   notebook
DK0911_SRM_cov_DK0911_on_DK0911.ipynb,

Pst_104E_SRM_cov_Pst_104E_on_Pst_104E.ipynb,   and

DK0911_Pst104E_comperative_coverage_analysis.ipynb reference (xxx).

Cross isolate genome wide presence-absence analysis

We aimed at determining presence absence polymorphism at the whole genome level. We first

mapped short reads derived from one *P. striiformis* f. sp*. tritici* isolate ('query') against the

complete genome (ph) of the other isolate ('subject') using BWA-MEM v0.7.15-r1142-dirty using

standard parameters. We then calculated sequencing depths in 1 kb sliding windows as

described above. We defined regions that fell below 10% normalized sequencing depth in the

cross mapping but not self mapping experiments as absent in the 'query' isolate. We used

bedtools (Dale et al. 2011; Quinlan & Hall 2010) to identify genes which are completely

contained within these regions and defined them as 'variable genes'. We estimated the

distribution of the expected number of variable genes using a permutation test randomly

reshuffling the absent regions five thousand times. We used this random distribution to calculate

a p-value testing the hypothesis if the observed number of variable genes is different from the

observed random distribution at a p-value cut-off of 0.05. We performed a similar analysis for

TEs with the important difference to determine variable TE content at single base pair resolution

and with two thousand five hundred permutations. In addition, we performed similar tests on the

variable TE content using reciprocal whole genome alignments as input. We aligned each

genome against each other with the mummer package v4. Initially, alignments with nucmer

used -l 20 -c 65 --max-match. Alignments were filtered with delta-filter requiring no uniqueness -

u 0 (see notebook Mummer_DK0911_Pst_104E.ipynb for more detail). We converted mummer

delta files into bed files for input in the presence absence analysis as described above. Details

on   this   analysis   can   be   found   in   jupyter   notebooks

DK0911_SRM_cov_Pst_104E_on_DK0911.ipynb,

Pst_104E_SRM_cov_Pst_104E_on_Pst_104E.ipynb, Mummer_DK0911_Pst_104E.ipynb, and

DK0911v04_Pst104E_presence_absence.ipynb.

Ortholog analysis

We performed orthology analysis orthofinder/2.2.6 (Emms & Kelly 2018) of all non-redundant

protein sets with publicly available *Pucciniomycotina* genomes. Protein sets were downloaded

from MycoCosm (Grigoriev et al. 2014) (see Supplemental Data 5 for details). *Zymoseptoria*

tritici and *Verticillium dahliae* protein sets were also included as out-group.

We defined singletons between the two isolates as proteins that were in orthogroups that did not

contain a member from the other isolate. We performed enrichment analysis interrogating if

different gene categories are enriched for singletons over others. For this we used Fisher's

exact test and bonferroni correction executed in python (McKinney 2010). Detailed analysis can

be   found   in   jupyter   notebook   DK0911_vs_Pst104E_synteny.ipynb,

Pst104E_vs_DK0911_synteny.ipynb,

DK0911_vs_Pst104E_reciprocal_gene_pair_analysis.ipynb,   and

DK0911v04_Pst104E_presence_absence_orthology.ipynb.

Synteny analysis

We performed detailed synteny analysis to identify the best reciprocal gene pairs between the

two isolates. In addition, we investigated if certain gene categories are more syntenic than

others considering parameters like maximum synteny block size and number of allowed gaps

within a synteny block.

We identified best reciprocal gene pairs as follows. For each orthogroup we identified the

reciprocal gene pairs or groups belonging to each isolate. If there was more than one gene pair

candidate we identified the pair(s) for which each downstream and upstream gene was

conserved in terms of orthogroup identity as well. In the case of multiple candidate gene pairs

we assigned gene pair status to pairs that formed the longest synteny block when allowing for a

maximum synteny gap number of eight.

We identified synteny blocks using MCScanX (Wang et al. 2012) inspired by previous analysis

(Zhao & Schranz 2019). We ran MCScanX (Wang et al. 2012) and Diamond (Buchfink et al.

2015) within the Synnet.sh script with following parameters. MCScanX was run fixing the -w flag

to 0, the -s flag to 3, and varying the -m flag with the following values: 0, 1, 2, 3, 4, 5, 6, 7, 8, 9,

10, 15, 25, 35. These settings allow for the identification of synteny blocks involving at least

three genes with a varying number of allowed gaps within the synteny block. Diamond was run

with the following flags: --max-hsps 1 -k 25 -f 6 -b 5. Resulting *.collinearity files were parsed

and further analysed as described above. Detailed description of downstream analysis can be

found in in jupyter notebook DK0911_vs_Pst104E_reciprocal_gene_pair_analysis.ipynb and

DK0911v04_Pst104E_presence_absence_orthology.ipynb.

Gene pair variation analysis

We assessed the variation between the gene pairs of the two isolates using three approaches.

We calculated the Levenshtein distance based on the codon based CDS alignments and on the

protein alignments of each gene pair. We calculated the dN:dS ratios by using these two
alignment sets with yn00 paml version 4.9 (Yang & Nielsen 2000). The protein sequences of
each gene pair was aligned using muscle v3.8.31 (Edgar 2004), and codon-based alignments
were generated using PAL2NAL v14 (Suyama et al. 2006). The Levenshtein distance was
calculated in python using the distance module v0.1.3. Analysis details can be found in jupyter
notebook DK0911_vs_Pst104E_reciprocal_gene_pair_analysis.ipynb.

Telomere analysis
We estimated telomere length and telomere patterns using computel (Nersisyan & Arakelyan
2015) providing a chromosome number of 18 (-nchr) and the genome size (-lgenome) of the
primary assemblies.

Genome architecture analysis
We analyzed various features between certain gene families and in regards to TEs using
bedtools (Quinlan & Hall 2010) and pybedtools (Dale et al. 2011) as described previous
(Schwessinger et al. 2018). See jupyter notebook DK0911_v04_effector_analysis.ipynb.

Mitochondrial genome assembly and annotation
To assemble the Pst-DK0911 mitochondrial genome, we extracted mitochondrial reads from
whole genome PacBio reads of Pst-DK0911 by mapping them to the closely related *Phakopsora*
meibomiae (soybean rust pathogen) mt-genome (Stone et al. 2010) using bwa version 0.7.15-
r1140 (Li 2013). The reads were *de novo* assembled with Canu v1.6 (Koren et al. 2017). Canu
produced a single circular contig of ~124 kb. The circularity of the contig was confirmed by dot
plot using Gepard v-1.40 (Krumsiek et al. 2007). This produced a typical dot plot with identical
repeats at both ends, indicating a circular topology. The repeated sequences at the ends of the
contig were removed and the remainder circularized using circlator v-1.4.0 (Hunt et al. 2015).
The final circularized contig was of 101,813 bp. The contig was polished/error corrected with
two iterations of long reads using PacBio's GenomicConsensus Arrow (2019) and five iterations
of Illumina short reads using pilon v-1.22 (Walker et al. 2014). The polished mt-genome was
annotated with two independent mt-genome annotation tools, GeSeq web browser,
https://chlorobox.mpimp-golm.mpg.de/geseq.html (Tillich et al. 2017) using the *Phakopsora* mt-
genome as a reference and MITO web server using genetic code 4 http://mitos.bioinf.uni-
leipzig.de/index.py (Bernt et al. 2013). Annotations using these web browsers were checked
individually to confirm gene boundaries as well as intron-exon boundaries by aligning the

predicted genes with their orthologs in closely-related fungal species. Annotated tRNA genes

were further evaluated and compared using tRNAscan-SE v2.0.3 (Lowe & Eddy 1997) and

ARAGORN v1.2.38 (Laslett & Canback 2004). Introns were annotated using RNAweasel (Lang

et al. 2007).

Results

<u>A high quality partially haplotype phased genome for Pst-DK0911, a Warrior (PstS7) isolate</u>

We aimed to generate a haplotype-phased genome of the isolate Pst-DK0911 representing the

PstS7 lineage (Ali et al. 2017) generally known as "Warrior" (Hovmøller et al. 2016) using

PacBio long-reads (Supplemental Data 6, Table 1) assembled with FALCON-Unzip (Chin et al.

2016), which is designed to output haplotype phase blocks (haplotigs) for regions where

homologous chromosomes vary significantly. The primary genome assembly of Pst-DK0911 is

highly contiguous with 94 contigs and a contig N50 of 1.54 Mb, whereas the haplotigs are far

more fragmented consisting of 1176 contigs. The 94 primary contigs totalled 74 Mb in size and

the 1176 haplotigs constituted a size of 52 Mb (Table 1). Here haplotigs represent genome

regions where the two haploid genome copies are significantly different. The genome was

annotated based on new and previously published expression data from several different

developmental and infection stages of *P. striiformis* f. sp*. tritici* on wheat (Dobon et al. 2016;

Schwessinger et al. 2018). The Pst-DK0911 genome appears to be complete as measured by

the identification of benchmarking conserved single copy orthologs (BUSCOs, basidiomycota

odb9) (Simão et al. 2015). We could identify complete BUSCO hits for 97% (1290/1335) of the

benchmarking set with only 2% (25/1335) missing. In addition, the assembly has a low level of

duplicated BUSCOs (< 5%) indicating that it is completely phased and that primary contigs do

not contain extensive duplicated regions or unassigned haplotigs. This makes our Pst-DK0911

genome assembly comparable with previous long-read genome assemblies for *P. striiformis* f.

sp*. tritici* (Schwessinger et al. 2018; Xia, Wang, Yin, Cornejo, Scot H. Hulbert, et al. 2018). In

total we annotated 15,070 genes on primary contigs and 10,870 genes on haplotigs. The overall

gene length (~1550 bp on primary contigs and ~1420 bp on haplotigs) and average number of

introns per genes (~3.3 and ~3.1 for primary- and haplotigs, respectively) is similar between

primary contigs and haplotigs and near identical to our previous long-read assembly of Pst-

104E (Table 1) (Schwessinger et al. 2018).

Potential molecular functions of all protein coding genes were assessed by searches against

several databases (see Methods section). About 50% of the proteome contained at least one

functional domain (Supplemental Data 6, Table 2). We predicted candidate effectors with the
machine learning program EffectorP 2.0 (Sperschneider et al. 2018) applied to the secretome
identified with SignalP 3.0 (Bendtsen et al. 2004). We identified 819 candidate effectors on
primary contigs and 557 on haplotigs.

<u>A high quality mitochondrial genome for Pst-DK0911</u>
Using PacBio long-reads we were able to assemble a high-quality, circular mitochondrial
genome for Pst-DK0911 with a length of 101,813 bp (map of *P. striiformis* f. sp*. tritici* mt-
genome, Supplemental Figure 1). The Pst-DK0911 mt-genome contains 14 protein coding
genes that encode components of the electron transport chain, including three copies of the
ATP synthase complex (*atp6*, *atp8* and *atp9*), seven subunits of the nicotinamide adenine
dinucleotide ubiquinone oxidoreductase complex (*nad1*, *nad2*, *nad3*, *nad4* ,*nad4L*, *nad5* and
nad6), one cytochrome b (*cob*), and three subunits of cytochrome c oxidase (*cox1*, *cox2* and
cox3). In addition, the mt-genome encodes a ribosomal protein (*rps3*) which is a frequent
component of fungal mt-genomes (Korovesi et al. 2018; Losada et al. 2014), and a set of 24
tRNAs. All coding and non-coding genes are transcribed clockwise by the plus strand only. The
protein coding regions are separated by large (800-2,100 bp) regions containing abundant AT-
rich repeats.

<u>The difference in genome size between Pst-DK0911 and Pst-104E, representing two different *P.*</u>
<u>*striiformis* f. sp*. tritici* lineages, is explained by the difference in TE content and suggests a</u>
<u>recent TE burst</u>

The overall genome composition and coding capacity (Table 1) of Pst-DK0911 is very similar to
that of previous long-read assemblies (Schwessinger et al. 2018; Xia, Wang, Yin, Cornejo, Scot
H. Hulbert, et al. 2018). At the same time the cumulative size of the primary contigs of Pst-
DK0911 is 74 Mb whereas it is 83 Mb for Pst-104E. We investigated if this difference might be
linked to differences in transposable element (TE) content. When analysing repetitive elements,
approximately 56% of the Pst-DK0911 genome was annotated as encoding TEs, shared equally
between Class I and Class II elements (Supplemental Figure 2). In the case of Class I
retrotransposons, LTR Gypsy and Copia elements were by far the most abundant superfamilies.
TEs belonging to the TIR order were the most abundant Class II DNA transposons. The overall
distribution of TEs into superfamiles and their relative percentage identity compared to the
family consensus sequence in Pst-DK0911 was very similar to what have previously been

observed in Pst-104E (Schwessinger et al. 2018) (Supplemental Figure 2 and 3 A). Yet TE
genome coverage was distinct between the two genomes contributing to the difference in
primary contig size (approximation of genome size) between Pst-DK0911 (Table 1) and Pst-
104E (Schwessinger et al. 2018). TE content of Pst-DK0911 was 41 Mb when compared to 47
Mb in Pst-104E, which accounts for a 14% increase in TE content and for 54% of the observed
genome size difference between the two genomes. Indeed, there appears to be a specific
expansion of TEs in Pst-104E that is absent in Pst-DK0911 (Supplemental Figure 3 B). About
40% of the difference in TE coverage in Pst-104E relative to Pst-DK0911 is explained by an
expansion of TEs with an average sequence identity relative to the family consensus sequence
ranging from 82 to 95% (Supplemental Figure 3B). Here sequence conservation (identity) can
be used as a proxy for TE age (Maumus & Quesneville 2014), so more recent TE duplication
events display a greater sequence conservation than older TE copies, indicating that the
expansion of TEs in Pst104E has happened in recent history. It is also the relatively young TE
families that demonstrate the largest difference in coverage between Pst-104E and Pst-
DK0911, these belong mainly to the Class I LTR Gypsy and unclassified Class II elements
(Supplemental Figure 3 C and D).
498
499 <u>Comparatively low heterozygosity levels in Pst-DK0911 suggest long-term asexual reproduction</u>
500 <u>contributes positively to the accumulation of genetic variations</u>
501
In addition to a substantially smaller genome size, the Pst-DK0911 genome was phased to a
much lower level than Pst-104E, reaching levels of 70% phasing compared with over 90% for
Pst-104E (Schwessinger et al. 2018). This suggests that Pst-DK0911 is less heterozygous than
Pst-104E, because lower levels of heterozygosity prevent the phasing of haplotypes during the
assembly process (Chin et al. 2016). Hence, we tested heterozygosity levels in four
independent ways. First, we determined the number of heterozygous single nucleotide
polymorphisms (SNPs) when mapping Illumina short reads of the same isolate against its
primary contigs only. Here Pst-104E had about 7.1 SNPs per kb (~ 0.71% variation) which is
consistent with previous reports of isolates belonging to the PstS0 and PstS1/2 lineage
(Schwessinger et al. 2018; Xia, Wang, Yin, Cornejo, Scot H. Hulbert, et al. 2018; Cantu et al.
2013, 2011; Hubbard et al. 2015; Radhakrishnan et al. 2019; Bueno-Sancho et al.; Cuomo et al.
2016; Zheng et al. 2013). In contrast Pst-DK0911 had only 1.6 SNPs per kb (~ 0.16% variation),
which is dramatically less and consistent with our observation of reduced phasing in the Pst-
DK0911 genome assembly. Secondly, we made use of the partially phased assemblies to

estimate larger scale variations. When comparing Pst-DK0911 with Pst-104E we consistently
observed that the cumulative variation size in different size bins is always larger in Pst-104E
(Figure 1). In total, the larger scale structural variants covered 5.10 Mb (~ 6.39% variation) in
Pst-104E and 2.00 Mb (~ 2.66% variation) in Pst-DK0911. Third, we analyzed the normalized
read mapping coverage of primary contigs and haplotigs to estimate the level of heterozygosity
and of hemizygosity versus the level of collapsed regions in the two genome assemblies. We
normalized the read mapping coverage to one ('haploid coverage') using completely phased
regions of each genome as reference when mapping reads against both primary contigs and
haplotigs (Figure 2 B and D). When mapping against the primary contigs only (Figure 2A),
nearly all regions of the Pst-DK0911 genome displayed two times ('diploid') coverage,
suggesting a high level of similarity between the two haplotypes, such that reads from one
haplotype are able to map to the other haplotype. In contrast, Pst-104E also showed a peak at
1x coverage which suggests the presence of hemizygous regions in its genome. Indeed, when
mapping reads against primary contigs and haplotigs, Pst-104E showed a single strong
coverage peak for all regions analyzed (Figure 2 B-D) including regions in primary contigs that
lack an alternate haplotig. This supports the conclusion that Pst-104E contains hemizygous
regions in its genome that are significantly different between the two haploid nuclei. In contrast,
nearly all genome regions in Pst-DK0911 which lack a corresponding haplotig display 2x
'diploid' coverage indicating that these regions are collapsed in this genome assembly and
present in both haploid nuclei. Fourth, we analyzed the nucleotide variation of high confidence
gene pairs (alleles) within each assembly (primary contigs versus linked haplotigs). A total of
52% (7472/14321) of all possible high quality gene pairs in Pst-104E displayed variation on the
CDS level (Levenshtein distance greater than zero) compared to 40% (4344/10870) in Pst-
DK0911. The distribution of nucleotide variation of high quality gene pairs varied significantly
between Pst-104E and Pst-DK0911 (median: 0.0053 vs. 0.0012, Mann-Whitney U test, p-value
~ 3.13e-93) with the first quartiles being more extended in Pst-104E compared to Pst-DK0911,
while following quantiles are more extended in Pst-DK0911 (Figure 3). This suggests two
different potential sources of gene pair variation. Overall Pst-104E is clearly more heterozygous
than Pst-DK0911 in its genome architecture.
545

<u>Reduced telomere length in Pst-104E might be caused by long-term asexual reproduction</u>
It is well known that telomere ends shorten with increasing mitotic cell divisions due to the
incomplete replication of chromosome ends in mammalian cells (Aubert & Lansdorp 2008) but
this phenomenon is less studied in filamentous fungi (Wang et al. 2014). We tested if telomere

length is different in Pst-104E compared to Pst-DK0911. Our hypothesis was that if Pst-104E
has been asexual for a longer time period and hence undergone a greater number of
successive mitotic cell divisions, it should have shorter telomeres than Pst-DK0911. We
approximated telomere length via searching the canonical 'TTAGGG' telomere repeat and
variants thereof in whole genome Illumina short-read data (see Methods). We estimated 55
telomere repeats in Pst-DK0911 and 18 in Pst-104E, which suggests significant telomere
shortening in Pst-104E. Consistently, the canonical 'TTAGGG' repeat accounts only for 76% of
all patterns in Pst-104E versus 84% in Pst-DK0911 suggesting that Pst-104E has accumulated
more variations at its telomeres over time. Overall, this observation of relatively short telomeres
in Pst-104E compared to Pst-DK0911 is consistent with the increased heterozygosity in Pst-
104E.

<u>Inter-lineage genome variation is dominated by variation in transposable elements</u>
Next we tested for variation and presence-absence polymorphisms between the Pst-104E and
Pst-DK0911 genomes using a conservative coverage based read mapping approach. We
defined unique regions (highly variable or absent) in one genome if they fell below a 10%
normalized coverage threshold in cross-mapping (e.g. Pst-104E reads onto the Pst-DK0911
genome), but not self-mapping (e.g. Pst-104E reads on to the Pst-104E genome) experiments.
In total, we identified 6 Mb of sequence that was unique to Pst-DK0911 and 21 Mb to Pst-104E.
We then considered how many genes are completely contained in these unique regions. We
identified 724 variable genes that were in the Pst-DK0911 genome and not covered by Pst-
104E reads and hence absent in Pst-104E. Conversely, we identified 2394 variable genes of
Pst-104E that were not covered by Pst-DK0911 reads and hence missing from Pst-DK0911.
When analyzing if any specific gene group is enriched or depleted within this variable gene set,
we found that BUSCOs are significantly depleted in both cases (Multiple testing corrected
Fisher's exact test p-value ~ 3.1 e-11 for Pst-DK0911 and p-value ~ 8.2 e-9 for Pst-104E as
reference), which is expected for genes that belong to the highly conserved core genome. We
could not find any significant enrichment signal for candidate effectors or secreted proteins in
either Pst-104E or Pst-DK0911. On the contrary, genes encoding for secreted proteins
appeared to be depleted from the Pst-104E variable gene set (multiple testing corrected
Fisher's exact test p-value ~ 2.2 e-5).
We next tested if the unique regions were randomly distributed across the genomes or if they
contained a different number of genes than expected by chance. To do this, we performed a
permutation test by randomly reshuffling the unique regions to establish the expected

distribution of variable genes when the unique regions are randomly distributed. In both cases
BUSCO genes were significantly depleted from unique regions (Supplemental Figure 4). In the
case of Pst-DK0911, candidate effectors and genes encoding for secreted proteins were also
depleted from unique regions. In contrast, more genes than expected by chance were fully
contained within the unique genome regions of Pst-DK0911 and hence absent from Pst-104E.
The reverse analyses did not reveal any strong signal for any gene group except for genes
encoding for BUSCOs and secreted proteins which were significantly depleted from unique
regions in Pst-104E. Next we tested if TEs contribute to sequence variation between the two
genomes using a similar approach (see methods for details). Indeed, TEs were significantly
enriched in unique regions identified by our coverage based read mapping approach
(Supplemental Figure 5 A and B). As read mapping is known to suffer in repetitive regions, we
performed identical tests based on whole genome alignments allowing for multi-mapping with a
percentage identity coverage cut-off of 95%. Consistently, this second independent method also
identified that TEs are highly enriched in unique regions in both genomes (Supplemental Figure
6 A and B). Overall, these analyses suggest that sequence variation between the two isolates is
mostly caused by variation in TE sequences.

<u>Inter-lineage gene content variation does not favor candidate effectors</u>
Similar to previous observations in *P. striiformis* f. sp*. tritici* (Schwessinger et al. 2018; Xia,
Wang, Yin, Cornejo, Scot H. Hulbert, et al. 2018) and other rust fungi (Duplessis et al. 2011;
Miller et al. 2018), candidate effectors did not show any obvious genome compartmentalization
in Pst-DK0911 (Supplemental Figure 7). Candidate effectors were not located in gene sparse
regions or linked to TEs. Yet similar to Pst-104E, candidate effectors appeared to be slightly
closer to BUSCOs than to other genes (Supplemental Figure 7). This is not consistent with the
'two-speed' model of genome organisation (Dong et al. 2015; Frantzeskakis, Kusch, et al. 2019)
and indicates that rust fungi do not employ this mechanism to increase effector diversity within
lineages as effectors are not located in gene sparse regions close to TEs. Another leading
hypothesis in the field of plant pathology is that presence-absence polymorphisms within a
species is more common for candidate effectors than for other genes (Frantzeskakis et al.). We
tested this hypothesis by performing gene family (orthology) analysis with OrthoFinder (Emms &
Kelly 2018) using 59 fungal proteomes representing 38 fungal species with a focus on rust fungi
(see Supplemental Data 5 for details). Overall, candidate effectors did not appear to be
significantly enriched in the singleton gene pool of either isolate. When identifying genes without
orthologs in one of the two isolates, a total of 1275 and 1404 proteins were singletons in Pst-

DK0911 and in Pst-104E, respectively. As expected BUSCOs are depleted from the singleton

sets in both genomes (multiple testing error corrected Fisher's exact test p-value ~ 3.0 e-33 for

Pst-DK0911 and p-value ~ 5.3 e-31 for Pst-104E). When we investigated the EffectorP

predicted candidate effector subset in Pst-104E we saw a weak signal of enrichment in the

singleton gene set (Multiple testing error corrected Fisher's exact test p-value ~ 1.0 e-4). In

contrast, genes belonging to the secretome were depleted from the singleton gene set in Pst-

DK0911 (multiple testing error corrected Fisher's exact test p-value ~ 4.6 e-07).

<u>Candidate effectors fall into conserved and hypervariable categories</u>

While candidate effectors were not specifically enriched in presence-absence polymorphisms

between Pst-DK0911 and Pst-104E, we tested if some of the candidate effectors displayed

signs of positive selection. First, we investigated the variation between syntenic orthologs on the

nucleotide and amino acid level. About 44% and 57% of all effector candidates displayed

variation at the nucleotide level, which is slightly below the genome average of 65% in Pst-

DK0911 and 70% in Pst-104E, respectively. The variation between candidate effector genes

was higher than BUSCOs but similar to the overall gene pool (Figure 4 A, see Supplemental

Data 7-10 for statistical testing and gene category annotation). This trend was very similar at the

protein level with 36% and 52% of candidate effectors displaying variation in Pst-DK0911 and

Pst-104E, respectively. The variation in BUSCOs was significantly reduced compared to all

other gene groups, yet similar between all other gene groups. Lastly, we calculated the dN:dS

ratio for all syntenic ortholog pairs. The percentage of candidate effectors with a dN:dS ratio

greater than zero was lower than the genome average in both isolates (23% vs 35% in Pst-

DK0911 and 33% vs 42% in Pst-104E). At the same time the candidate effector population that

displayed greater than zero dN:dS ratio displayed a marked increase dN:dS ratios (Mann-

Whitney U test, all genes vs candidate effectors, p-value ~ 7.1 e-13 (Pst-104E), p-value ~ 2.2 e-

2 (Pst-DK0911)). Indeed, 51 and 133 candidate effectors displayed a dN:dS ratio greater than

one and hence a strong signal for positive selection in Pst-DK0911 and Pst-104E, respectively.

<u>Candidate effectors do not show strong signals for altered synteny</u>

While candidate effectors are not compartmentalized in rust fungi, we aimed to explore other

signals of specific genome organization in regard to candidate effectors in *P. striiformis* f. sp*.*

tritici. The availability of two highly contiguous partially phased genomes enabled us to ask the

question if candidate effectors displayed an altered synteny pattern similar to *R* genes in plants.

R genes in plants are the least syntenic genes in land plants (Zhao & Schranz 2019). If this

were the case for effectors, it could indicate that candidate effectors are in genomic regions with
higher recombination rates or general higher plasticity leading to disruption of gene synteny.
Using a similar approach as described for syntenic gene orthologs in plants (Zhao & Schranz
2019), we first tested how sensitive different gene groups are towards allowing a varying
number of gene gaps within synteny blocks using primary contigs as queries against the
complete assembly as subject. As expected (Zhao & Schranz 2019), BUSCOs were the least
affected by the number of gaps (Figure 5). The number of BUSCOs within a synteny block
plateaued at about 97% already at a maximum gap size of two or three in Pst-DK0911 or Pst-
104E, respectively (Figure 5 A). In contrast, for all other gene groups the percentage of genes
contained in synteny blocks kept on increasing until the maximum tested allowed gap size of 35.
Over 94% and 91% of all candidate effectors were contained within synteny blocks with a gap
size of 8 or more in Pst-DK0911 or Pst-104E, respectively. This is more than the genome
average of 88% and 86%, respectively. Hence overall candidate effectors do not appear to be
less syntenic than the genome average. This observation was further supported by the fact that
we observed very little difference in the mean best synteny block length at various numbers of
allowed gaps within the alignments (Figure 5 B). While BUSCOs appeared to have slightly
longer mean synteny block length, the variation of synteny block length was significant in each
gene group (see Supplemental Figure 8) not allowing for any generalization. Overall, these
observations are consistent with the fact that candidate effectors are also not enriched in inter-
lineage singletons and appear to be mostly surprisingly conserved at the gene level.
672

Discussion

Genomic insights into the genetic effects of sexual reproduction in *P. striiformis* f. sp*. tritici*
In the early 2010s, *P. striiformis* f. sp*. tritici* isolates from the 'Warrior' (PstS7) lineage arrived in
Europe severely impacting wheat production by causing stripe rust on previously resistant
wheat varieties (Hovmøller et al. 2016; Hubbard et al. 2015). The Pst-DK0911 isolate analyzed
in this manuscript belongs to the PstS7 lineage and was captured in Denmark in June 2011
(Hovmøller et al. 2016). The PstS7 lineage is genetically distinct from the pre-existent globally
important lineages such as PstS0 and PstS1/2, which are part of presumed asexual lineages at
least tracing back to isolates in Europe in the 1950s (Thach et al. 2016) and Africa in the 1980s
(Walter et al. 2016), respectively. Pst-DK0911 in contrast is closely related to sexual *P.*
striiformis f. sp*. tritici* populations in the Himalayan regions, which represent the center of
genetic diversity for *P. striiformis* f. sp*. tritici* (Ali et al. 2017; Ali, Gladieux, Leconte, et al. 2014).

Pst-DK0911 is one of very few reported *P. striiformis* f. sp*. tritici* isolates that has undergone the
complete sexual cycle under laboratory conditions (Rodriguez-Algaba et al. 2014). Self-crosses
of Pst-DK0911 suggest that this isolate is largely homozygous, in terms of having two
recognized alleles, for several recognized effector loci as none of the tested progeny isolates
were able to infect wheat differential lines that were resistant to the parental isolate. Hence, we
hypothesize that Pst-DK0911 originated from a highly inbred population that potentially might
have undergone several rounds of selfing or crosses between highly related genotypes leading
to relatively low heterozygosity. Indeed, our data supports this hypothesis as Pst-DK0911
displays the lowest heterozygosity at the SNP level of any *P. striiformis* f. sp*. tritici* isolate
analyzed so far. In addition, Pst-DK0911 displays about half the inter-haplotype structural
variation when compared to Pst-104E (Figure 1), which belongs to the long-term asexual
lineage PstS0. This observation is consistent with the insight gained from *Puccinia coronata* f.
sp. *avenae* where the genome of an isolate closely related to sexual populations displays
reduced heterozygosity by comparison to the genome of an isolate more closely related to long-
term asexual populations (Miller et al. 2018). It is likely that repeated sexual cycles of *P.*
striiformis f. sp*. tritici* lead to reduced heterozygosity as the physiology of sexual reproduction
favors inbreeding within one population. In addition, sexual recombination allows for gene
conversion and unequal crossover at homologous sides including transposable elements
(Underwood & Choi 2019; Liu et al. 2018). Interestingly, the variation analysis of high-quality
gene-pairs in Pst-DK0911 revealed that its genome harbors discontinuous populations of gene-
pairs. In general, variation in gene-pairs in dikaryotic rust fungi can be caused by two major
factors. The first would be continuous accumulation of mutations during asexual reproduction
cycles leading to linear accumulation of variation between gene-pairs. The second being
variation that has been introduced at a given point in time, e.g. mating of two genetically distinct
parents. The gene pair variation in Pst-DK0911 appears to be shaped by both sources of
variation and is best explained by an initial cross of genetically distinct parents, several rounds
of selfing of closely related individuals on barberry, followed by a limited number of asexual
reproduction cycles on wheat.
714
715   <u>The high quality mitochondrial genome of *P. striiformis* f. sp*. tritici* DK0911 reveals long low-</u>
716   <u>complexity AT regions</u>
717   The mitochondrial genome of Pst-DK0911 consists of extensive intergenic regions interspersed
718   with long low-complexity AT repeats. Comparative analysis of *P. striiformis* f. sp*. tritici* mt-
719   genome with rust and other fungi reveled that these repeats are unique to *P. striiformis* f. sp*.

tritici mt-genome. These regions are difficult to assemble with short Illumina reads alone but can be assembled with long PacBio reads that easily span the repeat regions, partially explaining the absence of a high-quality mitochondrial reference for *P. striiformis* f. sp. *tritici* so far. The overall GC content of the *P. striiformis* f. sp. *tritici* mt-genome is 31% rising to 35.8% in the protein-coding regions. This is similar to other fungal mt-genomes (Torriani et al. 2014; Stone et al. 2010), although the *P. striiformis* f. sp. *tritici* mt-genome is comparatively large. The size of fungal mt-genomes can vary by as much as 10-fold; for example, the smallest reported fungal mt-genome is just 20,063 bp for *Candida glabrata* (Koszul et al. 2003)and the largest 230,000 bp for *Rhizoctonia solani* (Losada et al. 2014). The genetic content of these mt-genomes is largely conserved and consists of 14 genes with the order of the genes being highly variable between different fungi. This variability in gene order is the highest in basidiomycete fungi (Aguileta et al. 2014). The gene order of *P. striiformis* f. sp. *tritici* mt-genome is similar to the close relative basidiomycete *Phakopsora sp.*, but the order of tRNA genes varies (Stone et al. 2010). Overall, this high-quality reference genome will enable the analysis of mitochondrial inheritance in *P. striiformis* f. sp. *tritici* in future studies.

The potential effects of long-term asexual reproduction on genome expansion, heterozygosity and fitness in *P. striiformis* f. sp. *tritici*

In contrast to Pst-DK0911 (Rodriguez-Algaba et al. 2014), isolates belonging to the long-term asexual lineages of PstS0 and PstS1/2 are often compromised in teliospore production on wheat (Ali et al. 2010), which is the entry point into the sexual infection cycle on barberry. It was hypothesized that this reduction is caused by counterselection of genes contributing to sexual reproduction in these two lineages given their likely exclusive asexual reproduction over multiple decades (Rodriguez-Algaba et al. 2014). Interestingly, our read cross-mapping presence-absence analysis identified a non-random distribution of genes located in highly variable regions of Pst-DK0911 compared to Pst-104E (Supplemental Figure 4). These and other highly variable genes might be involved in sexual reproduction in Pst-DK0911. In the future, it will be interesting to investigate expression patterns of these genes during teliospore production on wheat and during the sexual cycle of *P. striiformis* f. sp. *tritici* on barberry.

Overall, our comparative analysis suggests that long-term asexual reproduction in Pst-104E has led to a continuous accumulation of mutations, structural variations and a high level of heterozygosity between the two dikaryotic nuclei. The observed gene-pair variation pattern in Pst-104E is distinct compared to Pst-DK0911 and best explained by the continuous accumulation of mutations during asexual reproduction without other major events introducing

genetic diversity. Of course, we cannot exclude a somatic hybridization event in Pst-104E's
distant past similar to what has been recently suggested for *P. striiformis* (Lei et al. 2016a) and
clearly demonstrated at the whole genome level for the *Puccinia graminis* f.sp. *tritici* isolate
Ug99 (Li et al. 2019). In either case, the extensive relative telomere shortening in Pst-104E
does support the hypothesis of an extensive number of mitotic cell divisions without interruption
by meiosis as telomere shortening is an indication of aging in other organisms (Aubert &
Lansdorp 2008). Long-term asexual reproduction and independent accumulation of mutations in
both nuclei might have also impacted Pst-104E's overall fitness. This effect of accumulation of
mutations in asexual 'diploids' are often referred as the Meselson effect (Weir et al. 2016) and
might at least in part explain the rapid replacement of the PstS0 lineage by novel incursion e.g.
in Australia in the early 2000s (Wellings 2007).
Another striking fact is the relative genome size increase in Pst-104E which is partially (~54%)
explained by the expansion of the absolute TE content in this isolate. The contribution of TE
expansion to genome size increase is well documented for plants, animals, and fungi. The
overall TE family identity profile (TE 'age') is not different between Pst-104E and Pst-DK0911
(Supplemental Figure 2 A). At the same time, most of the TE content difference between the
two isolate coincides with the major TE burst at around 80% TE family identity (Supplemental
Figure 2 A and B). The difference in TE content could be either explained by selective purging
of TEs via unequal crossovers during meiosis (Shirleen Roeder 1983) in Pst-DK0911 in its
recent past during sexual reproduction cycles. This hypothesis is most parsimonious with the
observed data. Alternatively, the extended asexual reproduction cycles in Pst-104E might have
led to an increase in TE content in the absence of TE purging and continuous TE activity during
infection. The latter has been recently shown for *Zymoseptoria tritici* in which specific TE
families are expressed during the infection of wheat (Fouché et al. 2019). The upregulation of
specific TE families during infection was isolate specific, which could explain the difference of
relative abundance of some TE families over others. Indeed, TEs are the most variable genomic
regions between Pst-DK0911 and Pst-104E as measured at the superfamily level
(Supplemental Figure 4). In the future, it will be important to explore the expression of TEs in
several *P. striiformis* f. sp. *tritici* isolates during the infection of wheat. In addition, it would be
fascinating to perform whole genome comparative analysis of a time series of PstS0 isolates
available in the Stubbs collection ranging from the 1950s to the 1990s (collected by R.W.
Stubbs, Wageningen University) and later on expanded until today by the Global Rust
Reference Center (Thach et al. 2015). These analyses could directly reveal the impact of long-

term asexual reproduction on genome architecture, heterozygosity, and TE movement and
plasticity.
789
<u>Effector evolution in *P. striiformis* f. sp*. tritici* and beyond</u>
Clearly candidate effectors are not compartmentalized or linked to TEs in *P. striiformis* f. sp*.*
tritici (Supplemental Figure 6) (Schwessinger et al. 2018; Xia, Wang, Yin, Cornejo, Scot H.
Hulbert, et al. 2018). The availability of two haplotype-aware assemblies allowed us to test if
candidate effectors behave differently compared to other genes in terms of synteny (Figure 5).
Overall, candidate effectors did not behave strikingly differently from all genes (Figure 5). Yet
comparison between candidate effectors and secreted proteins revealed that fewer candidate
effectors, and especially EffectorP predicted candidates, are within a synteny block at any given
allowed gap-size (Figure 5A). This might suggest that these genes have a higher turn-over in
regions of the genome that have increased recombination rates. Future studies capturing more
of the genetic diversity of *P. striiformis* f. sp*. tritici* with haplotype aware assemblies will answer if
this trend of reduced synteny of candidate effectors holds true in general. Overall, candidate
effectors were well conserved between our two isolates and not enriched in presence absence
polymorphisms. Yet we could observe two distinct populations of candidate effectors. About
70% of all candidate effectors were invariable between the two isolates. These might be core
effectors required for plant infection of this obligate biotroph with a focus on wheat or not under
selection by the plant immune system within the experienced environments. At the same time
the remaining set of candidate effectors displayed an increased level of variation and dN:dS
ratios, which is likely caused by specific selection pressures exerted by the plant immune
system (Figure 4). This was especially visible in Pst-104E (Figure 4 A). This is consistent with
the fact that Pst-104E is more heterozygous with more distinct candidate effector genes
reducing the reciprocal mapping rate compared to Pst-DK0911. Overall the difference effector
repertoire between Pst-104E and Pst-DK0911 might explain the differences in virulence profiles
or aggressiveness observed between these two *P. striiformis* f. sp*. tritici* lineages (Ali et al.
2017).
815
<u>Evolution of wheat rust fungi</u>
Genome compartmentalization of effectors is absent in rusts (Duplessis et al. 2011; Miller et al.
2018; Schwessinger et al. 2018; Xia, Wang, Yin, Cornejo, Scot H. Hulbert, et al. 2018) and
indeed in most other obligate biotrophic plant pathogens including mildews of wheat and barley
(Frantzeskakis, Németh, et al. 2019; Müller et al. 2019; Fletcher et al. 2019; Frantzeskakis et al.

2018). This poses the question of what other genetic mechanisms lead to variation in candidate
effector complements within and between *P. striiformis* f. sp*. tritici* lineages. It has been shown
recently that candidate effectors are enriched in recombination hotspots in *Zymoseptoria tritici*
(Grandaubert et al. 2019) and *Blumeria graminis* f. sp*. tritici* (Müller et al. 2019). Similarly,
recombination within sexual populations in the Himalayan region could lead to reshuffling of
candidate effector loci in *P. striiformis* f. sp*. tritici.* This might bring together the most suitable
candidate effectors for infection of wheat lines currently grown on large acreages. Indeed, we
hypothesis that *P. striiformis* f. sp*. tritici* undergoes a two-step selection process. First,
candidate effector allele combinations must be suited to the infection of modern wheat varieties
in contrast to strains only adapted to local barberry populations and the proximal grasses, which
are likely regular hosts of *P. striiformis* f. sp*. tritici* within the region. Second, once a *P. striiformis*
f. sp*. tritici* lineage is adapted to modern wheat varieties it must overcome the specific
resistance genes within currently grown wheat varieties (Ali et al. 2017). This second step of
adaptation is driven by genetic variation introduced during the asexual reproduction cycle of
wheat. The mutations in recognized effectors can be caused by SNPs, small insertions and
deletions, larger structural variations and TE movement. Because both nuclei evolve
independently during asexual reproduction, recognized effectors that are heterozygous to begin
with are highly likely to be lost during this process. This effect is illustrated by the two cloned
recognized effectors of *P. graminis* f. sp*. tritici*, AvrSr35 and AvrSr50, which were both
heterozygous in the isolates they were identified in (Chen et al. 2017; Salcedo et al. 2017).

A third genetic mechanism of introducing variation is hybridization between distinct species or
lineages (Stukenbrock 2016). Hybridization leads to novel allele combinations and previously
heterozygous recessive alleles to be unearthed. Interspecies hybridization is important for the
emergence of new plant pathogens as demonstrated for *Blumeria graminis* f. sp. *triticalae*,
which arose from a cross between *B. graminis* f. sp. *tritici* and *B. graminis* f. sp. *secali* (Menardo
et al. 2016). A special form of hybridization is the swapping of whole nuclei between different
dikaryotic rust isolates (Park & Wellings 2012). This process is referred to as somatic
hybridization and has long been speculated upon based on low resolution genetic markers or
novel virulence profiles emerging during co-infection of asexual hosts (Park & Wellings 2012;
Lei et al. 2016b). Yet only very recently, somatic hybridization has been conclusively
demonstrated to be important for wheat rust evolution under natural conditions. A recent study
used completely haplotype phased assemblies to show that the hypervirulent *P. graminis* f. sp.
tritici Ug99 isolate is a product of nuclei swapping between isolates of an old South African and
a poorly described Iranian *P. graminis* f. sp*. tritici* lineage (Li et al. 2019). It is likely that similar

somatic hybridization events occur in *P. striiformis* f. sp*. tritici.* Somatic hybridization of *P.*
striiformis f. sp*. tritici* circumvents the first adaptation process on wheat as each haploid nucleus
is likely already adapted to wheat. In addition, it can rapidly generate novel effector allele
combinations and drive rapid *R* gene adaptation by introducing heterozygosity to the effector
gene profile. In addition, the impact of somatic hybridization on generating hypervirulent wheat
rust strains increases with the number of genetically distinct lineages within a given location.
This furthers the biosecurity risk of higher rates of novel pathotype incursions in wheat growing
areas globally.

Conclusions
The future of wheat rust and *P. striiformis* genomics lies in generating complete haplotype
phased genomes that assort each set of chromosomes into specific karyotypes. This will enable
us to predict the potential of generating novel allele combinations during the asexual
reproduction cycle via mutations and somatic hybridization. In addition, haplotype phased
genomes will facilitate the cloning of recognized effectors. In combination, this will enable us to
better detect specific recognized effector allele pair combinations using modern long-read DNA
sequence approaches and enable us to better model the durability of certain *R* gene stacking
approaches. Lastly, it will be important to survey *P. striiformis* populations in the Himalayan
regions, which are not adapted to modern wheat varieties but to locally growing wheat and
grasses. This will provide insight into the true genetic diversity of *P. striiformis* and enable us to
catalog the entire genetic diversity of effectors. In addition, we might identify effectors that are
important for the infection of wheat or those that are negatively selected for by modern wheat
varieties.

Acknowledgments
The research on Pst-DK0911 conducted at Aarhus University was funded by the Danish
Innovation Fund (grant no. 11-116241, RUSTFIGHT). This work was supported by a Australian
Research Council DECRA (DE150101897) and Future Fellowship (FT180100024) to BS.

Data availability
All raw sequencing data and the assembled genomes can be found on NCBI under the
Bioproject accession number PRJNA588102. Additional, annotation files and axillary genome
files can be found on zenodo 10.5281/zenodo.3556840. Supplementary data files can be found
on zenodo 10.5281/zenodo.3556860. Supplementary figures can be found on zenodo

10.5281/zenodo.3556929. All code related to this project can be found on github

https://github.com/Team-Schwessinger/DK_0911-1 and has a zenodo doi of

10.5281/zenodo.3556925.

Authors contribution

The experimental work on Pst-DK0911 was made by Y-J.C. and C.K.S. under the supervision of

A.F.J and M.S.H. Preliminary bioinformatic analyses of short read Illumina genomic and

transcriptomic data for Pst-DK0911 was made by Y-J.C and J.K.V. Provision of a genetically

pure stock of Pst-DK0911 and virulence phenotyping of Pst-DK0911 was made by M.S.H. B.S.

conducted all bioinformatic analysis except for the mitochondrial genome assembly. R. T.

contributed to bioinformatic analysis. R. N. analyzed the mitochondrial genome. All authors

contributed intellectually to this study. B.S wrote the manuscript with contributions from Y-J.C,

A.F.J, R. N. and J. P. R. All authors comment on drafts of this manuscript and approved the final

submission.

904
905
References

Aguileta G et al. 2014. High Variability of Mitochondrial Gene Order among Fungi. Genome Biol
Evol. 6:451–465. doi: 10.1093/gbe/evu028.

Ali S et al. 2016. CloNcaSe: Estimation of sex frequency and effective population size by
clonemate re-sampling in partially clonal organisms. Mol Ecol Resour. n/a-n/a. doi:
10.1111/1755-0998.12511.

Ali S, Gladieux P, Rahman H, et al. 2014. Inferring the contribution of sexual reproduction,
migration and off-season survival to the temporal maintenance of microbial populations: a case
study on the wheat fungal pathogen Puccinia striiformis f.sp. tritici. Mol Ecol. 23:603–617. doi:
10.1111/mec.12629.

Ali S, Gladieux P, Leconte M, et al. 2014. Origin, Migration Routes and Worldwide Population
Genetic Structure of the Wheat Yellow Rust Pathogen Puccinia striiformis f.sp. tritici. PLoS
Pathog. 10:e1003903. doi: 10.1371/journal.ppat.1003903.

Ali S et al. 2017. Yellow Rust Epidemics Worldwide Were Caused by Pathogen Races from
Divergent Genetic Lineages. Front. Plant Sci. 8. doi: 10.3389/fpls.2017.01057.

Ali S, Leconte M, Walker A-S, Enjalbert J, de Vallavieille-Pope C. 2010. Reduction in the sex
ability of worldwide clonal populations of Puccinia striiformis f.sp. tritici. Fungal Genetics and
Biology. 47:828–838. doi: 10.1016/j.fgb.2010.07.002.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool.
J. Mol. Biol. 215:403–410. doi: 10.1016/S0022-2836(05)80360-2.

Anikster Y. 1986. Teliospore Germination in Some Rust Fungi. Phytopathology. 76:1026. doi:
10.1094/Phyto-76-1026.

Aubert G, Lansdorp PM. 2008. Telomeres and aging. Physiol. Rev. 88:557–579. doi:
10.1152/physrev.00026.2007.

Bairoch A, Apweiler R. 2000. The SWISS-PROT protein sequence database and its supplement
TrEMBL in 2000. Nucleic Acids Res. 28:45–48.

Bao W, Kojima KK, Kohany O. 2015. Repbase Update, a database of repetitive elements in
eukaryotic genomes. Mobile DNA. 6:11. doi: 10.1186/s13100-015-0041-9.

Beddow JM et al. 2015. Research investment implications of shifts in the global geography of
wheat stripe rust. Nature Plants. 15132. doi: 10.1038/nplants.2015.132.

Bendtsen JD, Nielsen H, von Heijne G, Brunak S. 2004. Improved prediction of signal peptides:
SignalP 3.0. J. Mol. Biol. 340:783–795. doi: 10.1016/j.jmb.2004.05.028.

Bernt M et al. 2013. MITOS: Improved de novo metazoan mitochondrial genome annotation.
Molecular Phylogenetics and Evolution. 69:313–319. doi: 10.1016/j.ympev.2012.08.023.

Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using DIAMOND.
Nature Methods. 12:59–60. doi: 10.1038/nmeth.3176.

Bueno-Sancho V et al. Pathogenomic analysis of wheat yellow rust lineages detects seasonal
variation and host specificity. Genome Biol Evol. doi: 10.1093/gbe/evx241.

Cantu D et al. 2013. Genome analyses of the wheat yellow (stripe) rust pathogen Puccinia
striiformis f. sp. tritici reveal polymorphic and haustorial expressed secreted proteins as
candidate effectors. BMC Genomics. 14:270. doi: 10.1186/1471-2164-14-270.

Cantu D et al. 2011. Next generation sequencing provides rapid access to the genome of
Puccinia striiformis f. sp. tritici, the causal agent of wheat stripe rust. PLoS ONE. 6:e24230. doi:
10.1371/journal.pone.0024230.

Chen J et al. 2017. Loss of AvrSr50 by somatic exchange in stem rust leads to virulence for
Sr50 resistance in wheat. Science. 358:1607–1610. doi: 10.1126/science.aao4810.

Chen W, Wellings C, Chen X, Kang Z, Liu T. 2014. Wheat stripe (yellow) rust caused by
Puccinia striiformis f. sp. tritici. Molecular Plant Pathology. 15:433–446. doi:
10.1111/mpp.12116.

Chin C-S et al. 2016. Phased diploid genome assembly with single-molecule real-time
sequencing. Nat Meth. 13:1050–1054. doi: 10.1038/nmeth.4035.

Cuomo CA et al. 2016. Comparative Analysis Highlights Variable Genome Content of Wheat
Rusts and Divergence of the Mating Loci. G3: Genes, Genomes, Genetics. g3.116.032797. doi:
10.1534/g3.116.032797.

Dale RK, Pedersen BS, Quinlan AR. 2011. Pybedtools: a flexible Python library for manipulating
genomic datasets and annotations. Bioinformatics. 27:3423–3424. doi:
10.1093/bioinformatics/btr539.

Dobon A, Bunting DCE, Cabrera-Quio LE, Uauy C, Saunders DGO. 2016. The host-pathogen
interaction between wheat and yellow rust induces temporally coordinated waves of gene
expression. BMC Genomics. 17:380. doi: 10.1186/s12864-016-2684-4.

Dong S, Raffaele S, Kamoun S. 2015. The two-speed genomes of filamentous pathogens: waltz
with plants. Curr. Opin. Genet. Dev. 35:57–65. doi: 10.1016/j.gde.2015.09.001.

Duan X et al. 2010. Puccinia striiformis f.sp. tritici presents high diversity and recombination in
the over-summering zone of Gansu, China. Mycologia. 102:44–53. doi: 10.3852/08-098.

Duplessis S et al. 2011. Obligate biotrophy features unraveled by the genomic analysis of rust
fungi. PNAS. 108:9166–9171. doi: 10.1073/pnas.1019315108.

Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high
throughput. Nucleic Acids Res. 32:1792–1797. doi: 10.1093/nar/gkh340.

Ellis JG, Lagudah ES, Spielmeyer W, Dodds PN. 2014. The past, present and future of breeding
rust resistant wheat. Front. Plant Sci. 5:641. doi: 10.3389/fpls.2014.00641.

Emms DM, Kelly S. 2018. OrthoFinder2: fast and accurate phylogenomic orthology analysis
from gene sequences. bioRxiv. 466201. doi: 10.1101/466201.

Fletcher K et al. 2019. Genomic signatures of heterokaryosis in the oomycete pathogen Bremia
lactucae. Nature Communications. 10:2645. doi: 10.1038/s41467-019-10550-0.

Flutre T, Duprat E, Feuillet C, Quesneville H. 2011. Considering Transposable Element
Diversification in De Novo Annotation Approaches. PLOS ONE. 6:e16526. doi:
10.1371/journal.pone.0016526.

Fouché S et al. Stress-Driven Transposable Element De-repression Dynamics and Virulence
Evolution in a Fungal Pathogen. Mol Biol Evol. doi: 10.1093/molbev/msz216.

Frantzeskakis L et al. Rapid evolution in plant–microbe interactions – a molecular genomics
perspective. New Phytologist. 0. doi: 10.1111/nph.15966.

Frantzeskakis L et al. 2018. Signatures of host specialization and a recent transposable element
burst in the dynamic one-speed genome of the fungal barley powdery mildew pathogen. BMC
Genomics. 19:381. doi: 10.1186/s12864-018-4750-6.

Frantzeskakis L, Németh MZ, et al. 2019. The Parauncinula polyspora Draft Genome Provides
Insights into Patterns of Gene Erosion and Genome Expansion in Powdery Mildew Fungi. mBio.
10. doi: 10.1128/mBio.01692-19.

Frantzeskakis L, Kusch S, Panstruga R. 2019. The need for speed: compartmentalized genome
evolution in filamentous phytopathogens. Molecular Plant Pathology. 20:3–7. doi:
10.1111/mpp.12738.

Garrison E, Marth G. 2012. Haplotype-based variant detection from short-read sequencing.
arXiv:1207.3907 [q-bio]. http://arxiv.org/abs/1207.3907 (Accessed September 15, 2017).

Gladieux P et al. 2014. Fungal evolutionary genomics provides insight into the mechanisms of
adaptive divergence in eukaryotes. Molecular Ecology. 23:753–773. doi: 10.1111/mec.12631.

Grabherr MG et al. 2011. Trinity: reconstructing a full-length transcriptome without a genome
from RNA-Seq data. Nat Biotechnol. 29:644–652. doi: 10.1038/nbt.1883.

Grandaubert J, Dutheil JY, Stukenbrock EH. 2019. The genomic determinants of adaptive
evolution in a fungal pathogen. Evolution Letters. 3:299–312. doi: 10.1002/evl3.117.

Grigoriev IV et al. 2014. MycoCosm portal: gearing up for 1000 fungal genomes. Nucleic Acids
Res. 42:D699-704. doi: 10.1093/nar/gkt1183.

Haas BJ et al. 2008. Automated eukaryotic gene structure annotation using EVidenceModeler
and the Program to Assemble Spliced Alignments. Genome Biology. 9:R7. doi: 10.1186/gb-
2008-9-1-r7.

Heitman J. 2015. Evolution of sexual reproduction: A view from the fungal kingdom supports an
evolutionary epoch with sex before sexes. Fungal Biology Reviews. 29:108–117. doi:
10.1016/j.fbr.2015.08.002.

Hoff KJ, Lange S, Lomsadze A, Borodovsky M, Stanke M. 2016. BRAKER1: Unsupervised
RNA-Seq-Based Genome Annotation with GeneMark-ET and AUGUSTUS. Bioinformatics.
32:767–769. doi: 10.1093/bioinformatics/btv661.

Hovmøller MS et al. 2016. Replacement of the European wheat yellow rust population by new
races from the centre of diversity in the near-Himalayan region. Plant Pathol. 65:402–411. doi:
10.1111/ppa.12433.

Hovmøller MS, Justesen AF. 2007. Rates of evolution of avirulence phenotypes and DNA
markers in a northwest European population of Puccinia striiformis f. sp. tritici. Molecular
Ecology. 16:4637–4647. doi: 10.1111/j.1365-294X.2007.03513.x.

Hovmøller MS, Rodriguez-Algaba J, Thach T, Sørensen CK. 2017. Race Typing of Puccinia
striiformis on Wheat. In: Wheat Rust Diseases: Methods and Protocols. Periyannan, S, editor.
Methods in Molecular Biology Springer: New York, NY pp. 29–40. doi: 10.1007/978-1-4939-
7249-4_3.

Hovmøller MS, Yahyaoui AH, Milus EA, Justesen AF. 2008. Rapid global spread of two
aggressive strains of a wheat rust fungus. Molecular Ecology. 17:3818–3826. doi:
10.1111/j.1365-294X.2008.03886.x.

Hubbard A et al. 2015. Field pathogenomics reveals the emergence of a diverse wheat yellow
rust population. Genome Biology. 16:23. doi: 10.1186/s13059-015-0590-8.

Huerta-Cepas J et al. 2016. eggNOG 4.5: a hierarchical orthology framework with improved
functional annotations for eukaryotic, prokaryotic and viral sequences. Nucleic Acids Res.
44:D286–D293. doi: 10.1093/nar/gkv1248.

Hunt M et al. 2015. Circlator: automated circularization of genome assemblies using long
sequencing reads. Genome Biology. 16:294. doi: 10.1186/s13059-015-0849-0.

Jin Y, Szabo LJ, Carson M. 2010. Century-Old Mystery of Puccinia striiformis Life History
Solved with the Identification of Berberis as an Alternate Host. Phytopathology. 100:432–435.
doi: 10.1094/PHYTO-100-5-0432.

Jones P et al. 2014. InterProScan 5: genome-scale protein function classification.
Bioinformatics. 30:1236–1240. doi: 10.1093/bioinformatics/btu031.

Kim D, Langmead B, Salzberg SL. 2015. HISAT: a fast spliced aligner with low memory
requirements. Nat Meth. 12:357–360. doi: 10.1038/nmeth.3317.

Koren S et al. 2017. Canu: scalable and accurate long-read assembly via adaptive k-mer
weighting and repeat separation. Genome Res. gr.215087.116. doi: 10.1101/gr.215087.116.

Korovesi AG, Ntertilis M, Kouvelis VN. 2018. Mt-rps3 is an ancient gene which provides insight
into the evolution of fungal mitochondrial genomes. Molecular Phylogenetics and Evolution.
127:74–86. doi: 10.1016/j.ympev.2018.04.037.

Koszul R et al. 2003. The complete mitochondrial genome sequence of the pathogenic yeast
Candida (Torulopsis) glabrata. FEBS Letters. 534:39–48. doi: 10.1016/S0014-5793(02)03749-3.

Krumsiek J, Arnold R, Rattei T. 2007. Gepard: a rapid and sensitive tool for creating dotplots on
genome scale. Bioinformatics. 23:1026–1028. doi: 10.1093/bioinformatics/btm039.

Kurtz S et al. 2004. Versatile and open software for comparing large genomes. Genome Biol.
5:R12. doi: 10.1186/gb-2004-5-2-r12.

Lang BF, Laforest M-J, Burger G. 2007. Mitochondrial introns: a critical view. Trends in
Genetics. 23:119–125. doi: 10.1016/j.tig.2007.01.006.

Laslett D, Canback B. 2004. ARAGORN, a program to detect tRNA genes and tmRNA genes in
nucleotide sequences. Nucleic Acids Res. 32:11–16. doi: 10.1093/nar/gkh152.

Lechner M et al. 2011. Proteinortho: Detection of (Co-)orthologs in large-scale analysis. BMC
Bioinformatics. 12:124. doi: 10.1186/1471-2105-12-124.

Lei Y et al. 2016a. Virulence and Molecular Characterization of Experimental Isolates of the
Stripe Rust Pathogen (Puccinia striiformis) Indicate Somatic Recombination. Phytopathology™.
107:329–344. doi: 10.1094/PHYTO-07-16-0261-R.

Lei Y et al. 2016b. Virulence and Molecular Characterization of Experimental Isolates of the
Stripe Rust Pathogen (Puccinia striiformis) Indicate Somatic Recombination. Phytopathology.
107:329–344. doi: 10.1094/PHYTO-07-16-0261-R.

Li F et al. 2019. Emergence of the Ug99 lineage of the wheat stem rust pathogen through
somatic hybridisation. bioRxiv. 692640. doi: 10.1101/692640.

Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.
arXiv:1303.3997 [q-bio]. http://arxiv.org/abs/1303.3997 (Accessed September 14, 2017).

Liu H et al. 2018. Tetrad analysis in plants and fungi finds large differences in gene conversion
rates but no GC bias. Nat Ecol Evol. 2:164–173. doi: 10.1038/s41559-017-0372-7.

Lorrain C, Santos KCG dos, Germain H, Hecker A, Duplessis S. 2019. Advances in
understanding obligate biotrophy in rust fungi. New Phytologist. 222:1190–1206. doi:
10.1111/nph.15641.

Losada L et al. 2014. Mobile elements and mitochondrial genome expansion in the soil fungus
and potato pathogen Rhizoctonia solani AG-3. FEMS Microbiol Lett. 352:165–173. doi:
10.1111/1574-6968.12387.

Lowe TM, Eddy SR. 1997. tRNAscan-SE: A Program for Improved Detection of Transfer RNA
Genes in Genomic Sequence. Nucleic Acids Res. 25:955–964. doi: 10.1093/nar/25.5.955.

Markell SG, Milus EA. 2008. Emergence of a Novel Population of Puccinia striiformis f. sp. tritici
in Eastern United States. Phytopathology. 98:632–639. doi: 10.1094/PHYTO-98-6-0632.

Maumus F, Quesneville H. 2014. Ancestral repeats have shaped epigenome and genome
composition for millions of years in *Arabidopsis thaliana*. Nature Communications. 5:4104. doi:
10.1038/ncomms5104.

Mboup M et al. 2012. Genetic structure and local adaptation of European wheat yellow rust
populations: the role of temperature-specific adaptation. Evolutionary Applications. 5:341–352.
doi: 10.1111/j.1752-4571.2011.00228.x.

McKinney W. 2010. Data Structures for Statistical Computing in Python. In: pp. 51–56.
http://conference.scipy.org/proceedings/scipy2010/mckinney.html (Accessed September 15,
2017).

Menardo F et al. 2016. Hybridization of powdery mildew strains gives rise to pathogens on novel
agricultural crop species. Nat Genet. 48:201–205. doi: 10.1038/ng.3485.

Miller ME et al. 2018. De Novo Assembly and Phasing of Dikaryotic Genomes from Two
Isolates of Puccinia coronata f. sp. avenae, the Causal Agent of Oat Crown Rust. mBio.
9:e01650-17. doi: 10.1128/mBio.01650-17.

Müller MC et al. 2019. A chromosome-scale genome assembly reveals a highly dynamic
effector repertoire of wheat powdery mildew. New Phytologist. 221:2176–2189. doi:
10.1111/nph.15529.

Nattestad M, Schatz MC. 2016. Assemblytics: a web analytics tool for the detection of variants
from an assembly. Bioinformatics. 32:3021–3023. doi: 10.1093/bioinformatics/btw369.

Nersisyan L, Arakelyan A. 2015. Computel: Computation of Mean Telomere Length from Whole-
Genome Next-Generation Sequencing Data. PLoS One. 10. doi: 10.1371/journal.pone.0125201.

Park RF, Wellings CR. 2012. Somatic Hybridization in the Uredinales. Annual Review of
Phytopathology. 50:219–239. doi: 10.1146/annurev-phyto-072910-095405.

Pertea M et al. 2015. StringTie enables improved reconstruction of a transcriptome from RNA-
seq reads. Nat Biotech. 33:290–295. doi: 10.1038/nbt.3122.

Petersen TN, Brunak S, von Heijne G, Nielsen H. 2011. SignalP 4.0: discriminating signal
peptides from transmembrane regions. Nat Meth. 8:785–786. doi: 10.1038/nmeth.1701.

Quesneville H et al. 2005. Combined Evidence Annotation of Transposable Elements in
Genome Sequences. PLOS Computational Biology. 1:e22. doi: 10.1371/journal.pcbi.0010022.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. 26:841–842. doi: 10.1093/bioinformatics/btq033.

Radhakrishnan GV et al. 2019. MARPLE, a point-of-care, strain-level disease diagnostics and
surveillance tool for complex fungal pathogens. BMC Biology. 17:65. doi: 10.1186/s12915-019-
0684-y.

Rapilly F. 1979. Yellow Rust Epidemiology. Annu. Rev. Phytopathol. 17:59–73. doi:
10.1146/annurev.py.17.090179.000423.

Rawlings ND, Barrett AJ, Finn R. 2016. Twenty years of the MEROPS database of proteolytic
enzymes, their substrates and inhibitors. Nucleic Acids Res. 44:D343–D350. doi:
10.1093/nar/gkv1118.

Rodriguez-Algaba J, Walter S, Sørensen CK, Hovmøller MS, Justesen AF. 2014. Sexual
structures and recombination of the wheat rust fungus Puccinia striiformis on Berberis vulgaris.
Fungal Genetics and Biology. 70:77–85. doi: 10.1016/j.fgb.2014.07.005.

Salcedo A et al. 2017. Variation in the AvrSr35 gene determines Sr35 resistance against wheat
stem rust race Ug99. Science. 358:1604–1606. doi: 10.1126/science.aao7294.

Savary S et al. 2019. The global burden of pathogens and pests on major food crops. Nature
Ecology & Evolution. 1. doi: 10.1038/s41559-018-0793-y.

Schwessinger B et al. 2018. A Near-Complete Haplotype-Phased Genome of the Dikaryotic
Wheat Stripe Rust Fungus Puccinia striiformis f. sp. tritici Reveals High Interhaplotype Diversity.
mBio. 9:e02275-17. doi: 10.1128/mBio.02275-17.

Schwessinger B. 2017. Fundamental wheat stripe rust research in the 21(st) century. New
Phytol. 213:1625–1631. doi: 10.1111/nph.14159.

Schwessinger B, Rathjen JP. 2017. Extraction of High Molecular Weight DNA from Fungal Rust
Spores for Long Read Sequencing. In: Wheat Rust Diseases.Methods in Molecular Biology
Humana Press, New York, NY pp. 49–57. doi: 10.1007/978-1-4939-7249-4_5.

Seudre O et al. 2018. Why outcross? The abandon-ship hypothesis in a facultative
outcrossing/selfing fungal species. Fungal Genetics and Biology. 120:1–8. doi:
10.1016/j.fgb.2018.08.005.

Shirleen Roeder G. 1983. Unequal crossing-over between yeast transposable elements. Mol
Gen Genet. 190:117–121. doi: 10.1007/BF00330332.

Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. 2015. BUSCO:
assessing genome assembly and annotation completeness with single-copy orthologs.
Bioinformatics. 31:3210–3212. doi: 10.1093/bioinformatics/btv351.

Singh RP et al. 2016. Disease Impact on Wheat Yield Potential and Prospects of Genetic
Control. Annual Review of Phytopathology. 54:303–322. doi: 10.1146/annurev-phyto-080615-
095835.

Sørensen CK, Hovmøller MS, Leconte M, Dedryver F, de Vallavieille-Pope C. 2014. New Races
of Puccinia striiformis Found in Europe Reveal Race Specificity of Long-Term Effective Adult
Plant Resistance in Wheat. Phytopathology. 104:1042–1051. doi: 10.1094/PHYTO-12-13-0337-
R.

Sørensen CK, Thach T, Hovmøller MS. 2016. Evaluation of Spray and Point Inoculation
Methods for the Phenotyping of Puccinia striiformis on Wheat. Plant Disease. PDIS-12-15-1477-
RE. doi: 10.1094/PDIS-12-15-1477-RE.

Sperschneider J et al. 2015. EffectorP: predicting fungal effector proteins from secretomes
using machine learning. New Phytol. n/a-n/a. doi: 10.1111/nph.13794.

Sperschneider J, Dodds PN, Gardiner DM, Singh KB, Taylor JM. 2018. Improved prediction of
fungal effector proteins from secretomes with EffectorP 2.0. Mol. Plant Pathol. 19:2094–2110.
doi: 10.1111/mpp.12682.

Steele KA, Humphreys E, Wellings CR, Dickinson MJ. 2001. Support for a stepwise mutation
model for pathogen evolution in Australasian Puccinia striiformis f.sp. tritici by use of molecular
markers. Plant Pathology. 50:174–180. doi: 10.1046/j.1365-3059.2001.00558.x.

Stone CL, Buitrago MLP, Boore JL, Frederick RD. 2010. Analysis of the complete mitochondrial
genome sequences of the soybean rust pathogens Phakopsora pachyrhizi and P. meibomiae.
Mycologia. 102:887–897. doi: 10.3852/09-198.

Stukenbrock EH. 2016. The Role of Hybridization in the Evolution and Emergence of New
Fungal Plant Pathogens. Phytopathology. PHYTO-08-15-0184-RVW. doi: 10.1094/PHYTO-08-
15-0184-RVW.

Suyama M, Torrents D, Bork P. 2006. PAL2NAL: robust conversion of protein sequence
alignments into the corresponding codon alignments. Nucleic Acids Res. 34:W609–W612. doi:
10.1093/nar/gkl315.

Taylor JW, Hann-Soden C, Branco S, Sylvain I, Ellison CE. 2015. Clonal reproduction in fungi.
PNAS. 112:8901–8908. doi: 10.1073/pnas.1503159112.

Testa AC, Hane JK, Ellwood SR, Oliver RP. 2015. CodingQuarry: highly accurate hidden
Markov model gene prediction in fungal genomes using RNA-seq transcripts. BMC Genomics.
16:170. doi: 10.1186/s12864-015-1344-4.

Thach T, Ali S, Justesen A f., Rodriguez-Algaba J, Hovmøller M s. 2015. Recovery and
virulence phenotyping of the historic 'Stubbs collection' of the yellow rust fungus Puccinia
striiformis from wheat. Ann Appl Biol. 167:314–326. doi: 10.1111/aab.12227.

Thach T, Ali S, de Vallavieille-Pope C, Justesen AF, Hovmøller MS. 2016. Worldwide population
structure of the wheat rust fungus Puccinia striiformis in the past. Fungal Genetics and Biology.
87:1–8. doi: 10.1016/j.fgb.2015.12.014.

Tillich M et al. 2017. GeSeq – versatile and accurate annotation of organelle genomes. Nucleic
Acids Res. 45:W6–W11. doi: 10.1093/nar/gkx391.

Torriani SFF et al. 2014. Comparative analysis of mitochondrial genomes from closely related
Rhynchosporium species reveals extensive intron invasion. Fungal Genetics and Biology.
62:34–42. doi: 10.1016/j.fgb.2013.11.001.

Underwood CJ, Choi K. 2019. Heterogeneous transposable elements as silencers, enhancers
and targets of meiotic recombination. Chromosoma. 128:279–296. doi: 10.1007/s00412-019-
00718-4.

Walker BJ et al. 2014. Pilon: An Integrated Tool for Comprehensive Microbial Variant Detection
and Genome Assembly Improvement. PLOS ONE. 9:e112963. doi:
10.1371/journal.pone.0112963.

Walter S et al. 2016. Molecular markers for tracking the origin and worldwide distribution of
invasive strains of Puccinia striiformis. Ecol Evol. n/a-n/a. doi: 10.1002/ece3.2069.

Wang N et al. 2014. Novel Telomere-Anchored PCR Approach for Studying Sexual Stage
Telomeres in Aspergillus nidulans. PLOS ONE. 9:e99491. doi: 10.1371/journal.pone.0099491.

Wang Y et al. 2012. MCScanX: a toolkit for detection and evolutionary analysis of gene synteny
and collinearity. Nucleic Acids Res. 40:e49–e49. doi: 10.1093/nar/gkr1293.

Weir W et al. 2016. Population genomics reveals the origin and asexual evolution of human
infective trypanosomes. eLife Sciences. 5:e11473. doi: 10.7554/eLife.11473.

Wellings CR. 2011. Global status of stripe rust: a review of historical and current threats.
Euphytica. 179:129–141. doi: 10.1007/s10681-011-0360-y.

Wellings CR. 2007. Puccinia striiformis in Australia: a review of the incursion, evolution, and
adaptation of stripe rust in the period 1979–2006. Aust. J. Agric. Res. 58:567–575.

Xia C, Wang M, Yin C, Cornejo OE, Hulbert Scot, et al. 2018. Genome sequence resources for
the wheat stripe rust pathogen (Puccinia striiformis f. sp. tritici) and the barley stripe rust
pathogen (Puccinia striiformis f. sp. hordei). MPMI. doi: 10.1094/MPMI-04-18-0107-A.

Xia C, Wang M, Yin C, Cornejo OE, Hulbert Scot H., et al. 2018. Genomic insights into host
adaptation between the wheat stripe rust pathogen (Puccinia striiformis f. sp. tritici) and the
barley stripe rust pathogen (Puccinia striiformis f. sp. hordei). BMC Genomics. 19:664. doi:
10.1186/s12864-018-5041-y.

Yang Z, Nielsen R. 2000. Estimating synonymous and nonsynonymous substitution rates under
realistic evolutionary models. Mol. Biol. Evol. 17:32–43.

Yin Y et al. 2012. dbCAN: a web resource for automated carbohydrate-active enzyme
annotation. Nucleic Acids Res. 40:W445-451. doi: 10.1093/nar/gks479.

Zeyl C. 2009. The role of sex in fungal evolution. Current Opinion in Microbiology. 12:592–598.
doi: 10.1016/j.mib.2009.09.011.

Zhao T, Schranz ME. 2019. Network-based microsynteny analysis identifies major differences
and genomic outliers in mammalian and angiosperm genomes. PNAS. 201801757. doi:
10.1073/pnas.1801757116.

Zheng W et al. 2013. High genome heterozygosity and endemic genetic recombination in the
wheat stripe rust fungus. Nat Commun. 4:2673. doi: 10.1038/ncomms3673.

2019. *PacificBiosciences/GenomicConsensus.* Pacific Biosciences
https://github.com/PacificBiosciences/GenomicConsensus (Accessed November 4, 2019).

2017a. *rtg-tools: RTG Tools: Utilities for accurate VCF comparison and manipulation.* Real Time
Genomics https://github.com/RealTimeGenomics/rtg-tools (Accessed September 15, 2017).

2017b. *vcflib: a simple C++ library for parsing and manipulating VCF files, + many command-*
line utilities. vcflib https://github.com/vcflib/vcflib (Accessed September 15, 2017).

1236 **Tables:**
1237

	Primary assembly		Haplotype assembly
	Primary contigs with haplotigs	Primary contigs without haplotigs	Haplotigs
Number of contigs	86	8	1176
Bases	73,945,211	482,406	52,126,201
TEs Coverage	56.02%	80.90%	51.80%
Number of genes	15003	67	10870
Average gene length	1553	1032	1419
Average introns per gene	3.31	1.73	3.12
Number of genes per 10kb	2.02	1.39	2.08

	BUSCOs	1290 (S: 1184, D: 50, F: 43)	2 (S: 2)	886 (S: 727, D: 77, F: 56)
Number of BUSCOs per 10kb		0.17	0.04	0.17
Number of candidate effectors[a]		816	3	557
Number of candidate effectors[a] per 10kb		0.11	0.06	0.11

Table 1: Summary of Pst-DK0911 genome assembly and annotation

Summary statistics for the genome assembly according to the three different contig categories as described in the main text.

For the BUSCO annotation we used the following abbreviattions. S single copy, D duplicated, F fragmented.

[a]Candidate effectors have been predicted based on the machine learning algorithm EffectorP2 only.

In the BUSCO column the abbreviations are as follows: S single-copy, D duplicated, F fragmented

Figure legends

Figure 1: Reduced inter haplotype structural variations in Pst-DK0911

Summary of interhaplotype variation between primary contigs and their respective haplotigs, analyzed using Assemblytics. Each plot indicates the number of bases that are spanned by the specific variation category. Pst-DK0911 displays less variation then Pst-104E in all categories.

Figure 2: Low level of hemizygous regions in Pst-DK0911

Normalized Illumina short-read coverage plots for the color-coded (Pst-104E blue and Pst-DK-0911 brown) genome regions in primary contigs when mapping against primary contigs only (A) or when mapping against primary contigs and haplotigs (B, D, and F). The same of haplotigs in panel C. Coverage was normalized to one in regions of the genome that are fully phased. The Pst-DK0911 contains significantly less homozygous regions when compared to Pst-104E as it only has a single 2x coverage peak when mapping against primary contigs only (A). In addition, nearly all regions in primary contigs that do not possess a corresponding haplotig have 2x coverage suggesting that these regions are collapsed regions in the assembly and not hemizygous (E).

Figure 3: Different level of allelic gene pair variation in Pst-DK0911

Boxenplot of levenshstein edit distance of coding regions of high-quality allelic gene pairs in Pst-104E and Pst-DK0911. For values greater zero, n[Pst-104E]=7472 and n[Pst-DK0911]=4344.

Figure 4: A subset of effector candidates displays patterns of positive selection

Violinplots of the pairwise comparison of syntenic orthologs in Pst-104E and Pst-DK0911 split according to gene classifications. (A) Shows the Levenshtein distance at the coding sequence level (CDS). (B) Shows the Levenshtein distance at the protein level. (C) Shows the dN:dS ratio. n indicates number of orthologs genes or proteins with any difference.

max indicates the total number of orthologous genes or proteins

% indicates the percentage of orthologs genes or proteins with any variation greater zero.

See Supplemental Data 7 and 8 for detailed statistical analysis of the complete data set.

Figure 5: Candidate effectors are mostly syntenic between the two wheat rust isolates

(A) Shows the percentage of genes that are syntenic at increasing number of allowed gaps within the synteny block. (B) Shows the mean synteny block length at increasing number of allowed gaps within the synteny block.

The minimum synteny block size is three. Data points only include genes on the primary contig of the query genome. Data points represent different gene classifications.

Supplemental Figure 1: Map of the Pst-DK0911 mt-genome

The outer ring represents genes (coding and non-coding) and their respective positions in the genome. The yellow colour represents genes of nicotinamide adenine dinucleotide ubiquinone oxidoreductase complex, the pink colour represents genes of cytochrome c oxidase complex, the light green colour represents cytochrome b gene, dark green represents ATP synthase genes, brown represents ribosomal protein gene, navy blue represents tRNAs genes and red represents rRNAs genes. The inner ring representing GC content. The gaps in the inner ring representing AT abundant repeat.

Supplemental Figure 2: *P. striiformis* f. sp*. tritici* characteristic transposable element distribution in Pst-DK0911

Repetitive element annotation on primary contigs (A) and on haplotigs (B). Top panels show percentages of genome coverage for all repetitive elements and different subcategories. These include TEs of class I (RNA retrotransposons) and class II (DNA transposons), simple sequence repeats (SSR), and unclassifiable repeats (no Cat). Middle and bottom panels show percentages of genome coverage of class I and class II TEs categorized to class, order, and superfamily levels wherever possible. Repetitive elements were identified using the REPET pipeline, and classifications were inferred from the closest BLAST hit (see Materials and Methods). The TE order is color coded in each ClassI and ClassII TE plot.

Supplemental Figure 3: Specific TE expansion in Pst-104E absent from Pst-DK0911

(A) Mean identity distribution relative to the consensus sequence of TE families as a proxy of TE age in Pst-104E and Pst-DK0911. (B) Relative difference in TE coverage accumulation (0-1) in Pst-104E compared to Pst-DK0911 versus TE identity distribution as proxy of TE age. Genome coverage per TE superfamily elements in Pst-DK0911 (C) and Pst-104E (D) with a minimum mean identity of 80% and more than 50 copies. These TEs are most likely to contribute to current genome evolution. Bars are color-coded according to the order level.

Supplemental Figure 4: Short read cross-mapping based presence-absence polymorphisms between Pst-104E and Pst-DK0911 is not specifically enriched for candidate effectors

Expected (violinplots) and observed (red line) number of gene completely contained in low coverage regions in cross-mapping experiments for Pst-DK0911 (A) or Pst-104E (B) as reference. Expected distributions were generated by randomly reshuffling the observed low coverage regions 5000 times and measuring the number of fully contained genes in the random low coverage regions. P-values are two-sided and calculated by comparing the observed number of contained genes to the expected distribution (see methods for details).

Supplemental Figure 5: Short read cross-mapping based variation and presence-absence polymorphisms between Pst-DK0911 and Pst-104E is enriched for TEs

Expected (violinplots) and observed (red line) number of bases pair regions of TEs contained in low coverage regions in cross-mapping experiments for Pst-DK0911 (A) or Pst-104E (B) as reference. Expected distributions were generated by randomly reshuffling the observed low coverage regions 2500 times and measuring the number of fully contained genes in the random low coverage regions. P-values are two-sided and calculated by comparing the observed number of contained genes to the expected distribution (see methods for details). Literally, all TEs are significantly enriched for being contained in regions highly variable between the two *P. striiformis* f. sp*. tritici* strains.

Supplemental Figure 6: Whole genome alignment-based variation and presence-absence polymorphisms between Pst-DK0911 and Pst-104E is enriched for TEs

Expected (violinplots) and observed (red line) number of bases pair regions of TEs contained in low coverage regions in whole genome alignment experiments for Pst-DK0911 (A) or Pst-104E (B) as reference. Expected distributions were generated by randomly reshuffling the observed low coverage regions 2500 times and measuring the number of fully contained genes in the random low coverage regions. P-values are two-sided and calculated by comparing the observed number of contained genes to the expected distribution (see methods for details). Literally, all TEs are significantly enriched for being contained in regions highly variable between the two *P. striiformis* f. sp*. tritici* strains.

Supplemental Figure 7: Candidate effector are not found in gene spares regions and are not linked to TEs in Pst-DK0911

(A) Nearest-neighbor gene distance density hexplots for three gene categories, including all genes, BUSCOs, and candidate effectors. Each subplot represents a distance density hexplot with the log10 3'-flanking and 5'-flanking distance to the nearest-neighboring gene plotted along the x axis and y axis, respectively. (B) Violin plots for the log10 distance to the most proximal transposable element for genes in each category without allowing for overlap. (C) Violin plots for the log10 distance to the most proximal gene in the same category for subsamples of each category equivalent to the smallest category size (n = 814). (D) Violin plots for the minimum distance (log10) of candidate effectors and BUSCOs to each other or a random subset of genes (n = 814). The P values for panels B, C, and D were calculated using the Wilcoxon rank-sum

1359   test after correction for multiple testing (Bonferroni; alpha = 0.05) on the linear distance in
1360   bases.
1361
Supplemental Figure 8: Candidate effectors synteny block size distribution reflects the
genome average.
Violinplots of the mean synteny block length in (A) Pst-104E and (B) Pst-DK0911 and split
according to gene classifications. The maximum allowed gap number is eight. Minimum synteny
block length is three.
n indicates numbers of orthologs within a synteny block.
max indicates the total number of orthologs.
% indicates the percentage of orthologs within a synteny block.

Self mapping