Mapping Human Pluripotent Stem Cell Derived Erythroid Differentiation by Single-Cell Transcriptome Analysis

Zijuan Xin [1,2,#,a], Wei Zhang[1,2,#,b], Shangjin Gong[1,2,c], Junwei Zhu[1,d], Yanming Li[1,e], Zhaojun Zhang[1,3,4,*,f], Xiangdong Fang [1,2,3,4,*,g]

[1] *CAS Key Laboratory of Genome Science and Information, Beijing Institute of Genomics, Chinese Academy of Sciences, Beijing 100101, China*

[2] *College of Life Sciences, University of Chinese Academy of Sciences, Beijing 100049, China*

[3] *Sino-Danish College, University of Chinese Academy of Sciences, Beijing 100190, China*

[4] *Institute for Stem Cell and Regeneration, Chinese Academy of Sciences, Beijing 100101, China*

[#] Equal contribution.

[*] Corresponding authors.

E-mail: fangxd@big.ac.cn (Fang X), zhangzhaojun@big.ac.cn (Zhang Z)

Running title: *Xin Z and Zhang W / scRNA-seq Analysis of iPSC-derived RBC differentiation system*

[a] ORCID: 0000-0001-9418-4848.

[b] ORCID: 0000-0002-0519-3935.

[c] ORCID: 0000-0002-9811-5302.

[d] ORCID: 0000-0001-9766-3334.

[e] ORCID: 0000-0002-8213-9166.

[f] ORCID: 0000-0003-0490-6507.

[g] ORCID: 0000-0002-6628-8620.

Total word counts: 6,288

Total figures: 6

Total supplementary figures: 7

Total supplementary tables: 7

Abstract

There is currently an imbalance between the supply and demand of functional red

blood cells (RBCs) in clinical applications, and this imbalance can be addressed by

regenerating RBCs with a variety of in vitro methods. Induced pluripotent stem cells

(iPSCs) can address the low supply of cord blood and the ethical issues in embryonic

stem cell research and provide a promising strategy to eliminate immune rejection.

However, no complete single-cell level differentiation pathway exists for the

iPSC-derived RBC differentiation system. In this study, we used iPSC line BC1 to

establish an RBC regeneration system and used the 10× Genomics single-cell

transcriptome platform to map the cell lineage and differentiation trajectories on day

14 (D14) of the regeneration system. We found iPSC differentiation was not

synchronized during embryoid body (EB) culture, and the D14 cells in the system

mainly consisted of mesodermal and various blood cells, similar to yolk sac

hematopoiesis. During asynchronous EB differentiation, iPSCs undergo three

bifurcations before they enter erythroid differentiation, and the driver genes of each

bifurcation were identified. The key roles of cell adhesion and estradiol in RBC

regeneration were observed.   This study provides systematically theoretical guidance

for the optimization of the iPSC-derived RBC differentiation system.

KEYWORDS: scRNA-seq; iPSCs; Hematopoiesis; Erythropoiesis; Differentiation

trajectory

Introduction

The substantial gap between the supply and demand of blood has always been a serious problem in clinical practice [1]. Artificial blood is an important means of solving this worldwide shortage, and obtaining functional red blood cells (RBCs) is the key to artificial blood generation [2]. RBC regeneration *in vitro* has been an important research direction in the blood research field for many years [3, 4]. In the current study, the primitive materials that can be used for RBC regeneration are mainly cord blood hematopoietic stem cells (HSCs) [5], embryonic stem cells (ESCs) [6, 7], and induced pluripotent stem cells (iPSCs) [6, 8]. The regeneration technology of cord blood HSCs is relatively mature, but there are problems that come with this technology, such as difficulty in obtaining materials, significant differences among individuals, high price of this technology, *etc.*, and ESCs are subject to ethical restrictions [9]. Human iPSCs can be differentiated into a variety of cell types *in vitro*, providing a model for basic research and a source of clinically relevant cells [10-12]. Although the derivation of iPSC lines for patients has now become a routine technique, how to accurately differentiate iPSCs into the desired cell types is still a major challenge [13, 14].

Decades of research have shown that *in vitro* hematopoietic differentiation is closely related to in vivo development [15]. Hematopoiesis originates from the FLK1+ (KDR) lateral mesoderm, and some of the specialized cells can be induced to form vascular hemangioblasts by the transcription factor (TF) ETV2 [16, 17]. These cells have the ability to differentiate into blood and vascular precursor cells, during which ETV2 is regulated by the BMP and WNT signaling pathways [18, 19]. At this time, the primitive RBCs that express embryonic globin (ε-globin; HBE) are produced, which is the first wave of hematopoiesis in the yolk sac [20]. Hemangioblasts differentiate into endothelial cells and specialize in hematopoietic endothelial cells that produce erythroid-myeloid progenitors (EMPs). EMPs can differentiate into most of the myeloid and into definitive RBCs that can simultaneously express embryonic (ε-globin; HBE), fetal (γ-globin, HBG1 and HBG2) and adult (β-globin, HBB) globin;

this differentiation represents the second wave of hematopoiesis in the yolk sac [21].

HSCs can also be directly produced through endothelial-hematopoietic transition

(EHT) induced by GATA2 and RUNX1 [22-24]. This occurs in the

aorta-gonad-mesonephros (AGM) region that maintains lifelong hematopoietic

differentiation and the hematopoietic cycle [25], and these cells can differentiate into

definitive RBCs that express only adult globin and are regulated by signaling pathways

such as VEGF and HIF [26, 27]. Hematopoietic differentiation is the differentiation

process of HSCs to form various types of blood cells [28]. After differentiation and

activation of HSCs, it is necessary to go through the cell fate determination program

regulated by GATA1, KLF1 and other TFs to enter the erythroid differentiation

process [29]. The generation of RBCs is a complex multistep process involving the

differentiation of early erythroid progenitor cells into definitive RBCs, which require

spatiotemporal specific completion of globin synthesis and assembly [30, 31], iron

metabolism [32, 33], heme synthesis [34], cell denucleation and other processes,

eventually forming intact functional RBCs [35]. However, the molecular and cellular

mechanisms involved in these processes are still poorly understood.

Based on the existing molecular mechanisms of hematopoietic development and

erythroid differentiation, scientists simulate embryonic hematopoiesis [36] used the

spin EB method to regenerate HSPCs *in vitro*, and then induced the differentiation of

RBCs from iPSCs. However, the differentiation efficiency and denucleation rate of the

existing system is low, and the expression of adult globin is limited; these drawbacks

limit the clinical availability [37]. At the same time, iPSC differentiation is a complex

process, and flow cytometry and immunostaining have been used to determine the cell

types during iPSC differentiation culture. However, these methods are limited by the

number of fluorescent probes used, and it is not possible to solve important problems,

such as the cell composition and differentiation path of the iPSC differentiation system,

under high-resolution conditions. Recently, single-cell transcriptome technology has

been able to efficiently capture and identify rare and transient cell types, determine the

spatial or temporal localization of cells, and reconstitute gene regulatory networks,

helping scientists understand the mechanisms by which development and cell fate are

determined [38]. In recent years, there have been many single-cell hematopoiesis

studies *in vitro*. By single-cell sequencing of embryoid bodies, researchers found that

naïve H9 ESCs have a stronger hematopoietic capacity than primer H9 ESCs [39] and

illuminated the heterogeneity of pluripotent stem cell-derived endothelial cell

differentiation [40]. Single-cell sequencing of D29 erythrocytes belonging to the

iPSC-derived erythroid differentiation system revealed that cells expressing β-globin

showed reduced transcripts encoding ribosomal proteins and increased expression of

ubiquitin-proteasome system members [41]. The lack of a complete single-cell

transcriptome map as iPSCs differentiate into RBCs does not allow scientists to

systematically guide iPSCs to efficiently produce functional RBCs. Therefore, in this

study, we established an iPSC-derived erythroid differentiation system and obtained a

dynamic transcriptional map of cell differentiation through high-resolution single-cell

transcriptomics sequencing. The cell map of the *in vitro* iPSC-derived RBC

differentiation system and the *in vitro* differentiation trajectory of the intact iPSCs to

RBCs were mapped for the first time. iPSCs undergo three bifurcations before they

enter erythroid differentiation: the bifurcation between hematopoiesis and

angiogenesis, the bifurcation between myeloid and lymphoid lineage, and the

bifurcation between granulocyte-mononuclear and megakaryocyte-erythroid lineage.

Cells undergo two transformations: during transformation one, cells actively express

cell adhesion molecules, while during transformation two, cells downregulate these

adhesion molecules. During erythroid differentiation, estradiol promotes erythroid

progenitor cell proliferation and inhibits erythroid differentiation. This study has

important guiding significance for analyzing the basic fate of cells and the gene

network structure, as well as for designing more effective hematopoietic and

erythropoiesis strategies for regenerative medicine.

Results

iPSC cell line-derived HSPC production and erythropoiesis

Depending on the spin EB culture method, we used the iPSC cell line BC1 to collect the suspension cells (SCs) through the cell strainer in the system after 14 days of culture [42, 43] (**Figure 1**A). The established culture system can effectively produce SCs. The cell count analysis showed that the number of SCs on day 14 (D14) was approximately 10-fold higher than the number of iPSCs on D0 (Figure 1B). Flow cytometry analysis showed that approximately 25.9±3.05% of the SCs expressed CD34, and approximately 27.08±1.63% of the SCs expressed CD45, indicating that we collected a certain percentage of HSPCs from the SCs (Figure 1C). We performed colony-forming unit (CFU) assays on the collected SCs on D14. After 14 days of culture, blood lineage colonies were observed under the microscope, indicating that the SCs have the potential to differentiate into various blood lineage cells; and the CFU-GEMM colony showed a slight rust red color, indicating that the erythroid cell colonies begin to express hemoglobin (Figure 1D). Subsequently, we attempted to induce HSPCs in SCs for erythroid differentiation. By assessing the changes in the expression of the cell surface markers CD71 and CD235a at different time points (Figure 1E), we found that on D14 of the EB stage, the proportion of CD71 and CD235a double-positive cells in SCs exceeded 30%, indicating that some cells in the system entered the erythroid differentiation stage. On D28, more than 95% of cells expressed both CD71 and CD235a (Figure 1E). Through Wright-Giemsa staining analysis, we observed that most of the cells were at the end of erythroid differentiation, and some cells were denucleated RBCs. In summary, we successfully obtained HSPCs by spin EB technology used the iPSC cell line BC1 and induced HSPCs to enter the erythroid differentiation process, eventually producing denucleated RBCs.

Analysis of the cell composition of the iPSC-derived RBC differentiation system

In this system, D14 is the intermediate time point. At this time, the sample contains both EBs and SCs, and the SCs contain a certain proportion of CD34+, CD45+ and CD71+ CD235a+ cells, including various cell types; in addition, the cell number is

large and the differentiation system is stable. Therefore, to establish the cell lineage

and developmental trajectory of the iPSC-derived RBC differentiation system, we

selected EB cells (EBCs) and SCs on D14 of culture and sequenced them using a 10×

Genomics platform. After sequencing and data processing, we detected 470,711,092

reads and 4,441 cells, with an average of 105,992 reads per cell; these reads

corresponded to 5,888 unique molecular identifier (UMI) counts and 1,644 genes.

After quality control analysis of the data by Seurat [44], we finally obtained 3,215

cells with high-quality transcripts, including 704 SCs and 2,511 EBCs (Figure S1A).

The t-SNE analysis at the sample level revealed that the cells were mainly divided

into upper and lower parts. EBCs were mainly distributed in the upper part, and the

corresponding highly expressed genes were mainly related to mesoderm

differentiation. SCs were mainly distributed in the lower part, and the corresponding

highly expressed genes mainly regulate hematopoietic differentiation and immune

function (**Figure 2**A, S1B and C). We also performed bulk RNA-seq on SCs cultured

at D14 and the iPSC line BC1. After comparing them with published HSPC

transcriptome data from human cord blood, we found that SCs are significantly

different from iPSCs, and the hematopoietic lineage-related gene expression patterns

of which are similar to HSPCs and exhibit HSPC-like characteristics (Figure S1D and

E). Subsequently, the intersection of the top 2,000 genes in EBCs and SCs was

selected for further principal component analysis (PCA) and t-SNE analysis. The

results showed that the differentiation process of cells was not synchronized. Seurat

analysis divided our samples into ten clusters, including three progenitor cells clusters

(cluster 9 mainly include pluripotent stem cells and high gene expression of *ZFP42*;

cluster 1, 2 mainly include mesoderm cells and high gene expression of *PDGFRA*);

CD33 cells cluster (cluster 0, high gene expression of *MNDA* and *S100A8*);

mesenchymal cells (cluster 3, high gene expression of *LUM* and *DCN*); HSPCs

(cluster 4, high gene expression of *RUNX1* and *MYB*); hemangioblasts (cluster 5, high

gene expression of *KDR* and *CD34*) [45]; vascular endothelial cells (VECs) (cluster 6,

high gene expression of *MALAT1* and *WSB1*); megakaryocyte and erythroid cells

(MEs) (cluster 7, high gene expression of *GATA1* and *HBA1*) [46]; and CD14 cells

(cluster 8, high gene expression of *LYZ* and *CD14*) [47] (Figure 2B, C and S2C). The

CD33 cells cluster had the largest number of cells, 695, which accounted for 22% of

the total cells in the sample, and the progenitor cells-9 cluster had the fewest cells,

only 115, accounting for 4%, indicating that the D14 cells in the system almost

entered the differentiation process. Combined with the sample distribution results,

SCs were mainly distributed in blood cell clusters, such as clusters 0, 4, 7, and 8, and

EBCs were scattered and distributed in almost every cluster (Figure S2B); however,

the MEs cluster was almost entirely made of SCs, which may be because the RBCs

had low adhesion and were more easily released from the EBs (Figure 2A and B).

To further determine the function of the cell cluster, we performed functional

annotations based on the top 50 genes expressed in each cluster. Gene annotation

analysis revealed that progenitor cells-9 cluster genes were particularly involved in

the function of embryonic morphogenesis ($P = 1.7E\text{-}05$), and the results also indicated

that the progenitor cells-9 cluster had the differentiation potential to form each germ

layer cell, such as kidney, podosoma, epithelial cells and sensorium. The progenitor

cells-1 cluster was characterized by gastrula cells, and highly expressed genes were

enriched in gastrulation ($P = 6.01E\text{-}03$). Gene annotation analysis results of the

progenitor cells-1 cluster showed that these cells respond to growth factor stimulation

and highly expressed the genes related to organ growth and tissue morphogenesis.

Genes that were highly expressed in the progenitor cells-2 cluster were enriched in

mesoderm formation ($P = 1.55E\text{-}05$), which was related to cardiac development and

muscle formation, indicating that the progenitor cell-2 cluster was lateral

mesoderm-like mesoderm. Genes that were highly expressed in the mesenchymal

cells cluster were enriched in the connective tissue development pathway ($P =$

$1.40E\text{-}06$). Hemangioblasts cluster had the characteristics of hematopoietic

endothelium (HE), and highly expressed genes were enriched in the pathways of

endothelial development ($P = 1.02E\text{-}08$) and endothelial cell migration ($P =$

$2.04E\text{-}04$), suggesting that the hemangioblasts cluster simultaneously undergoes EHT

and highly expresses cell adhesion-related genes. Hemangioblasts further

differentiated into VECs and HSPCs. Genes that were highly expressed in VECs were

mainly enriched in the function of blood vessel development (P = 3.94E-03). Genes

that were highly expressed in HSPCs were enriched in the pathways of lymphocyte

differentiation (P = 3.94E-03), erythrocyte differentiation (P =9.58E-03), and myeloid

cell differentiation (P = 1.49E-03), suggesting that HSPCs have the potential to

differentiate into myeloid, lymphoid and erythroid cells. Genes that were highly

expressed in the CD33 cells, MEs and CD14 cells clusters were obviously enriched in

the pathways of granulocyte activation (P = 5.32E-35), oxygen transport (P =

2.18E-08), and phagocytosis (P = 9.33E-05), respectively, and these clusters all

differentiated from HSPCs (Figure S3).

A higher resolution clustering of the scRNA-seq data

To further determine the process of cell differentiation, we performed a higher

resolution clustering of the scRNA-seq data, and the results showed that clusters 0, 1,

4, and 8 had subclusters. Among them, the CD33 cells cluster had three subclusters,

which were recorded as 0a, 0b, and 0c. Compared with clusters 0b and 0c, cluster 0a

had increased *CD14* and *CTSS* gene expression, indicating that cluster 0a was a

monocyte-like cells cluster [47]. Compared with clusters 0a and 0c, cluster 0b had

increased *ELANE* and *MPO* gene expression, suggesting that cluster 0b was a

granulocyte-macrophage progenitor (GMP)-like cells cluster. Compared with clusters

0a and 0b, cluster 0c had increased *NFC1* and *DEFA3* gene expression and was a

granulocyte-like cells cluster (**Figure 3**A and B). Cluster 1 had two subclusters that

were marked as clusters 1a and 1b. Cluster 1a had high expression levels of the *PITX2*

and *PRRX1* genes, indicating that this cluster contained mesodermal cells; compared

with cluster 1a, cluster 1b showed more stem cell characteristics with high expression

of the *PDGFRB* and *ID3* genes [48, 49] (Figure 3A and C). The HSPCs cluster was

divided into 2 subclusters that were marked as clusters 4a and 4b. Cluster 4a was

characterized as a common myeloid progenitor (CMP)-like cluster and had the

potential for myeloid cell differentiation and had high *PRG3* and *PRG2* gene

expression; cluster 4b was characterized as a myeloid-lymphoid progenitor

(MLP)-like cluster with high *EGR1* and *TRDC* gene expression [50] and was

enriched in the pathways related to lymphocyte differentiation [50]. In addition, the

CD14 cells cluster was also divided into two clusters, namely, the macrophages

cluster (cluster 8a, high gene expression of *CD36* and *CD163*) and the dendritic cells

cluster (cluster 8b, high gene expression of *S100A8*) [51] (Figure S4). These results

indicated that the cells in our system originated from cluster 9, differentiated into the

KDR+ lateral mesoderm and formed endothelial progenitor cells (EPCs) under BPM4

factor induction; then by VEGF induction, endothelial cells undergo EHT, allowing

the cells to differentiate into HSPCs. Then, HSPCs differentiate into various blood

cells, mainly myeloid cells, similar to yolk sac hematopoiesis [45, 52, 53] (Figure

S5A).

Dynamic regulation of cell adhesion molecules in hematopoietic development

To explain how endothelial characteristics lost during the formation of HSPCs, we

performed differentially expressed gene (DEG) analysis between HSPCs and

hemangioblasts and found that with the formation of HSPCs, the expression levels of

TFs, such as GATA2 and TAL1, were significantly increased, and the expression

levels of endothelial characteristic genes, such as *TIE1* and *KDR*, were significantly

decreased (Figure 3D). Then, 466 genes that were significantly higher in HSPCs than

in hemangioblasts were intersected with the 1,639 human TFs retrieved from the TF

database [54], and 24 TFs were obtained. An interactive network analysis of these 24

TFs revealed that 4 of them were involved in erythroid differentiation, 4 TFs were

involved in lymphoid differentiation, 8 TFs were involved in myeloid differentiation,

and the others were potentially involved in hematopoietic differentiation (Figure 3E).

At the same time, compared with HSPCs, the functional annotations of the top 50

genes that were highly expressed in hemangioblasts showed significant enrichment in

hematopoietic differentiation pathways, such as the myeloid and lymphoid

differentiation pathways. The top 50 highly expressed genes in the hemangioblasts

were significantly enriched in the pathways of endothelial development and

PI3K-AKT signaling, and a number of genes were also found to be enriched in

pathways associated with cell adhesion; these genes were most highly expressed in

hemangioblasts (Figure 3F and S5C). Making use of these genes for interaction

network analysis, the inner circle on the interaction network map showed genes

enriched in the pathway, and the outer circle was genes that interacted with the inner

circle genes. We can see that the outer circle genes were characteristic of HE genes,

such as *KDR*, *NOTCH4*, and *ERG*, indicating that the outer circle genes were

necessary for the formation of hemangioblasts but were significantly lost during

hematopoietic differentiation [55, 56]. In addition, we found from the bulk RNA-seq

data that relative to iPSCs, SCs also downregulated the expression of cell

adhesion-related genes (Figure S5E), which may be the direct cause of SCs separation

from EBs.

Cell differentiation trajectory of the iPSC-derived RBC differentiation system

To further explain the differentiation trajectory of cells and the key factors affecting

cell differentiation, we performed a pseudotime analysis of scRNA-seq data using R

package Monocle [57]. The results also suggested that cell differentiation originated

from the progenitor cell clusters from hemangioblasts then experienced the first

bifurcation hemangioblasts. At the first bifurcation, cells differentiated into VECs and

HSPCs; then, at the second bifurcation, HSPCs differentiated into lymphoid cells

(Lyms) (small part cells) and CMPs (most part cells), and at the third bifurcation,

CMPs differentiated into MEs and granulocytes and monocytes (GMs) (**Figure 4**A

and B). Pseudotime heatmap analysis with 743 genes showed that all cells were

divided into three clusters, marked as early mesoderm cells, EPCs, and blood cells.

We compared the genes corresponding to the three clusters with the human TF

database. The *TBX2* and *PITX1* genes corresponding to the early mesodermal cells

cluster were involved in the differentiation of mesoderm, which guides mesodermal

organ formation and somite formation (Figure 4C and D). The *ETV2* and *ERG* genes

corresponding to the EPCs cluster could guide the cells to enter the hematopoietic

lineage, and the cluster also significantly expressed endothelial characteristic genes,

such as *TIE1* and *KDR* (Figure 4C and E). The TFs, including GATA2, corresponding

to blood cells cluster marked the initiation of hematopoiesis and then formed various

blood lineage cells under the guidance of EGR1, SPI1, GATA1, *etc.*, and the cluster

also   significantly   expressed   *F2R*,   *HBA1*   (the   characteristic   genes   of

megakaryocyte-erythroid cells) and *LYZ*, *MNDA*, and *ELANE* (the characteristic genes

of granulocyte-macrophage cells) (Figure 4C and F). In addition, pseudotime analysis

showed that the expression of the MEP characteristic genes were close to the

expression time of the HE characteristic gene *CD34*, and this phenomenon was

similar to the gene expression in early yolk sac hematopoiesis [58, 59] (Figure 4E and

F).

Functional annotation of the genes corresponding to each cluster indicated that the

early mesoderm cells cluster had the potential to differentiate into various mesoderm

organs, and the corresponding pathways were embryonic morphogenesis ($P =$

6.31E-34) and heart development ($P = 2.14$E-28). The EPCs cluster could further

differentiate into blood vessel and blood cells, and the corresponding genes were

significantly enriched in blood vessel development ($P = 1.35$E-46) and regulation of

the MAPK cascade ($P = 6.92$E-21). Many genes were also significantly enriched in

the pathways related to cell adhesion, such as KEGG pathway focal adhesion ($P =$

3.31E-22). And the blood cells underwent hematopoietic differentiation, the

corresponding genes were enriched in myeloid leukocyte activation ($P = 1.20$E-97),

hemostasis ($P = 2.69$E-28), cytokine production ($P = 2.04$E-24) and other pathways

(Figure S6). The above results suggest that the system underwent two key

transformations during the asynchronous differentiation process: (1) cells exited

pluripotent stem cells during germ layer differentiation, entered the lateral mesoderm

and were driven by the *ETV2* gene to differentiate into hematopoietic lineage [45]; (2)

HE cells entered hematopoietic differentiation under the guidance of GATA2 to form

various blood cells [60].

Fate-determining TFs drive cell bifurcation

To identify the key factors that drove cells into each bifurcation, we used a branched

heatmap to show the gene expression dynamics of these bifurcations. At the first

bifurcation, 606 genes divided cells into four clusters: mesodermal cells cluster, blood

vessel cluster, HSPCs cluster and blood lineage cells cluster. Driven by the SOX18

TF, the blood vessel cluster entered the vascular branch, and the direction of cell

differentiation was controlled by the SOX7 transcription suppressor factor [61]. The

HSPCs cluster driven by GATA2 and RUNX1 was further differentiated into a blood

lineage cells cluster under the induction of lineage-specific TFs, such as SPI1 and

SOX4 (**Figure 5**A). Functional annotation of genes in each cluster showed that the

genes in mesodermal cells were riched in skeletal system development (P = 5.01E-21)

and ossification (P = 1.35E-13). The blood vessel cluster genes were enriched in

angiogenesis (P = 6.17E-09), and the HSPCs cluster genes were enriched in ribosome

assembly (P = 6.92E-26) and *RUNX1* regulates transcription of genes involved in

differentiation of HSCs (P = 3.55E-05). The blood lineage cells cluster genes were

enriched in lymphocyte activation (P = 2.34E-13) and myeloid cell differentiation (P

= 5.62E-09).

To identify the driving factors for the differentiation of lymphoid and myeloid, we

constructed a differentiation trajectory for the HSPCs cluster and used 336 genes to

generate a pseudotime heatmap to show the gene expression dynamics of these

branches. Cells in the HSPCs cluster were divided into four clusters: multipotent

progenitors (MPPs), common lymphoid progenitors (CLPs), GMPs, and

megakaryocyte-erythroid progenitors (MEPs) (Figure 5B and C). In MPPs, the genes

GATA2, *RUNX1* and *CD34* were highly expressed, and functional annotation revealed

that the genes were significantly enriched in ribosome biogenesis (P = 1.70E-14) and

hematopoiesis (P = 9.55E-04). The CLPs cluster that driven by TFs such as IKZF1

and EGR1 was highly expressed *JAML*, *AIF1* and other lymphoid characteristic

genes, and functional annotation revealed a significant enrichment for lymphocyte

activation (P = 1.95E-55) and cytokine production (P = 8.51E-32). GMPs and MEPs

are both myeloid cells, under SPI1 and GATA1 induction, myeloid-specific genes

such as *LYZ, CLC*, *CD36* and *KLF1* were highly expressed. Functional annotation

revealed that the enriched genes were involved in myeloid leukocyte activation (P =

1.58E-48) and hemostasis (P = 5.13E-25) (Figure 5C, D and S6).

To identify the driving factors for the differentiation of GMPs and MEPs, we

constructed a differentiation trajectory for the CD33 cells cluster, MEs cluster and

CD14 cells cluster and used 633 genes to perform a pseudotime heatmap to show the

gene expression dynamics of these cells. Cells were divided into three clusters: MEs,

CMPs and GMs. MEs driven by GATA1 were highly expressed in the RBC-specific

gene *HBA1* and in the platelet-specific gene *GP9* [29]. Functional annotation of this

cluster was significantly enriched in platelet activation ($P = 8.51E-24$) and regulation

of ion transport ($P = 1.95E-10$). CMPs highly expressed *SPI1* and *MYB*, and

functionally annotated genes from this cluster were significantly enriched in response

to interleukin-1 ($P = 2.14E-04$) and the CMYB pathway ($P = 3.80E-04$). GMs driven

by MAFB had high expression of the granulocyte-specific gene *MNDA* and

monocyte-specific cell marker *CD14* (Figure 5E-G). Functional annotation of this

cluster was significantly enriched in myeloid leukocyte activation ($P = 1.23E-30$) and

regulation of the inflammatory response ($P = 6.76E-14$) (Figure S6).

Pseudotime trajectory of erythroid cells in the iPSC-derived RBC differentiation

system

To accurately elucidate the molecular activities in erythroid differentiation, we

performed pseudotime analysis of MEs cluster alone. Simultaneously, pseudotime

heatmap analysis with 200 genes showed that cells could be divided into two

subclusters, marked as MEPs and erythroid cells (Erys). The TF GATA1 was highly

expressed in the MEPs cluster and Erys cluster highly expressed of the erythrocyte

characteristic TF KLF1, while high expression of *GFL1B*, *NFE2*, *MAX* and other

genes played an important role in the differentiation and maturation of RBCs (**Figure**

6A, B). At the same time, the pseudotime trajectory analysis of the characterizing

genes showed that in the process of erythroid differentiation, the expression levels of

CD34 and *TIE1* (endothelial marker genes) were downregulated and decreased

rapidly during the differentiation process. *GATA1* was closely related to the early

differentiation of erythroid cells and was rapidly upregulated during the

differentiation process but was rapidly downregulated at the end of differentiation;

consistent with previous studies, *TAL1* and *GATA1* also showed similar trends [62].

KLF1 and *ALAS2* were also key factors during erythroid differentiation, and the

expression level gradually increased with the differentiation process; *HBA1* and *HBB*

encoded adult-type globin, *HBA1* was highly expressed throughout the erythroid

differentiation process, and the expression levels of both *HBA1* and *HBB* gradually

increased. *FTH1* and *FTL* encoded ferritin, which were highly expressed throughout

the differentiation process and were gradually upregulated throughout the erythroid

differentiation process, which may be related to the synthesis of heme (Figure 6C).

Functional annotation results revelated that genes corresponding to MEPs were

enriched for the functions of embryonic hemopoiesis (P = 2.95E-05) and response to

estradiol (P = 7.72E-04). Functional annotation results of genes corresponding to Erys

showed significant enrichment in oxygen transport (P = 5.84E-13), erythrocyte

differentiation (P = 1.73E-07) and other RBC characteristic pathways. The target gene

of estradiol is highly expressed at the beginning of the erythroid differentiation

process, and then the expression level is reduced. The results indicate that estradiol

may promote the proliferation of erythroid progenitor cells with erythroid

differentiation but simultaneously inhibits erythroid differentiation (Figure 6E). The

interaction network analysis of the corresponding genes in the O_2/CO_2 exchange in

erythrocytes, heme metabolic process and iron ion homeostasis pathways showed that

ALAS2, the hub gene in the interaction network, played a key role in the process of

erythroid differentiation and maturation [63] (Figure 6F). In addition, we mapped the

TFs of these 200 genes and the genes enriched in the pathways involved in erythroid

differentiation (Figure S7B and C). In addition, the β-chain globin expression profile

and the qPCR results corresponding to the erythroid cells in the data showed that the

erythroid cells produced by our iPSC system are mainly derived from yolk sac

hematopoiesis, and interestingly, during the hemangioblast formation period of D4

EBs [64], RBCs expressed *HBG*, which indicates that RBC production occurs before

HSPCs are formed during embryo culture (Figure S7D).

Our results showed that iPSCs entered erythroid differentiation, undergo two

transformations, exit pluripotency, and high expression of *ETV2* and *KDR* under

BMP4 and VEGF stimulation to induce cells to differentiate into hemangioblasts; this

process is accompanied by cell adhesion activation [45, 55]. In addition, HSPCs enter

the erythroid differentiation process accompanied by the loss of cell adhesion and the

fluctuation of estradiol target genes expression. During this process, the HSPCs would

be induced by IKZF1 to undergo lymphoid differentiation, and induced by SPI1 to

undergo myeloid differentiation; subsequently CMPs would be induced by MAFB to

enter granulocyte-mononuclear differentiation, and induced by GATA1 to enter

megakaryocyte-erythroid differentiation; lastly, MEPs would be induced by KLF1 to

enter erythroid differentiation (Figure 6G). We confirmed the expression trend of cell

adhesion-related genes (*KDR*, *FN1*, *CD34*, *CDH5*, *CD99* and *ESAM*) in EB and SC

differentiation (Figure S7E, left) and estradiol target genes (*ALAS2*, *GATA1* and

KLF1) in erythroid differentiation (Figure S7E, right) by qPCR analysis. The

experimental results were consistent with the data analysis.

Discussion

With the rapid development of single-cell technology, single-cell transcriptome

studies on the development of the hematopoietic lineage have been extensively

reported, and the known hematopoietic theory has been revised and supplemented, but

most studies were focused on bone marrow, peripheral blood, cord blood and other

systems [65, 66]. In this study, we performed a single-cell transcriptomics study on

the RBC regeneration system derived from iPSCs. The cell map and cell

differentiation trajectory of the system were mapped for the first time, and the whole

transcriptome dynamics from iPSCs to the erythroid lineage were delineated.

Through the analysis of the cell map, we found that in the process of

hematopoiesis, there are mainly mesoderm and various blood cells, which means that

cell differentiation is not synchronized. The process of RBC regeneration by iPSCs

through the embryoid stage is similar to yolk sac hematopoiesis, and these results are

consistent with those of previous research [20, 21]. The D14 erythrocytes are

composed of the primitive RBCs produced by the first wave of hematopoiesis and the

definitive RBCs produced by the second wave of yolk sac hematopoiesis. Similar to

the recent single-cell sequencing results of D29 iPS-derived erythroid differentiation,

it was shown that the globin expression profile of iPSC-derived definitive

erythrocytes corresponds to the globin expression profile of yolk sac erythroid

progenitor cells [41]. Both results suggest that iPSC-derived erythropoiesis is

accompanied by globin gene expression switching, and this system can be used as a

favorable model for studying the globin conversion mechanism [30].

According to the results of cell mapping analysis, cell differentiation originated

from iPSCs. Under the action of the cytokine BMP4, iPSCs form KDR+ lateral

mesoderm by means of germ layer differentiation and produce EPCs. Subsequently,

under the action of VEGF, the EPCs are transdifferentiated to form HE cells, and

HSPCs are released. Then, HSPCs are further induced to differentiate into blood

lineage cells. Our system can form various blood cells by hierarchical induction,

especially RBCs, and this type of blood cell formation is very close to the embryonic

hematopoiesis model *in vivo*, indicating that our system is a good model for

simulating *in vivo* hematopoietic differentiation and development [67].

By studying the differentiation trajectory of the iPSC-derived RBC differentiation

system, we found that the iPSCs differentiated enter hematopoietic lineage under the

driving force of *ETV2* and underwent three bifurcations, then enter erythroid

differentiation process. During the process of the bifurcation of hematopoiesis and

angioblasts, the cells enter the hematopoietic lineage driven by TFs, such as GATA2

and RUNX1; in the process of bifurcation of the myeloid and lymphoid lines, the cells

enter the myeloid lineage driven by TF SPI1; and during the process the bifurcation of

the GMPs and MEPs, the cells enter the megakaryocyte-erythroid lineage driven by

GATA1, and then erythrocytes are formed under the driving force of KLF1. These

series of fate-determining TFs can effectivly guide the regulation of iPSC-derived

RBC differentiation process and provide estimable advice for cell regeneration in

other blood lineages. In addition, pseudotime clustering analysis also suggested that

the pluripotent stem cells to erythroid cell production underwent two major

transformations, including *ETV2*-directed withdrawal of iPSC pluripotency into the

hematopoietic lineage, followed by GATA1-directed HSPC erythroid differentiation.

We noted that cell adhesion molecules are actively expressed during the

transformation one process but are downregulated during transformation two, possibly

related to cell adhesion for the formation of the gastrula [55]. We found that during

iPSC-derived erythroid differentiation, estradiol promoted the proliferation of

erythroid progenitor cells and inhibited erythroid differentiation. We were advised

that we could add a cell adhesion activator to stimulate the production of

hemangioblasts during the first four days of culturing EBs and then could add

estradiol to the medium 6 days before inducing SC differentiation to increase the

efficiency of RBC regeneration. In addition, a number of TFs that play a key role in

the differentiation and maturation of RBCs, such as *NEF2* and *MAX*, have also been

discovered. This information provides a powerful theoretical guide for optimizing our

iPSC-derived RBC regeneration system. With the continuous development of

single-cell technology, we will be able to excavate more potential mechanisms

regulating RBC development and maturation and will be able to achieve the

development and differentiation of RBCs in our system, thereby obtaining more

mature regenerated functional RBCs *in vitro* for use in clinical applications.

Materials and methods

iPSC and EB culture and erythropoiesis

iPSCs, which the original source was Linzhao Cheng's lab, were cultured in

vitronectin (Life Technologies)-coated culture dishes with Essential 8 medium

(Gibco). When the cells reached 70% to 80% confluence (usually around 3~4 days),

they were digested to suspension cells using 0.5 mM EDTA (Invitrogen) and passaged

according to a ratio of 1 to 4.

EB culture was performed during D0 to D11 as previously described [68]. From

D11 to D14 of EB culture, TPO was replaced by 3 U/ml erythropoietin (EPO)

(PeproTech). On D14, SCs were collected by a 70 μm cell strainer. Subsequently, the

SC solution was centrifuged at 300 g for 5 minutes to remove the supernatant, and

serum-free medium (SFM) containing 50 μg/ml stem cell factor (SCF) (PeproTech),

100 μg/ml IL-3 (PeproTech), 2 U/ml EPO and 1% penicillin/streptomycin (P/S) (Life

Technologies) was used to resuspend the SC precipitates. The cell concentration was

adjusted to 5 x 10^5/ml and cultured in a 6-well plate at a total volume of 2 ml per well.

All cells were cultured for 6 days at 37 °C under 5% CO_2, and the medium was

changed every three days. On D20, the medium was replaced with SFM containing 2

U/ml EPO and 1% P/S, and the cell concentration was adjusted appropriately. The

number of cells in each well did not exceed 2 x 10^6, and the total volume of the

medium per well was 2 ml. The cells were cultured at 37 °C under 5% CO_2, and the

medium was changed every three days.

Colony formation unit assay

Incubation of D14 SCs used MethoCult™ H4434 Classic Methylcellulose Medium

for Human Cells (StemCell). Each well of a 12-well plate contained 8,000-10,000

cells and 0.6 ml of medium. Other operations were performed as described in the

instructions. Cells were incubated for 12-14 days, and the clones were observed under

the microscope.

Flow cytometry analysis

First, ~1 × 10^5 cells were collected from the culture, centrifuged at 300 g for 5

minutes to remove the medium, and washed twice with 2 ml 1× Dulbecco's

phosphate buffered saline (DPBS) (Gibco) solution containing 2% FBS (Gibco) and 2

mM EDTA. After centrifugation at 300 g for 5 minutes to remove the supernatant, the

cells were resuspended in 50 µl DPBS buffer, and the antibodies (0.1 µl for anti-CD71

and 0.5 µl for anti-CD235a) were added, followed by incubation at 4 °C for 10

minutes in the dark. The cells were washed twice by adding 2 ml of 1× DPBS buffer

after the incubation and resuspended in 200 µl of DPBS buffer to prepare a cell

suspension for the assay. The BD FACSAria II instrument was used for flow

cytometry analysis, and the data analysis was performed with Flowjo software

(Version 7.6, Three Star). The antibodies (Miltenyi Biotec) used in this study included

anti-human CD34-FITC, anti-human CD45-PE, anti-human CD71-PE, and

anti-human CD235a-APC.

Wright-Giemsa staining

~1 × 10^5 cells were taken, and the medium was removed by centrifugation at 300 g for

5 minutes; the cells were resuspended in 20 µl of DPBS buffer. The cells were placed

onto one of two clean slides, and the other slide was used to push the slide at a 30 to

45-degree angle to spread the cells evenly. The pushed film was placed in a 37 °C

incubator for 1 to 3 hours. Then, 200 μL of Wright-Giemsa staining A solution

(BASO, BA-4017) was placed onto the cell-containing slide, the slide was placed at

room temperature for 1 minute and was immediately covered with 400 μL of B

solution. After the slide was incubated at room temperature for 8 minutes, the slides

were gently rinsed with fluid distilled water along the area around the slide without

cells. The slides were rinsed for an additional 1 minute until the water ran clear and

transparent. Then, the slides were placed at room temperature until dry, and a

microscope was used to observe and record cell morphology.

Total RNA extraction and qPCR

Total RNA was extracted from the iPSC line BC1, D4 EBs, D14 SCs, and D20

erythroid cells, and the total RNA was reverse transcribed into cDNA using the

PrimeScript RT Reagent Kit with gDNA Eraser (TaKaRa). The diluted cDNA was

used as a template for qPCR. The instrument used for qPCR was the Bio-Red CFX96

Real-Time PCR Detection System. All primers used in qPCR are listed in

Supplementary Table S7.

scRNA-seq data analysis and bulk RNA-seq data analysis

D14 EBs were dissociated with Accutase solution (Sigma). Single-cell suspensions at

800 cells/μL were subjected to Chromium 10× Genomics library construction. Raw

gene expression matrices generated by CellRanger (version 2.0.0) were imported into

Seurat (version 2.4) [44]. After removing genes expressed in fewer than 5 cells, in

cells with more than 5% mitochondria reads or in cells with less than 200 unique

genes, two Seurat objects were integrated and clustered into 10 and 14 clusters based

on fifteen principal components with the resolution parameter set at 0.2 and 0.8. Each

cluster was identified according to the marker genes found by the Seurat

"FindAllMarkers" function. Particular clusters were imported into Monocle2 [57] for

pseudotime ordering. For bulk RNA-seq data, FASTQC and Trimmomatic were used

to remove adaptor sequences and low-quality reads. Then, clean data were mapped to

the reference genome (GRCh38) by HISAT2 [69] to generate a gene expression

matrix, which was then imported into DEGseq [70] R Package to identify DEGs.

Heatmap analysis and PCA were performed by pheatmap and PRINCOMP, respectively. All data analyses were completed in the R 3.5.2 environment. We used Metascape [71] to perform functional annotation analysis. Human TF sequences were downloaded from http://humantfs.ccbr.utoronto.ca/download.php. The network was visualized by GeneMANIA [72].

Data availability

The single-cell RNA-seq of D14 EBCs and SCs, and the bulk RNA-seq of BC1 and D14 SCs is available in the Genome Sequence Archive (GSA) http://bigd.big.ac.cn/gsa/s/ZVZPnz5q

Authors' contributions

XF and ZZ designed and guided the study. WZ and JZ completed the experiments. ZX and SG completed the data analysis. YL supported the experiments. ZX and WZ drafted the manuscript, XF and ZZ edited the manuscript. All authors read and approved the final manuscript.

Competing interests

The authors have declared no competing interests.

Acknowledgments

This research was supported by the Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No. XDA16010602), the National Key Research and Development Program of China (Grant Nos. 2016YFC0901700, 2016YFC0901603, 2017YFC0907400, 2018YFC0910700, 2017YFC0907403 and 2018YFC0910402) and the National Natural Science Foundation of China (Grant Nos. 81670109, 81870097, 81700097 and 81700116). We were grateful to Linzhao Cheng for agreeing to use iPSC line BC1 and Qianfei Wang for providing the iPSCs.

References

[1] Williamson LM, Devine DV. Challenges in the management of the blood supply. The Lancet 2013;381:1866-75.

[2] Moradi S, Jahanian-Najafabadi A, Roudkenar MH. Artificial blood substitutes: First steps on the long route to clinical utility. Clin Med Insights Blood Disord 2016;9:33-41.

[3] Batta K, Menegatti S, Garcia-Alegria E, Florkowska M, Lacaud G, Kouskoff V. Concise review: Recent advances in the in vitro derivation of blood cell populations. Stem Cells Transl Med 2016;5:1330-7.

[4] Christaki EE, Politou M, Antonelou M, Athanasopoulos A, Simantirakis E, Seghatchian J, et al. Ex vivo generation of transfusable red blood cells from various stem cell sources: A concise revisit of where we are now. Transfus Apher Sci 2019;58:108-12.

[5] Lopez-Yrigoyen M, Yang CT, Fidanza A, Cassetta L, Taylor AH, McCahill A, et al. Genetic programming of macrophages generates an in vitro model for the human erythroid island niche. Nat Commun 2019;10:881.

[6] Shen J, Zhu Y, Lyu C, Feng Z, Lyu S, Zhao Y, et al. Sequential cellular niches control the generation of enucleated erythrocytes from human pluripotent stem cells. Haematologica 2019.

[7] Choi KD, Vodyanik MA, Togarrati PP, Suknuntha K, Kumar A, Samarjeet F, et al. Identification of the hemogenic endothelial progenitor and its direct precursor in human pluripotent stem cell differentiation cultures. Cell Rep 2012;2:553-67.

[8] Liu Y, Wang Y, Gao Y, Forbes JA, Qayyum R, Becker L, et al. Efficient generation of megakaryocytes from human induced pluripotent stem cells using food and drug administration-approved pharmacological reagents. Stem Cells Transl Med 2015;4:309-19.

[9] Sun S, Peng Y, Liu J. Research advances in erythrocyte regeneration sources and methods in vitro. Cell Regen (Lond) 2018;7:45-9.

[10] Park YJ, Cha S, Park YS. Regenerative applications using tooth derived stem cells in other than tooth regeneration: A literature review. Stem Cells Int 2016;2016:9305986.

[11] Jiang L, Jones S, Jia X. Stem cell transplantation for peripheral nerve regeneration: Current options and opportunities. Int J Mol Sci 2017;18.

[12] Illich DJ, Demir N, Stojkovic M, Scheer M, Rothamel D, Neugebauer J, et al. Concise review: Induced pluripotent stem cells and lineage reprogramming: Prospects for bone regeneration. Stem Cells 2011;29:555-63.

[13] Wu SM, Hochedlinger K. Harnessing the potential of induced pluripotent stem cells for regenerative medicine. Nat Cell Biol 2011;13:497-505.

[14] Bilic J, Izpisua Belmonte JC. Concise review: Induced pluripotent stem cells versus embryonic stem cells: Close enough or yet too far apart? Stem Cells 2012;30:33-41.

[15] Kauts ML, Rodriguez-Seoane C, Kaimakis P, Mendes SC, Cortes-Lavaud X, Hill U, et al. In vitro differentiation of gata2 and ly6a reporter embryonic stem cells

corresponds to in vivo waves of hematopoietic cell generation. Stem Cell Reports
2018;10:151-65.

[16] Koyano-Nakagawa N, Kweon J, Iacovino M, Shi X, Rasmussen TL, Borges L, et
al. Etv2 is expressed in the yolk sac hematopoietic and endothelial progenitors and
regulates lmo2 gene expression. Stem Cells 2012;30:1611-23.

[17] Kataoka H, Hayashi M, Nakagawa R, Tanaka Y, Izumi N, Nishikawa S, et al.
Etv2/er71 induces vascular mesoderm from flk1+pdgfralpha+ primitive mesoderm.
Blood 2011;118:6975-86.

[18] Kirmizitas A, Meiklejohn S, Ciau-Uitz A, Stephenson R, Patient R. Dissecting
bmp signaling input into the gene regulatory networks driving specification of the
blood stem cell lineage. Proc Natl Acad Sci U S A 2017;114:5814-21.

[19] Hubner K, Grassme KS, Rao J, Wenke NK, Zimmer CL, Korte L, et al. Wnt
signaling positively regulates endothelial cell fate specification in the fli1a-positive
progenitor population via lef1. Dev Biol 2017;430:142-55.

[20] Palis J. Primitive and definitive erythropoiesis in mammals. Front Physiol
2014;5:3.

[21] McGrath KE, Frame JM, Fromm GJ, Koniski AD, Kingsley PD, Little J, et al. A
transient definitive erythroid lineage with unique regulation of the beta-globin locus in
the mammalian embryo. Blood 2011;117:4600-8.

[22] Lie ALM, Marinopoulou E, Li Y, Patel R, Stefanska M, Bonifer C, et al. Runx1
positively regulates a cell adhesion and migration program in murine hemogenic
endothelium prior to blood emergence. Blood 2014;124:e11-20.

[23] de Pater E, Kaimakis P, Vink CS, Yokomizo T, Yamada-Inagawa T, van der
Linden R, et al. Gata2 is required for hsc generation and survival. J Exp Med
2013;210:2843-50.

[24] Huang K, Du J, Ma N, Liu J, Wu P, Dong X, et al. Gata2(-/-) human escs
undergo attenuated endothelial to hematopoietic transition and thereafter granulocyte
commitment. Cell Regen (Lond) 2015;4:4.

[25] Ivanovs A, Rybtsov S, Welch L, Anderson RA, Turner ML, Medvinsky A.
Highly potent human hematopoietic stem cells first emerge in the intraembryonic
aorta-gonad-mesonephros region. J Exp Med 2011;208:2417-27.

[26] Stratman AN, Davis MJ, Davis GE. Vegf and fgf prime vascular tube
morphogenesis and sprouting directed by hematopoietic stem cell cytokines. Blood
2011;117:3709-19.

[27] Guitart AV, Subramani C, Armesilla-Diaz A, Smith G, Sepulveda C, Gezer D, et
al. Hif-2alpha is not essential for cell-autonomous hematopoietic stem cell
maintenance. Blood 2013;122:1741-5.

[28] Upadhaya S, Sawai CM, Papalexi E, Rashidfarrokhi A, Jang G, Chattopadhyay
P, et al. Kinetics of adult hematopoietic stem cell differentiation in vivo. J Exp Med
2018;215:2815-32.

[29] Palii CG, Cheng Q, Gillespie MA, Shannon P, Mazurczyk M, Napolitani G, et al.
Single-cell proteomics reveal that quantitative changes in co-expressed
lineage-specific transcription factors determine cell fate. Cell Stem Cell
2019;24:812-20.

[30] Qiu C, Olivier EN, Velho M, Bouhassira EE. Globin switches in yolk sac-like
primitive and fetal-like definitive red blood cells produced from human embryonic
stem cells. Blood 2008;111:2400-8.
[31] Yu X, Azzo A, Bilinovich SM, Li X, Dozmorov M, Kurita R, et al. Disruption of
the mbd2-nurd complex but not mbd3-nurd induces high level hbf expression in
human erythroid cells. Haematologica 2019.
[32] Chou AC, Broun GO, Jr., Fitch CD. Abnormalities of iron metabolism and
erythropoiesis in vitamin e-deficient rabbits. Blood 1978;52:187-95.
[33] Anderson GJ, Powell LW, Halliday JW. Transferrin receptor distribution and
regulation in the rat small intestine. Effect of iron stores and erythropoiesis.
Gastroenterology 1990;98:576-85.
[34] Rutherford TR, Weatherall DJ. Deficient heme synthesis as the cause of
noninducibility of hemoglobin synthesis in a friend erythroleukemia cell line. Cell
1979;16:415-23.
[35] Ji P, Murata-Hori M, Lodish HF. Formation of mammalian erythrocytes:
Chromatin condensation and enucleation. Trends Cell Biol 2011;21:409-15.
[36] Murry CE, Keller G. Differentiation of embryonic stem cells to clinically
relevant populations: Lessons from embryonic development. Cell 2008;132:661-80.
[37] Mazurier C, Douay L, Lapillonne H. Red blood cells from induced pluripotent
stem cells: Hurdles and developments. Curr Opin Hematol 2011;18:249-53.
[38] Tang F, Lao K, Surani MA. Development and applications of single-cell
transcriptome analysis. Nat Methods 2011;8:S6-11.
[39] Han X, Chen H, Huang D, Chen H, Fei L, Cheng C, et al. Mapping human
pluripotent stem cell differentiation pathways using high throughput single-cell
rna-sequencing. Genome Biol 2018;19:47.
[40] Swiers G, Baumann C, O'Rourke J, Giannoulatou E, Taylor S, Joshi A, et al.
Early dynamic fate changes in haemogenic endothelium characterized at the
single-cell level. Nat Commun 2013;4:2924.
[41] Vanuytsel K, Matte T, Leung A, Naing ZH, Morrison T, Chui DHK, et al.
Induced pluripotent stem cell-based mapping of beta-globin expression throughout
human erythropoietic development. Blood Adv 2018;2:1998-2011.
[42] Cheng L, Hansen NF, Zhao L, Du Y, Zou C, Donovan FX, et al. Low incidence
of DNA sequence variation in human induced pluripotent stem cells generated by
nonintegrating plasmid expression. Cell Stem Cell 2012;10:337-44.
[43] Chou BK, Mali P, Huang X, Ye Z, Dowey SN, Resar LM, et al. Efficient human
ips cell derivation by a non-integrating plasmid from blood cells with unique
epigenetic and gene expression signatures. Cell Res 2011;21:518-29.
[44] Satija R, Farrell JA, Gennert D, Schier AF, Regev A. Spatial reconstruction of
single-cell gene expression data. Nat Biotechnol 2015;33:495-502.
[45] Koyano-Nakagawa N, Garry DJ. Etv2 as an essential regulator of mesodermal
lineage development. Cardiovasc Res 2017;113:1294-306.
[46] Tusi BK, Wolock SL, Weinreb C, Hwang Y, Hidalgo D, Zilionis R, et al.
Population snapshots predict early haematopoietic and erythroid hierarchies. Nature
2018;555:54-60.

[47] Stansfield BK, Ingram DA. Clinical significance of monocyte heterogeneity. Clin
Transl Med 2015;4:5.

[48] Scialdone A, Tanaka Y, Jawaid W, Moignard V, Wilson NK, Macaulay IC, et al.
Resolving early mesoderm diversification through single-cell expression profiling.
Nature 2016;535:289-93.

[49] Pijuan-Sala B, Griffiths JA, Guibentif C, Hiscock TW, Jawaid W, Calero-Nieto
FJ, et al. A single-cell molecular map of mouse gastrulation and early organogenesis.
Nature 2019;566:490-5.

[50] Min IM, Pietramaggiori G, Kim FS, Passegue E, Stevenson KE, Wagers AJ. The
transcription factor egr1 controls both the proliferation and localization of
hematopoietic stem cells. Cell Stem Cell 2008;2:380-91.

[51] Villani AC, Satija R, Reynolds G, Sarkizova S, Shekhar K, Fletcher J, et al.
Single-cell rna-seq reveals new types of human blood dendritic cells, monocytes, and
progenitors. Science 2017;356.

[52] Nostro MC, Cheng X, Keller GM, Gadue P. Wnt, activin, and bmp signaling
regulate distinct stages in the developmental pathway from embryonic stem cells to
blood. Cell Stem Cell 2008;2:60-71.

[53] Metcalf D. Hematopoietic cytokines. Blood 2008;111:485-91.

[54] Lambert SA, Jolma A, Campitelli LF, Das PK, Yin Y, Albu M, et al. The human
transcription factors. Cell 2018;172:650-65.

[55] Adams JC, Watt FM. Regulation of development and differentiation by the
extracellular matrix. Development 1993;117:1183-98.

[56] Angelos MG, Abrahante JE, Blum RH, Kaufman DS. Single cell resolution of
human hematoendothelial cells defines transcriptional signatures of hemogenic
endothelium. Stem Cells 2018;36:206-17.

[57] Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The
dynamics and regulators of cell fate decisions are revealed by pseudotemporal
ordering of single cells. Nat Biotechnol 2014;32:381-6.

[58] Dzierzak E, Philipsen S. Erythropoiesis: Development and differentiation. Cold
Spring Harb Perspect Med 2013;3:a011601.

[59] Palis J, Robertson S, Kennedy M, Wall C, Keller G. Development of erythroid
and myeloid progenitors in the yolk sac and embryo proper of the mouse.
Development 1999;126:5073-84.

[60] Gao X, Johnson KD, Chang YI, Boyer ME, Dewey CN, Zhang J, et al. Gata2
cis-element is required for hematopoietic stem cell generation in the mammalian
embryo. J Exp Med 2013;210:2833-42.

[61] Downes M, Koopman P. Sox18 and the transcriptional regulation of blood vessel
development. Trends Cardiovasc Med 2001;11:318-24.

[62] Rylski M, Welch JJ, Chen YY, Letting DL, Diehl JA, Chodosh LA, et al.
Gata-1-mediated proliferation arrest during erythroid maturation. Mol Cell Biol
2003;23:5031-42.

[63] To-Figueras J, Ducamp S, Clayton J, Badenas C, Delaby C, Ged C, et al. Alas2
acts as a modifier gene in patients with congenital erythropoietic porphyria. Blood
2011;118:1443-51.

[64] Fehling HJ, Lacaud G, Kubo A, Kennedy M, Robertson S, Keller G, et al.
Tracking mesoderm induction and its specification to the hemangioblast during
embryonic stem cell differentiation. Development 2003;130:4217-27.
[65] Dzierzak E, Bigas A. Blood development: Hematopoietic stem cell dependence
and independence. Cell Stem Cell 2018;22:639-51.
[66] Wilson NK, Göttgens B. Single-cell sequencing in normal and malignant
hematopoiesis. HemaSphere 2018;2.
[67] Yvernogeau L, Gautier R, Khoury H, Menegatti S, Schmidt M, Gilles JF, et al.
An in vitro model of hemogenic endothelium commitment and hematopoietic
production. Development 2016;143:1302-12.
[68] Liu Y, Wang Y, Gao Y, Forbes JA, Qayyum R, Becker L, et al. Efficient
generation of megakaryocytes from human induced pluripotent stem cells using food
and drug administration-approved pharmacological reagents. STEM CELLS
Translational Medicine 2015;4:309-19.
[69] Kim D, Langmead B, Salzberg SL. Hisat: A fast spliced aligner with low
memory requirements. Nat Methods 2015;12:357-60.
[70] Wang L, Feng Z, Wang X, Wang X, Zhang X. Degseq: An r package for
identifying differentially expressed genes from rna-seq data. Bioinformatics
2010;26:136-8.
[71] Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, et al.
Metascape provides a biologist-oriented resource for the analysis of systems-level
datasets. Nat Commun 2019;10:1523.
[72] Warde-Farley D, Donaldson SL, Comes O, Zuberi K, Badrawi R, Chao P, et al.
The genemania prediction server: Biological network integration for gene
prioritization and predicting gene function. Nucleic Acids Res 2010;38:W214-20.

Figure legends

Figure 1   Identification of iPSC-derived HSPCs and erythroid cells.

A. Schematic diagram of the protocol used to generate HSPCs and erythroid cells

from iPSCs *in vitro*. **B.** Histogram of iPSC counts on D0 and collected SCs on D14

per round-bottom 96-well plate (error bar = mean ± sd). **C.** Percentage of CD34+ and

CD45+ cells in the total SCs (mean ± sd). **D.** Representative image of CFU-GEMM

grown from collected SCs on D14. **E.** Representative results of CD71+ and CD235a+

cells expressed on D14, D17, D23 and D28 during the erythroid differentiation of

SCs. F. Representative image of erythroid cells on D28.

826
827 **Figure 2   Cell lineage map of the iPSC-derived RBC regeneration system.**

828 **A**. t-SNE plot of single-cell sample profiles. **B**. t-SNE plot of single-cell cluster

829 distribution. **C**. Heatmap shown the expression pattern of the top 50 DEGs in each

830 cell cluster; all genes used are listed in Supplementary Table S2. **D**. The dotplot of

831 marker genes of each cluster.

832
Figure 3   Subcluster analysis and DEG analysis.

A . t-SNE plots of single-cell subcluster distribution. **B**. Heatmap shown the DEG

pattern of the subcluster from the CD33 cells cluster. Top 50 DEGs of each subcluster

were shown, and genes were listed in Supplementary Table S4. **C**. Heatmap shown

the DEG pattern of subclusters from progenitor cells-1 cluster. Top 50 DEGs of each

subcluster were shown, and genes were listed in Supplementary Table S4. **D**.

Volcanic plot of DEGs from DEG analysis between HSPCs and hemangioblasts ($P <$

0.05, -1< Log2 FC <1). **E**. GO analysis of DEGs from DEG analysis between HSPCs

and hemangioblasts. Each functional term was shown in Supplementary Table S3. **F**.

Network shown the high expression of TFs in HSPCs.

843
844 **Figure 4   Cell pseudotime trajectory of the iPSC-derived RBC regeneration**

845 **system.**

846 **A**. Differentiation trajectory of iPSC-derived erythrocyte regeneration system by

847 monocle. **B**. Heatmap shown the gene expression dynamics during iPSC

differentiation. Genes were listed in Supplementary Table S3. **C-E**. Pseudotime analysis of marker genes of each cluster from early mesodermal cells cluster (**C**), HE cells cluster (**D**) and blood cells cluster (**E**).

Figure 5 Driving branching TFs of cell pseudotime trajectory.
A. Branched heatmap of differentiation branch point 1 from Figure 4a. Genes were listed in Supplementary Table S5. **B**. Differentiation trajectory of HSPCs. **C**. Heatmap shown the gene expression dynamics in HSPCs. Genes were listed in Supplementary Table S5. **D**. Pseudotime analysis of the marker gene of each cluster in C. **E**. Differentiation trajectory of cells from the CD33 cells cluster, MEs cluster and CD14 cells cluster in Figure 2B. **F**. Heatmap shown the gene expression dynamics in cells from E. Genes were listed in Supplementary Table S5. **G**. Pseudotime analysis of the marker genes of each cluster in F.

Figure 6 Differentiation trajectory from MEPs to erythroid cells.
A. Differentiation trajectory of MEs. **B**. Heatmap shown the gene expression dynamics in MEs. Genes were listed in Supplementary Table S5. **C**. Pseudotime analysis of marker genes of each cluster in B. **D**. Functional annotation analysis of MEPs cluster in B. Each functional term was shown in Supplementary Table S5. **E**. Pseudotime analysis of genes in the response to the estradiol functional term from D. **F**. Network shown the genes involved in RBC development from the Erys cluster in B.

Supplementary material

Figure S1 Quality control for scRNA-seq data and bulk RNA-seq data analysis.
A. Distribution of genes, transcripts and mitochondrial read proportion detected in each cell. **B.** Heatmap shown the expression pattern of the top 50 DEGs in each single-cell sample. Genes were listed in Supplementary Table S1. **C.** Functional annotation analysis of the top 50 DEGs in each single-cell sample. Each functional term was shown in Supplementary Table S1. **D.** Heatmap of hematopoietic lineage

related gene expression patterns for four bulk RNA-seq datasets. **E.** PCA of four bulk

RNA-seq datasets.

Figure S2 Overview of the 3,215 single cells from EBCs and SCs.

A. Distribution of cells in each cluster. **B.** Distribution of EBCs and SCs in each

cluster. **C.** Identification of marker genes for each cluster.

Figure S3 Functional annotation analysis of the top 50 marker genes in each

cluster, each functional term was shown in Table S3.

Figure S4 Subclusters of the CD33 cells cluster, progenitor cells-1 cluster, HSPCs

cluster and CD14 cells cluster.

A, **B, E, F.** Violin plots shown the expression distributions of marker genes across

subclusters: CD33 cells cluster (**A**) progenitor cells-1 cluster (**B**), HSPCs cluster (**E**)

and CD14 cells cluster (**F**). Heatmaps shown the DEG pattern of each subcluster from

the HSPCs cluster (**C**) and CD14 cells cluster (**D**). Gene sets of each cluster were

listed in Supplementary Table S4.

Figure S5 iPSC differentiation process and DEG analysis results

A. The process of iPSCs as they differentiate into erythroid cells. **B.** Venn diagram

shown the highly expressed TFs in HSPCs. **C.** t-SNE plots shown the cell adhesion

molecules. **D.** Network shown the high expression of TFs in HSPCs. **E.** Functional

annotation analysis of genes downregulation in SCs compared with that in iPSCs.

Each functional term was shown in Supplementary Table S3.

Figure S6 Functional annotation analysis for the heatmap genes in Figures 4 and

5. Each functional term was shown in Table S6.

Figure S7 Function of cell adhesion and estradiol during erythroid

differentiation.

A. Functional annotation analysis of the Erys cluster in Figure 6B. Each functional

term was shown in Supplementary Table S6**. B, C.** The network shown the TFs in

Figure 6B (**B**) and genes of the erythrocyte differentiation functional term from A and

Figure 6D (**C**). **D.** The β-chain globin expression profile. **E.** qPCR of cell adhesion

molecules and the effects of estradiol throughout erythroid cell maturation.

914
915   **Table S1** Heatmap genes for Figure S1B and functional term for Figure S1C

916
917   **Table S2** Heatmap genes for Figure 2C

918
919   **Table S3** Functional terms for Figure S3, 3F and S5E

920
921   **Table S4** Genes used for heatmap of subclusters 0a, 0b, 0c, 1a, 1b, 4a, 4b, 8a and 8b

922
923   **Table S5** Genes used for Figure 4C, 5A, 5C, 5F and 6B

924
925   **Table S6** Functional terms for Figure 5A, 5C, 5F and 6B

926
927   **Table S7** qPCR primer for globin, cell adhesion and estradiol related genes

Cluster
0 1 2 3 4 5 6 7 8 9

Cluster 0 (pink, top left):
- 28 Neutrophil degranulation
- 28 Neutrophil activation involved in immune response
- 28 Granulocyte activation
- 28 Myeloid leukocyte mediated immunity
- 30 Regulated exocytosis
- 11 Defense response to bacterium
- 13 Cytokine production
- 6 Neutrophil chemotaxis
- 6 Granulocyte chemotaxis
- 6 Granulocyte migration

x-axis: 0 5 10 15 20 25 30 35 40

Cluster 1 (gold, top right):
- 10 Focal adhesion
- 11 PI3K-Akt signaling pathway
- 6 Signaling by PDGF
- 4 Chondrocyte morphogenesis
- 6 Organ growth
- 9 Cell morphogenesis involved in differentiation
- 8 Tissue morphogenesis
- 5 Skeletal system morphogenesis
- 7 Cellular response to growth factor stimulus
- 3 Gastrulation

x-axis: 0 5 10 15 20 25

Cluster 2 (olive, second left):
- 15 Extracellular structure organization
- 10 Tissue morphogenesis
- 10 Cell morphogenesis involved in differentiation
- 10 Supramolecular fiber organization
- 9 Blood vessel morphogenesis
- 4 Mesoderm formation
- 6 Muscle tissue morphogenesis
- 4 Regulation of cell adhesion
- 4 Actin-mediated cell contraction
- 3 Cellular response to growth factor stimulus

x-axis: 0 2 4 6 8 10 12 14

Cluster 3 (green, second right):
- 8 Connective tissue development
- 9 Regulation of cytoskeleton organization
- 6 Cytoskeleton-dependent intracellular transport
- 5 Positive regulation of binding
- 3 Regulation of protein localization to cell surface
- 3 Establishment or maintenance of apical/basal cell polarity
- 8 Regulation of cell cycle process
- 7 Cell division
- 5 Reproductive structure development
- 4 Extracellular matrix organization

x-axis: 0 2 4 6 8 10

Cluster 4 (cyan, third left):
- 9 Protein targeting to ER
- 9 Eukaryotic translation initiation
- 9 Ribosome
- 8 Homeostasis of number of cells
- 8 Lymphocyte differentiation
- 10 Regulation of cytokine production
- 6 Regulation of hemopoiesis
- 3 Erythrocyte differentiation
- 3 Hematopoietic progenitor cell differentiation
- 4 Myeloid cell differentiation

x-axis: 0 2 4 6 8 10 12 14 16

Cluster 5 (turquoise, third right):
- 7 Endothelium development
- 6 Cell adhesion moleculesnt
- 7 Regulation of angiogenesis
- 4 Regulation of cell junction assembly
- 3 Regulation of endothelial cell differentiation
- 5 Endothelial cell migration
- 5 Cell-substrate adhesion
- 3 Vascular smooth muscle contraction
- 3 cAMP-mediated signaling
- 3 Rap1 signaling pathway

x-axis: 0 2 4 6 8 10 12 14

Cluster 6 (cyan, fourth left):
- 7 Integrin cell surface interactions
- 5 Signaling by PDGF
- 8 Extracellular matrix organization
- 6 Focal adhesion
- 7 PI3K-Akt signaling pathway
- 7 Signaling by receptor tyrosine kinases
- 6 Blood vessel development
- 4 Regulation of Wnt signaling pathway
- 3 Response to glucose
- 4 Response to oxygen levels

x-axis: 0 2 4 6 8 10 12

Cluster 7 (blue/purple, fourth right):
- 14 Regulated exocytosis
- 5 Hydrogen peroxide catabolic process
- 4 Oxygen transport
- 5 Detoxification
- 5 Protein heterooligomerization
- 6 Reactive oxygen species metabolic process
- 7 Response to toxic substance
- 7 Protein complex oligomerization
- 7 Regulation of protein kinase activity
- 6 Cytokine-mediated signaling pathway

x-axis: 0 2 4 6 8 10 12 14

Cluster 8 (magenta, bottom left):
- 11 Cellular response to molecule of bacterial origin
- 12 Myeloid leukocyte activation
- 6 I-kappaB kinase/NF-kappaB signaling
- 4 Mononuclear cell migration
- 4 Interleukin-1 production
- 6 Activation of innate immune response
- 6 Phagocytosis
- 5 Cytokine secretion
- 4 Interleukin-6 production
- 3 Monocyte chemotaxis

x-axis: 0 2 4 6 8 10 12

Cluster 9 (magenta/pink, bottom right):
- 8 Kidney development
- 6 Morphogenesis of embryonic epithelium
- 10 Epithelial cell differentiation
- 9 Tissue morphogenesis
- 8 Embryonic morphogenesis
- 7 Regulation of cell adhesion
- 6 Sensory organ development
- 4 Regulation of animal organ morphogenesis
- 3 Limb morphogenesis
- 6 Response to growth factor

x-axis: 0 2 4 6 8

y-axis (left): Number of Gene

x-axis (both bottom): -log10(*p* value)