Information content of cheek and lip colour in relation to the timing of ovulation in women

Lucie Rigaill [1]

[1] Kyoto University Primate Research Institute KUPRI, Inuyama, Japan

lucie.rigaill@gmail.com

Abstract

Various animal species have evolved a sexual communication system with female displaying

and male discriminating information about the timing of ovulation through sexual signals. More

research is now investigating the potential ovulatory signalling function of female red skin

colour in human and non-human primates. However, to date it is still challenging to draft

satisfying hypotheses of the evolution and function of female red skin colour, due to

methodological discrepancies between human and non-human primate studies. The present

study used a within-individual design and objective methods to analyse the relationship

between fine-scale variation in cheek and lip colour (luminance and redness) and the estimated

day of ovulation in 15 cycling women. Lip, but not cheek, colour, appeared to contain

information about the timing of ovulation, with lips getting darker around the timing of

ovulation. This study adds to the growing evidence that female red skin colour may play a role

in sexual signalling in human and non-human primates but also underlines variation in trait

forms and functions at the species-level.

Keywords: Red skin colour, Information content, Ovulatory signalling, Lips, Humans, Primates

Introduction

Do females display visual information about their reproductive status that can modulate mate attraction and mating strategies? This question is central to sexual selection theory. The number of possible offspring by females is constrained by physiological factors such as menstrual cyclicity, gestation and post-partum amenorrhea, and reproductive senescence. Reproduction is usually more costly for females due to the production of larger gametes, extended maternal care, and male monopolization [1,2]. Thus, in several species, females appear to have evolved traits which are attractive to males and can act as probabilistic signals of ovulation to maximize reproduction while balancing its associated costs. Across human and non-human primates (hereafter, primates), there is evidence that male behaviour is modulated by female traits, suggesting common evolutionary pathways and underlying mechanisms for sexual signalling [3,4].

Among the different female traits, there is a growing interest in the potential role of red skin colour in primate sexual communication. Intra-cycle variation in oestrogens induces ovulation and also affects some chromatic (redness) and achromatic (luminance) parameters of skin colour [5,6]. Circulating oestrogens bind to receptors in the skin, causing an increase in blood flow and consequently a decrease in the perceived skin luminance (i.e., darkening of the skin) [7,8]. Increase in blood flow can modulate the ratio of oxygenated/deoxygenated blood in the skin vessels which may influence perceived redness. Primate studies provided evidence that facial skin colour varies across the cycle in mandrills [9], informs about the probability of ovulation in rhesus macaques [10,11], and simultaneously conceals ovulation and advertises pregnancy in Japanese macaques [12,13]. Moreover, this colourful trait appears to be attractive to males suggesting a role in mate attraction, at least in macaques [14].

Skin and lip colour also correlate with female attractiveness in human [15–18]. Skin colour may thus be involved into human mate attraction, although the colour of the stimuli was

artificially manipulated in these studies which may impair the ecological validity of the results.

Studies of the potential signalling function of red skin colour in women yielded mixed results.

Oberzaucher et al. [19] described that cheeks were redder around ovulation compared to the

end of the cycle while Samson et al [20] did not replicate these findings. Recently, Burris et al.

[21] found no intra-cycle variation in cheek colour. Interpretation of these findings is

constrained by some methodological limitations. While most studies used photography to

analyse skin colour, which provides accurate measurements, they did not always correct

lighting conditions across photography sessions using colour standards that may have altered

measurements [22,23]. Human studies usually rely on a restricted data set to assess intra-cycle

difference: e.g. 2 samples from mid and late cycle, which underestimate or overestimate cycle

effects, with the exception of [21]. Finally, human studies often failed to confirm ovulation with

sex hormones profiles (with the exception of [21]) especially when studying perception of intra-

cycle variation. Taken together, these methodological limitations constrain our understanding

of the potential role of female red skin colour in sexual signalling in humans.

To determine if women display a colourful trait that may play a role in ovulatory

signalling, there is a fundamental need for studies of the relationship between fine-scale

variation in trait expression and the probability of ovulation. Such studies would benefit from

objective and quantitative methods, inspired by primate studies, using standardized

photography, regular biological sampling, and hormonal estimation of the ovulation date [23–

26]. Following this premise, this study aims at investigating whether cheek and lip colour

contain information about intra-cycle variation in the probability of ovulation in women. If

cheek or lip colour is related to the timing of ovulation within a cycle, I predict that cheeks or

lips will be darker/redder around this timing.

Methods

Ethic statement

This study was approved by the Human Research Ethics Committee of Kyoto University

Primate Research Institute (KUPRI, project number 2017-13). All participants signed a written

agreement concerning their participation in the study, confidentiality, and the use of their

photographs for research purposes and illustrations. Participants received a financial

compensation in return for their participation (5,000 JPY ~ 45 USD).

Participants

In total, 18 women participated in the study (mean age = 28.3 ± SD 4.3 years, "Asian" = 11,

"Caucasian" = 7, additional information on the supplementary material). All participants were

naturally cycling and not taking hormonal contraceptives for at least 3 months. I collected

digital photographs and saliva samples every two days, excluding weekends and national

holidays, for the duration of one complete menstrual cycle (i.e., from the beginning of their

menstruations until the next ones) between September and June 2018-19, to limit the effect of

tanning. Sampling occurred at a fixed time of the day for each participant to control for possible

diurnal variation in sex hormones concentration [27]. I collected a total of 204 photographs and

saliva samples (mean par participant = 11.3 ± SD 1.5, range = 8-13).

Assessment of cheek and lip colour

Participants removed face and lip make-up a minimum of 30 min before sampling. A single

female experimenter (LR) photographed the participants, as the sex of the photographer can

influence participant facial temperature and potentially skin colour [28]. Sampling was

conducted in a unique location, i.e., an office room with closed curtains, ceiling lights turned

on, and a constant 24° C temperature. Participants sat in front of a beige background and

adopted a neutral expression. The camera was place 2m from the participant's chair and held at

the same height of the participant's face. The experimenter used a Nikon 5000D camera with a

12.3 megapixel CMOS censor and a Nikon AF-S NIKKOR 18-55mm f/3.5-5.6G lens.

Photographs were collected in NEF format, i.e., Nikon's RAW, with the flash disabled, the

shutter speed and aperture size determined automatically by the camera, and a 55mm focal to

limit distortion [29]. The experimenter manually set the white balance using a X-Rite White

Balance Card (GretagMacbeth ColorChecker) and a Gretag X-Rite Color Checker (colour

standards). This technique standardizes photographs such that colour measurements are

comparable across all photographs [23–25]. Photographs were converted to uncompressed 16-

bit TIFF files. Cheek colour was measured from a pair of points located on the area described

in [21]. Lip colour was measured from the whole lip skin (supplementary material, Figure S1).

CIELAB values were extracted using Colourworker software (Chrometics Ltd. available at:

http://www.colourworker.com/) which provided luminance (L*) and red-green ratio (a*,

hereafter redness) values calculated from the estimated reflectance spectrum according to the

standard CIE (Commission Internationale de l'Éclairage) equations [30].

Determination of the estimated ovulation

Participants followed the sampling guideline recommended by Salimetrics (Salimetrics, LLC,

USA). Samples were collected during the photo session, labelled and stored within 15 min at -

80°C until analyses at KUPRI. Saliva samples were analysed for oestradiol (17ß-estradiol) and

progesterone (4-pregenene-3,20-dione) using Salimetrics enzyme immunoassays kits. The

oestradiol assay had a sensitivity of 0.1 pg/ml with an intra-assay coefficient of variation (CV)

of 6.2 % and inter-assay CV of 10.8 %. The progesterone assay had a sensitivity of 5 pg/ml

with an intra-assay CV of 6.6 % and inter-assay CV of 9.7 %.

The onset of the luteal phase was defined as the sample with a progesterone

concentration at least 2 standard deviations greater than the mean of the 2-3 preceding baseline

values [26]. Ovulation was considered to have occurred when a mid-cycle peak of oestradiol

was detected around the onset of the luteal phase. Because sampling occurred every 2-3 days,

I considered the day of peak oestradiol as the most likely day of ovulation and labelled it as *day*

0. The day directly preceding the estimated ovulation day was labelled as *day -1*, the day

directly following it as *day +1*, and so on. Ovulation could not be determined (i.e., abnormal

hormone variations) for 3 out of the 18 participants.

Statistical analyses

I used photographs of 15 participants showing ovulatory cycles to analyse intra-cycle variation

in cheek and lip luminance and redness (N = 165 photographs, mean per participant = 11.0 ±

SD 1.4, range = 8-13).

This study tests for a possible quadratic effect of the days relative to ovulation on cheek

and lip colour: higher effect toward the estimated ovulation and lower effect toward the

beginning and end of the cycle. I thus constructed general linear mixed-effects models (LMMs)

for cheek and lip colour (luminance and redness respectively) which included the linear (DAY)

and quadratic (DAY^2) effects of the days relative to the estimated ovulation as fixed effects,

and fitted by maximum likelihood using lme4 [31] and lmerTest [32] packages in R version

3.6.0 [33]. Inspection of the cumulative distribution functions revealed good fits to the normal

distribution for luminance and to the lognormal distribution for redness. Prior to modelling, the

days relative to the estimated ovulation were standardized to mean = 0 and SD = 1 to improve

model performance and interpretability [34]. On one hand, we can expect that the relationship

between the timing of ovulation and colour would be similar across participants; I thus

constructed random intercept models (RIM) with random intercepts for participant identity. On

the other hand, this relationship may vary between participants; I thus constructed uncorrelated

random slope models (RSM) with random slopes for the linear and quadratic terms. I also

constructed the respective RIM and RSM null models in which the predictor variables were

removed but the random effect structure was maintained. This resulted in 4 candidate models

for each colour/skin combination: RIM-full, RIM-null, RSM-full, and RSM-null. I ensured that

all relevant model assumptions were met by visually inspecting histograms of the residuals and

plots of residuals against fitted values. I used an information-theory approach to objectively

compare and rank the 4 candidate models in terms of how well they fitted the existing data thus

assessing the likelihood that one or more models among the candidates is/are best supported by

the data [35,36]. I used the function model.sel of the MuMIn package [37] to rank models based

on the Akaike's information criterion corrected for small sample size (AICc values). I reported

the weight of the models which indicates to what extent one candidate model is more likely

than another to provide a reasonable explanation of the variance in the data. I extracted weighted

parameter estimates (β), standard errors (s.e.), and 95% confidence intervals (95% CIs) of

model intercepts and predictors from conditional averaging of the 4 candidate models (function

model.avg).

Results

RIMs generally performed better than their respective RSMs (Table 1). Concerning cheek

colour, the RIM-null for luminance had a weight of 0.75, while the RIM-full for redness had a

weight of 0.51 (vs. 0.37 for its RIM-null), suggesting some weak evidence for variation in cheek

redness only (Table 1). Concerning lip colour, the RIM-full for luminance had a weight of 0.53

(vs. 0.31 for RIM-null) and the RIM-null for redness had a weight of 0.50 (vs. 0.37 for its RIM-

null), suggesting some weak evidence for variation in lip luminance only (Table 1). The timing

of ovulation explained little to none of the variance in cheek redness (Table 2, Figure 1). Lip

luminance varied according to the timing of ovulation with lips being darker around ovulation

(ß = 0.21 ± 0.09 s.e., 95% CI = 0.03; 0.39, Table 2, Figure 2).

Table 1. Characteristics of candidate models for cheek and lip redness and luminance. AICc:

corrected Akaike's information criterion, weight: model probabilities

	logLik	AICc	weight		logLik	AICc	weight
Cheek luminance				*Cheek redness*			
RIM - null	-247.751	501.7	0.75	RIM - full	279.799	-549.2	0.51
RIM - full	-247.292	505.0	0.14	RIM - null	277.351	-548.6	0.37
RSM - null	-247.751	505.9	0.09	RSM - full	279.799	-544.9	0.06
RSM - full	-247.292	509.3	0.02	RSM - null	277.621	-544.9	0.06
Lip luminance				*Lip redness*			
RIM - full	-272.607	555.6	0.53	RIM-null	221.984	-437.8	0.60
RIM - null	-275.332	556.8	0.30	RIM-full	223.354	-436.3	0.29
RSM - null	-274.316	559.0	0.10	RSM-null	222.055	-433.7	0.08
RSM - full	-272.345	559.4	0.08	RSM-full	223.356	-432.0	0.03

175　**Table 2.** Model averaged parameters estimates (ß) ± standard errors (s.e.) and confidence

176　intervals (95% CIs) from conditional averaging of all candidate models. Result for which CI

177　does not include zero are presented in bold.

	Intercept	DAY	DAY^2
Cheek luminance	26.99 ± 0.50 (**26.01; 27.97**)	-0.06 ± 0.07 (-0.21; 0.08)	0.03 ± 0.07 (-0.11; 0.17)
Cheek redness	2.05 ± 0.02 (**2.01; 2.10**)	0.00 ± 0.00 (-0.00; 0.01)	0.01 ± 0.00 (-0.00; 0.01)
Lip luminance	20.77 ± 0.32 (**20.13; 21.40**)	0.01 ± 0.09 (-0.18; 0.19)	0.21 ± 0.09 (**0.03; 0.39**)
Lip redness	2.20 ± 0.03 (**2.14; 2.27**)	0.01 ± 0.00 (-0.00; 0.01)	0.00 ± 0.00 (-0.00; 0.01)

179　**Figure 1**. Relationship between cheek (a) luminance and (b) redness and the days relative to

180　the estimated ovulation. Raw data are presented as grey circle. The red line presents the global

181　model prediction; the black lines show prediction for each participant.

Figure 2. Relationship between lip (a) luminance and (b) redness and the days relative to the estimated ovulation. Raw data are presented as grey circle. The red line presents the global model prediction; the black lines show prediction for each participant.

Discussion

Using objective methods to analyse fine-scale and intra-individual changes in cheek and lips colour according to the timing of ovulation, I found that lip luminance contains information about the timing of ovulation in women. Cheek colour may not be related to the timing of ovulation in agreement with a previous study [21].

Different facial features may respond differently to variation in the probability of ovulation. Intra-cycle variation in colour may be less cryptic for the lip skin as a result of its higher vascularization and the finer cellular layers [38] or due to tissue-specific differences in

oestrogen receptor expression or sensitivity. The latter has been already suggested to explain

the differential effect of cycle phase on facial and hindquarters colour in Japanese macaques

[13]. Further studies on tissue-specific expression and sensitivity of oestrogen receptors would

help to clarify this question. It is also possible that hormonal contraceptives have on-going and

long-term effect on red skin colour (e.g., female odours [39,40]), a question that I could not

assess in the present study and requires further investigation. Alternatively, colour changes in

the facial skin may be more condition-dependent and related to individual health thus masking

a potential cycle effect [15,16,41].

So, can lip colour act as an ovulatory signal? The present study suggest that lip colour

contains information about the timing of ovulation. To determine whether this information is

conveyed, further studies should assess men and women responses toward intra-cycle variation

in lip colour to determine whether these changes are perceptible. However, caution should be

used when designing such experiments. Studies should use an intra-individual design, i.e., using

individual pictures rather than pictures from different participants or composite stimuli, as the

differences in skin colour appeared to be relatively greater between than within participant in

the present study. Studies should also collect and test multiple samples per individual in order

to detect fine-scale variation, and confirm ovulation using hormonal data or ovulation detection

kit. Such experimental design should reduce the risk of over- or under-estimating a potential

cycle effect. However, whether variation in this trait is detectable under "optimal" conditions,

i.e., artificial environments and controlled settings, does not necessarily entail consequences on

human sexual communication and mate preferences since laboratory studies can have low

ecological validity. At best, studies on perception could provide further evidence that lip and/or

cheek colour has the potential to play a role in ovulatory signalling.

Humans may have inherited the biological bases for female red skin colour and sexual

signalling from a primate ancestor. Female red skin colour has been suggested to play a role

into the sexual communication of some primate species, i.e., signalling [9–11] or concealing

ovulation [13], advertising pregnancy [9,12]. The differences in red skin traits (face, lips, and

sexual skin) and signalling functions probably result from the different socio-environmental

constraints on mating across species. Concealing the reproductive status may have evolved in

species facing higher male monopolization or pair-bonding, lower intra-sexual competition, and

higher costs on signalling; while higher infanticide risks and intra-sexual competition, limited

mating opportunities, and lower costs on signalling may have favoured exaggerated or

multimodal signalling [42,43]. However, these proposed effects do not appear to fully explain

the inter-species differences observed in female red skin colour function, as species with similar

socio-ecology express traits that may be involved in either ovulatory signalling or concealing

(e.g. [11,13]). More studies using longitudinal and ecologically valid designs are needed to

better understand the possible evolutionary pathways and underlying mechanisms leading to

the evolution of female red skin colour in primate species including humans.

235
Acknowledgments

I acknowledge the Human Research Ethics Committee of Kyoto University Primate Research

Institute for permission to conduct my research. I would like to sincerely thank C. Neumann

for his insightful comments on the study and his valuable advice with statistical analyses, J.

Duboscq and C. Garcia for comments on an earlier version of the manuscript, and K. Mouri for

her general support in the lab.

Funding

LR is funded by the Japanese Society for the Promotion of Science (project 18H05810 JSPS).

References

1.   Galef BG. 1983 Costs and Benefits of Mammalian Reproduction. In *Symbiosis in Parent-Offspring Interactions* (eds LA Rosenblum, H Moltz), pp. 249–278. Boston, MA: Springer US. (doi:10.1007/978-1-4684-4565-7_11)

2.   Gittleman JL, Thompson SD. 1988 Energy Allocation in Mammalian Reproduction. *Integr Comp Biol* **28**, 863–875. (doi:10.1093/icb/28.3.863)

3.   Campbell C. 2007 Primate sexuality and reproduction. In *Primates in perspective*, pp. 423–437. New York: Campbell CJ, Fuentes A, MacKinnon KC, Panger M, Bearder SK.

4.   Dixson AF. 2012 *Primate sexuality: comparative studies of the prosimians, monkeys, apes and human beings. 2nd edition*. Oxford University Press. New York.

5.   Rhodes L, Argersinger ME, Gantert LT, Friscino BH, Hom G, Pikounis B, Hess DL, Rhodes WL. 1997 Effects of administration of testosterone, dihydrotestosterone, oestrogen and fadrozole, an aromatase inhibitor, on sex skin colour in intact male rhesus macaques. *Journal of Reproduction and Fertility* **111**, 51–57. (doi:10.1530/jrf.0.1110051)

6.   Thornton MJ. 2002 The biological actions of estrogens on skin. *Experimental Dermatology* **11**, 487–502. (doi:10.1034/j.1600-0625.2002.110601.x)

7.   Pelletier G, Ren L. 2004 Localization of sex steroid receptors in human skin. *Histol. Histopathol.* **19**, 629–636. (doi:10.14670/HH-19.629)

8.   Thornton Mj, Taylor AH, Mulligan K, Al-Azzawi F, Lyon CC, O'Driscoll J, Messenger AG. 2003 The Distribution of Estrogen Receptor β Is Distinct to That of Estrogen Receptor α and the Androgen Receptor in Human Skin and the Pilosebaceous Unit. *Journal of Investigative Dermatology Symposium Proceedings* **8**, 100–103. (doi:10.1046/j.1523-1747.2003.12181.x)

9.   Setchell JM, Wickings EJ, Knapp LA. 2006 Signal content of red facial coloration in female mandrills (*Mandrillus sphinx*). *Proceedings of the Royal Society B: Biological Sciences* **273**, 2395–2400. (doi:10.1098/rspb.2006.3573)

10.   Dubuc C, Brent LJN, Accamando AK, Gerald MS, MacLarnon A, Semple S, Heistermann M, Engelhardt A. 2009 Sexual skin color contains information about the timing of the fertile phase in free-ranging *Macaca mulatta*. *International Journal of Primatology* **30**, 777–789. (doi:10.1007/s10764-009-9369-7)

11.   Higham JP, Brent LJN, Dubuc C, Accamando AK, Engelhardt A, Gerald MS, Heistermann M, Stevens M. 2010 Color signal information content and the eye of the beholder: a case study in the rhesus macaque. *Behavioral Ecology* **21**, 739–746. (doi:10.1093/beheco/arq047)

12.   Rigaill L, MacIntosh AJJ, Higham JP, Winters S, Shimizu K, Mouri K, Furuichi T, Garcia C. 2015 Multimodal advertisement of pregnancy in free-ranging female Japanese macaques (*Macaca fuscata*). *PLOS ONE* **10**, e0135127. (doi:10.1371/journal.pone.0135127)

13.   Rigaill L, Higham JP, Winters S, Garcia C. 2019 The redder the better? Information content of red skin coloration in female Japanese macaques. *Behavioral Ecology and Sociobiology* **73**, 103. (doi:10.1007/s00265-019-2712-x)

14.   Higham JP, Winters S. 2015 Color and mate choice in non-human animals. In *Handbook of Color Psychology*, pp. 502–530. Cambridge: A. Elliot, M. Fairchild, & A. Franklin.

15.   Re DE, Whitehead RD, Xiao D, Perrett DI. 2011 Oxygenated-Blood Colour Change Thresholds for Perceived Facial Redness, Health, and Attractiveness. *PLOS ONE* **6**, e17859. (doi:10.1371/journal.pone.0017859)

16.   Pazda AD, Thorstenson CA, Elliot AJ, Perrett DI. 2016 Women's Facial Redness Increases Their Perceived Attractiveness: Mediation Through Perceived Healthiness. *Perception* **45**, 739–754. (doi:10.1177/0301006616633386)

17.   Young SG. 2015 The effect of red on male perceptions of female attractiveness:
Moderation by baseline attractiveness of female faces. *European Journal of Social*
Psychology **45**, 146–151. (doi:10.1002/ejsp.2098)
18.   Stephen ID, McKeegan AM. 2010 Lip colour affects perceived sex typicality and
attractiveness of human faces. *Perception* **39**, 1104–1110. (doi:10.1068/p6730)
19.   Oberzaucher E, Katina S, Schmehl SF, Holzleitner IJ, Mehu-Blantar I, Grammer K.
2012 The myth of hidden ovulation: Shape and texture changes in the face during the
menstrual cycle. *Journal of Evolutionary Psychology* **10**, 163–175.
(doi:10.1556/JEP.10.2012.4.1)
20.   Samson N, Fink B, Matts P. 2011 Does a woman's skin color indicate her fertility
level? Preliminary findings. *Swiss Journal of Psychology* **70**, 199–202. (doi:10.1024/1421-
0185/a000057)
21.   Burriss RP, Troscianko J, Lovell PG, Fulford AJC, Stevens M, Quigley R, Payne J,
Saxton TK, Rowland HM. 2015 Changes in Women's Facial Skin Color over the Ovulatory
Cycle are Not Detectable by the Human Visual System. *PLOS ONE* **10**, e0130093.
(doi:10.1371/journal.pone.0130093)
22.   Rowland HM, Burriss RP. 2017 Human colour in mate choice and competition.
Philosophical Transactions of the Royal Society B: Biological Sciences **372**.
(doi:10.1098/rstb.2016.0350)
23.   White T, Dalrymple R, Noble D, O'Hanlon J, Zurek D, Umbers K. 2015 Reproducible
research in the study of biological coloration. *Faculty of Science, Medicine and Health -*
Papers , 51–57. (doi:10.1016/j.anbehav.2015.05.007)
24.   Bergman TJ, Beehner JC. 2008 A simple method for measuring colour in wild
animals: validation and use on chest patch colour in geladas (*Theropithecus gelada*).
Biological Journal of the Linnean Society **94**, 231–240. (doi:10.1111/j.1095-
8312.2008.00981.x)
25.   Stevens M, Stoddard MC, Higham JP. 2009 Studying primate color: towards visual
system-dependent methods. *International Journal of Primatology* **30**, 893–917.
(doi:10.1007/s10764-009-9356-z)
26.   Hodges JK, Heistermann M. 2011 Field endocrinology: monitoring hormonal changes
in free-ranging primates. In *Field and laboratory methods in primatology: a practical guide*,
Joanna M. Setchell and Deborah J. Curtis.
27.   Bao AM, Liu RY, Someren E van, Hofman MA, Cao YX, Zhou JN. 2003 Diurnal
rhythm of free estradiol during the menstrual cycle. *European Journal of Endocrinology* **148**,
227–232. (doi:10.1530/eje.0.1480227)
28.   Hahn AC, Whitehead RD, Albrecht M, Lefevre CE, Perrett DI. 2012 Hot or not?
Thermal reactions to social contact. *Biology Letters* , rsbl20120338.
(doi:10.1098/rsbl.2012.0338)
29.   Třebický V, Fialová J, Kleisner K, Havlíček J. 2016 Focal Length Affects Depicted
Shape and Perception of Facial Images. *PLOS ONE* **11**, e0149313.
(doi:10.1371/journal.pone.0149313)
30.   Vorobyev M, Gumbert A, Kunze J, Giurfa M, Menzel R. 1997 Flowers Through
Insect Eyes. *Israel Journal of Plant Sciences* **45**, 93–101.
(doi:10.1080/07929978.1997.10676676)
31.   Bates D, Mächler M, Bolker B, Walker S. 2015 Fitting linear mixed-effects models
using lme4. *Journal of Statistical Software* **67**. (doi:10.18637/jss.v067.i01)
32.   Kuznetsova A, Brockhoff PB, Christensen RHB. 2017 lmerTest Package: Tests in
Linear Mixed Effects Models. *Journal of Statistical Software* **82**, 1–26.
(doi:10.18637/jss.v082.i13)
33.   R Core Team. 2019 R: A language and environment for statistical computing. See

https://www.R-project.org/.
34.   Harrison XA, Donaldson L, Correa-Cano ME, Evans J, Fisher DN, Goodwin CED,
Robinson BS, Hodgson DJ, Inger R. 2018 A brief introduction to mixed effects modelling and
multi-model inference in ecology. *PeerJ* **6**, e4794. (doi:10.7717/peerj.4794)
35.   Burnham KP, Anderson DR, Huyvaert KP. 2011 AIC model selection and multimodel
inference in behavioral ecology: some background, observations, and comparisons. *Behav*
Ecol Sociobiol **65**, 23–35. (doi:10.1007/s00265-010-1029-6)
36.   Symonds MRE, Moussalli A. 2011 A brief guide to model selection, multimodel
inference and model averaging in behavioural ecology using Akaike's information criterion.
Behav Ecol Sociobiol **65**, 13–21. (doi:10.1007/s00265-010-1037-6)
37.   Bartoń K. 2017 MuMIn: Multi-Model Inference. R package version 1.40.0.
38.   Crouzet C, Fournier H, Papon X, Hentati N, Cronier P, Mercier Ph. 1998 Anatomy of
the arterial vascularization of the lips. *Surgical and Radiologic Anatomy* **20**, 273–278.
(doi:10.1007/BF01628490)
39.   Crawford JC, Boulet M, Drea CM. 2010 Smelling wrong: hormonal contraception in
lemurs alters critical female odour cues. *Proceedings of the Royal Society of London B:*
Biological Sciences , rspb20101203. (doi:10.1098/rspb.2010.1203)
40.   Kuukasjärvi S, Eriksson CJP, Koskela E, Mappes T, Nissinen K, Rantala MJ. 2004
Attractiveness of women's body odors over the menstrual cycle: the role of oral
contraceptives and receiver sex. *Behavioral Ecology* **15**, 579–584.
(doi:10.1093/beheco/arh050)
41.   Thorstenson CA, Pazda AD, Elliot AJ, Perrett DI. 2017 Facial Redness Increases
Men's Perceived Healthiness and Attractiveness. *Perception* **46**, 650–664.
(doi:10.1177/0301006616680124)
42.   Rooker K, Gavrilets S. 2018 On the evolution of visual female sexual signalling.
Proceedings of the Royal Society B: Biological Sciences **285**, 20172875.
(doi:10.1098/rspb.2017.2875)
43.   Fitzpatrick CL, Servedio MR. 2018 The evolution of male mate choice and female
ornamentation: a review of mathematical models. *Curr Zool* **64**, 323–333.
(doi:10.1093/cz/zoy029)
374