Characterization of transcription activation domain of EcaICE1 and its interaction with EcaSIZ1 in *Eucalyptus camaldulensis*

Ziyang Zhang[1,2*], Ling Cheng[3*], Weihua Zhang[4*], Jianlin Hu[1,2], Yan Liu[1,2], Yuanzhen LIN[1,2,§]

1 College of Forestry and Landscape Architecture, South China Agricultural University, Guangzhou 510642, China;

2 Guangdong Key Laboratory for Innovative Development and Utilization of Forest Plant Germplasm, Guangzhou 510642, China;

3 Guangzhou Genedenovo Biotechnology Company Limited, Guangzhou 510642, China;

4 Guangdong Academy of Forestry, Guangzhou 510640, China

[*] Ziyang Zhang, Ling Cheng and Weihua Zhang have contributed equally to this work.

[§]Corresponding author: Tel/Fax:86-020-85280256;E-mail:yzhlin@scau.edu.cn

Abstract: SUMOylation in plants is associated with biotic and abiotic stress responses, flowering and other

aspects of development, and ICE1 protein SUMOylation by SUMO E3 ligase SIZ1 plays important roles in

plant cold tolerance. Here, we reported the subcellular localization of EcaICE1 and its interaction with

EcaSIZ1 in *Eucalyptus camaldulensis*. The genes *EcaICE1* and *EcaSIZ1* were isolated by homologous

cloning. The subcellular localization analysis showed that EcaICE1 was located in nucleus . Bimolecular

fluorescence complementation (BiFC) analysis revealed that EcaICE1 could interact with EcaSIZ1 in the

nucleus of *Nicotiana benthamiana* leaves. Moreover, yeast two-hybrid assay confirmed that the amino acid

region from position 84 to 126 in EcaICE1 was critical for the strong transactivation activity of EcaICE1

and that the C terminal region from position 361 to 557 in EcaICE1 was the key region for its interaction

with EcaSIZ1 by using different truncated lengths of non-transactivation activity of EcaICE1 as the bait

protein. Collectively, our results showed that EcaICE1 may have a SUMOylation pathway similar to

Arabidopsis thaliana.

Key words: *Eucalyptus camaldulensis*; EcaICE1; EcaSIZ1; protein interaction; SUMOylation

Low temperature is one of the important environmental factors that could restrict the growth and

development, geographical distribution and production of plants, and low temperature stress could reduce

production in agriculture and forest (Janmohammadi et al. 2015). In order to overcome low temperature stress, a

series of complex response mechanisms can be activated at physiological and molecular levels for plants (Shi et

al. 2018). Researches reveal that a large number of cold-regulated genes (*CORs*) are induced after plant cold

acclimation. These *CORs* genes usually contain *DRE*/*CRT* cis-acting elements and can be combined by

transcription factors CBFs or DREB1s family (Stockinger et al. 1997). CBFs or DREB1s are key transcription

factors of low temperature signaling pathway and play important roles in enhancing plant cold tolerance

(Chinnusamy et al. 2007). However, *CBFs* or *DREB1s* are also induced by low temperature. ICE1 gene has been

firstly verified from *Arabidopsis thaliana*, which can specifically bind to the cis-acting element of MYC

promoter of *CBF3* and then induce the expression of CBF3 downstream gene (Chinnusamy et al. 2003). After

that, a lot of studies reveal that ICE-CBF-COR is the important signal network for plants to adapt to the cold

stress (Shi et al. 2018). In *Arabidopsis thaliana*, ICE-CBF-COR pathway is positively or negatively controlled

by some regulators at transcriptional, post-transcriptional and post-translational levels. SIZ1 (Miura et al. 2007)

and OST1 (Ding et al. 2015) are positive regulators, while HOS1 (Dong et al. 2006) and MPK3/6 (Li et al. 2017)

are negative regulators. The Ub E3 ligase HOS1, can ubiquitinate and degrade ICE1 protein, thereby increase the

cold sensitivity of the transgenic plants (Dong et al. 2006). On the contrary, SUMO E3 ligase SIZ1, can enhance

the stability of ICE1 protein by SUMOylation and promote the expression of CBF3, and then causes enhanced

cold tolerance (Miura et al. 2007). OST1 can interact with ICE1, competed with HOS1, enhancing the stability

and transcriptional activity of ICE1, further resulting in increased cold tolerance (Ding et al. 2015). Recently, the

protein kinase MPK3/6 can interact with and phosphorylate ICE1, enhancing the degree of ubiquitination of

ICE1, but the interaction between HOS1 and ICE1 is not affected, suggesting that the degradation of ICE1

protein may be performed by the unknown E3 ligase (Li et al. 2017). Although ICE1 gene had been isolated

from *A. thaliana* (Chinnusamy et al. 2003), *Brassica juncea* (Wang et al. 2005), *Populus suaveolens* (Lin et al.

2007), wheat (Badawi et al. 2008), banana (ZHAO et al. 2013), *E. camaldulensis* (Lin et al. 2014), *Pyrus*

ussuriensis (Huang et al. 2015) and *Hevea brasiliensis* (Deng et al. 2017), the understanding of its positive and

negative regulations is still limited in woody plants.

Eucalyptus species are the important commercial tree species worldwide. The molecular regulation

mechanism of low temperature stress of *Eucalyptus* could have the potential to increase *Eucalyptus* plantation

range. *Eucalyptus* CBF genes has been studied (Kayal et al. 2006; Gamboa et al. 2007; Navarro et al. 2009,

2011), but the knowledge about ICE1 or ICE1-like modules is still rather limited. Our precious studies have

shown that the overexpression of *EcaICE1* from *E. camaldulensis* can enhance the cold tolerance and the

expression level of downstream genes in transgenic tobaccos (Lin et al. 2014). However, the studies of

subcellular localization, transcription activity and upstream regulators of EcaICE1 remain uncertain. In this study,

we analyzed the subcellular localization and the transcription activation region of EcaICE1. Furthermore, the

protein interactions between EcaICE1 and EcaSIZ1 were verified by Bimolecular fluorescence complementation

(BiFC) and Yeast two-hybrid (Y2H) assays.

Methods and Materials

Plant materials

The 30-day-old rooting tissue culture plantlets of *E. camaldulensis* cv. 103 were used in this study. Tissue culture

condition was performed as described previously (Lin et al. 2014). Tobacco (*Nicotiana benthamiana*) plants

were maintained in a growth chamber of 25 ◦C under long day conditions (16/8 h light/dark photoperiod), and

5-week-old plants were selected for further subcellular localization and BiFC analysis.

RNA isolation and Gene cloning

Total RNA was extracted as described previously (Lin et al. 2014). 1 μg DNase I-treated RNA was used for

synthesizing the first strand cDNA according to the manufacturer's instructions (PrimeScript II 1st Strand

cDNA Synthesis Kit; Takara, Dalian, China). The primers of *EcaICE1* and *EcaSIZ1* (listed as table 1) were

designed by homology cloning techniques, and then used for amplifying the aim genes with the first strand

cDNA as the template. The PCR process as follows: 95 °C for 4 min, followed by 35 cycles of 95 °C for 10 s, 58

°C for 10 s and 72 °C for 20 s.

Bioinformatics analysis of gene sequence

The coding sequence (CDS) of *EcaICE1* and *EcaSIZ1* were predicted using FGENESH 2.6 software

(http://linux1.softberry.com/berry.phtml?topic=fgenesh&group=programs&subgroup=gfind), and further

confirmed by BLASTP program on the NCBI website (http://blast.ncbi.nlm.nih.gov/Blast.cgi). The nuclear

localization signals were predicted using ProtComp 9.0 (http://linux1.softberry.com/berry.phtml?topic=

protcomppl&group=programs&subgroup=proloc), and the protein secondary domains were searched by Motif

scan (https://myhits.isb-sib.ch/cgi-bin/motif_scan). Finally, sequence alignments with other plants were

performed with ClUSTALX software.

Subcellular localization of EcaICE1

92 The complete Open Reading Frame (ORF) of *EcaICE1* was recombinated to the 5′-terminus of the yellow

93 fluorescent protein (YFP) of the pEarleyGate101 vector with CaMV 35S promoter using primers (listed as table

94 S1). The recombinant vector *35S::EcaICE1-YFP* was transferred into *Agrobacterium tumefaciens* strain GV3101,

95 further injected into the abaxial surfaces of 5-week-old tobacco with an incubator for 2-3 d. The YFP signal in

96 the leaves of tobacco was examined using confocal microscopy (Olympus BX61, Tokyo, Japan).

Bimolecular fluorescence complementation (BiFC) assay

To generate the constructs for BiFC assays, the full-length CDS of *EcaICE1* and *EcaHOS1* (without their stop

codons) were subcloned into pUC-pSPYNE or pUCpSPYCE vectors as described previously (Walter *et al*.

2004). Expressions of target genes alone were used as negative controls. The resulting recombinant vectors were

used for transient assays of tobacco leaves as described earlier. The transformed tobacco leaves were then

incubated at 22 °C for 24–48 h. The YFP signal was observed using a florescence microscope (Zeiss Axioskop 2

plus). The primers for BiFC assay were listed in Supporting Information Table S2.

Yeast two-hybrid (Y2H) assay

Yeast two-hybrid assays were performed using the Matchmaker™ gold Yeast two-Hybrid Systems (Clontech).

Different truncated CDS of *EcaICE1* without transcriptional activation activity and EcaSIZ1 were subcloned into

pGBKT7 and pGADT7 vectors to fuse with the DNA-binding domain (DBD) and activation domain (AD),

respectively, to create different baits and preys (primers are listed in Supplemental table S3). Then, different

pairs of bait and prey recombinants were co-transformed into yeast strain gold Y2H by the lithium acetate

method, and yeast cells were grown on SD/-LW medium (minimal media double dropouts, SD medium with

-Leu/-Trp) according to the manufacturer's protocol (Clontech) for 3 days. Transformed colonies were plated

onto SD/-LWHA medium (minimal media quadruple dropouts, SD medium with-Leu/-Trp/-Ade/-His) containing

125-μM Aureobasidin A (AbA), to test for possible interactions among EcaICE1 and EcaSIZ1, according to the

yeast cell growth status.

Transcriptional activation assay

The full-length ORF of *EcaICE1* and *EcaSIZ1* were recombinated into pGBKT7 vector, respectively, and

co-transformed into yeast strain gold Y2H. Then we found that both EcaICE1 and EcaHOS1 had strong

transcriptional activation activity (results are listed in Supplemental Figure S4). So we further performed

transcriptional activation assay for EcaICE1. Different truncated coding regions of EcaICE1 (EcaICE1$_{T1\sim T6}$)

were recombinated into pGBKT7 vector, then followed Y2H assay, adding 0, 125, 250, and 500 μM AbA, to

further discover the critical region of transactivation activity for EcaICE1. The primers were listed in Supporting

Information Table S3.

RESULT

The gene amplification and sequencing analysis of EcaICE1 and EcaSIZ1

The multiple alignments of plant ICE1 protein sequences (Figure 1) shows that the *EcaICE1* protein has a highly

conserved MYC-like bHLH domain (basic helix-loop-helix) and a zipper structure at the C-terminus, and an

S-rich (Serine-rich) acidic domain at the N-terminus. In addition, the SUMO binding site is also present in

EcaICE1 and the other two *Eucalyptus* ICE1, and it is identical to ICE1 from *A. thaliana* and *Capsella*

buras-pastoris, but slightly different from the other woody plants. Furthermore ,the 363-381 position of EcaICE1

protein is a nuclear localization signal, as shown in the NLS（Nuclear localization sequence）box, which means

that it may be located in the nucleus. However, it is interesting that only *Eucalyptus* ICE1 proteins have a Q-rich

(Glutamine-rich) domain, suggesting that *Eucalyptus* ICE1 proteins might have different characteristics with the

other plants.

The amplified *EcaSIZE1* cDNA is 2847bp with a full ORF (2616 bp) encoding 872 amino acids. BLAST

analysis illustrates that EcaSIZ1 shares a high sequence identity with other plant SIZ1-like proteins, such as *E.*

grandis (99%, XP_010043707), *Ziziphus jujube* (72%, XP_015880489), *Malus domestica* (72%,

XP_008385287), *Citrus sinensis* (70%, XP_006488140), *P. trichocarpa* (71%, XP_024454498), and *A. thaliana*

(68%, OAO90655), respectively. The multiple alignments of plant SIZ1 protein sequences (Figure 2)

demonstrate that EcaSIZ1 protein has a conserved MIZ1/SP zinc finger domain, SAP, PINIT, SXS and

PHD, similar to other plant HOS1 proteins. These results show that EcaSIZ1 is the SIZ1 protein from *E.*

camaldulensis.

Subcellular localization of EcaICE1

Based on the multiple alignment analysis (Figure 1), there was one potential NLS sequence in the bHLH domain

of EcaICE1. To investigate the subcellular localization of EcaICE1 in plant cells, the full-length coding region of

EcaICE1 was fused to the N–terminus of the YFP reporter gene under the control of the 35S promoter

(*35S::EcaICE1-YFP*). By transient analysis using tobacco leaf epidermis, the yellow fluorescence of EcaICE1

fused-proteins was localized exclusively in the nucleus (Figure 3), indicating that EcaICE1 was a nuclear

localized protein, similar to other plant ICE1(Huang et al. 2015; Deng et al. 2017), consistent with the classical

role of bHLH as transcriptional regulators in plants (Gabriela et al. 2003).

EcaICE1 could interact with EcaSIZ1

Previous report showed that SIZ1 could interact with ICE1 to regulate its transcriptional activity in *A. thaliana*,

which plays an important role in response to cold stress (Miura et al. 2007). In order to investigate whether

EcaSIZ1 could also interact with EcaICE1 in *E. camaldulensis*, the protein-protein interactions between

EcaICE1 with EcaHOS1 were first analysed using BiFC assay in tobacco leaves. The YFP fluorescent signal was

observed at the nucleus when EcaICE1-pSPYCE was co-transfected with EcaHOS1-pSPYNE (Figure 4). In

contrast, no YFP fluorescent signal was observed in the negative controls including EcaICE1-pSPYCE

co-expressed with unfused pSPYNE or EcaHOS1-pSPYNE co-expressed with unfused pSPYCE. These results

showed that EcaICE1 could interact with EcaSIZ1 to form heterodimers at the nucleus (Figure 4).

Transcriptional activation domain of EcaICE1

To further discover the protein-protein interaction region between EcaICE1 with EcaHOS1, we performed the

Y2H assay. It was surprised that not only EcaICE1 but also EcaHOS1 had strong transactivation activity when

their full-length sequences were fused into the pGBKT7 vector and co-transfected with pGADT7 empty vector,

respectively (Figure S4). In order to remove the effects of strong transactivation activity on protein-protein

interaction, we further characterized which region of EcaICE1 acts as transcriptional transactivation, Y2H assays

were carried out using intact or truncated EcaICE1 as an effector (Figure 5A). The transfected yeast cells

harboring either the full-length EcaICE1 (pGBKT7-EcaICE1) or the truncated version (EcaICE1$_{T1}$(1-41a);

EcaICE1$_{T2}$(1-83a)) grew well on the selection medium SD/-LWHA, suggesting that the N-terminal 1-83 residues

are not necessary for transactivation activity of the EcaICE1. In contrast, when the N-terminal amino acids of the

other truncated version (EcaICE1$_{T3}$(1-125aa); EcaICE1$_{T4}$ (1-184aa); EcaICE1$_{T5}$(1-316aa); EcaICE1$_{T6}$(1-360aa))

were deleted, no interactions were detected (Figure 5B). Taken together, these results demonstrate that amino

acids from positions 84 to 125 in EcaICE1 are critical for the transactivation activity of EcaICE1, which contains

Q-rich domain (Figure 1), in agreement with transactivation domain mode in Sp1 (Courey et al. 1989).

Protein-protein interaction region between EcaICE1 with EcaSIZ1

Now the truncated EcaICE1 proteins without transcriptional activation activity (EcaICE1$_{T3-T6}$) were fused

to the GAL4 activation domain of vector pGBKT7 and co-transformed with pGADT7-EcaHOS1 into yeast strain

separately to exam the protein-protein interaction region between EcaICE1 with EcaHOS1. The results showed

(Figure 6) that all the truncated EcaICE1$_{T3}$ could interact with EcaHOS1, indicating that the C-terminus protein

of EcaICE1(361-557aa) is indispensable for its interaction with EcaHOS1 protein, similar to *A. thaliana* (Miura

et al. 2007).

DISCUSSION

It is reported that China is the world's second largest wood consumer and the first largest wood

importer (Lin 2019). At present, Chinese annual wood consumption has exceeded 600 million cubic meters,

and its wood demand has become more than 50% dependent on imports. *Eucalyptus*, the main trees of

man-made forest in South China even whole China, plays an important role in Chinese wooden market (Li

et al. 2017). However, low temperature is a primary disadvantage factor, which causes the biomass to fall,

and limits the development of its health (Janmohammadi et al. 2015). As the result, it is important to reveal

the molecular mechanism of cold stress in *Eucalyptus* for improving the cold tolerance on the inheritance

level as well as expanding the growing in northern region. In *Arabidopsis*, the cold signaling pathways,

ICE-CBF-COR pathway (Ding et al. 2015; Shi et al. 2018) and ICE-CBF pathway is positively correlated

with response to low temperature stress (Lin et al. 2014), had been studied clearly while the research of

subcellular localization, transcriptional activation and interactions with upstream regulators in *E.*

camaldulensis has still uncertain. Therefore, we further analyze the subcellular localization, transcriptional

activation of EcaICE1, and EcaSIZ1 (the homologous gene of SIZ1) was cloned to determine the protein

interactions between EcaICE1 and EcaSIZ1 as well as the key region by BiFC and yeast two-hybrid system

technology for exploring the SUMOylation pathway of EcaICE1.

According to the multiple comparison of EcaICE1 protein sequence, these is a nuclear localization

sequence signal in its bHLH-ZIP domain, indicated that EcaICE1 may be located in nucleus. Further, the

analysis of subcellular localization demonstrates that EcaICE1 is really located in nucleus, which is similar

to *P. ussuriensis* (Huang et al. 2015) and *H. brasiliensis* (Deng et al. 2017) as well as according with the

characteristics of transcription factor family of plant bHLH (Gabriela et al. 2003). Above all, EcaICE1,

located in nucleus, may play a regulatory role in the molecular mechanism of cold stress. What is more, we

further found that the EcaICE1 84th to 95th amino acid of N-terminal, which is the Glutamine-rich domain,

was identical with its homologous gene of ICE1 in *E. grandis* and *E. globulus*, and this domain only was

found in the ICE1 homologous protein in *Eucalyptus*. Interestingly, it was found that EcaICE1 protein had

strong transcriptional activation activity. Similar results were also reported on the ICE1 in *P. ussuriensis*

(Huang et al. 2015) and *H. brasiliensis* (Deng et al. 2017) while ICE1 protein did not have transcriptional

activation activity in *Arabidopsis*. The 46th to 95th and the 1th to 402th amino acid of N-terminal of

protein were verified that those are the key region of transcriptional activation activity in PuICE1 of *P.*

ussuriensis (Huang et al. 2015) and HbICE1 of *H. brasiliensis* (Deng et al. 2017), respectively.

Unfortunately, neither of them further explored the structural characteristics of the transcriptional activation

region. This study indicated that 84th to 125th amino acid of N-terminal of protein was the critical region

of transcriptional activation in EcaICE1 of *E. camaldulensis*, including the Q-rich domain as well as

according with the transcriptional activation structure of classical transcription factor Sp1 (Courey et al.

1989). It means that Q-rich domain may be the key transcriptional activation region of EcaICE1 protein.

SUMOylation is a posttranslational regulatory process in eukaryotes, including cold tolerance (Miura

et al. 2007), ABA response (Miura et al. 2009), flowering time (Bo et al. 2010) and drought tolerance

(Miura et al. 2013) and most of them are mainly devoting to *A. thaliana*. Studies in *A. thaliana* have shown

that the SUMOylation pathway is mainly mediated by the SUMO E3 ligase SIZ1. Currently, SIZ1 genes

have been isolated from plants such as rice (PARK et al. 2010), *Dendrobium nobile* (Liu et al. 2015), apple

(Zhang et al. 2016), soybean (Cai et al. 2017) and tomato (Zhang et al. 2017), but no study on cloning the

gene SIZ1 and its interaction with ICE1 in forest trees. The coding sequence of *EcaSIZ1*, which was

isolated in this study, is similar to the cloned SIZ1 protein in the above plants, and also has similar protein

secondary domains such as MIZ1/SP zinc finger domain, SAP domain, PINIT domain, SXS domain and

PHD domain. Among them, MIZ1/SP zinc finger domain is associated with SUMO E3 ligase activity, and

The SXS domain is associated with binding of SUMO (Miura et al. 2007). The SAP domain promotes

binding to DNA, PINIT domain is related to intracellular retention. The above plant SIZ1 proteins are

located in the nucleus and EcaSIZ1 protein should also have this conserved domain, so it indicated that

EcaSIZ1 may be located in the nucleus too. PHD domain is a plant homologous domain, which only exists

in SIZ1 proteins in plant (Park et al. 2010). Meanwhile, EcaSIZ1 protein shared more than 70% identity

with other woody plants at the SIZ1 homologous protein level, indicating that *EcaSIZ1* is the SIZ1 gene

from *E. camaldulensis*, and its encoded protein may have SUMO E3 ligase activity and mediate

SUMOylation in the nucleus.

BiFC assay showed that EcaICE1 and EcaSIZ1 had the protein interaction in plant nucleus, but which

region of EcaICE1 was the key region of protein interaction is still unknown. Therefore, using EcaICE1

with different truncated lengths without transcriptional activation activity as decoy protein, the 361th to

557th amino acid region at the C-terminal of EcaICE1 protein where was the key region of its interaction

with EcaSIZ1 was further identified by yeast hybrid technology. The protein sequence of EcaICE1 in this

region contained a SUMO domain, which was completely consistent with the result of *A. thaliana*,

suggesting that the process of ICE1 SUMOylation, mediated by SIZ1, might be similar to *A. thaliana*

(Miura et al. 2007). As for whether EcaSIZ1 has SUMO E3 ligase activity and whether it could couple with

EcaICE1 forming the SUMO coupling for playing a role in cold toleration in plant, need further

experiments to confirm.

CONCLUSIONS

In this study, we cloned the gene *EcaICE1* and *EcaSIZ1* from *E. camaldulensis*, and the amplification

of *EcaICE1* was the same as the previous result, and *EcaSIZ1* was highly conserved with other plant SIZ1

genes. EcaICE1 was similar to other ICE1 in plant, located in nucleus, and it had a strong transcriptional

activation activity. Moreover, the 84-126 amino acid regions of N-terminal domain of EcaICE1 protein,

were the key region of transcriptional activation activity of EcaICE1. In addition, the C-terminal region

from position 361 to 557 in EcaICE1 was the key region for its interaction with EcaSIZ1. According to this

study, we made a foundation on SUMOylation pathway and the molecular regulation mechanism of

EcaICE1.

Acknowledgements

This work has been supported by NSFC project 31470673. We also wish to express our appreciation to the

anonymous reviewers and technical editors of the Journal of Trees for their comments and corrections of the

article.

References

Badawi M, Reddy YV, Agharbaoui Z, Tominaga Y, Danyluk J, Sarhan F, Houde M (2008) Structure and

functional analysis of wheat ICE (Inducer of CBF Expression) genes. Plant Cell Physiol 49:

1237-1249

Bo JJ, Hua JY, Jiyoung L, Kenji M, Yul YC, Woe-Yeon K, Michael VO, Youbong H, Somers DE, Ilha L

(2010) The SUMO E3 ligase, AtSIZ1, regulates flowering by controlling a salicylic acid-mediated

floral promotion pathway and through affects on FLC chromatin structure. Plant J 53: 530-540

Cai B, Kong X, Zhong C, Sun S, Zhou XF, Jin YH, Wang Y, Li X, Zhu Z, Jin JB (2017) SUMO E3

Ligases GmSIZ1a and GmSIZ1b regulate vegetative growth in soybean: functional characterization of

soybean SUMO E3 ligase. J Integr Plant Biol 59: 2-14

Chinnusamy V, Ohta M, Kanrar S, Lee B, Hong X, Agarwal M, Zhu J (2003) ICE1: a regulator of

cold-induced transcriptome and freezing tolerance in *Arabidopsis*. Gene Dev 17: 1043-1054

Chinnusamy V, Zhu JH, Zhu JK (2007) Cold stress regulation of gene expression in plants. Trends Plant

Sci 12: 444-451

Courey AJ, Holtzman DA, Jackson SP, Tjian, R (1989) Synergistic activation by the glutamine-rich

domains of human transcription factor Sp1. Cell 59(5):827-836

Deng XM, Wang JX, Li Y, Wang J, Tian W (2017) Characterization of a cold responsive HbICE1 gene

from rubber trees. Trees 31: 137-147

Ding Y, Li H, Zhang X, Xie Q, Gong Z, Yang S (2015) OST1 Kinase modulates freezing tolerance by

enhancing ICE1 stability in *Arabidopsis*. Dev Cell 32: 278-289

Dong C, Agarwal M, Zhang Y, Xie Q, Zhu J (2006) The negative regulator of plant cold responses, HOS1,

is a RING E3 Ligase that mediates the ubiquitination and degradation of ICE1. Pro Natl Acad Sci

USA 103: 8281-8286

Gabriela TO, Enamul H, Quail PH (2003) The *Arabidopsis* basic/helix-loop-helix transcription factor

family. Plant Cell 15: 1749-1770

Halabian R, Fathabad ME, Masroori N, Roushandeh AM, Saki S, Amirizadeh N, Najafabadi AJ,

Gharehbaghian A, Roudkenar MH (2009) Expression and purification of recombinant human

coagulation factor VII fused to a histidine tag using Gateway technology. Blood transfus 7(4):

305-312

Huang X, Li K, Jin C, Zhang S (2015) ICE1 of *Pyrus ussuriensis* functions in cold tolerance by enhancing

PuDREBa transcriptional levels through interacting with PuHHP1. Sci Rep 5: 17620

Janmohammadi M, Zolla L, Rinalducci S (2015) Low temperature tolerance in plants: Changes at the

protein level. Phytochemistry 117: 76-89

Li C, Wu ZH, Shang XH. Research progress on soil microbial diversity of *Eucalyptus* plantations (2017)

Eucalypt Sci & Technol 34(03):51-54 [in Chinese]

Li H, Ding Y, Shi Y, Zhang X, Zhang S, Gong Z, Yang S (2017) MPK3- and MPK6-mediated ICE1

phosphorylation negatively regulates ICE1 stability and freezing tolerance in *Arabidopsis*. Dev Cell 43:

630-642

Lin YZ (2019) Research methodologies for genotype by environment interactions in forest trees and their

applications. Scientia Silvae Sinicae 55(5): 142-151 [in Chinese]

Lin YZ, Zhang ZY, Liu CX, Zhu BQ, Chen XY (2007) In silico cloning and analyzing of PsICE1 from

Populus suaveolens, a freezing-resistant transcription factor. Mol Plant Breeding 5(3): 424-430 [in

Chinese]

Lin YZ, Zheng HQ, Zhang Q, Liu CX, Zhang ZY (2014) Functional profiling of EcaICE1 transcription

factor gene from *Eucalyptus camaldulensis* involved in cold response in tobacco plants. J Plant

Biochem Biot 23: 141-150

Liu F, Wang X, Su M, Yu M, Zhang S, Lai J, Yang C, Wang Y (2015) Functional characterization of

DnSIZ1, a SIZ/PIAS-type SUMO E3 ligase from *Dendrobium*. BMC Plant Biol 15: 225

Miura K, Jin JB, Lee J, Yoo CY, Stirm V, Miura T, Ashworth EN, Bressan RA, Yun D, Hasegawa PM

(2007) SIZ1-Mediated sumoylation of ICE1 controls CBF3/DREB1A expression and freezing

tolerance in *Arabidopsis*. Plant Cell 19: 1403-1414

Miura K, Lee J, Jin JB, Yoo CY, Miura T, Hasegawa PM, Estelle M (2009) Sumoylation of ABI5 by the

Arabidopsis SUMO E3 ligase SIZ1 negatively regulates abscisic acid signaling. Pro Natl Acad Sci

USA 106: 5418-5423

Miura K, Okamoto H, Okuma E, Shiba H, Kamada H, Hasegawa PM, Murata Y (2013) SIZ1 deficiency

causes reduced stomatal aperture and enhanced drought tolerance via controlling salicylic acid-

induced accumulation of reactive oxygen species in *Arabidopsis*. Plant J 73: 91-104

Park HC, Kim H, Koo SC, Park HJ, Cheong MS, Hong H, Baek D, Chung WS, Kim DH, Bressan RA, LEE

SY, Bohnert HJ, Yun DJ (2010) Functional characterization of the SIZ/PIAS-type SUMO E3 ligases,

OsSIZ1 and OsSIZ2 in rice. Plant Cell Environ 33: 1923-1934

Shi Y, Ding Y, Yang S (2018) Molecular regulation of CBF signaling in cold acclimation. Trends Plant Sci

23: 623-637

Stockinger EJ, Gilmour SJ, Thomashow MF (1997) *Arabidopsis thaliana CBF1* encodes an AP2

domain-containing transcriptional activator that binds to the C-Repeat/DRE, a Cis-Acting DNA

regulatory element that stimulates transcription in response to low temperature and water deficit. Pro

Natl Acad Sci USA 94: 1035-1040

Walter M, Chaban C, Schütze K, Batistic O, Weckermann K, Näke C, Blazevic D, Grefen C, Schumacher

K, Oecking C, Harter K, Kudla J (2004) Visualization of protein interactions in living plant cells using

bimolecular fluorescence complementation. Plant J 40(3): 428-438

Wang X, Sun X, Liu S, Liu L, Liu X, Sun X, Tang K (2005) Molecular cloning and characterization of a

novel ice gene from *Capsella bursa-pastoris*. Mol Biol 39: 18-25

Zhang R, Guo Y, Li Y, Zhou L, Hao Y, You C (2016) Functional identification of MdSIZ1 as a SUMO E3

ligase in apple. J Plant Physiol 198: 69-80

Zhang S, Zhuang K, Wang S, Lv J, Ma N, Meng Q (2017) A novel tomato SUMO E3 ligase, SlSIZ1,

confers drought tolerance in transgenic tobacco. J Integr Plant Biol 59: 102-117

Zhao ML, Wang JN, Shan W, Fan JG, Kuang JF, Wu KQ, LI XP, Chen WX, He FY, Chen JY, Lu WJ

(2013) Induction of jasmonate signalling regulators MaMYC2s and their physical interactions with

MaICE1 in methyl jasmonate-induced chilling tolerance in banana fruit. Plant Cell Environ 36: 30-51

Table 1 Primers used in the experiments

primers	Primer sequences(5′→3′)
EcaICE1-F	ATGGTTCTGGGTCCCAGCG
EcaICE1-R	TCACACCATCCCATGGAAGC
EcaSIZ1-F	ATGGATTTGGTCTCTAGTTGCAAG
EcaSIZ1-R	CTATTCAGAATCTGATTCAATAGAAAGGTACA

Figure 1 Amino acid alignment of the EcaICE1 protein with twelve plant ICE1 proteins.

The predicted protein domains were shown. The *Arabidopsis thaliana* AtICE1 (Accession: AAP14668), *Capsella bursa*

CbICE1 (Accession: AAS7935), *Camellia sinensis* CsICE1 (Accession: GQ229032), *Eucalyptus camaldulensis* EcaICE1

(Accession: MH899180), *E. globulus* EglICE1 (Accession: AEF33833), *E. grandis* EgrICE1 (Accession: XP_010066419),

Glycine max GmICE1 (Accession: ACJ39211), *Hevea brasiliensis* HbICE1 (Accession: JT926796), *Malus domestica*

MdICE1 (Accession: ABS50251), *Populus suaveolens* PsICE1 (Accession: ABF48720), *P. trichocarpa* PtICE1 (Accession:

ABN58427) and *Pyrus ussuriensis* PuICE1 (Accession: APC57593) proteins are included.

Figure 2 Multi- alignment of amino acid of SIZ1 from *Eucalyptus camaldulensis* **and other plants and their conserved domains**

Arabidopsis thaliana: AtSIZ1, OAO90655; *Citrus sinensis*: CsSIZ1, XP_006488140; *Eucalyptus camaldulensis*: EcaSIZ1, MH899182; *E. grandis*: EgrSIZ1, XP_010043707; *Malus domestica*: MdSIZ1, XP_008385287; *Ziziphus jujuba*: ZjSIZ1, XP_015880489; *Populus trichocarpa*: PtrSIZ1, XP_024454498.

367 **Figure 3 Subcellular localization of EcaICE1 in leaves of *Nicotiana benthamiana***

368 YFP: the yellow fluorescence signals under YFP-excited; Bright field: figure under bright field; Merge: The

369 merge of YFP and bright field . Bar is 20 μm.

Figure 4 The protein interaction of EcaICE1 and EcaSIZ1 identified by bimolecular fluorescence complementation methods

YFP: the yellow fluorescence signals under YFP-excited; Bright field: figure under bright field; Merge: The

merge of YFP and bright field . Bar is 20 μm.

Figure 5 Transcriptional activation region analysis of EcaICE1 protein

BD: pGBKT7 vetor; AD: pGADT7 vetor; SD/-LW: SD/-Leu-Trp: double dropouts, SD medium with -Leu/-Trp;

SD/-LWHA: quadruple dropouts, SD medium with-Leu-Trp-His-Ade;10^{0}、10^{-1}、10^{-2}: dilutions with 1, 10 and 100

times respectively.

Figure 6 The interaction identification of EcaICE1 and EcaSIZ1 by yeast two-hybrid methods

BD: pGBKT7 vetor; AD: pGADT7 vetor; SD/-LW: SD/-Leu-Trp: double dropouts, SD medium with -Leu/-Trp;

SD/-LWHA: quadruple dropouts, SD medium with-Leu-Trp-His-Ade;10^0、10^{-1}、10^{-2}: dilutions with 1, 10 and

100 times respectively.

389