Original Manuscript

Title: Is vectorial transmission of *Trypanosoma cruzi* an efficient route to

support high infection rates in sylvatic hosts?

Juan Manuel Cordovez[1,4*], Mauricio Santos-Vega[1], Diana Erazo[1], Camilo Sanabria[2] ,

Lina María Rendón[3], Felipe Guhl[3]

[1] Departamento de Ingeniería Biomédica, Grupo de Investigación en Biología

Matemática y Computacional BIOMAC, Universidad de los Andes, Bogotá Colombia.

[2] Departamento de Matemáticas, Grupo de Investigación en Biología Matemática y

Computacional BIOMAC, Universidad de los Andes, Bogotá Colombia.

[3] Departamento de Ciencias Biológicas, Centro de Investigaciones en Microbiología y

Parasitología Tropical CIMPAT, Universidad de los Andes, Bogotá Colombia.

[4] Simon A. Levin Mathematical, Computational and Modeling Sciences Center,
Arizona State University, Tempe, Arizona
15

18 *Corresponding author: jucordov@uniandes.edu.co

Abstract

Chagas disease is caused by the parasite *Trypanosoma cruzi* and it is transmitted to humans by the triatomine bug *Rhodnius prolixus*. The main insect vector in the Andean countries presents sylvatic and domestic cycles involving humans, insects and reservoirs (e.g small mammals). It is commonly assumed that vectorial transmission is the main route for parasite spread between hosts. Recent studies have reported high percentages (21-80%) of infected opossums (*Didelphis marsupialis*) in the sylvatic cycle, raising the question of whether such a high proportion of infected could be only maintained by vectorial transmission, a seemingly inefficient pathway. To address this question, we formulated a mathematical model that describes the sylvatic transmission dynamics considering vectors and hosts and parametrized with field data. Our results show that vectorial transmission it is not sufficient to explain such high percentages of infected host-mammals reported in the literature. Here we propose oral transmission as an alternate route of transmission that may increase the number of infected individuals found in field studies.

Introduction

The transmission of the parasite *Trypanosoma cruzi*, etiological agent of Chagas disease, involves several pathways and results in 6 million infected people in Latin America [1]. Human infections are caused by multiple routes, the main suggested mechanism is vectorial transmission that occurs when triatomine insects feed on host blood (Sylvatic mammals and humand after a short period, the vector defecates releasing large amounts of parasites in the skin close to the wound allowing the parasite to reach the bloodstream [2-4]. Vertical transmission occurs in humans from an infected mother to a child; however, the ability of the parasite to cross the placenta of sylvatic reservoirs has not been fully demonstrated yet. Oral transmission has been proved to cause more aggressive clinical symptoms in humans and to have a high mortality rate (8-35% compared to 5-10% by vectorial transmission) only two weeks post infection [5]. In sylvatic mammals, oral transmission has been reported when mammals feed on *Rhodnius prolixu*s infected with *T. cruzi* were ingested [6]. Recent studies in central Brazil have demonstrated that both vertical and oral transmission are not a rare event in this biological system. In fact, a recent study in the Pantanal Region of Brazil, have demonstrated that both the vertical an oral transmission are likely to occur, depending on the encounter possibilities of the mammals and vectors [7].

In sylvatic mammals, particularly of the family Didelphidae, it has also been suggested that spraying from anal glands could be playing an important role in the transmission of *T. cruzi*. Opossums, mainly of the species *D. marsupialis*, have been proposed not only as a reservoir but also as a *T. cruzi* vector, since the parasite can multiply extracellularly in the anal glands of the animal [8-9]. This variety of transmission mechanisms and their relative importance in human infection compared to reservoir infection, suggests that parasites have different transmission cycles in the environment, sylvatic or domestic, that can be connected or isolated depending on the feeding behaviour of insect vectors.

When insect vectors breed and feed inside the houses, a domestic cycle is occurring, involving human and domestic mammals as reservoirs [4,10]. On the other hand, the sylvatic cycle involves triatomine bugs living in the wild feeding on sylvatic mammals such as opossums and rodents that act as reservoirs. The connection between the two cycles occurs when insects migrate from the sylvatic to the domestic habitat attracted by light sources and domestic species presence combined with the increase in the number of domestic animals like dogs [11-14]. It must be considered that the ecological

interactions and encounters between vectors and reservoirs depend on the faunal
composition, which is directly related with the landscape structure [15].

The domestic and peri domestic cycle has been the focus of many reports [12, 16-17]
and control programs [4, 18-20], probably because is the easiest one to intervene and
involves humans directly. However, understanding the sylvatic cycle is crucial because
it is the source of infected insects that ultimately invade the houses. Furthermore,
control programs in Colombia that used pyrethroid insecticides, that succeeded in
countries like Chile and Uruguay in eliminating insects from houses [21], have shown
limited effectiveness in Colombia due to a strong re-infestation phenomenon that
occurs weeks after the application of the insecticide [22]. In addition, vector prevention
and control activities have not been very efficient in endemic areas due to the lack of
knowledge about the biological characteristics of the vector populations present in
each region, leading to uncertainty about the most appropriate control measures in
each transmission scenario [23]. Thus, we investigate in more detail the sylvatic cycle
dynamics for the Colombian endemic department of Casanare a territory of high
interest involving both Chagas disease transmission cycles and high *T. cruzi* natural
infection in *R. prolixus* [24-27].

In rural areas of Colombia, particularly in the Orinoco region where the Casanare
department is located, *R. prolixus* is considered the main vector of *T. cruzi*
transmission. Its main habitat are palm trees, particularly the species *Attalea*
butyracea, in a landscape where houses are scarce and well spread. This palm tree
species is widely distributed and establishes a large faunistic reserve, where
mammalian reservoirs are typically found in the crowns [23]. In this natural habitat,
where triatomines and mammals coexist and interact, the vectorial pathway of parasite
transmission between them has been demonstrated by many studies [2]. Few authors
had documented the existence of both oral and congenital pathways [28-29]. This
raises the difficulty of knowing whether vector transmission is acting as a sole route,
or if multiple transmission modes act simultaneously.

Regardless of the transmission route, the question of how severe is the infection in
sylvatic reservoirs has appeared repeatedly and several studies have reported the
proportion of different infected mammals. This is important because high levels of
reservoirs infection suggest higher probabilities for human infection. For example, bats
can play an important role in transmission scenarios, because their high mobility allows
the parasite to migrate between the sylvatic and domestic habitat. Studies carried out

in Casanare in different bat species reported infection indexes of T. cruzi from 6.5% to
51% [26, 30]. Moreover, Didelphis marsupialis has been considered one of the main
reservoirs of the parasite and previous studies suggest infection rates ranging from 5
to 90% [31-32]. More recent studies have reported 80% and 89% infected mammals
in an area with similar ecological characteristics to the department of Casanare [33-
34. Finally, the results from our studies showed an infection rate of 21% in *D.*
marsupialis. Hypothetically if vectorial transmission is acting alone, host reservoirs will
become infected only if after a blood meal the insect defecates on the skin and then
the parasites find their way into the bloodstream, and insects would become infected
if they suck blood from an infected mammal. This poses the question of whether
vectorial transmission is sufficient to sustain more than 21% of infected hosts in a
population, as reported before.

Several mathematical models of Chagas disease have been proposed in the last two
decades to study its epidemiology and more recently its ecology. Among the first
models we can find general analysis of vector and host dynamics [35], the
incorporation of acute and chronic stages [36] or models accounting for congenital
transmission [37], spatially explicit models [38], age-structure models [39], and
stochastic models [40]. On the ecological side, more recent models have investigated
the role of dogs [41] and synantropic animals in human infection [42], and the sylvatic
cycle dynamics in different habitats varying in their host communities [43]. However,
none of these models has evaluated if the assumed mechanisms of transmission could
explain the sylvatic host infection that is observed in the wild.

Here we investigate if vectorial transmission per se is capable to maintain the
transmission observed in the data by using a mathematical modeling approach. By
exploring the model, we will address if the well-known vectorial transmission is enough
to support the high infection rates among sylvatic reservoirs and, if so, to propose
entomological control strategies that would be adequate to reduce the risk of infection
to humans.

Methods and Results

We formulate different epidemiological models to recreate the sylvatic transmission
cycle and establish the interactions between vectors and reservoirs. In the model, we
consider hosts (H) and vectors (V) to represent the population of *D. marsupialis* and
R. prolixus respectively, and the model was set to represent the dynamics in the region

of Casanare, Colombia composed by large palm plantations. The results are
normalized per palm as a study unit. The total number of hosts, N_H, is divided into
susceptible (S_H) and infected class (I_H). Similarly, the total population of vectors N_V, is
divided into susceptible (S_V) and infected (I_V). Our first model depicted in figure 1A, the
vectors become infected at a rate (β_{HV}) that reflects the transmission of the parasite
from an infected host to a susceptible vector due to biting. Then the host become
infected at a rate ($\beta_{VH,0}$) that reflects the transmission of the parasite from an infected
vector to a susceptible host due to biting. Likewise, ($\beta_{VH,1}$) is the transmission rate by
ingestion (consumption or predation) of the infected vector by the reservoir. In this way
β_{HV}, $\beta_{VH,0}$, and $\beta_{VH,1}$ consider triatomine – host contact rate by the probability of
infection from the one to the other. The second model, in figure 1B, considers the
same routes of transmission as the previous model and includes a new rate reflecting
the possibility of susceptible host acquiring the parasite from an infected host (β_{HH}).
159
160

161
162 **Figure 1**. A) Shows the Vector-host model. Vector-host model with transmission
163 between hosts
164
165 The two models are summarized by the equations:
166
167
$$\frac{dS_V}{dt} = \mu_V N_V - \beta_{HV} I_H S_V - \mu_V S_V$$

168
169
$$\frac{dI_V}{dt} = \beta_{HV} I_H S_V - \mu_V I_V$$

170
171
$$\frac{dS_H}{dt} = \mu_H N_H - (\beta_{VH,0} + \beta_{VH,1}) I_V S_H - \mu_H S_H$$

$\dfrac{dI_H}{dt} = \{(\beta_{VH,0} + \beta_{VH,1})I_V + \beta_{HH}I_H\}S_H - \mu_H I_H$

Where the model with no transmission between hosts (Figure 1A) is obtained by

making $\beta_{HH} = 0$. Having no vital dynamics, $N_V = S_V + I_V$ and $N_H = S_H + I_H$ are

constant, the equations collapse into:

$\dfrac{dI_V}{dt} = \beta_{HV}I_H(N_V - I_V) - \mu_V I_V$

$\dfrac{dI_H}{dt} = \{(\beta_{VH,0} + \beta_{VH,1})I_V + \beta_{HH}I_H\}(N_H - I_H) - \mu_H I_H$

The Next Generation Matrix (G) for this model is:

$$G = \begin{bmatrix} 0 & \dfrac{N_V \beta_{HV}}{\mu_H} \\ \dfrac{N_H(\beta_{VH,0} + \beta_{VH,1})}{\mu_V} & \dfrac{N_H \beta_{HH}}{\mu_H} \end{bmatrix}$$

And the adjacency matrix $(S(G))$ is then:

$$S(G) = \begin{bmatrix} 0 & 1 \\ 1 & \epsilon \end{bmatrix}$$

Where $\epsilon = 1$ or $\epsilon = 0$, depending whether there is transmission between hosts. The

spectral radius ρ of $S(G)$ is either 1 if $\epsilon = 0$, or $(1 + \sqrt{5})/2$ if $\epsilon = 1$. Now if $\mu(G)$ is the

critical virulence [44] of our epidemiological system and R_0 its basic reproductive

number:

$$\mu(G) \leq R_0 \leq \rho\,\mu(G)$$

Therefore, with no transmission between hosts

$$R_0 = \mu(G) = \sqrt{\frac{N_V N_H \beta_{HV}(\beta_{VH,0} + \beta_{VH,1})}{N_V \mu_H}}$$

204 And with it

$$\mu(G) \leq R_0 \leq \frac{1 + \sqrt{5}}{2} \mu(G) \simeq 1.618\,\mu(G)$$

206 Where

$$\mu(G) = \max\left\{\sqrt{\frac{N_V N_H \beta_{HV}(\beta_{VH,0} + \beta_{VH,1})}{\mu_V \mu_H}}, \frac{N_H \beta_{HH}}{\mu_H}\right\}$$

209 In order to have an endemic equilibrium, where the parasite invaded the ecosystem, it

210 is necessary to have $R_0 > 1$. In the first model (Figure 1A), in this equilibrium, if it exists,

211 the number of infected hosts corresponds to:

$$I_H^* = \frac{N_V N_H \beta_{HV}(\beta_{VH,0} + \beta_{VH,1}) - \mu_V \mu_H}{\beta_{HV}\{N_V(\beta_{VH,0} + \beta_{VH,1}) + \mu_H\}}$$

$$I_H^* = N_H\left(1 - \frac{1}{R_0^2}\right)\left(\frac{N_V(\beta_{VH,0} + \beta_{VH,1})}{N_V(\beta_{VH,0} + \beta_{VH,1}) + \mu_V}\right)$$

$$I_H^* \leq N_H\left(1 - \frac{1}{R_0^2}\right)$$

220 In the second model (Figure 1B), we know that the parasite will not invade the

221 ecosystem if

$$\mu(G) < \frac{2}{1 + \sqrt{5}} = \frac{\sqrt{5} - 1}{2} \simeq 0.618$$

For the numerical exploration of the model we used parameters from the literature. To

determine their maximum values for transmission coefficients, which are the most

uncertain parameters, we combined expert knowledge, field and lab measures from

several sources (Table I). Death rate parameters were estimated based on the average

of the life expectancy (1/life expectancy) from literature reports.

Finally, to relax the assumption of the transmission following the mass action law an

alternate way to model the transmission would be incorporate saturation effects in the

transmission. The most common way to include the effects of saturation in a biological

model is to consider an analogous to the Michaelis-Menten equation in the definition

of both transmission coefficients. Here, we redefined the transmission rates in our

model to include saturation in the following manner:

$$\beta_{HV} = \frac{b_{HV}N_H}{a_{HV}N_H}; \ \beta_{VH} = \frac{b_{VH}N_V}{a_{VH}N_V}$$

where b is the maximum transmission coefficient and a reflects the number of hosts

necessary to reach the saturation level (when $N_H = a$, the transmission coefficient

takes the value of $b/2$). However, even if we consider a more realistic interaction

between reservoirs and vectors the model could predicts reproductive numbers higher

than 1 for some combinations of the transmission coefficients and does not explain the

high levels of infection found in the field studies.

Table I. Parameters used in the mathematical model of the sylvatic cycle of Chagas

disease.

Terminology	Definition	Value	References
N_V	Total number of vectors[a]	9.6/palm	[26]
N_H	Total number of hosts[a]	1.5/100ha	[26]
$\beta_{HV}*$	Max transmission rate of vector by contact with infected host[b]	1	[55]
$\beta_{VH,0}*$	Max transmission rate of host due to an infected vector (biting)[ab]	1×10^{-2}	[45]

$\beta_{VH,1}$*	Transmission rate of host due to an infected vector (oral)[a][b]	1 x 10^{-1}	[45]
β_{HH}*	Transmission rate of host by contact with infected host[b]	1 x 10^{-1}	[55]
μ_V	Death rate for vectors[a]	1.7/year	[46]
μ_H	Death rate for host[a]	0.666/year	[47,48,49]

* Transmission rates are reported to the nearest order of magnitude.

[a] Parameter was calculated using literature reports.

[b] Parameter was calculated using expert knowledge.

A sensitivity analysis for global reproductive number R0 was performed using the Latin

Hypercube method (LH) to estimate each parameter contribution. Negative Partial

Rank Correlation Coefficients (PRCC) indicate a decrease in R0 and PRCC positive

values indicate an increment in R0. Thus, increases in N_v, N_H, β_{HV}, $\beta_{VH,0}$, and $\beta_{VH,1}$

produce a rise in R0 (Fig. 2). The positive effect (PRCC) of these parameters ranges

between 0.61 and 0.65 for the model without oral transmission and when it is included,

the contribution of vector-host transmission rates ($\beta_{VH,0}$, $\beta_{VH,1}$) is shared by both

parameters ($\beta_{VH,0}$ PRCC = -0.30 and $\beta_{VH,1}$ PRCC = - 0.33). On the other hand, vector

and host death rates have significant effect on lowering R0 (model without oral

transmission μ_V, PRCC = -0.61 and μ_H PRCC = - 0.58, including oral transmission μ_V,

PRCC = -0.61 and μ_H PRCC = - 0.58).

Figure. 2. Latin Hypercube Sampling for the global reproductive number R0. PRCC:

Partial Rank Correlation Coefficient. Black colored circles correspond to the model

without oral transmission and gray to the model including oral transmission. For

information about each parameter's explanation see Table 1. Note that parameters

related to host-vector (and vice versa) infection rates and populations (N_v, N_H, β_{HV},

$\beta_{VH,0}$, and $\beta_{VH,1}$) have a positive contribution to R0. On the contrary, vector and host

death rates (μ_V and μ_H) have a negative effect in R0.

Plugging the parameter values into the maximum critical virulence equation $\mu(G)$ and

considering the maximum probability value in each transmission rate (β) we got that μ

(G) max is between 0.3584 and 0.0224. Since none of these values are greater than

0.618, an endemic infection with the parasite cannot occur, interestingly this indicates

that another route of transmission, apart from the vector route, is needed to explain

high percentages of host infection rates. Figure 3 show a numerical exploration of the

model considering parameter ranges for β_{HV} and $\beta_{VH,0}$, here is evident that for every

combination of parameters a model that just incorporates vectorial transmission is not

capable to exhibit an R0 bigger than one.

Figure 3. R0 values depicted in z depending on β_{HV} (here in x) and $\beta_{VH,0}$ (here in y) we used for the analyses the palm as the study unit. In this case the max R0 is lower than 0.4.

In addition, Figure 4 shows values of R0 resulted from simulations of the model with a range of values of the oral transmission parameter $\beta_{VH,1}$ assuming that β_{HV} and $\beta_{VH,0}$ are at the maximum value found in figure 2, we found that $\beta_{VH,1}$ has to be 0.07 or bigger in order to reach and R_0 above 1 (Figure 4).

Figure 4. The x axis represents n $\beta_{VH,1}$ values range and their corresponding Ro. We

assume maximum values for β_{HV} (1) and $\beta_{VH,0}$ (0.01). In this case, $\beta_{VH,1}$ must be larger

than 0.07, for Ro to have a value above 1.

Discussion

To the date, studies that investigate in detail the dynamics of the sylvatic cycle of *T.*

cruzi transmission are still rare. Nonetheless, it has been shown for the endemic region

of Casanare a stable sylvatic transmission, where *R. prolixus* individuals were captured

in palm trees (*A. butyracea*) [25-27]. These studies have reported infections rates in

mammals ranging from (21-89%) [26, 34, 50, 51]. This is remarkable fact, given that

vectorial route has been considered inefficient since the parasite faces great

challenges to infiltrate the host bloodstream via vectorial route [52] , thus this route its

unable to explain the observed reservoir prevalence reported in the literature.

T. cruzi vectorial transmission has been suggested to be among one the most

inefficient ways for parasites to infect susceptible hosts, although the number of

infected hosts it is high on the field fluctuating between 40 to 90% [31-33]. Our results

from model simulations only considering vectorial transmission show that the basic

reproductive number R_0 is always less than 1. This implies that an additional

transmission route is needed to guarantee an endemic state (R0>1). Although, there

are no reports in the literature for natural occurring populations where the transmission

of the parasite is not supported due to low transmission rates. Perhaps for other vector

populations, like *Triatoma dimidiata*, this could be the case and it would be interesting

to verify it in the field.

One of our main goals with this study was to establish if vectorial transmission per se

was able to explain the high-reported levels of infected hosts. Our results demonstrate

that even if we simulate the system at their maximal critical virulence $\mu(G)$ over the

highest possible value combinations of transmission rates the system never reaches

high number of infected hosts. The incorporation of a new route of transmission, such

as the oral transmission, let the system reach the high proportion of infected hosts

reported in field studies [47,52-54]. However, these high proportion of infected host

could also be obtained with higher levels of transmission rates, in particular increasing

the transmission rate from vectors to host β_{VH}, even though we believe that the high

values needed are outside of the biological range we have no report to compare to and

thus this is mainly speculative.

Results from our sensitivity analysis suggest there is no single variable or parameter

that by itself explains the dynamics in the systems. Instead, we were able to identify a

subset of factors that together help to explain the temporal variation in the system.

First, in the absence of oral transmission the dynamics is explained by the transmission

rate and the population size and the incorporation of the oral transmission add a

significant positive effect that help to reach a higher number of infected hosts. In

addition, it is important to note that the direction of change in the fraction of infected

hosts after a change in the parameter is given by the sign of its condition number: a

same direction change (e.g. increase parameter-increase i_H) is a positive condition

number, while an opposite direction change (e.g. increase parameter-decrease i_H)

produces negative condition numbers. Using the parameter values in table 1, we found

that for increasing growth rates increases the fraction of infected hosts. However, the

effect of changing N_V is an order of magnitude lower than changing N_H in the same

proportion (same relative change). For death rates, we have an opposite-direction

effect as it was expected, meaning that if we have more vector or host, the contact

between both would be lower accordingly to the mass action law, so the net flow of

individuals from susceptible to infected populations would be lower. Again, we saw that

vectors have an effect almost an order of magnitude higher than the hosts. Thus, we

could expect that these parameters become an interesting target for disease control

initiatives.

Importantly, our model is implemented using the palm as spatial unit; the choice is

based in its fundamental role in the ecology of both vector and host populations. A

mathematical model with a different spatial resolution (i.e. a village) faces the

challenge of vector and host mobility. In addition, a temporal model often assumes

complete mixing in the spatial component and that is an important aspect to study any

host – vector disease model. Here restricting the model to the palm for the analysis

simplify and constrain the model results. Extrapolating our results to villages with

multiple palms has to be done carefully because hosts often visit multiple palms and

insects could move also between palms and houses. We believe that our results

should apply to higher levels of aggregation such as villages with high and

homogeneous palms density with easy access between palms to guaranteed

population mixing. However, including the palm distribution in the model implies a

different theoretical approach that although is an important hypothesis it is out of the

scope of this paper.

From a biological perspective, the ability of the model to capture important disease

dynamics is what makes it useful for testing potential control strategies and studying

T. cruzi transmission in sylvatic host species different from *D. marsupialis*. For

example, if we increase death rate of vectors and simultaneously decrease the

transmission rates to a point where they cancel out is possible to eradicate the disease.

This is an important result, because control strategies often target one parameter at a

time (i.e. increase vector mortality - house spraying, decrease contact rates - improve

house materials), but it seems more reasonable to intervene all of them at the same

time taking care in shifting the disease balance from endemic to temporal. In fact, it

has been proposed that using a particular control method does not exclude using

another one [10]. One way to achieve this at the household level is to combine

spraying, which increases insect mortality, with presence of non-reservoirs peri-

domestic species, such as chickens, providing another vector feeding. However, there

is potential negative effect because the latter could increase vector carrying capacity.

In addition to the previous analysis the model could be used to further refine the range

of unknown parameters. For example, the transmission rates, β_{HV} and β_{VH}, are difficult

to estimate, however if one has reports of the densities for host and vectors, proportion

of infected host and vectors, and a good estimate of death rates, then it is easy to

calculate the transmission rates. This technique could be implemented combining
fieldwork and the mathematical expressions to make the model adequate to a certain
region, and thus a useful disease control tool.
390

Conclusions

In Latin America, the transmission dynamics of Chagas disease vary significantly
between regions and the Orinoco epidemiological scenario involves a unique mixture
of factors that requires interdisciplinary approaches. Computational models, along with
biological knowledge, are a great tool to test hypotheses and forecast epidemic events,
becoming great allies in understanding transmission mechanisms and designing
control strategies.
398

Acknowledgements

[funding sources should not be included here or in the manuscript file, only during

manuscript submission]

References

1. World Health Organization 2015. Chagas disease in Latin America: an
epidemiological update based on 2010 estimates. *Weekly epidemiological*
record Relevé épidémiologique hebdomadaire No. 6, 90: 33-40.
2. Catala S, Crocco LB, Morales GF 1997. *Trypanosoma cruzi* transmission risk
index (TcTRI): an entomological indicator of Chagas disease vectorial
transmission to humans. *Acta Trop 68*: 285-295.
3. Rassi AJ RA, Marin-Neto JÁ 2010. Chagas disease. Seminar. *Lancet 375*:
1338-1402.
4. Coura JR, Junqueira AC, Fernandes O, Valente SA, Miles MA 2002.
Emerging Chagas disease in Amazonian Brazil. *Trends Parasitol 18*: 171-
176.
5. Sanchez LV, Ramírez JD 2012. Congenital and oral transmission of American
trypanosomiasis: an overview of physiopathogenic aspects. *Parasitology 139*:
1-13.
6. Thomas ME, Rasweiler IV JJ, D'Alessandro A 2007. Experimental
transmission of the parasitic flagellates *Trypanosoma cruzi* and *Trypanosoma*

rangeli between triatomine bugs or mice and captive neotropical bats. *Mem Inst Oswaldo Cruz 102*: 559-565.

7. Herrera HM, Rocha FL, Lisboa CV, Rademaker V, Mourao GM, Jansen AM 2011. Food web connections and the transmission cycles of Trypanosoma cruzi and Trypanosoma evansi (Kinetoplastida, Trypanosomatidae) in the Pantanal Region, Brazil.   : 380-387.

8. Jansen A, Santos de Pinho A, Varella LC, Cupolillo E, Mangia RH, Fernandes O 1999. The Sylvatic Cycle of *Trypanosoma cruzi*: a Still Unsolved Puzzle. *Mem Inst Oswaldo Cruz 94*: 203-204.

9. Jansen A, Carreira J, Deane M 1988. Infection of a mammal by monogenetic insect Trypanosomatids (Kinetoplastida, Trypanosomatidae). *Mem Inst Oswaldo Cruz 81*: 271-272.

10. Guhl F 2009. Enfermedad de Chagas: Realidad y perspectivas. *Biomedica 20*: 228-234.

11. Cohen JE, Gurtler RE 2001. Modeling household transmission of American trypanosomiasis. *Science 293*: 694-698.

12. Fitzpatrick S, Feliciangeli MD, Sanchez-Martin MJ, Monteiro FA, Miles MA 2008. Molecular genetics reveal that silvatic *Rhodnius prolixus* do colonise rural houses. *PLoS Negl Trop Dis 2*: e210.

13. Sanchez-Martin MJ, Feliciangeli MD, Campbell-Lendrum D, Davies CR 2006. Could the Chagas disease elimination programme in Venezuela be compromised by reinvasion of houses by sylvatic *Rhodnius prolixus* bug populations? *Trop Med Int Health 11*: 1585-1593.

14. Minoli SA, Lazzari CR 2006. Take-off activity and orientation of triatomines (Heteroptera: Reduviidae) in relation to the presence of artificial lights. *Acta Trop 97*: 324-330.

15. Jansen AM, Xavier SCC, Roque ALR 2015. The multiple and complex and changeable scenarios of the *Trypanosoma cruzi* transmission cycle in the sylvatic environment. *Acta Trop 151*: 1-15.

16. Miles MA, Feliciangeli MD, de Arias AR 2003. Science, medicine, and the future - American trypanosomiasis (Chagas' disease) and the role of molecular epidemiology in guiding control strategies. *Brit Med J 326*: 1444-1448.

17. Feliciangeli MD, Dujardin JP, Bastrenta B, Mazzarri M, Villegas J, Flores M, Muñoz M 2002. Is Rhodnius robustus (Hemiptera: Reduviidae) responsible for Chagas disease transmission in Western Cenezuela? *Trop Med Int Health 7*: 280-287.

18. Dias JCP 2007. Southern Cone Initiative for the elimination of domestic
populations of *Triatoma infestans* and the interruption of transfusional Chagas
disease. Historical aspects, present situation, and perspectives. *Mem Inst*
Oswaldo Cruz 102: 11-18.

19. Dias JCP, Silveira AC, Schofield CJ 2002. The impact of Chagas disease
control in Latin America - A review. *Mem Inst Oswaldo Cruz 97*: 603-612.

20. Campbell-Lendrum DH, Angulo VM, Esteban L, Tarazona Z, Parra GJ
Restrepo M, Restrepo BN, Guhl F, Pinto N, Aguilera G, Wilkinson P, Davies
CR 2007. House-level risk factors for triatomine infestation in Colombia. *Int J*
Epidemiol 36: 866-872.

21. WHO/TDR UW 1999. Chile and Brazil to be certified free of transmission of
Chagas disease. TDR News.

22. Guhl F 1998. Estado actual del control de la enfermedad de Chagas en
Colombia. In: Guhl F JCE, editor. Curso-taller control de tripanosomiasis
americana y leishmaniasis: aspectos biológicos, genéticos y moleculares.
Bogotá: Corcas Editores. pp. 47-81.

23. Guhl F, Pinto N, Aguilera G 2009. Sylvatic triatominae: a new challenge in
vector control transmission. *Mem Inst Oswaldo Cruz 104*: 71-75.

24. Guhl F, Aguilera G, Pinto N, Vergara D 2007. Actualización de la distribución
geográfica y ecoepidemiología de la fauna de triatominos (Reduviidae:
Triatominae) en Colombia. *Biomédica 27*(Supl. 1): 143-62.

25. Angulo V, Esteban L, Luna KP 2012. *Attalea butyracea* próximas a las
viviendas como posible fuente de infestación domiciliaria por *Rhodnius prolixus*
(Hemiptera: Reduviidae) en los Llanos Orientales de Colombia. *Biomédica*
32(2).

26. Rendón LM, Guhl F, Cordovez JM, Erazo D 2015. New scenarios of
Trypanosoma cruzi transmission in the Orinoco region of Colombia. *Mem Inst*
Oswaldo Cruz 110(3): 283-288.

27. Urbano P, Poveda C, Molina J 2015. Effect of the physiognomy of *Attalea*
butyracea (Arecoideae) on population density and age distribution of *Rhodnius*
prolixus (Triatominae). *Parasit Vectors 8*: 199.

28. Benchimol Barbosa PR 2006. The oral transmission of Chagas' disease: an
acute form of infection responsible for regional outbreaks. *Int J Cardiol 112*:
132-133.

29. Brutus L, Schneider D, Postigo J, Romero M, Santalla J, Chippaux JP 2008.
Congenital Chagas disease: diagnostic and clinical aspects in an area without
vectorial transmission, Bermejo, Bolivia. *Acta Trop 106*: 195-199.

30. Ramirez et al. 2014

31. Fernandes O, Mangia RH, Lisboa CV, Pinho AP, Morel CM, Zingales B,
Campbell DA, Jansen AM 1999. The complexity of the sylvatic cycle of
Trypanosoma cruzi in Rio de Janeiro state (Brazil) revealed by the non-
transcribed spacer of the mini-exon gene. *Parasitology 118 (Pt 2)*: 161-166.

32. Grisard EC, Carvalho-Pinto CJ, Scholz AF, Toma HK, Schlemper BR Jr,
Steindel M 2000. *Trypanosoma cruzi* infection in *Didelphis marsupialis* in
Santa Catarina and Arvoredo Islands, southern Brazil. *Mem Inst Oswaldo*
Cruz 95: 795-800.

33. Sanchez JL CJ, Vallejo GA, Lozano LE, Jaramillo JC, Guhl F 2000.
Diagnóstico molecular de *Trypanosoma cruzi* en reservorios y humanos en
un área endémica del departamento del Tolima. Curso-Taller Internacional:
"Biología, Epidemiología y Control de la Tripanosomiansis Americana y
Leshmaniasis": Mayo 29 Junio 23, pp. 63-68.

34. Pinto N MD, Herrera C, Vallejo G, Naranjo JM, Guhl F 2005. Comprobación
del ciclo selvatico de *Rhodnius prolixus* en reductos de *Attalea Butyracea* en
el departamento de Casanare. *Biomédica 25*: 159.

35. Velasco-Hernandez JX 1991. An epidemiological model for the dynamics of
Chagas' disease. *Biosystems 26*: 127-134.

36. Das P, Mukherjee D 2006. Qualitative study of a model of Chagas' disease.
Math Comput Model 43: 413-422.

37. Raimundo SM, Massad E, Yang HM 2010. Modelling congenital transmission
of Chagas' disease. *Biosystems 99*: 215-222.

38. Slimi R, El Yacoubi S, Dumonteil E, Gourbiere S 2009. A cellular automata
model for Chagas disease. *Appl Math Model 33*: 1072-1085.

39. Inaba H, Sekine H 2004. A mathematical model for Chagas disease with
infection-age-dependent infectivity. *Math Biosci 190*: 39-69.

40. Castañera MB, Aparicio JP, Gurtler RE 2003. A stage-structured stochastic
model of the population dynamics of *Triatoma infestans*, the main vector of
Chagas disease. *Ecol Model 162*: 33-53.

41. Fabrizio et al. 2016

42. Petersen et al. 2015

43. Erazo, D., & Cordovez, J. (2016). Modeling the effects of palm-house proximity
on the theoretical risk of Chagas disease transmission in a rural locality of the
Orinoco basin, Colombia. Parasites & Vectors, 9(1), 592.
https://doi.org/10.1186/s13071-016-1884-8

44. Cordovez JM and Sanabria C 2014. Environmental Changes Can Produce
531   Shifts in Chagas Disease Infection Risk. *Environmental Health Insights* 8(S2):
532   43-48 doi: 10.4137/EHI.S16002.

533   45. Rabinovich J, Solarz ND, Gürtler R, Wisnivesky-Colli C 1990. Probability of
534   transmission of Chagas disease by *Triatoma infestans* (Hemiptera:
535   Reduviidae) in an endemic area of Santiago del Estero, Argentina. *Bull World*
536   *Heal Organ 68*: 737-746.

537   46. Chaves LF, Hernandez MJ, Revilla TA, Rodriguez DJ, Rabinovich JE 2004.
538   Mortality profiles of *Rhodnius prolixus* (Heteroptera: reduviidae), vector of
539   Chagas disease. *Acta Trop 92(2)*: 119-125.

540   47. Schweigmann NJ, Pietrokovsky S, Bottazzi V, Conti O, Bujas MA,
541   Wisnivesky-Colli C 1999. Prevalence of *Trypanosoma cruzi* infection in
542   opossum (*Didelphis albiventris*) in Santiago del Estero, Argentina. *Rev*
543   *Panam Salud Publica 6*: 371-377.

544   48. Reyes AA 2009. Fauna mammalia asociada a los focos de leishmaniasis
545   neotropical. Situación en Venezuela. *Boletín de Malariología y Salud*
546   *Ambiental 1*: 35-52.

547   49. Telford S, Tonn R 1982. Dinámica de *Trypanosoma cruzi* en poblacionesde
548   un reservorio primario, *Didelphis marsupialis* en los llanos altos de
549   Venezuela. *Bol Oficina Sanit Panam 93*: 341-364.

550   50. Freire-de-Lima L, da Fonseca LM, da Silva VA, da Costa KM, Morrot A,
551   Freire-de-Lima CG, Previato JO, Mendonça-Previato L. Modulation of Cell
552   Sialoglycophenotype: A Stylish Mechanism Adopted by Trypanosoma cruzi to
553   Ensure Its Persistence in the Infected Host. *Front Microbiol.* 2016 May
554   11;7:698. doi: 10.3389/fmicb.2016.00698. PMID: 27242722; PMCID:
555   PMC4862976.

556   51. Pinho A., Cupolillo E., Mangia R., Fernandes O. & Jansen A. M.
557   (2000). *Trypanosoma cruzi* in the sylvatic environment: distinct transmission
558   cycles involving two sympatric marsupials. *Trans. R. Soc. Trop. Med.*
559   *Hyg.* **94:** 509-514.

560   52. Deane M., Lenzi H. & Jansen A. (1986). Double development cycle
561   of *Trypanosoma cruzi* in the opossum. *Parasitol. Today.* **2:** 146-147.

562   53. Conti O, Schweigmann NJ, Pietrokovsky S, Bottazzi V, Wisnivesky-Colli C.
563   Search for *Trypanosoma cruzi* in the anal glands of wild *Didelphis*
564   *albiventris* from Santiago del Estero, Argentina. Mem Inst Oswaldo Cruz
565   1995; 90(6):687

566   54. Jansen AM, Madeira FB, Deane MP. *Trypanosoma cruzi* infection in the

567   opossum *Didelphis marsupialis:* absence of neonatal transmission and

568   protection by maternal antibodies in experimental infections. Mem Inst

569   Oswaldo Cruz 1994;89:4145.

574 **Figure legends**

575 F**igure 1**. A) Shows the Vector-host model. Vector-host model with transmission
576 between hosts

577
Figure. 2. Latin Hypercube Sampling for the global reproductive number R0. PRCC:
Partial Rank Correlation Coefficient. Black colored circles correspond to the model
without oral transmission and gray to the model including oral transmission. For
information about each parameter's explanation see Table 1. Note that parameters
related to host-vector (and vice versa) infection rates and populations (N_v, N_H, β_{HV},
$\beta_{VH,0}$, and $\beta_{VH,1}$) have a positive contribution to R0. On the contrary, vector and host
death rates (μ_V and μ_H) have a negative effect in R0.

585
586 **Figure 3.** R0 values depicted in z depending on β_{HV} (here in x) and $\beta_{VH,0}$ (here in y)
587 we used for the analyses the palm as the study unit. In this case the max R0 is lower
588 than 0.4.

589
590 **Figure 4.** The x axis represents n $\beta_{VH,1}$ values range and their corresponding Ro. We
591 assume maximum values for β_{HV} (1) and $\beta_{VH,0}$ (0.01). In this case, $\beta_{VH,1}$ must be larger
592 than 0.07, for Ro to have a value above 1.

593
594

595 **Tables**

596 **Table I.** Parameters used in the mathematical model of the sylvatic cycle of Chagas
597 disease.
598