Developing synergistic drug combinations to restore antibiotic sensitivity in drug-resistant

Mycobacterium tuberculosis

(Running title: Restoring sensitivity in drug-resistant tuberculosis)

Charles Omollo,[a,b,c,d,#] Vinayak Singh,[a,b,c,d] Elizabeth Kigondu,[a,b,c,*] Antonina Wasuna,[a,b,c] Pooja

Agarwal,[c,d] Atica Moosa,[c,d] Thomas R. Ioerger,[e] Valerie Mizrahi,[c,d,f] Kelly Chibale,[a,b,d] and

Digby F. Warner[c,d,f,#]

a. Department of Chemistry, University of Cape Town, Rondebosch 7701, South Africa;

b. South African Medical Research Council Drug Discovery and Development Research Unit,

University of Cape Town, Rondebosch 7701, South Africa;

c. SAMRC/NHLS/UCT Molecular Mycobacteriology Research Unit, DSI/NRF Centre of

Excellence for Biomedical Tuberculosis Research, Department of Pathology, University of Cape

Town, Rondebosch 7701, South Africa;

d. Institute of Infectious Disease & Molecular Medicine, University of Cape Town, Rondebosch

7701, South Africa;

e. Texas A&M University, Department of Computer Science, College Station, TX, 77843, USA

f. Wellcome Centre for Infectious Diseases Research in Africa, University of Cape Town,

Rondebosch 7701, South Africa.

Current address: Centre for Traditional Medicine & Drug Research, Kenya Medical Research

Institute, Nairobi, Kenya

[#]For   correspondence: charles.omollo@uct.ac.za;   digby.warner@uct.ac.za

ABSTRACT

Tuberculosis (TB) is a leading global cause of mortality owing to an infectious agent, accounting

for almost one-third of antimicrobial resistance (AMR) deaths annually. We aimed to identify

synergistic anti-TB drug combinations with the capacity to restore therapeutic efficacy against

drug-resistant mutants of the causative agent, *Mycobacterium tuberculosis*. We investigated

combinations containing the known translational inhibitors, spectinomycin (SPT) and fusidic

acid (FA), or the phenothiazine, chlorpromazine (CPZ), which disrupts mycobacterial energy

metabolism. Potentiation of whole-cell drug efficacy was observed in SPT-CPZ combinations.

This effect was lost against an *M. tuberculosis* mutant lacking the major facilitator superfamily

(MFS) efflux pump, Rv1258c. Notably, the SPT-CPZ combination restored SPT efficacy against

an SPT-resistant mutant carrying a g1379t point mutation in *rrs*, encoding the mycobacterial 16S

ribosomal RNA. Combinations of SPT with FA, which targets the mycobacterial elongation

factor G, exhibited potentiating activity against wild-type *M. tuberculosis.* Moreover, this

combination produced a marginal potentiating effect against both FA-monoresistant and SPT-

monoresistant mutants. Finally, combining SPT with the frontline anti-TB agents, rifampicin

(RIF) and isoniazid, resulted in enhanced activity *in vitro* and *ex vivo* against both drug-

susceptible *M. tuberculosis* and a RIF-monoresistant *rpoB* S531L mutant.These results support

the utility of novel potentiating drug combinations in restoring antibiotic susceptibility of *M.*

tuberculosis strains carrying genetic resistance to any one of the partner compounds.

INTRODUCTION

The increasing prevalence of multidrug-resistant tuberculosis (MDR-TB) – defined as resistance to the frontline anti-TB agents, isoniazid (INH) and rifampicin (RIF) – necessitates the urgent development and implementation of new anti-mycobacterial drugs and therapeutic strategies (1, 2). A number of anti-TB compounds are currently in the drug discovery pipeline, with several others in advanced preclinical development (3). However, with the exception of bedaquiline (BDQ) and delamanid, no TB-specific drugs have been introduced into clinical use within the past 40 years (4). Therefore, new options need to be explored to address the problem of drug resistance.

Novel combination regimens comprising standard anti-TB agents and repurposed drugs represent a logical approach, especially where the new partner drug has already been approved for other clinical indications (5-7). Recent advances in understanding the physicochemical properties that determine drug distributions within complex tissue and cellular (micro)environments (8, 9), together with improved methods for rapid selection of multiple potentially synergistic drug partners *in vitro* and *in vivo* (8, 10-12), suggest the potential for rational development of novel combination approaches. This is important since it might address the long-held concern that developing combinations should be avoided owing to the complexities inherent in ensuring simultaneous and sustained delivery of the optimal partner compounds to the same target site (11). The impact of pre-existing drug-resistance on the utility of new drug combinations presents an additional challenge, and is of particular concern when these combinations comprise current frontline anti-TB agents. To minimize the risks of exposing an individual to effective monotherapy, the likely pre-existence of resistance to individual drugs must be recognized, and informed combination approaches for drug therapies designed. These

should incorporate multiple attributes beyond simply selecting individual molecules based on

their biological activities as single agents (13).

In this study, we employed spectinomycin (SPT) as an anchor compound in combination

with other experimental antibiotics and existing frontline anti-TB agents. SPT is an

aminocyclitol antibiotic that inhibits protein synthesis by disrupting mRNA interactions with the

30S ribosome (14). Unlike other aminocyclitol antibiotics (including gentamycin and

kanamycin), SPT is not ototoxic (15) and has been used extensively in treating *Neisseria*

gonorrhoeae infections in patients who cannot tolerate first-line treatments (10). From the

perspecective of new regimen design, SPT has been shown in combination screens against *M.*

tuberculosis to synergize with several different classes of antimycobacterial compounds, both *in*

vitro and in a macrophage model (10). Unfortunately, a key liability undermining its utility as a

single agent is that SPT is subject to active efflux by *M. tuberculosis* – an observation which

motivated an elegant medicinal chemistry solution in the development of the spectinamides as

derivative "efflux-resistant" anti-TB antibiotics (16-18). Spectinamides are also known to

synergize with a variety of antibiotic classes (11) with lead spectinamide molecules, such as

1599, shown to be active against MDR *Mtb* strains and to synergize with existing and

experimental anti-TB drugs *in vivo* (19, 20). However, developing suitable spectinamide

formulations for therapeutic delivery remains an obstacle to the advancement of these

compounds as novel anti-TB agents (21, 22).

We investigated the utility of potentiating combinations, anchored by SPT, to circumvent

drug resistance and, potentially, restore (partial) susceptibility where genetically drug-resistant

mutants pre-exist for one of the partner compounds. We applied combination screens utilizing (i)

chlorpromazine (CPZ), a phenothiazine whose complex and unresolved mechanism of action

involving disruption of the mycobacterial electron transport chain (23) has been implicated in

efflux pump inhibition (24); and (ii) fusidic acid (FA), a translational inhibitor with demonstrated

(albeit moderate) activity *in vitro* (25, 26). FA was selected owing to its potential for

repositioning as anti-TB agent and because it possesses a unique mechanism of action:

specifically, inhibition of bacterial protein synthesis by binding to elongation factor G (EF-G)

(27). The antimicrobial-potentiating effect of FA with other antibiotics, including the frontline

anti-TB drug, ethambutol (EMB), as well as its lack of cross-resistance to other antimicrobial

classes, provided additional motivation for our choice of FA (12, 28). By testing these

combinations against both drug-susceptible *M. tuberculosis* H37Rv and selected drug-resistant

mutants, we explored new potentiating combinations and demonstrated the utility of developing

potent combinations against bacilli carrying pre-existing genetic resistance to either of the

partner drugs. In addition, this work revealed that the inclusion of SPT as third agent with the

existing first-line anti-TB drug combination of RIF and NIH restores activity *in vitro* against

defined pre-MDR mutants of *M. tuberculosis.*

MATERIALS AND METHODS

Chemicals and reagents

All chemicals and solvents were purchased from Sigma-Aldrich. Working solutions of all

antimicrobial agents were prepared in dimethyl sulfoxide (DMSO).

Bacterial strains and culture conditions

The laboratory strain, *M. tuberculosis* H37Rv, its derivative mutants, and a reporter strain which

has been used previously in high-throughput antimicrobial drug screening and constitutively

expresses green fluorescent protein (GFP), H37Rv pMSP*::eGFP* (29), were maintained as

freezer stocks. Strains were inoculated in standard Middlebrook 7H9 medium supplemented with

10% oleic acid-albumin-dextrose-catalase (OADC) (Difco) an incubated as stationary cultures at

37°C for approximately 3 days, sub-cultured, and incubated until culture density was

approximatly OD$_{600}$ ~0.5. A second reporter mutant, *M. tuberculosis*

H37Rv::(pSMYC::mCherry) (30), that constitutively expresses the mCherry fluorophore, was

grown in media containing 50 mg/L hygromycin. Cell suspensions were diluted to give an

expected final concentration of 10^5 cells/ml at the time of inoculation into the microplate for the

minimum inhibitory concentration (MIC) assays.

Drug susceptibility assays

Resazurin was used to determine the susceptibility of drugs against *M. tuberculosis* strains. The

MIC$_{90}$ for all drugs was determined by broth microdilution assay in 96-well microtiter plates

(31). Briefly, 2-fold serial dilutions of compounds were performed on clear, round-bottom 96-

well plates using 7H9-OADC medium. *M. tuberculosis* cultures, grown to to an OD$_{600}$ of 0.5 (~

10^8 cells/ml) and diluted 1,000-fold, were added at equal volume for a total volume of 100 µl per

well. The plates were sealed in zip-lock bags and incubated at 37 °C for 7 days, consistent with

EUCAST guidelines (32) and published literature (33) recommending that MIC plates should be

read after 7 and 14 days post inoculation. Resazurin dye was added and the plates incubated for a

further 24 h. Fluorescence readings, at excitation and emission wavelengths of 540 and 590 nm

respectively, were recorded using a SpectraMax i3x plate reader (Molecular Devices), and the

MIC$_{90}$, the lowest drug concentration to inhibit growth by more than 90%, was determined from

the dose-response curve.

Checkerboard assays

2D Checkerboard

Standard "two-dimensional" (2D) drug-drug interactions were determined by checkerboard

titration in a 96-well plate **(Fig. S1a)**. The 2D microdilution was carried out as described (34),

with slight modification. Briefly, the first drug (A) to be serially diluted was dispensed (2 µl)

along the x-axis (columns 3 to 11, row B) at a starting concentration 100 times higher than the

final concentration in the well, and (2 µl) per well of the second drug (B) was serially dispensed

along the y-axis (from row B to H) at a starting concentration 100 times higher than the final

concentration in the 96-well microtitre plate. The first column (column 1) and last column

(column 12) contained drug-free controls (with 1% DMSO as a diluent) and a control drug

concentration giving maximum inhibition, respectively. The second column from B2-H2 and

first row from A3-A11 contained individual drugs, thus providing the MIC for each drug alone in

each assay (each plate). The plates were placed in zip-lock bags and incubated at 37 °C for 7

days. Resazurin dye was then added and the plates incubated for a further 24 h. A visible colour

change from blue to pink indicated growth of bacteria, and the visualized MIC was defined as

the lowest concentration of drug that prevented growth (at which the colour change did not

occur) (31). Fluorescence readings (excitation: 544 nm, emission: 590 nm) were obtained using

BMG Labtech POLARstar Omega plate reader (BMG Labtech, Offenburg, Germany). The

mean fluorescence value for the "maximum inhibition" wells (column 12) was subtracted from

all other wells to control for background fluorescence. Percent inhibition was defined as 1-(test

well fluorescence units/mean fluorescence units of maximum inhibition wells) × 100 on day 8

after incubation. The lowest drug concentration effecting inhibition of 90% was considered the

$MIC_{90.}$ In addidition, synergy was interpreted according to the sum of fractional inhibitory

concentration (ΣFIC). The fractional inhibitory concentration for each compound was calculated

as follows (35): FIC_A _(MIC of compound A in the presence of compound B)/(MIC of

compound A alone), where FIC_A is the fractional inhibitory concentration of compound A.

Similarly, the FIC for compound B was calculated. The ΣFIC was calculated as $FIC_A + FIC_B$.

Synergy was defined by values of $\Sigma FIC \leq 0.5$, antagonism by $\Sigma FIC > 4.0$, and no interaction by

ΣFIC values from 0.5 to 4.0 (36).

3D Checkerboard

In the three-drug ("three-dimensional", 3D) combinations **(Fig. S1b)**, microdilutions for the first

two drugs were initially set up principally following the standard 2D checkerboard assay

protocol described above. The third drug (2 µl) was then added at a starting concentration 100

times higher than the final concentration in the well as an overlay at five sub-inhibitory

concentrations ranging from 1/32 to 1/2 of the single-drug MIC. Well A2 on all plates contained

the third drug only, providing the single-drug MIC for the third drug in each assay (set of 5

plates). After inoculation with a log-phase culture, OD_{600} of 0.5 ($\sim 10^8$ cells/ml) of *M.*

tuberculosis, 50 µl to each well, the plates were placed in zip-lock bags and incubated for 7 days

at 37°C before addition of resazurin. The plates were further incubated for 24 h and the results

were read in the BMG Labtech plate reader (excitation 544 nm; emission 590 nm) (33). The %

inhibition was calculated as described earlier.

Macrophage assays

Cell culture and maintenance

Human promonocytic THP-1 cells were maintained in RPMI-1640 medium (Sigma)

supplemented with 10% fetal bovine serum (FBS, Invitrogen) at an initial density of 8×10^5

cells/ml at 37^oC in an humidified, 5% CO_2 atmosphere. Prior to plating of the cells, viability was

assessed by trypan blue exclusion method (37). Maturation of THP-1 cells into macrophages was

induced by adding 200 nM PMA (phorbol 12-myristate 13-acetate; Sigma) in cell culture

medium for 24 h. Differentiated macrophages were then washed three times with pre-warmed

phosphate buffer saline (PBS) to remove the PMA and replenished with cell culture medium.

Infection of macrophages and drug treatment

To check the efficacy of drugs in macrophages, 5×10^4 THP-1 cells/well (100 µl final volume) in

96-well flat-bottomed tissue culture plates were differentiated into macrophages. To infect

macrophages, an exponentially growing *M. tuberculosis* H37Rv::(pSMYC::mCherry) culture

was harvested by centrifugation and washed twice with PBS. The pellet was resuspended in PBS

and passed through a 5 µm filter to generate a suspension of single-cell bacilli. The bacterial

suspension density was estimated by measuring OD at 600 nm, corresponding an $OD_{600}\sim0.5$ to

1×10^8 CFU/ml. Infection medium comprised cell culture medium containing a number of

bacteria required to achieve a multiplicity of infection (MOI) of 5:1 (5 bacilli for every THP-1

cell) (38). The macrophage cells were overlaid with infection medium and incubated at 37°C in

5% CO_2 for the phagocytic period of 3 hours. Cells were then washed gently and thoroughly

with prewarmed PBS to remove extracellular bacteria. The cells from three wells were lysed by

adding triton X-100 (0.05% in PBS) and the lysate was plated onto 7H10 to score colony

forming units (CFUs) for untreated day zero. The used as untreated control were refreshed with

cell culture medium. The cells in remaining wells were treated with the indicated antibiotic either

alone or in combination at $1\times$ MIC_{90} or $5\times$ MIC_{90}, as determined in liquid culture. Hygromycin at

50 mg/L was added into the cell culture medium for all wells to maintain the plasmid expressing

mCherry .

Fluorescence measurement

The fluorescence of the mCherry reporter (excitation: 590 nm, emission: 610 nm) was measured

at different time points on a BMG Labtech plate reader.

CFU enumeration

To estimate the numbers of live bacilli after drug treatment, untreated and drug-treated *M.*

tuberculosis-infected cells were lysed in triton X-100 (0.05% in PBS) on days 2, 4 and 6, and

serial dilutions of the cell lysate were plated onto 7H10 agar. Colonies were counted after 3-4

weeks of incubation at 37 $^{\circ}$C.

Statistical Analyses

Statistical analyses were performed using Prism 9.0.0.121 (GraphPad). Means were compared

via ANOVA, with post-test evaluation using Dunnett's or Bonferroni's test. *P* values are

abbreviated as follows: *, *P<0.05*; **, *P<0.01*; ****P<0.001*

RESULTS

CPZ potentiates SPT activity by inhibiting Rv1258c-mediated efflux. The combination of

SPT and CPZ was previously reported as synergistic against *M. smegmatis* (39). When tested

against wild-type *M. tuberculosis* H37Rv **(Fig. 1a),** the same combination yielded a ΣFIC value

of 0.09 **(Table 1)**, confirming strong synergy (36). We investigated whether this effect resulted

from CPZ-mediated disruption of the activity of the Major Facilitator Superfamily (MFS) efflux

pump, Rv1258c, which has been implicated in innate resistance to SPT (17). We performed

checkerboard assays using the efflux-defective Δ*Rv1258c* ("tap") knockout mutant, which had

been used in the development of the spectinamides (SPD) (17), and its complemented derivative,

Δ*Rv1258c* pCRS4. Both mutants exhibited the same MIC$_{90}$ of 22 mg/L for CPZ **(Fig. 1 & Table**

1), whereas the Δ*Rv1258c* knockout mutant was hypersusceptible to SPT, returning an

approximately 6-fold lower MIC$_{90}$ of 3.9 mg/L **(Fig. 1b, Table 1 and Table S1)**, as observed

previously (40). Notably, the synergy detected on exposing wild-type *M. tuberculosis* to a

combination of CPZ and SPT **(Fig. 1a**) was eliminated in the Δ*Rv1258c* mutant **(Fig. 1b)** –

which yielded a ΣFIC value of 0.75 (**Table 1**) – but was restored in the complemented Δ*Rv1258c*

pCRS4 strain **(Fig. 1c),** with a ΣFIC of 0.12 **(Table 1)**. Previous studies have reported no

significant alteration in *Rv1258c* transcription in response to CPZ treatment (41). Therefore, our

observations suggested that CPZ treatment abrogated efflux-mediated intrinsic resistance to SPT

in wild-type *M. tuberculosis* in a manner dependent on Rv1258c, perhaps as a result of CPZ-

mediated inhibition of energy metabolism (23).

CPZ synergizes with compounds other than SPT, but SPT potentiation arises solely from

CPZ-mediated inhibition of Rv1258c. To determine whether the synergy observed was specific

for SPT or might apply to other antimycobacterial agents, CPZ was applied as the anchor

compound in combination assays with a panel of anti-TB antibiotics of different classes and

mechanisms of action (**Table S2**). Of the eight compounds tested with CPZ, four exhibited clear

synergy (ΣFIC ≤ 0.5): the frontline agents, RIF and INH, which returned ΣFICs of 0.37 and 0.5,

respectively, and bedaquiline (BDQ) and nalidixic acid, both of which gave ΣFICs ≤ 0.25. In

contrast, no potentiation was observed with KAN or the fluoroquinolones, ciprofloxacin (CIP)

and levofloxacin (LEV), all of which yielded ΣFICs > 0.5.

In a complementary approach, we also investigated if the potentiating effect observed

with the SPT-CPZ combination was unique to CPZ. To this end, we assayed SPT in combination

with an expanded panel of antimycobacterial agents (**Fig. S2** and **Table S3**). SPT was found to

synergize with only two of the eleven compounds tested: erythromycin (ERY), a macrolide

targeting the ribosome, and verapamil (VER), a cationic amphiphile which was originally

considered an *M. tuberculosis* efflux pump inhibitor but has been shown recently to disrupt

membrane function (42). RIF, the mycobacterial RNA polymerase inhibitor, was just beyond the

threshold determining synergistic activity.

To ascertain if inhibition of Rv1258c-mediated efflux resulted in the observed compound

synergies, the Δ*Rv1258c* mutant was tested for hypersensitivity to a corresponding panel of anti-

TB agents (**Fig. S3** and **Table S1**). The spectinamide 1599 had the same MIC_{90} for both wild-

type *M. tuberculosis* H37Rv and the Rv1258c mutant, reflecting its successful modification to

avoid Rv1258c-mediated efflux (43). Of the other 11 compounds tested, only SPT was

associated with hypersensitivity in the Rv1258c knock-out mutant, returning an MIC_{90} value of

0.39 mg/L compared to the MIC_{90} value of 62-125 mg/L against wild-type strain. In

combination, these results strongly support the inference that the synergy detected with the CPZ-

SPT combination arises from CPZ-mediated inhibition of Rv1258c.

The CPZ-SPT combination partially restores SPT sensitivity in an SPT-resistant mutant. A

spontaneous SPT-resistant mutant (SPT^R) carrying a g1379t point mutation in the mycobacterial

16S ribosomal RNA, *rrs*, was associated with >64-fold increase in the SPT MIC_{90} (**Table S1).** In

contrast, the activity of CPZ remained at the wild-type concentration for this strain, consistent

with a mechanism of action of CPZ that was independent of *rrs* inhibition (44). While the SPT^R

mutant was resistant to SPT at concentrations >248 mg/L in the absence of CPZ, combinations

utilizing CPZ at sub-MIC concentrations ([CPZ] ≤ 11.1 mg/L) restored SPT sensitivity, at least

partially (**Fig. 1d**). These results suggested the capacity for synergistic combinations to restore

drug activity against mutant strains genetically resistant to either of the partner compounds.

Assessing synergy with SPT and fusidic acid, two antibiotics acting on the mycobacterial

translational machinery. The systematic application of drug combinations can reveal

synergistic interactions. One form of synergy occurs when drug(s) which perturb normal cell

physiology trigger (compensatory) cellular responses that can, in turn, affect (potentiate) the

activities of other drugs (45). Nichols *et al*. demonstrated that the synergistic interaction between

sulfamethoxazole and trimethoprim was a result of the two drugs targeting tetrahydrofolate

biosynthesis (46). With this in mind, a combination comprising SPT and FA – translation

inhibitors which act at discrete steps of the elongation process – was tested against wild-type *M.*

tuberculosis and two resistant strains, FA[R] and SPT[R]. Isolation of spontaneous FA[R] mutants *in*

vitro yielded a single strain on $25\times$ MIC FA at a frequency $\sim10^{-8}$. Whole-genome sequencing

identified a c1384t (H462Y) substitution in *fusA1* (*Rv0684*), encoding the essential

mycobacterial elongation factor G (EF-G) (48). The histidine residue is highly conserved across

multiple bacterial species; therefore, using the *T. thermophilus* structure as template (49), it can

be inferred that *M. tuberculosis* H462 corresponds to *T. thermophilus* H458 (50), mutations of

which are likely to alter the FA-binding pocket (51). In MIC assays, the H462Y mutant

consistently yielded an $MIC_{90} \geq 25$ mg/L, confirming heritable FA[R] **(Table 2)**.

With these strains in hand, we evaluated the interaction between SPT and FA and,

furthermore, assessed whether this combination – which is synergistic against the parental, drug-

susceptible *M. tuberculosis* H37Rv – might counter pre-existing genentic resistance to either

compound. The combination of SPT and FA returned a ΣFIC value of 0.50 **(Table 2)** against

wild-type H37Rv; upon addition of FA at sub-MIC concentration ([FA] = 0.16 mg/L), the MIC_{90}

of SPT exhibited a $\sim$4-fold decrease from 62 mg/L to 16.3 mg/L **(Fig. 2a)**, p<0.001. FA at sub-

inhibitory concentration ([FA] = 0.32 and [FA] = 0.16 mg/L) enhanced the activity of SPT

([SPT] = 62 mg/L) against a SPT[R] mutant. The inhibitory effect was significantly different from

the results observed with similar concentrations of SPT in the absence of FA ([FA] = 0 mg/L),

p<0.001 **(Fig. 2b)**. Although the calculated sum FIC did not satisfy the criterion for "synergistic"

(ΣFIC$\leq$0.5) **(Table 2),** the effect was marked and reproducible in two independent biological

replicates **(Fig. 2b)**. Notably, the same combination did not return enhanced activity against the

FA^R mutant **(Fig 2c)**, strongly suggesting that FA was the major contributor to the SPT-FA

combination. A summary of the inhibitory effects of the CPZ-SPT and FA-SPT combinations is

presented in **Table 3**.

Confirmation that fluorescence intensities correlate with cell density readings. The

centrality of the resazurin microtiter assay (REMA) in determining the synergies reported in this

study demanded orthogonal evidence supporting the claimed results. To this end, a 96-well based

antimycobacterial assay was performed with a selected panel of drugs having different

mechanisms of action **(Fig. S4)**. The experiments were conducted using the H37Rv::GFP

reporter mutant in which expression of the fluorophore is constitutive (29). After 8 days of

incubation in the presence of 2-fold dilutions of the antimicrobial agents, GFP and resazurin

fluorescence intensities were determined, and the corresponding optical density read-outs

recorded in parallel. Strong agreement was discerned when comparing GFP and OD$_{600nm}$

methods with the standard resazurin read-out **(Fig. S4a & b)**, supporting the use of the resazurin

assay for both MIC and FIC determinations.

A three-drug combination comprising SPT, RIF and INH enhances *in vitro* activity against

***M. tuberculosis*.** The premise that synergistic combinations might be usefully applied to

overcome existing drug resistance was further explored using SPT in combination with the

frontline anti-TB agents, RIF and INH. In a 2D pairwise screening assay, we evaluated two-drug

permutations of RIF, INH and SPT. The RIF-INH combination showed no interaction **(Fig. S5a)**

while sub-MIC concentrations of INH (0.125 mg/L) and RIF (0.003 mg/L) halved the effective

SPT concentration from 125 mg/L to 62 mg/L **(Fig S5b & S5c)**. To leverage the potential effect

of SPT, a 3D combination assay was performed in which RIF and INH were titrated against

decreasing sub-MIC$_{90}$ concentrations of SPT (1/2×, 1/4×, 1/8×, 1/16× and 1/32× MIC$_{90}$), using

the format illustrated in **Fig. S1b**. When tested against drug-susceptible *M. tuberculosis* H37Rv,

RIF at sub-MIC ([RIF] = 0.003 mg/L) plus SPT at both $1/4\times$ and $1/2\times$ MIC resulted in an 8-fold

decrease in the effective concentration of INH from 0.25 mg/L to 0.03 mg/L **(Fig. S6)**. A kill

kinetics assay showed that the addition of SPT to the RIF-INH combination elicited ~1 $\log_{10}$ unit

reduction (p < 0.001) in the viable bacillary population following 8-day exposure to the three-

drug combination **(Fig. S7a)**. In contrast, when the $\Delta Rv1258c$ "tap" knock-out mutant was

tested, RIF at sub-MIC ([RIF] = 0.003 mg/L) plus $1/2\times$ MIC SPT resulted in only a ~3-fold

decrease in the effective concentration of INH – from 0.25 mg/L to 0.09 mg/L **(Fig. S7b)**, again

implicating Rv1258c in intrinsic antibiotic resistance in *M. tuberculosis*.

The RIF-INH-SPT combination is active in *M. tuberculosis*-infected macrophages and

against monoresistant pre-MDR strains.

Since *M. tuberculosis* survives and replicates in macrophages (52), the synergy of RIF-INH plus

SPT was evaluated against intracellular bacilli in *M. tuberculosis*-infected THP-1 cells. This

three-drug combination showed inhibitory activity at $1\times$ MIC_{90} of the combined drugs **(Fig. S8)**.

In contrast, the inhibitory effect was reduced when similar concentrations of each drug were

applied individually, or when the standard RIF-INH combination was used without SPT. The

intracellular activity of this triple combination was further confirmed by CFU enumeration **(Fig.**

S9), which revealed a 2-$\log_{10}$ decrease in CFU/ml when $1\times$ MIC_{90} RIF-INH-SPT was applied

compared to the untreated control.

To evaluate the efficacy of RIF-INH-SPT against known drug-resistant strains, the

combination was tested against two pre-MDR *M. tuberculosis* mutants: a RIF-monoresistant

mutant carrying the common *rpoB* S531L allele, and an INH-monoresistant strain harbouring a -

c15t mutation in the *inhA* promoter region that confers low-level INH resistance **(Fig. 3)**.

Duplicate checkerboard experiments showed that, for the $rpoB^{S531L}$ mutant, addition of 1/2× MIC

SPT to the RIF-INH plate resulted in a decrease in the effective concentrations of both RIF (20

to 10 mg/L) and INH (0.25 to 0.03 mg/L) (**Fig. 3a** and **3b**), indicating partial restoration of drug

susceptibility in the presence of SPT. Enhanced susceptibility was also observed for the INH^R

mutant, albeit to a lesser extent: a sub-MIC RIF concentration of 0.01 mg/L and INH at 2.5 mg/L

achieved ~80 % bacterial inhibition (**Fig. 3c and 3d**).

DISCUSSION

Notwithstanding recent promising claims (53), the bacterial capacity for acquisition of resistance

by multiple mechanisms means that it is difficult, perhaps even conceivably impossible, to

overcome antibiotic resistance sustainably (54). Different approaches can be used to circumvent

resistance transiently, ensuring antibiotic efficacy despite the pre-existence of resistant

organisms in an infecting population. Combination therapy represents one such approach.

In a previous study, Chen *et al*. demonstrated a synergistic interaction between SQ109

and RIF when tested against RIF^R isolates: at 0.5× MIC, SQ109 was able to increase RIF's

activity against *de facto* resistant organisms in a dose-dependent manner (34). Recently, Yang *et*

al reported the enhanced efficacy of the imipenem-colistin combination against multiple drug-

resistant *Enterobacter cloacae in vitro* and in an infection model (55). Relativly few studies have

been undertaken to illustrate the association between potentiating drug interactions and the

ability of the particular drug combination to overcome pre-existing genetic resistance. In a

clinical study, Ankomah *et al*. suggested that drugs acting synergistically can prevent treatment

failure even when bacteria resistant to one of these drugs are present at the beginning of therapy

(56). Our interaction studies between SPT and CPZ reaffirmed the susceptibility of SPT to

Rv1258c-mediated efflux, an observation which suggests that efforts to modify SPT – including

through novel chemical modifications that engineer resistance to efflux (17) – should be pursued.

We tested the susceptibility of the Δ*Rv1258c* mutant to a small panel of anti-

mycobacterial compounds with different mechanisms of action and observed that SPT alone was

associated with hypersusceptibility. A similar hypersusceptibility phenotype was achieved

against wild-type *M. tuberculosis* H37Rv via chemical potentiation of SPT using CPZ as the

combination agent. However, the potentiating effect of CPZ was not specific to SPT and was

instead observed for a handful of other agents. This suggests the likelihood that CPZ might

disable more than one intrinsic resistance mechanism in *M. tuberculosis*. Further work is

required to ascertain the precise mechanism for each compound, with evidence to date

implicating multiple potential efflux systems in intrinsic resistance to INH, RIF, and the

fluoroquinolones (57). For BDQ, the multi-substrate RND-family transporter, MmpS5-MmpL5,

appears a strong candidate based on previous reports (58).

Our results revealed synergy between FA and SPT against drug-susceptible bacteria via a

mechanism independent of the efflux inhibition seen with SPT-CPZ. Notably, the same FA-SPT

combination exerted an enhanced inhibitory effect against a genotypically confirmed SPT[R]

mutant compared to a FA[R] mutant. Although there is no definitive explanation for this finding,

we postulate that the relative potency of FA (~0.63 mg/L) against the drug-susceptible H37Rv

parent compared to that of SPT (~50 mg/L) could impact the FA-SPT combination against the

SPT[R] mutant, restoring susceptibility. A similar effect was not evident in a SPT-FA combination

against FA[R] mutant owing to the diminished activity of FA and high MIC_{90} of SPT. Of interest is

the impact of individual active drugs in driving synergy.

Other explanations for potentiation include the sustained drug pressure emanating from

the drug interactions which leads to an increased effective dose of the drug combination.

Moreover, some studies have demonstrated that the drug susceptibility of pathogens can be

significantly enhanced as a result of a reduced efflux pump efficiency either by genetic

manipulation (59), or addition of efflux pump inhibitors (60, 61). The clinical relevance of this

finding is that, despite the existence of bacterial resistance against a combination partner, it

would still be possible to achieve optimal therapeutic outcomes via the use of appropriate potent

drug combinations.

Previous work has demonstrated the potential of having three-drug combinations when

compared to individual or two-drug regimens (62, 63). Recently, Tekin *et al.* reported that

combinations of three different antibiotics can often overcome antimicrobial resistance to

antibiotics, even when none of the three antibiotics on their own — or even two of the three

together — is effective (64). In addition, based on drug interaction studies, Ramon-Garcia *et al.*

hypothesized that the synergistic activity of the triplet combination might have multiplicative

effects (10). Here, SPT was deployed as part of a three-drug regimen which also included RIF

and INH, the two drugs that form the cornerstone of TB treatment. Other studies have shown the

interaction between RIF and INH against *M. tuberculosis* to have no interaction or to be mildly

antagonistic (8, 65). The inclusion of SPT in this drug regimen was underpinned by reports that

24 out of 70 random combinations tested were synergistically active in *M. smegmatis* (10). This

suggests a large unexplored pool of synergistic combinations. Notably, SPT exhibited synergy

with several compounds both *in vitro* and *ex vivo* (10), even though the compound has high

MIC$_{90}$ against *M. tuberculosis* when administered on its own (17).

In the three-drug combination assay, synergy resulted when sub-inhibitory concentrations

(1/2× and 1/4× MICs) of SPT were titrated into media containing RIF and INH. This finding

correlated well with the results of time-kill kinetics. However, the time-kill assay suggested that

the inhibitory effect of the three-drug interaction was bacteriostatic ($0 \geq \log_{10}$ CFU <3 reduction)

and not bactericidal. This observation reveals that the combination of RIF and INH — two

bactericidal drugs that are most potent against actively dividing cells — shows bacteriostatic

effects. Furthermore, the inhibition of growth induced by a bacteriostatic drug, SPT, results in an

overall static effect when the drug is used in combination with a bactericidal drug. Other studies

have shown that, in similar interactions, the resulting effect achieves a more efficient clearance at

lower concentrations (40).

In attempting to exploit synergy for potential optimal treatment outcomes, an

investigation of the RIF-INH plus SPT interaction was performed in *rpoB* and *inhA* mutants. The

RIF^R *rpoB* mutant had an MIC value >2000 times the MIC$_{90}$ for drug-susceptible *M.*

tuberculosis. Notably, addition of 1/2× MIC of SPT restored partial drug efficacy against this

resistant mutant. As with the drug susceptible H37Rv strain, a mechanistic explanation for the

synergy observed using the RIF-INH plus SPT combination against the *rpoB* mutant is presently

lacking. INH targets mycobacterial cell envelope biosynthesis, possibly enhancing permeation of

SPT into the bacilli. However, access alone may not necessarily contribute to the synergistic

interaction. Chen *et al.* reported synergy between SQ109, a presumed cell envelope inhibitor,

and RIF (34). On the other hand EMB, which also affects mycobacterial cell envelope synthesis,

did not exhibit synergy with RIF (66).

Prior reports have shown RIF to be an efficient inducer of cytochrome P450 (CYP 450), a

superfamily of haem-containing enzymes involved in the biosynthesis of compounds such as

sterols, steroids, and fatty acids as well as detoxification of xenobiotics and chemicals (67). RIF

has been linked with the induction of CYP both in humans and in *M. tuberculosis* (68). The

elevated levels of CYP have been associated with drug resistance due to the enhanced rate of

elimination of the drugs by metabolism and detoxification pathways. INH, on the other hand,

inhibits CYP in *M. tuberculosis* (68). This ability of INH to inhibit CYP may contribute to

synergy in the RIF-INH plus SPT combination when the active form of INH is not rapidly

eliminated inside *M. tuberculosis* and when SPT acts by further reducing the activiy of CYPs.

There are prospects to combine SPT with RIF-INH in treatment regimens. SPT is given

by intramuscular injection to achieve therapeutic concentrations in serum of about 100 mg/L one

hour after a single 2 g dose. An over 4-fold increase in its effectiveness within a triple SPT-RIF-

INH combination , as indicated by these data, would potentially allow for oral formulation, a

critical delivery format when administering treatment to TB out-patients. In summary, these *in*

vitro and *ex vivo* results suggests that the RIF-INH plus SPT triple-combination may be an

effective therapeutic option for the treatment of both the drug susceptible and resistant *M.*

tuberculosis infections. They also reinforce a growing body of evidence supporting the utility of

drug potentiation strategies in improving treatment outcomes.

ACKNOWLEDGEMENTS

Research reported in this publication was supported by the Strategic Health Innovation

Partnerships (SHIP) Unit of the South African Medical Research Council (SAMRC) with funds

received from the South African Department of Science Innovation (to KC and DFW). We

acknowledge the support of the University of Cape Town, the SAMRC (to KC and VM), the

South African Research Chairs Initiative (to KC) and Department of Science Innovation

administered through the National Research Foundation, the Schlumberger Foundation Faculty

for the Future scheme (to EK and AW), and the Research Council of Norway (to DFW). We

thank Prof. J. A. Aínsa for providing the *M. tuberculosis* Δ*Rv1258c* ("tap") knockout mutant and

its complemented derivative, Prof. R. E. Lee for providing stocks of the spectinamide, 1599, Dr.

Krupa Naran for advice with checkerboard assays and Dr. Mandy Mason for critical review of

the manuscript.

FUNDING

The University of Cape Town, Novartis Research Foundation, South African Medical Research

Council, Strategic Health Innovations Partnerships Unit of the South African Medical Research

Council, South African Research Chairs Initiative of the Department of Science and Innovation

administered through the South African National Research Foundation, Carnegie Mellon

Scholarship Awards, and Research Council of Norway.

TRANSPARENCY DECLARATIONS

None   to   declare.

REFERENCES

1.   **Sullivan T, Ben Amor Y.** 2016. Global Introduction of New Multidrug-Resistant Tuberculosis Drugs-Balancing Regulation with Urgent Patient Needs. Emerg Infect Dis **22**.

2.   **Hoagland DT, Liu J, Lee RB, Lee RE.** 2016. New agents for the treatment of drug-resistant Mycobacterium tuberculosis. Adv Drug Deliv Rev **102:**55-72.

3.   **Evans JC, Mizrahi V.** 2018. Priming the tuberculosis drug pipeline: new antimycobacterial targets and agents. Curr Opin Microbiol **45:**39-46.

4.   **Zumla A, Nahid P, Cole ST.** 2013. Advances in the development of new tuberculosis drugs and treatment regimens. Nat Rev Drug Discov **12:**388-404.

5.   **Nzila A, Ma Z, Chibale K.** 2011. Drug repositioning in the treatment of malaria and TB. Future Med Chem **3:**1413-1426.

6.   **Oprea TI, Bauman JE, Bologa CG, Buranda T, Chigaev A, Edwards BS, Jarvik JW, Gresham HD, Haynes MK, Hjelle B, Hromas R, Hudson L, Mackenzie DA, Muller CY, Reed JC, Simons PC, Smagley Y, Strouse J, Surviladze Z, Thompson T, Ursu O, Waller A, Wandinger-Ness A, Winter SS, Wu Y, Young SM, Larson RS, Willman C, Sklar LA.** 2011. Drug Repurposing from an Academic Perspective. Drug Discov Today Ther Strateg **8:**61-69.

7.   **Zumla AI, Schito M, Maeurer M.** 2014. Advancing the portfolio of tuberculosis diagnostics, drugs, biomarkers, and vaccines. Lancet Infect Dis **14:**267-269.

8.   **Cokol M, Kuru N, Bicak E, Larkins-Ford J, Aldridge BB.** 2017. Efficient measurement and factorization of high-order drug interactions in Mycobacterium tuberculosis. Sci Adv **3:**e1701881.

9.   **Greenwood DJ, Dos Santos MS, Huang S, Russell MRG, Collinson LM, MacRae JI, West A, Jiang H, Gutierrez MG.** 2019. Subcellular antibiotic visualization reveals a dynamic drug reservoir in infected macrophages. Science **364:**1279-1282.

10. **Ramon-Garcia S, Ng C, Anderson H, Chao JD, Zheng X, Pfeifer T, Av-Gay Y, Roberge M, Thompson CJ.** 2011. Synergistic drug combinations for tuberculosis therapy identified by a novel high-throughput screen. Antimicrob Agents Chemother **55:**3861-3869.

11. **Bruhn DF, Scherman MS, Liu J, Scherbakov D, Meibohm B, Bottger EC, Lenaerts AJ, Lee RE.** 2015. In vitro and in vivo Evaluation of Synergism between Anti-Tubercular Spectinamides and Non-Classical Tuberculosis Antibiotics. Sci Rep **5:**13985.

12. **Yilancioglu K, Cokol M.** 2019. Design of high-order antibiotic combinations against M. tuberculosis by ranking and exclusion. Sci Rep **9:**11876.

13. **Brooks BD, Brooks AE.** 2014. Therapeutic strategies to combat antibiotic resistance. Adv Drug Deliv Rev **78:**14-27.

14. **Hoeksema H, Knight JC.** 1975. The production of dihydrospectinomycin by Streptomyces spectabilis. J Antibiot (Tokyo) **28:**240-241.

15. **Novak E, Gray JE, Pfeifer RT.** 1974. Animal and human tolerance of high-dose intramuscular therapy with spectinomycin. J Infect Dis **130:**50-55.

16. **Cully M.** 2014. Antibacterial drugs: Redesigned antibiotic combats drug-resistant tuberculosis. Nat Rev Drug Discov **13:**256.

17. **Lee RE, Hurdle JG, Liu J, Bruhn DF, Matt T, Scherman MS, Vaddady PK, Zheng Z, Qi J, Akbergenov R, Das S, Madhura DB, Rathi C, Trivedi A, Villellas C, Lee RB, Rakesh, Waidyarachchi SL, Sun D, McNeil MR, Ainsa JA, Boshoff HI, Gonzalez-Juarrero M, Meibohm B, Bottger EC, Lenaerts AJ.** 2014. Spectinamides: a new class of semisynthetic antituberculosis agents that overcome native drug efflux. Nat Med **20:**152-158.

18. **Liu J, Bruhn DF, Lee RB, Zheng Z, Janusic T, Scherbakov D, Scherman MS, Boshoff HI, Das S, Rakesh, Waidyarachchi SL, Brewer TA, Gracia B, Yang L, Bollinger J, Robertson GT, Meibohm B, Lenaerts AJ, Ainsa J, Bottger EC, Lee RE.** 2017. Structure-Activity Relationships of Spectinamide Antituberculosis Agents: A Dissection of Ribosomal Inhibition and Native Efflux Avoidance Contributions. ACS Infect Dis **3:**72-88.

19.   **Robertson GT, Scherman MS, Bruhn DF, Liu J, Hastings C, McNeil MR, Butler MM,**
Bowlin TL, Lee RB, Lee RE, Lenaerts AJ. 2017. Spectinamides are effective partner agents for
the treatment of tuberculosis in multiple mouse infection models. J Antimicrob Chemother
72:770-777.
20.   **Bruhn DF, Scherman MS, Liu J, Scherbakov D, Meibohm B, Böttger EC, Lenaerts AJ, Lee**
RE. 2015. In vitro and in vivo evaluation of synergism between anti-tubercular spectinamides
and non-classical tuberculosis antibiotics. Scientific reports **5:**13985.
21.   **Rathi C, Lukka PB, Wagh S, Lee RE, Lenaerts AJ, Braunstein M, Hickey A, Gonzalez-**
Juarrero M, Meibohm B. 2019. Comparative pharmacokinetics of spectinamide 1599 after
subcutaneous and intrapulmonary aerosol administration in mice. Tuberculosis (Edinb) **114:**119-
122.
22.   **Stewart IE, Lukka PB, Liu J, Meibohm B, Gonzalez-Juarrero M, Braunstein MS, Lee RE,**
Hickey AJ. 2019. Development and Characterization of a Dry Powder Formulation for Anti-
Tuberculosis Drug Spectinamide 1599. Pharm Res **36:**136.
23.   **Weinstein EA, Yano T, Li LS, Avarbock D, Avarbock A, Helm D, McColm AA, Duncan K,**
Lonsdale JT, Rubin H. 2005. Inhibitors of type II NADH:menaquinone oxidoreductase
represent a class of antitubercular drugs. Proc Natl Acad Sci U S A **102:**4548-4553.
24.   **Jaiswal I, Jain A, Verma SK, Singh P, Kant S, Singh M.** 2017. Effect of efflux pump
inhibitors on the susceptibility of Mycobacterium tuberculosis to isoniazid. Lung India **34:**499-
505.
25.   **Castanheira M, Watters AA, Bell JM, Turnidge JD, Jones RN.** 2010. Fusidic acid resistance
rates and prevalence of resistance mechanisms among Staphylococcus spp. isolated in North
America and Australia, 2007-2008. Antimicrob Agents Chemother **54:**3614-3617.
26.   **Collignon P, Turnidge J.** 1999. Fusidic acid in vitro activity. Int J Antimicrob Agents **12 Suppl**
2:S45-58.
27.   **Turnidge J.** 1999. Fusidic acid pharmacology, pharmacokinetics and pharmacodynamics. Int J
Antimicrob Agents **12 Suppl 2:**S23-34.
28.   **Fuursted K, Askgaard D, Faber V.** 1992. Susceptibility of strains of the Mycobacterium
tuberculosis complex to fusidic acid. APMIS **100:**663-667.
29.   **Srivastava V, Rouanet C, Srivastava R, Ramalingam B, Locht C, Srivastava BS.** 2007.
Macrophage-specific Mycobacterium tuberculosis genes: identification by green fluorescent
protein and kanamycin resistance selection. Microbiology **153:**659-666.
30.   **Carroll P, Schreuder LJ, Muwanguzi-Karugaba J, Wiles S, Robertson BD, Ripoll J, Ward**
TH, Bancroft GJ, Schaible UE, Parish T. 2010. Sensitive detection of gene expression in
mycobacteria under replicating and non-replicating conditions using optimized far-red reporters.
PLoS One **5:**e9823.
31.   **Montoro E, Lemus D, Echemendia M, Martin A, Portaels F, Palomino JC.** 2005.
Comparative evaluation of the nitrate reduction assay, the MTT test, and the resazurin microtitre
assay for drug susceptibility testing of clinical isolates of Mycobacterium tuberculosis. Journal of
Antimicrobial Chemotherapy **55:**500-505.
32.   **Schon T, Werngren J, Machado D, Borroni E, Wijkander M, Lina G, Mouton J, Matuschek**
E, Kahlmeter G, Giske C, Santin M, Cirillo DM, Viveiros M, Cambau E. 2020.
Antimicrobial susceptibility testing of Mycobacterium tuberculosis complex isolates - the
EUCAST broth microdilution reference method for MIC determination. Clin Microbiol Infect
doi:10.1016/j.cmi.2020.07.036.
33.   **Martin A, Camacho M, Portaels F, Palomino JC.** 2003. Resazurin microtiter assay plate
testing of Mycobacterium tuberculosis susceptibilities to second-line drugs: rapid, simple, and
inexpensive method. Antimicrob Agents Chemother **47:**3616-3619.
34.   **Chen P, Gearhart J, Protopopova M, Einck L, Nacy CA.** 2006. Synergistic interactions of
SQ109, a new ethylene diamine, with front-line antitubercular drugs in vitro. J Antimicrob
Chemother **58:**332-337.

35.   **Ramon-Garcia S, Martin C, Ainsa JA, De Rossi E.** 2006. Characterization of tetracycline
564   resistance mediated by the efflux pump Tap from Mycobacterium fortuitum. J Antimicrob
565   Chemother **57:**252-259.
566   36.   **Odds FC.** 2003. Synergy, antagonism, and what the chequerboard puts between them. J
567   Antimicrob Chemother **52:**1.
568   37.   **Strober W.** 2001. Trypan blue exclusion test of cell viability. Curr Protoc Immunol **Appendix
569   3:**Appendix 3B.
570   38.   **Stokes RW, Doxsee D.** 1999. The receptor-mediated uptake, survival, replication, and drug
571   sensitivity of Mycobacterium tuberculosis within the macrophage-like cell line THP-1: a
572   comparison with human monocyte-derived macrophages. Cell Immunol **197:**1-9.
573   39.   **Kigondu EM, Njoroge M, Singh K, Njuguna N, Warner DF, Chibale K.** 2014. Synthesis and
574   synergistic antimycobacterial screening of chlorpromazine and its metabolites. MedChemComm
575   **5:**502-506.
576   40.   **Lee A, Mao W, Warren MS, Mistry A, Hoshino K, Okumura R, Ishida H, Lomovskaya O.**
577   2000. Interplay between efflux pumps may provide either additive or multiplicative effects on
578   drug resistance. J Bacteriol **182:**3142-3150.
579   41.   **Li P, Gu Y, Li J, Xie L, Li X, Xie J.** 2017. Mycobacterium tuberculosis Major Facilitator
580   Superfamily Transporters. J Membr Biol **250:**573-585.
581   42.   **Chen C, Gardete S, Jansen RS, Shetty A, Dick T, Rhee KY, Dartois V.** 2018. Verapamil
582   targets membrane energetics in Mycobacterium tuberculosis. Antimicrobial agents and
583   chemotherapy **62**.
584   43.   **Lee R, Hurdle J, Liu J, Bruhn D, Matt T, Scherman M, Vaddady P, Zheng Z, Qi J,
585   Akbergenov R.** Rakesh. Waidyarachchi, SL**:**152-158.
586   44.   **Rubin H.** 2007. The respiratory chain of M. tuberculosis. The FASEB Journal **21:**A207-A207.
587   45.   **Bollenbach T.** 2015. Antimicrobial interactions: mechanisms and implications for drug discovery
588   and resistance evolution. Curr Opin Microbiol **27:**1-9.
589   46.   **Nichols RJ, Sen S, Choo YJ, Beltrao P, Zietek M, Chaba R, Lee S, Kazmierczak KM, Lee
590   KJ, Wong A, Shales M, Lovett S, Winkler ME, Krogan NJ, Typas A, Gross CA.** 2011.
591   Phenotypic landscape of a bacterial cell. Cell **144:**143-156.
592   47.   **Kigondu EM, Wasuna A, Warner DF, Chibale K.** 2014. Pharmacologically active metabolites,
593   combination screening and target identification-driven drug repositioning in antituberculosis drug
594   discovery. Bioorg Med Chem **22:**4453-4461.
595   48.   **DeJesus MA, Gerrick ER, Xu W, Park SW, Long JE, Boutte CC, Rubin EJ, Schnappinger
596   D, Ehrt S, Fortune SM, Sassetti CM, Ioerger TR.** 2017. Comprehensive Essentiality Analysis
597   of the Mycobacterium tuberculosis Genome via Saturating Transposon
598   Mutagenesis. mBio **8:**e02133-02116.
599   49.   **Gao YG, Selmer M, Dunham CM, Weixlbaumer A, Kelley AC, Ramakrishnan V.** 2009. The
600   structure of the ribosome with elongation factor G trapped in the posttranslocational state.
601   Science **326:**694-699.
602   50.   **Ramakrishnan G, Chandra NR, Srinivasan N.** 2015. Recognizing drug targets using
603   evolutionary information: implications for repurposing FDA-approved drugs against
604   Mycobacterium tuberculosis H37Rv. Mol Biosyst **11:**3316-3331.
605   51.   **Besier S, Ludwig A, Brade V, Wichelhaus TA.** 2003. Molecular analysis of fusidic acid
606   resistance in Staphylococcus aureus. Mol Microbiol **47:**463-469.
607   52.   **Takaki K, Davis JM, Winglee K, Ramakrishnan L.** 2013. Evaluation of the pathogenesis and
608   treatment of Mycobacterium marinum infection in zebrafish. Nature Protocols **8:**1114.
609   53.   **Ling LL, Schneider T, Peoples AJ, Spoering AL, Engels I, Conlon BP, Mueller A, Schäberle
610   TF, Hughes DE, Epstein S.** 2015. A new antibiotic kills pathogens without detectable resistance.
611   Nature **517:**455-459.
612   54.   **Wright GD.** 2016. Antibiotic Adjuvants: Rescuing Antibiotics from Resistance: (Trends in
613   Microbiology 24, 862-871; October 17, 2016). Trends Microbiol **24:**928.

55. **Yang H, Chen G, Hu L, Liu Y, Cheng J, Ye Y, Li J.** 2016. Enhanced efficacy of imipenem-
colistin combination therapy against multiple-drug-resistant Enterobacter cloacae: in vitro activity
and a Galleria mellonella model. J Microbiol Immunol Infect doi:10.1016/j.jmii.2016.01.003.
56. **Ankomah P, Johnson PJ, Levin BR.** 2013. The pharmaco -, population and evolutionary
dynamics of multi-drug therapy: experiments with S. aureus and E. coli and computer simulations.
PLoS Pathog **9:**e1003300.
57. **Rodrigues L, Cravo P, Viveiros M.** 2020. Efflux pump inhibitors as a promising adjunct therapy
against drug resistant tuberculosis: a new strategy to revisit mycobacterial targets and repurpose
old drugs. Expert Review of Anti-infective Therapy**:**1-17.
58. **Hartkoorn RC, Uplekar S, Cole ST.** 2014. Cross-resistance between clofazimine and
bedaquiline through upregulation of MmpL5 in Mycobacterium tuberculosis. Antimicrobial
agents and chemotherapy **58:**2979-2981.
59. **Lomovskaya O, Lee A, Hoshino K, Ishida H, Mistry A, Warren MS, Boyer E, Chamberland
S, Lee VJ.** 1999. Use of a genetic approach to evaluate the consequences of inhibition of efflux
pumps in Pseudomonas aeruginosa. Antimicrob Agents Chemother **43:**1340-1346.
60. **Markham PN, Neyfakh AA.** 1996. Inhibition of the multidrug transporter NorA prevents
emergence of norfloxacin resistance in Staphylococcus aureus. Antimicrob Agents Chemother
40:2673-2674.
61. **Markham PN.** 1999. Inhibition of the emergence of ciprofloxacin resistance in Streptococcus
pneumoniae by the multidrug efflux inhibitor reserpine. Antimicrob Agents Chemother **43:**988-
989.
62. **Cokol-Cakmak M, Bakan F, Cetiner S, Cokol M.** 2018. Diagonal Method to Measure Synergy
Among Any Number of Drugs. J Vis Exp doi:10.3791/57713.
63. **Cokol M, Weinstein ZB, Yilancioglu K, Tasan M, Doak A, Cansever D, Mutlu B, Li S,
Rodriguez-Esteban R, Akhmedov M, Guvenek A, Cokol M, Cetiner S, Giaever G, Iossifov I,
Nislow C, Shoichet B, Roth FP.** 2014. Large-scale identification and analysis of suppressive
drug interactions. Chem Biol **21:**541-551.
64. **Tekin E, Beppler C, White C, Mao Z, Savage VM, Yeh PJ.** 2016. Enhanced identification of
synergistic and antagonistic emergent interactions among three or more drugs. J R Soc Interface
13.
65. **Bhusal Y, Shiohira CM, Yamane N.** 2005. Determination of in vitro synergy when three
antimicrobial agents are combined against Mycobacterium tuberculosis. Int J Antimicrob Agents
26:292-297.
66. **Torella JP, Chait R, Kishony R.** 2010. Optimal drug synergy in antimicrobial treatments. PLoS
Comput Biol **6:**e1000796.
67. **Sarathy JP, Zuccotto F, Hsinpin H, Sandberg L, Via LE, Marriner GA, Masquelin T,
Wyatt P, Ray P, Dartois V.** 2016. Prediction of Drug Penetration in Tuberculosis Lesions. ACS
Infect Dis **2:**552-563.
68. **Zhang L, Yan K, Zhang Y, Huang R, Bian J, Zheng C, Sun H, Chen Z, Sun N, An R, Min F,
Zhao W, Zhuo Y, You J, Song Y, Yu Z, Liu Z, Yang K, Gao H, Dai H, Zhang X, Wang J,
Fu C, Pei G, Liu J, Zhang S, Goodfellow M, Jiang Y, Kuai J, Zhou G, Chen X.** 2007. High-
throughput synergy screening identifies microbial metabolites as combination agents for the
treatment of fungal infections. Proc Natl Acad Sci U S A **104:**4606-4611.

Figure 1. Inhibition of Rv1258c-mediated efflux of SPT by addition of CPZ. Combinations of
CPZ and SPT were applied in checkerboard assays against (*a*) wild-type *M. tuberculosis* H37Rv,
(*b*) the Δ*Rv1258c* deletion mutant, (*c*) the Δ*Rv1258c* pCRS4 complemented mutant, and (*d*) the
SPT[R] strain. Bacterial growth inhibition was assessed in two independent experiments by
fluorescence-based resazurin assay. The dashed horizontal line indicates 90% inhibition.

Figure 2. *In vitro* interaction between SPT and FA. Combinations of SPT and FA were applied in checkerboard assays against (*a*) wild-type *M. tuberculosis* H37Rv, (*b*) the SPT[R] mutant and (*c*) the FA[R] mutant. Bacterial viability was assessed by fluorescence-based resazurin assay. Dashed horizontal lines indicate 90% inhibition and data are the means and standard deviations of two independent biological replicates.

Figure 3. Activity against pre-MDR *M. tuberculosis* strains. *In vitro* activity of RIF-INH against
RIF[R] *M. tuberculosis rpoB*[S531L] mutant in the (*a*) absence or (*b*) presence of 1/2× MIC SPT, and
against the INH[R] *M. tuberculosis inhA* mutant in the (*c*) absence or (*d*) presence of 1/2× MIC SPT.
Bacterial viability was assessed in two independent experiments by fluorescence-based resazurin
assay. The dashed horizontal line indicates 90% inhibition.

Table 1 Investigation of potential synergies between SPT and CPZ against different *M.*
tuberculosis strains through the calculation of the FIC and sum of the FIC.

M. tuberculosis strain	Drug combination	MIC (mg/L)		ΣFIC*
		Alone	In Combination	
H37Rv	SPT	62	1.9	0.09
	CPZ	22.2	1.4	
Δ*Rv1258c*	SPT	3.9	1.0	0.76
	CPZ	22.2	11.1	
Δ*Rv1258c* pCRS4	SPT	31.0	1.9	0.12
	CPZ	22.2	1.4	
SPT^R	SPT	>248	15.5	0.56
	CPZ	22.2	11.1	

22   *The FIC of each drug was calculated as: (MIC of drug in combination) / (MIC of individual drug). The ΣFIC is
23   the sum of the FICs of the two drugs where ΣFIC ≤0.5 is synergistic; ≥4.0 is antagonistic; and any value in the
24   range $0.5<x<4.0$ is considered additive or no interaction (1).

25   **Table 2** Investigation of potential synergies between SPT and FA against different *M.*

26   *tuberculosis* strains through the calculation of the FIC and sum of the FIC

M. tuberculosis strain	Drug combination	MIC (mg/L)		ΣFIC
		Alone	**In Combination**	
H37Rv	SPT	62	16.3	0.5
	FA	0.63	0.16	
SPT^R	SPT	≥62	≥62	1.5
	FA	0.63	0.32	
FA^R	SPT	62	31.5	1.5
	FA	≥25	≥25	

27   *The FIC of each drug was calculated as: (MIC of drug in combination) / (MIC of individual drug). The ΣFIC is

28   the sum of the FICs of the two drugs where ΣFIC ≤0.5 is synergistic; ≥4.0 is antagonistic; and any value in the

29   range 0.5<*x*<4.0 is considered additive or no interaction (36).

Table 2 Summary of *in vitro* drug activities against *M. tuberculosis* strains

Drugs/Combination			
Strain	**CPZ**	**SPT**	**CPZ-SPT**
H37Rv	✔	✘	✅
Rv1258c	✔	✔	⊗
Rv1258c::WT	✔	✘	✅
SPT[R]	✔	✘	✅

Drugs/Combination			
Strain	**FA**	**SPT**	**FA-SPT**
WT H37Rv	✔	✘	✅
FA[R]	✘	✘	⊗
SPT[R]	✔	✘	✅

Legend:

✔	$MIC_{90} \leq 10$ mg/L
✘	$MIC_{90} > 10$ mg/L
✅	Synergy
⊗	No interaction