Females don't always sing in response to male song, but when they do, they sing to males with higher pitched songs

Alexander N. G. Kirschel[1,2,3], Zacharo Zanti[1], Zachary T. Harlow[3], Edgar E. Vallejo[4†], and Martin L. Cody[3], and Charles E. Taylor[3].

†Deceased 19th June 2019

[1]*Department of Biological Sciences, University of Cyprus, P.O. Box 20537, Nicosia 1678, Cyprus*

[2]*Edward Grey Institute, Department of Zoology, University of Oxford, South Parks Road, Oxford OX1 3PS, United Kingdom*

[3]*Department of Ecology and Evolutionary Biology, University of California, 621 Charles E. Young Dr. South, Los Angeles, CA 90095–1606*

[4]*Department of Information Technologies and Computer Science, ITESM-Campus Estado de México, Estado de México, Mexico*

Email of corresponding author: kirschel@ucy.ac.cy

Author contributions: ANGK, EEV, MLC and CET conceived of the study and ANGK designed the research; ANGK and ZTH collected field data, assisted by EEV; ANGK and ZZ analysed data; and ANGK wrote the manuscript, with contributions from MLC and CET.

Abstract

The long-held view that bird song is exclusively a male trait has been challenged

recently by a number of studies and reviews highlighting the prevalence of female

song. In spite of that, there remains a lack of knowledge on the function of female

song, with most evidence thus far focusing on females performing duets with males in

courtship displays, for joint territory defence, or for mate guarding purposes. Here we

show in a tracheophone suboscine passerine *Formicarius moniliger*, a sexually

monomorphic species in which both sexes sing, that females may participate in both

intrasexual and intersexual territory defence. Females sing more in response to

females than to males yet demonstrate an unexpected pattern of singing back to

playback of males singing higher frequency song than themselves. Unlike males, who

respond indiscriminately to playback of any song performed by either sex, females

appear to discern not only the sex, but also the size of the presumed intruder,

perceiving a strong negative relationship between body mass and frequency. Our

findings suggest females will only engage in territory defence with males when they

expect those males to be weaker than they are. While our results are consistent with

expectations of a shared ancestral function of song in territory defence, they also

suggest females may suffer greater costs in engaging in territorial disputes and thus

limit their vocal contribution according to the perceived threat.

Keywords: Animal communication, female song, intersexual interactions, sex

roles, intrasexual territoriality, Neotropical birds.

Introduction

Sexual dimorphism is manifested in animals in a variety of ways. Male elephant seals

can be 10 times the size of females (Mccann et al., 1989), and sport highly modified

proboscises; male deer grow antlers; male peacocks develop iridescent plumage and a

long train with eyespots (Yorzinski et al., 2013); all this in aid of attracting females

and fighting off rival males (Lande, 1980). Such dimorphism represents conventional

sex roles of males competing for mates and females investing in parental care (Kokko

and Jennions, 2008).

Yet many species are sexually monomorphic. No differences in body size; no

differences in colour or pattern. Genetic correlations in body size between the sexes

may constrain sexual dimorphism (Merilä et al., 1998). Also, strong stabilizing

selection acting on the sexes (Price and Grant, 1985), largely driven by them

occupying similar niches, might explain their lack of dimorphism (Székely et al.,

2007). Many such animals use acoustic communication for mate attraction and

territory defence. Insects, frogs and birds create a cacophony of sounds in natural

habitats as a result (Kirschel et al., 2009a), but except in some cases of sex-role

reversal (Goymann et al., 2004), song has typically been considered a sexually

divergent behaviour performed by males (Odom et al., 2014).

There are, however, some animals in which both sexes sing. For instance in

tropical birds, many females sing as well as males (Slater and Mann, 2004).

Females in these instances may at times sing loosely in association with the male or

at others sing a tightly-coordinated duet (Slater and Mann, 2004), the latter a

phenomenon which has received much attention in the literature (Langmore, 1998,

Slater and Mann, 2004, Hall, 2004). Duetting has been attributed to territory defence,

whereby the female sings to defend the territory jointly with her mate from a rival of

either sex (Fedy and Stutchbury, 2005, Hall, 2004), or she may sing to defend the

territory specifically from same sex rivals (Langmore, 1998, Cain and Langmore,

2015). Alternatively, duets could function in mate guarding, whereby either the male

or the female ensures their partner is not singing solo in an attempt to attract extra-

pair mating opportunities (Levin, 1996a, Levin, 1996b, Seddon and Tobias, 2006,

Tobias and Seddon, 2009).

The prevalence of female song in the tropics has been attributed to sex role

convergence in locations where birds are resident and defend territories year-round

(Slater and Mann, 2004), but this view has been challenged by studies suggesting

female song is ancestral in songbirds (Odom et al., 2014, Riebel et al., 2019). So

while songs may represent ornaments attractive to the opposite sex, their primary

function in females is thought to be in intrasexual competition in species defending

territories year-round from same-sex rivals (Tobias et al., 2011, Tobias et al., 2012b).

A further study across all birds identified year-round territoriality as the trait most

closely associated with duetting (Tobias et al., 2016). Such year-round territoriality is

particularly evident in insectivorous birds (Hau et al., 2000, Slater and Mann, 2004,

Tobias et al., 2011), with much previous work on duetting birds focusing on

insectivorous wrens and antbirds (Fedy and Stutchbury, 2005, Levin, 1985, Seddon

and Tobias, 2006). Female song is also common in many Australian passerines,

including in the insectivorous fairy-wrens, whistlers, shrike-thrushes, and bell birds,

which like the wrens and antbirds, have extended longevity when compared to most

temperate passerines, and defend territories year-round. In these taxa, the sexes are

either sexually dimorphic or dichromatic (e.g., antbirds, (Kirschel et al., 2019, Tobias

and Seddon, 2009), fairy-wrens and whistlers (Hall and Peters, 2008, van Dongen and

Mulder, 2008), sing distinctly different songs (e.g., wrens (Mennill and Vehrencamp,

2008), or both (again antbirds (Tobias and Seddon, 2009), and fairy-wrens (Hall and

Peters, 2008), suggesting songs may still serve different functions among the sexes.

Might there be situations in species where the sexes are monomorphic and sing

indistinguishable songs, and what might such cases reveal regarding sex roles?

Perhaps sexually monomorphic vocalisations could play a role in intersexual

competition (Tobias et al., 2012a). Here we examine such a case. We studied a

population of sexually monomorphic Mexican antthrush – a.k.a. Mayan antthrush

(Krabbe and Schulenberg 2019) - (*Formicarius moniliger*), for which previous studies

have shown both sexes sing (Blumstein et al., 2011, Kirschel et al., 2009b). Despite

evidence showing that individuals can reliably be recognized from the subtle spectral

and temporal idiosyncrasies in their songs, efforts thus far have not been able to

distinguish male from female song (Kirschel et al., 2011b).

109   Mexican antthrush is a tracheophone suboscine passerine (Tobias et al.,

2012a), a group whose simple stereotypic songs have allowed investigators to identify

songs to species (Trifa et al., 2008), and to track movements of individuals in space

and time using an acoustic sensor network (Collier et al., 2010). Previous work on

Mexican antthrush has shown how songs can reliably be assigned to individuals

(Kirschel et al., 2009b), and using such song classifications, resulting territory maps

have demonstrated little territory overlap between same sex rivals suggesting strong

intrasexual territoriality (Kirschel et al., 2011b).

117   We investigated how male and female Mexican antthrush perceive song

produced by possible territory intruders of the same or opposite sex using playback

experiments. We hypothesized that a vocal response to same sex playback represented

territory defence, though could also represent mate guarding. Conversely, a vocal

response to opposite sex playback could represent intersexual territory defence or

joint territory defence, if the mate also responded vocally, but potentially promiscuous

intentions if not. Our overall aim was to determine when each sex used songs in

response to possible intruders, thus informing us of the function of female song,

which remains understudied across birds (Odom and Benedict, 2018, Riebel et al.,

2019). Moreover, because of much evidence of a negative correlation between body

size and song frequency in birds (Ryan and Brenowitz, 1985a, Wallschlager, 1980),

we also tested whether such a relationship exists within antthrushes. If there is such a

relationship, might individuals perceive differences in body size? If larger birds sang

lower frequency song, responses might differ based on the size of the presumed

intruder. We tested this hypothesis by comparing responses to differences in

frequency between responder's song and playback stimulus, as well as differences in

body mass between responder and presumed intruder.

Materials and methods

Fieldwork

Fieldwork took place at the Estación Chajul in the Montes Azules Biosphere Reserve,

in south-eastern Chiapas, Mexico (16˚6′44′′N; 90˚56′27′′W), during 8-17 June 2007,

7-18 December 2008, 7-28 May 2009, and 11-31 May 2012. Montes Azules, also

known as Selva Lacandona, represents the largest expanse of pure tropical rainforest

in North America (ParksWatch, 2003). The study was focused within a 50-ha study

plot at an elevation range of 150-165 m, on the northern side of the Lacantún River

(Kirschel et al., 2011b).

We captured birds using target netting techniques and marked captured birds

with a unique combination of color rings to aid visual confirmation of their identity.

We deployed individual mist nets (Avinet or Ecotone, 30- or 36-mm mesh, 12 x 2.5

m) along 1 - 2 m-wide trails, with the bottom of each net set at ground level. We

target netted antthrushes using conspecific playback. Captured birds were also

weighed and measured and 50-100μL of blood was obtained via venepuncture of the

brachial vein for genetic sex identification. Individuals caught in 2012 also had a

radio-frequency identification (RFID) tag fitted on their backs with a harness as part

of a parallel study, and methods used are described therein (Kirschel et al.

unpublished ms). We recorded ringed birds singing along trails in the study area as

previously described and in accordance with methods used previously (Kirschel et al.,

2011b, Kirschel et al., 2009c). Specifically, we used a Marantz PMD 670 / 661

recorder with Sennheiser microphones ME-67/K6, MKH20 microphone with a

Telinga parabolic reflector, or MKH8050 housed in a Rycote windshield, as well as

an acoustic sensor network (see (Collier et al., 2010, Kirschel et al., 2011b).

Playback experiments

Fifty playback stimuli were prepared using Raven Pro 1.4 (Cornell Lab of

Ornithology) using songs recorded during previous field seasons (Kirschel et al.,

2011a), and additional recordings obtained during the 2012 field effort. The songs

used belonged to 24 ringed individuals and had been recorded while color-ring

combinations were confirmed, or were classified unambiguously as belonging to

specific individuals (Kirschel et al., 2011b, Kirschel et al., 2009c). Sex of individuals

whose songs were used in experiments had been previously identified genetically

(Blumstein et al., 2011, Kirschel et al., 2009b), or had been predicted based on

behaviour typical of one of the sexes during the 2012 field season and subsequently

confirmed genetically (see below). Each stimulus was prepared with songs that were

high-pass filtered at 400 Hz to remove background noise, maximum amplitude

normalized at 20,000 units (Raven's unit of amplitude measurement), and arranged

temporally with two songs of an individual within a one-minute stimulus with the

remainder silence and then looped continuously for the duration of the experiment.

Playback apparatus consisted of an Apple iPod mp3 player and a TivoliAudio PAL

loudspeaker, with output volume set at the same level for all experiments. Playbacks

were performed at different times of day and stimuli were randomly selected.

Experiments varied in length and depended on an approach by focal individuals.

Antthrushes are inconspicuous, terrestrial birds (Cody, 2000), which may approach

gradually and silently on the ground, possibly from far within their territory. In such

cases it could take them 10 - 20 minutes to approach to within a distance of the

loudspeaker where a recording of suitable quality might be obtained. Recordings were

thus obtained opportunistically when birds started vocalizing in response to the

playback and only those experiments where recordings were obtained are included in

analyses. Our aim was to document who sang and what sex they were, in response to

which individual playback and the sex, song characters, and body mass of the

individual whose song was used in the experiment. A parallel systematic study

performed in 2012 focused on approaches to specific stimuli (Kirschel et al.

unpublished ms). In field seasons prior to 2012, on some occasions we may not have

ringed both individuals of a focal pair. In such cases we still use the data of the

experiment from the one identified individual but do not consider any partner's

response in analyses.

Genetic sex identification

DNA was extracted from blood or feather samples collected from 21 individuals in

the field in 2012 using a QIAGEN DNEasy Blood and Tissue Kit (QIAGEN,

Valencia, CA) following manufacturer's protocols. Highly conserved primers (2550F

and 2718R, (Fridolfsson and Ellegren, 1999)) were used in polymerase chain reaction

(PCR) to amplify the differently sized introns of Z- and W- linked chromohelicase-

DNA binding protein 1 (CHD1) genes. PCR products were separated in a 1 or 2%

agorose gel run in TAE buffer, revealing one or two bands for males and females

respectively.

Song feature extraction

Eighty-three recordings from fieldwork in 2012 were imported into RAVEN Pro 1.4

(Cornell Lab of Ornithology, Ithaca, NY, USA) where song sequences were cut into

separate WAV files, which were then processed in a feature extraction program in

MATLAB 7 (MathWorks 2005). The procedure is described in detail in (Kirschel et

al., 2009c). Here, we follow the approach used in (Kirschel et al., 2011b), extracting

19 temporal and spectral features of song (See supplement).

Canonical discriminant analysis

The majority of songs used for this study had been identified to individual in previous

studies (Kirschel et al., 2011b, Kirschel et al., 2009c) or by sight or with radio

telemetry in 2012. However, there remained 39 recordings of antthrush songs for

which we did not know the singer. We followed (Kirschel et al., 2011b) in using

linear canonical discriminant analysis (CDA) to identify those songs to individuals.

The CDA was performed in STATA 11.2 (StataCorp 2009) on songs cut from

recordings obtained in May 2012 using the 19 variables extracted from the songs.

Following (Kirschel et al., 2011b), we used 1189 songs from recordings where the

singer could be unambiguously confirmed by sight or radio telemetry as training data

and predicted the singer of 619 songs on the remaining 39 recordings (see ESM for

details of post CDA classification filtering). The resultant songs classified to

individual were then pooled with the 2011 songs from 168 recordings previously

classified (Kirschel et al., 2011b, Kirschel et al., 2009c), and statistical analyses were

performed on a total of 3022 songs attributed to 24 individuals. Moreover, we tested

whether with the combined dataset of 50 individuals we could distinguish between

male and female song in a CDA with leave-one-out-classification using the mean

values of the 19 song features from the 230 recordings included in the study.

Statistical analyses

We first tested whether one of the sexes sings more than the other in response to

playback of either sex using a chi-squared test. We restricted this analysis to 158

experiments (78 male and 80 female song playbacks) in territories where both

members of the pair were ringed. To test for what determines when each sex sings, we

used a Bayesian Generalized Linear Mixed Model (GLMM) in 'blme' in R, fitted

with the binomial distribution and logit link function using the function bglmer. We

determined the effect on the binary response variable "female song response" or

"male song response" of the difference in peak frequency between the mean peak

frequency of the responder's song (from all recordings of the individual obtained that

year) and the peak frequency of the playback stimulus. Also included as fixed factors

in the model were the binary predictor of sex of the individual whose song was used

in the playback experiment, and its interaction with difference in peak frequency.

Subject of experiment and playback stimulus used were included as crossed random

effects. We then tested whether each sex responded to different frequencies of male or

female playback using Bayesian GLMM fitted with a Gaussian distribution. Because

the function blmer does not provide P values by default, these were computed using

the packages 'sjstats' and 'parameters' in R.

To determine whether there was a relationship between body mass and song

frequency we first calculated the mean peak frequency from all recordings of each

individual for each field trip. For this analysis, we calculated the mean peak frequency

only from recordings where the bird's ring combination was identified, or where the

bird was identified using radio telemetry (Kirschel et al. unpublished ms). We then

ran a GLMM in lme4 in R, fitted with a Gaussian distribution, with mean peak

frequency as the dependent variable and body mass (g) and sex as fixed effects and

individual included as a random effect, to account for individuals recorded (and body

mass measured) over several years. We then tested if the relative size influenced

responses based on sex of the presumed intruder. Specifically, we ran Bayesian

GLMMs to test whether the interaction between difference in body mass (centred and

standardized, (Schielzeth, 2010) between subject and playback singer and the latter's

sex affected male or female response levels, with individual subject and individual

whose song playback was used as crossed random effects.

Results

Genetic sex identification

Of the 21 birds we obtained samples from in 2012, 13 were male and 8 were female.

This information was combined with the data from previous genetic analyses

(Kirschel et al., 2011b), from which 12 sexed males and 11 females, were included as

playback stimuli singers (13 male, 11 female) and/or test subjects (21 male, 17

female) in experiments.

Canonical discriminant analysis

Based on the CDA trained using songs of 20 individuals, 619 songs from 39

recordings were classified. Of these, 88.7% were assigned to the predicted individual,

consistent with classification rates found previously for similar numbers of

individuals in a season (Kirschel et al., 2011b). By contrast, the CDA could only

classify 75.1% of male song and 55.9% of female song to the correct sex (See ESM

for detailed results of the CDA).

Playback experiments

Males responded with song significantly more than females did to both male (94.9%

vs 16.7% of experiments; $\chi^2 = 21.08$, $P < 0.0001$) and female playback (89.9% vs

26.6% of experiments; $\chi^2 = 24.59$, $P < 0.0001$). Males showed no difference in song

response levels based on sex of playback, and difference in peak frequency between

the subject and playback stimulus singer or their interactions (Table 1, Fig. 1). By

contrast, females sang more in response to female playback than male playback, but

not in relation to differences in peak frequency between subject and playback

stimulus singer (Table 1). There was, however, a significant negative interaction

between sex of playback and difference in peak frequency, indicating that females

increasingly respond to male song the lower their song is in frequency compared to

the playback stimulus (Fig. 1). Further, while there was no difference in the peak

frequency of female playback that males and females responded, females responded

to significantly higher frequency male playback than males did (Table S2).

Figure 1

Females (a) sing back in response to higher frequency male song than their own, and to males that are smaller than they are, but do not discriminate between frequencies or body mass in response to female song. Males (b) sing back in response to the vast majority of playback experiments (large circles) and do not discriminate between sexes or according to body size. Blue and green circles – vocal response to song playback of smaller and larger males respectively. Red and orange circles – vocal response to song playback of smaller and larger females respectively. Cyan and lime green dots – no vocal response to song playback of smaller and larger males respectively. Maroon and gold dots - no vocal response to song playback of smaller and larger females respectively. Dotted line represents y = x.

306 **Table 1:** Bayesian generalized linear mixed models with binomial distribution of the

307 effect of sex of singer on playback experiment, difference in peak frequency of songs

308 on the playback and responder's song, and the interaction of those effects, on vocal

309 responses of females (a) and males (b).

a) Female responses	Estimate	SE	Z	P
Intercept	-0.81	0.43	-1.89	0.058
Sex	-2.17	0.81	-2.67	**0.0076**
Peak frequency difference (standardized)	0.34	0.33	1.01	0.314
Sex/peak frequency difference interaction	-0.02	0.008	-3.12	**0.0018**
b) Male responses				
Intercept	3.95	1.45	2.73	**0.0063**
Sex	1.51	1.12	1.35	0.177
Peak frequency difference (standardized)	-0.004	0.008	-0.49	0.622
Sex-peak frequency difference interaction	0.02	0.015	1.49	0.137

While there was no difference in peak frequency according to sex (GLMM: t = 0.719,

P = 0.48) there was a significant negative relationship between body mass and peak

frequency (t = -3.29, P = 0.002), meaning larger individuals sang lower frequency

songs (Fig. 2). Yet, neither females nor males responded significantly differently to

song of each sex in relation to the body mass of the singer (Table S3). However, in

females specifically, there was a pattern of greater response to male song the lower

his body mass was in relation to the female's, but it was not significant.

Figure 2

There is a significant negative relationship between body mass and peak frequency,

with larger individuals singing lower frequency song, though no differences between

males (blue dots) and females (red dots). *Formicarius moniliger* illustration courtesy

of Del Hoyo et al. (2019).

Discussion

Female birds are widely recognized as being more discriminating than males in

response to playback stimuli (Searcy and Brenowitz, 1988, Seddon and Tobias, 2010).

But males are also more typically involved in territory defence and the cost of

responses to abnormal species-specific song or to heterospecific song would be lower

than in females (Searcy and Brenowitz, 1988). In this study, antthrush females

responded to playback less often than males did, supporting the view that males

devote more effort to territory defence than females do. Indeed, males appeared to

respond indiscriminately to song, whether performed by males or females and

irrespective of the body size of the presumed intruder. Such responses suggest either

heightened aggression towards intruders of both sex in males, promiscuous intentions

when a female trespasses into his territory, or an inability to distinguish song to sex.

By contrast, females responded significantly more often to female song than to male

song, suggesting they do discriminate between the songs of each sex, in spite of the

extent of overlap in song characteristics (Fig. 2), and our failure to reliably classify

songs to sex in a discriminant analysis. Their responses suggest they do identify song

to sex, consistent with the hypothesis that signals that may appear sexually

monomorphic to human observers are still likely to contain information on sex-

specific differences (Price, 2015). Females singing in response to female song is

consistent with both the function of intrasexual territory defence and mate guarding

(Langmore, 1998, Levin, 1996b, Tobias and Seddon, 2009, Cain and Langmore,

2015). Yet, on occasion, females responded to male song. They responded

significantly more to male song the lower the frequency of their song compared to the

male song. In other words, they respond to those males they perceived as singing

higher pitched songs than they did.

Frequency has been shown to be negatively correlated with body size in birds,

both among species (Wallschlager 1980, (Ryan and Brenowitz, 1985b), and within

species (Hall et al., 2013). Here, we tested whether there was a relationship between

frequency and body mass in individuals of Mexican antthrush. We found that larger,

heavier antthrushes do indeed sing lower frequency songs than lighter antthrushes,

while song frequency does not differ between the sexes. What this relationship

suggests is that females responding to males singing at higher frequencies are

responding to smaller males. Our test of that specific relationship found a pattern in

that direction, but it was not significant. Of course, there is variation in song

frequency that is not explained entirely by body mass (Fitch, 1999), and the birds

responded to what they heard and not what they saw. Other aspects of the playback

stimulus might provide further information to the receiver regarding the size of the

singer and the threat (or opportunity) represented. Indeed, the one female whose song

was higher in frequency than the playback in Fig. 1 was actually heavier than the

male whose song she responded to, and that male on average sang at higher

frequencies than that female, just not on the selected playback stimulus.

Not only did females respond to higher frequency male song then theirs, they

also sang back in response to higher frequency male song than males did, providing

further support that females responded to the smallest males. Might females be

attracted to males that are smaller than them? Or are they simply attracted to higher

frequency song? We suggest an alternative explanation. These paired females are not

responding to male song for mutual attraction purposes. Instead, we believe that

females will participate in intersexual territory defence by singing back only when

they perceive the intruding male to be smaller, and thus weaker, than they are.

Sexually monomorphic songs are expected to fulfil a similar function between the

sexes (Riebel et al., 2019), and thus could be used for both intrasexual and intersexual

territory defence. But in a species where the female sings a fraction of what the male

does, it seems she chooses carefully when she will use her song for the purpose of

intersexual territory defence.

We did not test whether female song was influenced by the male mate's

response and thus formed a coordinated territorial response to intruders (e.g., (Hall

and Peters, 2008)– males responded to almost every experiment rendering such a test

unworkable. In any case, such behaviour is more typical of duetting birds that

coordinate their songs into duets, which antthrushes do not do. Neither do female

antthrushes sing in tandem with the male or jam his song as a mate guarding strategy

(Tobias and Seddon, 2009). Instead, females sometimes sing solo (see also (Kirschel

et al., 2011b). We also caution that our results are based on a large number of

recordings of individuals identified based on song classifiers, and that we excluded

songs on recordings that we were unable to classify to specific individuals. What this

means is that there were individuals singing on a number of the recordings that we

could not classify as belonging to a known ringed individual. It does not mean that

those were not female or male partners of individuals we did identify singing on

recordings. Just that our CDA failed to identify them. Nevertheless, any situations

where we were unable to identify the individual singer reduced our sample size and

overall statistical power. We have no reason to assume any missing data would not be

representative of the patterns reported here. We also caveat that nonvocal responses

may play an important part in both territorial defence and mate guarding, so it would

be important to associate nonvocal responses such as approaches to playback stimuli

as well as vocal responses.

Female song has typically been thought of as a component of coordinated

duets in tropical birds (Fedy and Stutchbury, 2005). We have shown here that female

antthrushes sing independently of males and that their songs cannot reliably be

distinguished from male song. Rather than female song representing evidence for

convergence in sex roles, a notion that has been challenged by recent studies (Riebel

et al., 2019), our study on suboscine passerines is consistent with the premise that

female song is ancestral in songbirds (Odom et al., 2014), with females singing to

defend territories much in the way that males do. Nevertheless, with costs of

responding to the wrong stimulus potentially greater in females (Searcy and

Brenowitz, 1988), we find that females respond far less than males do and pick and

choose when they will respond with song to intruders, especially when they are males.

Acknowledgments

We thank T. Asef, A. Darrah, S. Michaelides, L. Powell, and the staff at the Estacion

Chajul, for assistance in the field and lab, with analyses, and with logistics. This work

was supported by a National Science Foundation Award 0410438 to C.E.T., by the

Consejo Nacional de Ciencia y Tecnologia (CONACYT) under Award Number

REF:J110.389⁄2006 and by Wolfram Research under Award Number L322 to E.E.V.

and a University of Oxford John Fell Fund grant to A.N.G.K. Research was

conducted under UCLA's Animal Research Committee protocols (ARC No. 2004-

094-12), and a SEMARNAT research permit (E.E.V).

References

BLUMSTEIN, D. T., MENNILL, D. J., CLEMINS, P., GIROD, L., YAO, K., PATRICELLI, G., DEPPE, J. L., KRAKAUER, A. H., CLARK, C., CORTOPASSI, K. A., HANSER, S. F., MCCOWAN, B., ALI, A. M. & KIRSCHEL, A. N. G. 2011. Acoustic monitoring in terrestrial environments using microphone arrays: applications, technological considerations and prospectus. *Journal of Applied Ecology,* 48**,** 758-767.

CAIN, K. E. & LANGMORE, N. E. 2015. Female and male song rates across breeding stage: testing for sexual and nonsexual functions of female song. *Animal Behaviour,* 109**,** 65-71.

CODY, M. L. 2000. Antbird guilds in the lowland Caribbean rainforest of southeast Nicaragua. *Condor,* 102**,** 784-794.

COLLIER, T. C., KIRSCHEL, A. N. G. & TAYLOR, C. E. 2010. Acoustic localization of antbirds in a Mexican rainforest using a wireless sensor network. *Journal of the Acoustical Society of America,* 128**,** 182-189.

FEDY, B. C. & STUTCHBURY, B. J. M. 2005. Territory defence in tropical birds: are females as aggressive as males? *Behavioral Ecology and Sociobiology,* 58**,** 414-422.

FITCH, W. T. 1999. Acoustic exaggeration of size in birds via tracheal elongation: comparative and theoretical analyses. *Journal of Zoology,* 248**,** 31-48.

FRIDOLFSSON, A. K. & ELLEGREN, H. 1999. A simple and universal method for molecular sexing of non-ratite birds. *Journal of Avian Biology,* 30**,** 116-121.

GOYMANN, W., WITTENZELLNER, A. & WINGFIELD, J. C. 2004. Competing females and caring males. Polyandry and sex-role reversal in African black coucals, Centropus grillii. *Ethology,* 110**,** 807-823.

HALL, M. L. 2004. A review of hypotheses for the functions of avian duetting. *Behavioral Ecology and Sociobiology,* 55**,** 415-430.

HALL, M. L., KINGMA, S. A. & PETERS, A. 2013. Male Songbird Indicates Body Size
with Low-Pitched Advertising Songs. *PLoS ONE,* 8.
HALL, M. L. & PETERS, A. 2008. Coordination between the sexes for territorial
defence in a duetting fairy-wren. *Animal Behaviour,* 76**,** 65-73.
HAU, M., WIKELSKI, M., SOMA, K. K. & WINGFIELD, J. C. 2000. Testosterone and
year-round territorial aggression in a tropical bird. *General and*
Comparative Endocrinology, 117**,** 20-33.
KIRSCHEL, A. N. G., BLUMSTEIN, D. T., COHEN, R. E., BUERMANN, W., SMITH, T. B.
& SLABBEKOORN, H. 2009a. Birdsong tuned to the environment: green
hylia song varies with elevation, tree cover, and noise. *Behavioral Ecology,*
20**,** 1089-1095.
KIRSCHEL, A. N. G., BLUMSTEIN, D. T. & SMITH, T. B. 2009b. Character
displacement of song and morphology in African tinkerbirds. *Proceedings*
of the National Academy of Sciences of the United States of America, 106**,**
8256-8261.
KIRSCHEL, A. N. G., CODY, M. L., HARLOW, Z. T., PROMPONAS, V. J., VALLEJO, E. E.
& TAYLOR, C. E. 2011a. Territorial dynamics of Mexican Ant-thrushes
Formicarius moniliger revealed by individual recognition of their songs.
IBIS 153**,** 255-268.
KIRSCHEL, A. N. G., CODY, M. L., HARLOW, Z. T., PROMPONAS, V. J., VALLEJO, E. E.
& TAYLOR, C. E. 2011b. Territorial dynamics of Mexican Ant-thrushes
Formicarius moniliger revealed by individual recognition of their songs.
Ibis, 153**,** 255-268.
KIRSCHEL, A. N. G., EARL, D. A., YAO, Y., ESCOBAR, I. A., VILCHES, E., VALLEJO, E.
E. & TAYLOR, C. E. 2009c. Using Songs to Identify Individual Mexican
Antthrush Formicarius Moniliger: Comparison of Four Classification
Methods. *Bioacoustics-the International Journal of Animal Sound and Its*
Recording, 19**,** 1-20.
KIRSCHEL, A. N. G., SEDDON, N. & TOBIAS, J. A. 2019. Range-wide spatial
mapping reveals convergent character displacement of bird song.
Proceedings of the Royal Society B-Biological Sciences, 286.
KOKKO, H. & JENNIONS, M. D. 2008. Parental investment, sexual selection and
sex ratios. *Journal of Evolutionary Biology,* 21**,** 919-948.
LANDE, R. 1980. Sexual Dimorphism, Sexual Selection, and Adaptation in
Polygenic Characters. *Evolution,* 34**,** 292-305.
LANGMORE, N. E. 1998. Functions of duet and solo songs of female birds. *Trends*
in Ecology & Evolution, 13**,** 136-140.
LEVIN, R. N. 1985. The Function of Vocal Duetting in the Bay Wren (Thryothorus-
Nigricapillus). *American Zoologist,* 25**,** A3-A3.
LEVIN, R. N. 1996a. Song behaviour and reproductive strategies in a duetting
wren,Thryothorus nigricapillus: I. Removal experiments. *Animal*
Behaviour, 52**,** 1093-1106.
LEVIN, R. N. 1996b. Song behaviour and reproductive strategies in a duetting
wren,Thryothorus nigricapillus: II. Playback experiments. *Animal*
Behaviour, 52**,** 1107-1117.
MCCANN, T. S., FEDAK, M. A. & HARWOOD, J. 1989. Parental Investment in
Southern Elephant Seals, Mirounga-Leonina. *Behavioral Ecology and*
Sociobiology, 25**,** 81-87.

MENNILL, D. J. & VEHRENCAMP, S. L. 2008. Context-Dependent Functions of
Avian Duets Revealed by Microphone-Array Recordings and Multispeaker
Playback. *Current Biology,* 18**,** 1314-1319.
MERILÄ, J., SHELDON, B. C. & ELLEGREN, H. 1998. Quantitative Genetics of
Sexual Size Dimorphism in the Collared Flycatcher,Ficedula Albicollis.
Evolution, 52**,** 870-876.
ODOM, K. J. & BENEDICT, L. 2018. A call to document female bird songs:
Applications for diverse fields. *The Auk,* 135**,** 314-325.
ODOM, K. J., HALL, M. L., RIEBEL, K., OMLAND, K. E. & LANGMORE, N. E. 2014.
Female song is widespread and ancestral in songbirds. *Nature
Communications,* 5.
PARKSWATCH. 2003. *Park Profile – Mexico, Montes Azules Biosphere Reserve*
[Online]. www.parkswatch.org Available:
http://www.parkswatch.org/parkprofiles/pdf/mabr_eng.pdf [Accessed
2004].
PRICE, J. J. 2015. Rethinking our assumptions about the evolution of bird song
and other sexually dimorphic signals. *Frontiers in Ecology and Evolution,*
3.
PRICE, T. D. & GRANT, P. R. 1985. The Evolution of Ontogeny in Darwin Finches -
a Quantitative Genetic Approach. *American Naturalist,* 125**,** 169-188.
RIEBEL, K., ODOM, K. J., LANGMORE, N. E. & HALL, M. L. 2019. New insights from
female bird song: towards an integrated approach to studying male and
female communication roles. *Biology Letters,* 15.
RYAN, M. J. & BRENOWITZ, E. A. 1985a. The Role of Body Size, Phylogeny, and
Ambient Noise in the Evolution of Bird Song. *American Naturalist,* 126**,**
87-100.
RYAN, M. J. & BRENOWITZ, E. A. 1985b. The Role of Body Size, Phylogeny, and
Ambient Noise in the Evolution of Bird Song. *The American Naturalist,*
126**,** 87-100.
SCHIELZETH, H. 2010. Simple means to improve the interpretability of
regression coefficients. *Methods in Ecology and Evolution,* 1**,** 103-113.
SEARCY, W. A. & BRENOWITZ, E. A. 1988. Sexual differences in species
recognition of avian song. *Nature,* 332**,** 152-154.
SEDDON, N. & TOBIAS, J. A. 2006. Duets defend mates in a suboscine passerine,
the warbling antbird (Hypocnemis cantator). *Behavioral Ecology,* 17**,** 73-
83.
SEDDON, N. & TOBIAS, J. A. 2010. Character displacement from the receiver's
perspective: species and mate recognition despite convergent signals in
suboscine birds. *Proceedings of the Royal Society B-Biological Sciences,*
277**,** 2475-2483.
SLATER, P. J. B. & MANN, N. I. 2004. Why do the females of many bird species
sing in the tropics? *Journal of Avian Biology,* 35**,** 289-294.
SZÉKELY, T., LISLEVAND, T. & FIGUEROLA, J. 2007. Sexual size dimorphism in
birds. *In:* FAIRBAIRN, D. J., BLANCKENHORN, W. U. & SZÉKELY, T. (eds.)
Sex, size, and gender roles : evolutionary studies of sexual size dimorphism.
Oxford: Oxford University Press.
TOBIAS, J. A., BRAWN, J. D., BRUMFIELD, R. T., DERRYBERRY, E. P., KIRSCHEL, A.
N. G. & SEDDON, N. 2012a. The importance of neotropical suboscine birds

as study systems in ecology and evolution. *Ornitologia Neotropical,* 23**,**
259-272.
TOBIAS, J. A., GAMARRA-TOLEDO, V., GARCIA-OLAECHEA, D., PULGARIN, P. C. &
SEDDON, N. 2011. Year-round resource defence and the evolution of male
and female song in suboscine birds: social armaments are mutual
ornaments. *Journal of Evolutionary Biology,* 24**,** 2118-2138.
TOBIAS, J. A., MONTGOMERIE, R. & LYON, B. E. 2012b. The evolution of female
ornaments and weaponry: social selection, sexual selection and ecological
competition. *Philosophical Transactions of the Royal Society B-Biological*
Sciences, 367**,** 2274-2293.
TOBIAS, J. A. & SEDDON, N. 2009. Signal Jamming Mediates Sexual Conflict in a
Duetting Bird. *Current Biology,* 19**,** 577-582.
TOBIAS, J. A., SHEARD, C., SEDDON, N., MEADE, A., COTTON, A. J. & NAKAGAWA,
S. 2016. Territoriality, Social Bonds, and the Evolution of Communal
Signaling in Birds. *Frontiers in Ecology and Evolution,* 4.
TRIFA, V. M., KIRSCHEL, A. N. G., TAYLOR, C. E. & VALLEJO, E. E. 2008. Automated
species recognition of antbirds in a Mexican rainforest using hidden
Markov models. *Journal of the Acoustical Society of America,* 123**,** 2424-
2431.
VAN DONGEN, W. F. D. & MULDER, R. A. 2008. Male and female golden whistlers
respond differently to static and dynamic signals of male intruders.
Behavioral Ecology, 19**,** 1025-1033.
WALLSCHLAGER, D. 1980. Correlation of Song Frequency and Body-Weight in
Passerine Birds. *Experientia,* 36**,** 412-412.
YORZINSKI, J. L., PATRICELLI, G. L., BABCOCK, J. S., PEARSON, J. M. & PLATT, M. L.
2013. Through their eyes: selective attention in peahens during courtship.
Journal of Experimental Biology, 216**,** 3035-3046.
576

577