The effects of arbuscular mycorrhizal fungi (AMF) and *Rhizophagus irregularis*

on soil microorganisms assessed by metatranscriptomics and metaproteomics

Lammel, D.R. [1,2], Meierhofer, D. [3], Johnston, P. [1,4], Mbedi, S. [2,5], Rillig, M.C. [1,2]

[1] Freie Universität Berlin, Institut für Biologie, D-14195 Berlin, Germany

[2] Berlin-Brandenburg Institute of Advanced Biodiversity Research (BBIB), D-14195 Berlin, Ge

[3] Max Planck Institute for Molecular Genetics, D-14195 Berlin, Germany

[4] Berlin Center for Genomics in Biodiversity Research (BeGenDiv), D-14195 Berlin, Germany

[5] Museum für Naturkunde, D-10115 Berlin, Germany

Running title: **AMF select and activate soil microorganisms in the hyphosphere**

Abstract

Arbuscular mycorrhizal fungi (AMF) form symbioses with approximately 80% of plant

species and potentially benefit their hosts (e.g. nutrient acquisition) and the soil

environment (e.g. soil aggregation). AMF also affect soil microbiota and soil

multifunctionality. We manipulated AMF presence (via inoculation of non-sterile soil

with *Rhizophagus irregularis* and using a hyphal compartment design) and used

RNA-seq and metaproteomics to assess AMF roles in soil. The results indicated that

AMF drove an active soil microbial community expressing transcripts and proteins

related to nine metabolic functions, including the metabolism of C and N. We suggest

two possible mechanisms: 1) the AMF hyphae produce exudates that select a

beneficial community, or, 2) the hyphae compete with other soil microbes for

available nutrients and consequently induce the community to mineralize nutrients

from soil organic matter. We also identified candidate proteins that are potentially

related to soil aggregation, such as Lpt and HSP60. Our results bridge microbial

ecology and ecosystem functioning. We show that the AMF hyphosphere contains an

active community related to soil respiration and nutrient cycling, thus potentially

improving nutrient mineralization from soil organic matter and nutrient supply to the

plants.

1. Introduction

Recent studies indicated that plant roots select beneficial microbial

communities related to nutrient cycling in the rhizosphere (Mendes et al., 2014,

2018b; Baltrus, 2017; Yan et al., 2017; Schmidt et al., 2019). Therefore, we

hypothesized that arbuscular mycorrhizal fungi (AMF) also select beneficial microbial

communities in their hyphosphere (the volume of soil influenced by AMF hyphae).

Previous studies have found that AMF affect the soil microbial community

structure at the taxonomic level (Turrini et al., 2018; Xu et al., 2018; Bhatt et al.,

2019). However, even though such diversity studies based on ribosomal DNA

regions (e.g. 16/18S rRNA or ITS) are robust methods to assess microbial

community structure, they usually fail to correlate to environmental processes (Hall et

al., 2018; Lammel et al., 2018; Zhang et al., 2019). More accurate information related

to ecosystem functioning, interactions and nutrient cycling can be obtained by

assessing directly the metabolically active microbial community by RNA-seq and

proteomics (Mendes et al., 2018a; Yao et al., 2018). Thus, learning about the

metabolically active microbial community can be particularly important for a better

understanding of arbuscular mycorrhizal fungi (AMF) interactions with the soil

microbiota.

AMF are soil fungi that form a symbiosis with approximately 80% of plant

species. During the symbiosis, plant roots supply these fungi with carbohydrates and

fatty acids, while the fungi induce metabolic changes in the plant (including improved

plant growth and better plant response to biotic and abiotic stress) (Kameoka et al.;

Delavaux et al., 2017; Powell and Rillig, 2018; Begum et al., 2019; Mateus et al.,

2019). In addition to intimately associating with the plant root, the fungi also grow in

the soil, foraging for mineral nutrients and water, and affecting soil properties

(Kameoka et al.; Rillig and Mummey, 2006; Pepe et al., 2017). Consequently, AMF

have three main roles in soil, they: 1) taking up mineral nutrients and water, therefore

changing soil nutrient and water dynamics; 2) interacting with and changing soil

microbial communities, and 3) interacting with and modifying soil structure, e.g. by

aggregating soil (Rillig and Mummey, 2006; Cavagnaro et al., 2015; Powell and

Rillig, 2018; Bhatt et al., 2019).

Thus, it is not only relevant to understand how AMF change the dynamics of

nutrients in the soil but also how AMF induce changes in the soil microbial

community itself (Turrini et al., 2018; Bhatt et al., 2019). The inoculation of AMF,

remarkably *R. irregularis,* affects soil microbial community structure at the taxonomic

level (Changey et al., 2019). Moreover, some bacterial taxa are likely associated with

AMF spores, for example, *Proteobacteria* and *Firmicutes* (Battini et al., 2016;

Changey et al., 2019). However, detailed information about the effects of AMF on the

functionally active microbial community is still unavailable.

AMF also affect soil structure and their presence and abundance in soil is

typically correlated with soil aggregation (Rillig and Mummey, 2006; Schlüter et al.,

2019, 2019). On the other hand, soil aggregation affects microbial communities in

soil (Rillig et al., 2017; Upton et al., 2019). Soil aggregation is likely influenced by

direct effects of the AMF, such as hyphal enmeshment of soil particles and

production of exo-polymeric substances (EPS), and indirect effects, such as AMF

effects on microbial communities and the soil food web that could then change the

soil structure and aggregates (Rillig and Mummey, 2006). EPS can also be produced

by the microbial community, and include mainly polysaccharides and

lipopolysaccharides, which are synthesized by proteins such as Wza, Wca, Kps, Alg,

and Sac, and have been reported to be potentially correlated to soil aggregation

(Cania et al., 2019).

Another possible mechanism for soil aggregation could be that specific AMF

proteins may act as binding agents for soil particles. In the 1990's, a monoclonal

antibody was produced (MAb32B11) for the purpose of quantifying *R. irregularis* in

soil and the protein attached to that antibody was called "glomalin" (name derived

from *Glomus*) and it correlated with soil aggregation (Wright et al., 1996). Later,

general "glomalin related soil protein" ("GRSP") extractions methods were proposed,

but such methods were not specific and very likely biased by substances other than

glomalin (Rosier et al., 2006; Gillespie et al., 2011). Thus, the initial hypotheses

linking specific AMF proteins and soil aggregation are not confirmed. Gadkar and

Rillig (2006) provided initial evidence that the protein that attaches to the *R.*

irregularis 90's antibody (that was first called glomalin) may be a HSP60 protein.

HSP60 proteins (GroEL) are mainly molecular chaperons, but they can also have

moonlight functions, and are widespread proteins present in all living organisms

(Henderson et al., 2013). It is still unclear whether this protein can directly interact

with soil particles, but a priority research question is to quantify the abundance of

HSP60 from *R. irregularis* in soil compared to the natural occurrence of HSP60 from

the overall soil microbial community.

The objective of this study was to use RNA-seq and proteomics to investigate

the hypothesis that the AMF hyphosphere selects for a beneficial microbial

community. We mainly investigated the AMF effects on: 1) the C, N and P soil

community metabolism at the gene/protein level; 2) the soil microbial community

structure itself (assessed by the taxonomic affiliation of the genes/proteins); and 3)

transcripts and proteins potentially related to soil aggregation (such as the above-

mentioned proteins Wza, Wca, Kps, Alg, Sac and HSP60).

2. Material and Methods

Compartmentalized experimental setup with hyphal compartment

To test the effects of AMF on soil, we used the hyphal compartment approach

(Johnson et al., 2001; Leifheit et al., 2014). We established two main treatments,

non-sterile soil control and the same soil inoculated with *R. irregularis* (DAOM

197198) with ten replicates (ten pots) for each treatment. In all pots we placed the

hyphal compartment cores, and then the control soils were further divided into two

treatments: static cores and cores that were rotated three times per week. The

rotation of the cores severs fungal hyphae thus avoiding the growth of AMF inside

the rotated cores (Leifheit et al., 2014). So the experiment was finally divided into

three operational treatments: control with static cores (C), control with rotated cores

(rC) and static cores in soil inoculated with the AMF *R. irregularis* (A). Due to

technical limitations of advanced molecular techniques (e.g. sequencing costs), we

randomly chose four samples of each treatment for RNA-seq and five samples of

each treatment for proteomics. We report all the non-molecular analysis results for all

the treatments and replicates in Supplementary Material 1.

The soil used in the experiment was collected at the agricultural experimental

station of the Humboldt University in Dahlem (Berlin, Germany). The top soil from a

meadow (0-30 cm) was taken and air-dried at room temperature. The soil was an

Albic Luvisol (World Reference Base for soil resources, 1998) and had the following

physical-chemical characteristics: 73.6% sand, 18.8% silt, 7.6% clay; pH 7.1 ($CaCl_2$)

(analyses conducted by LUFA Rostock Agricultural Analysis and Research Institute,

Germany); 1.87% C (total) and a C/N ratio 15.6 (analyzed on an Euro EA C/N

analyzer, HEKAtech GmbH, Wegberg, Germany) (Rillig et al., 2010). The soil was

sieved to 4mm and root and plant material removed. Twenty plastic pots (20cm

diameter x 20cm high) were filled with five liters of this soil and core compartments

inserted in each pot (details about the cores in Leifheit et al., 2014).

Seeds of clover (*Trifolium repens* L) were surface sterilized with 70% ethanol

for 30 secs, followed by 3% sodium hypochlorite for 5 min, and then rinsed five times

with autoclaved water. Ten of these seeds were sown evenly in each pot, but after

germination only five plants were kept per pot. Half of the pots (ten) received the

main treatment and were inoculated with *R. irregularis* and the other half, ten pots,

were kept as control without inoculation. The inoculum was prepared from *R.*

irregularis that was grown under *in vitro* culture conditions (Rillig et al., 2010). A piece

of solid media containing hyphae and spores ("a loopful") from the hyphae

compartment (containing 20 ± 7 spores) was inoculated nearby each seed in the

inoculated treatments and a "loopful" of sterile media was added nearby each seed

for the controls. The plants were grown in a greenhouse (22°C/ 18°C, day/ night) and

a photoperiod of 14h (a mix of natural light and supplementation of artificial light

when needed). An automatic irrigation system watered each pot with deionized water

once a day. It was programmed for 50ml/day for young plants and achieving 100

ml/day for larger plants. We fertilized the pots on days 15, 45 and 75 with 100 ml

Hoagland solution without P and N. On the same days we added 2 ml of full

Hoagland solution into the core compartments aiming to stimulate hyphal foraging

and growth into the cores.

Experiment harvest

After 105 days, the experiment was harvested. Pots were harvested one at a

time, to avoid delays in freezing the samples for molecular analysis. First, the shoots

of the plants were cut and placed in paper bags for later drying and biomass

determination. Then, 5 mm of the surface soil of the hyphal compartment cores were

removed (to avoid contamination from the surface) and the soil of each core was

transferred to a sterile 50ml conical tube and gently homogenized for 30s. From

these samples, two aliquots of 2ml soil were transferred to two sterile centrifuge

tubes and immediately frozen in liquid N_2 (to be later used in RNA extraction) and

5ml of soil transferred to a 5ml sterile centrifuge tube and frozen (to be later used for

protein extraction). All the spatulas and spoons were treated with 70% ethanol and

flamed between samples, and the plastic materials were previously autoclaved. The

samples were stored at -80°C. The remaining soil was air-dried for 48h at 40°C and

used for the soil aggregation and hyphal length analyses.

Plant and soil analyses

Plant shoots were dried in an oven for 48h at 60°C and then weighed. We

analyzed soil aggregation by measuring the newly formed aggregates and water

stable aggregates (WSA), and we also analyzed AMF and non-AMF hyphal length in

the soil.

The newly formed aggregates were measured by weighing the dry soil

particles >2mm and dividing this value by the total weight of the soil sample. For the

WSA measurement we used a wet sieving apparatus (Kemper and Rosenau, 1986).

The method was slightly modified as described by Leifheit et al. (2014): 4.0 g of dried

soil were rewetted by capillary action and sieved for 3 min using a 250 µm sieve. The

179 coarse matter was separated by crushing the aggregates that remained on the sieve

180 and the small particles washed out through the sieve. Coarse matter and soil were

181 dried at 60°C for 48 h and WSA calculated and corrected for coarse matter.

182  Hyphal length was determined by extraction from 4.0 g of soil, staining with

183 Trypan blue and followed by microscopy (Jakobsen et al., 1992; Rillig et al. 1999).

184 Dark to light blue stained aseptate hyphae with characteristic unilateral angular

185 projections ("elbows and coils") were considered mycorrhizal (Mosse, 1959), whereas

186 non-blue stained or blue stained hyphae with regular septation or straight growth

187 were considered non-mycorrhizal (Leifheit et al., 2014). We also counted the number

188 of AMF spores on the slides (Supp. Material 1).

190 *RNA extraction and sequencing*

191  RNA was extracted from 2g of the frozen soil using the RNeasy PowerSoil

192 Total RNA Kit (Qiagen, Germany). Since an important goal of RNA-seq was to

193 produce a broad and annotated library to allow protein identification, we adopted two

194 library strategies for mRNA enrichment. First, we targeted eukaryotic mRNA

195 enrichment in the samples and used 60% of the volume of each RNA extract for poly-

196 A enrichment using the kit NEXTflex Poly(A) Beads (Perkin Elmer, USA). Secondly,

197 we targeted the transcripts from the full soil diversity using rRNA depletion by

198 hybridization. For that, the four replicates of each treatment were combined in pairs,

199 using 20% of the volume of the RNA extracts from each sample, yielding two

200 samples for each treatment (we pooled samples to increase sequencing depth). The

201 samples were then treated with the kit RiboMinus Bacteria following the

202 manufacturer's instructions (ThermoFischer, USA).

The 18 mRNA samples, 12 samples poly-A enriched and six samples rRNA

depleted, were prepared for sequencing using NextFlex Rapid Directional RNA

SeqKit (Perkin Elmer, USA). All the samples were evaluated in an Agilent

TapeStation and quantified by QuBit (Invitrogen, USA). The samples of each library

were pooled in equimolar ratios for each library strategy. The libraries were then

pooled together using a six-fold higher concentration of the rRNA depletion library

compared to the poly-A library (since we expected higher diversity for the first). The

libraries were sent for sequencing to a commercial company using the Illumina HiSeq

X Ten platform. The sequences were deposited at the GenBank server

(https://www.ncbi.nlm.nih.gov/genbank - Bioproject number PRJNAxxxxx).

Soil RNA-seq analysis

The sequences from the HiSeq run were quality checked and demultiplexed

using the bcl2fastq software (Illumina, USA). All the sequences were then processed

together using the software Trinity and assembled and cleaned of rRNA (Grabherr et

al., 2011). The assemblage was annotated by Blast2go (Götz et al., 2008). The

sequences were also further classified according to the SEED system from MG-

RAST (Wilke et al., 2016). Lastly, the reads of each sample were mapped to the

sequences assembled by the software Trinity or to the genome of *R. irregularis* DAO

JGI v2 using the software Salmon (Grabherr et al., 2011; Morin et al., 2019). The

annotation table and the counting table of the mapped reads were further exported

for analysis in R.

Metaproteomics

Protein extraction was performed based on five grams of frozen soil for all

three conditions in biological quintuplicates by the addition of extraction buffer

containing 100 mM Tris-HCl pH 8.0, 5% w/v sodium dodecyl sulphate, and 10 mM

dithiothreitol. Samples were vortexed and boiled for 10 min at 100°C, centrifuged at

4,000 x g for 10 min and the supernatants were transferred into new protein low-

binding tubes. Proteins were then precipitated by the addition of chilled trichloroacetic

acid to a final concentration of 20% and kept at -80°C overnight. Proteins were

pelleted at 4,000 x g for 30 min, washed twice with cold acetone (-20°C), and air-

dried, as described in (Qian and Hettich, 2017).

Protein extracts were reduced and alkylated under denaturing conditions by

the addition of 200 µL of a buffer containing 3 M guanidinhydrochlorid, 10 mM tris(2-

carboxyethyl)phosphine (TCEP), 40 mM chloroacetamide (CAA), and 100 mM Tris

pH 8.5 to prevent proteolytic activities, briefly vortexed and boiled for 10 min at 95°C

at 1000 rpm. Pellet disintegration was achieved by a sonicator with five cycles of

P150W, C60%, A100% (Hielscher, Teltow, Germany). After centrifugation at 15,000

rcf at 4°C for 10 min, supernatants were transferred into new protein low binding

tubes. The lysates were loaded into the preOmics in-stage tip kit cartridges (iST kit

96x, Martinsried, Germany) to remove humics, centrifuged at 3,800 rcf for 3 min,

washed with 8 M urea by pipetting up and down ten times and centrifuged again. The

lysates were digested and purified according to the preOmics in-stage tip kit

(preOmics). In brief, 50 µl lysis buffer was added, diluted with resuspension buffer

and digested after 10 min at 37°C at 500 rpm overnight. After several washing steps,

peptides were eluted sequentially in three fractions using the SDB-RPS-1 and -2

buffers (Kulak et al., 2014) and the elution buffer provided by preOmics. Eluates were

first dried in a SpeedVac, then dissolved in 5% acetonitrile and 2% formic acid in

water, briefly vortexed, and sonicated in a water bath for 30 seconds for subsequent

analysis on a nanoLC-MS/MS.

LC-MS instrument settings for shotgun proteome profiling and data analysis

LC–MS/MS was carried out by nanoflow reverse-phase liquid chromatography

(Dionex Ultimate 3000, Thermo Scientific, Waltham, MA) coupled online to a Q-

Exactive HF Orbitrap mass spectrometer (Thermo Scientific). The LC separation was

performed using a PicoFrit analytical column (75 µm ID × 50 cm long, 15 µm Tip ID

(New Objectives, Woburn, MA) in-house packed with 3-µm C18 resin (Reprosil-AQ

Pur, Dr. Maisch, Ammerbuch-Entringen, Germany). Peptides were eluted using a

gradient from 3.8 to 50 % solvent B in solvent A over 121 min at 266 nL per minute

flow rate. Solvent A was 0.1 % formic acid and solvent B was 79.9 % acetonitrile, 20

% water, 0.1 % formic acid. An electrospray was generated by applying 3.5 kV. A

cycle of one full Fourier transformation scan mass spectrum (300–1750 m/z,

resolution of 60,000 at m/z 200, AGC target $1e^6$) was followed by 12 data-dependent

MS/MS scans (resolution of 30,000, AGC target $5e^5$) with a normalized collision

energy of 25 eV. In order to avoid repeated sequencing of the same peptides a

dynamic exclusion window of 30 sec. was used. In addition, only the peptide charge

states between two to eight were sequenced.

Label-Free Proteomics Data Analysis

Raw MS data were processed with MaxQuant software (v1.6.0.1) and

searched against the *Rhizophagus irregularis* UP000236242_747089 database,

released in May/2019, or the annotated RNA-seq assemblage. A false discovery rate

(FDR) of 0.01 for proteins and peptides, a minimum peptide length of 7 amino acids,

a precursor mass tolerance to 20 ppm for the first search and 4.5 ppm for the main

search were required. A maximum of two missed cleavages was allowed for the

tryptic digest. Cysteine carbamidomethylation was set as fixed modification, while N-

terminal acetylation and methionine oxidation were set as variable modifications.

The tables generated by MaxQuant were further processed using the online

server ANPELA ("analysis and performance assessment of label-free

metaproteomes"), data were log transformed and processed by the online pipeline,

including missing data imputation by the K-nearest Neighbor algorithm (KNN) and

mean normalization (MEA) (Tang et al., 2019).

Statistical analysis

Analyses were performed in R version 3.5. Plant biomass, water stable

aggregates and hyphal length were analyzed by ANOVA, followed by the Tukey

posthoc test. We also correlated variables using Spearman's rank correlation

coefficient (*p*). The distribution of the data and effect sizes were analyzed by the R-

DBR package (SI Figure S1).

The tables previously obtained from the RNA-seq and proteomics were

aggregated with the SEED functions for a first explanatory analysis. Differences

between the samples were analyzed by cluster analysis. The taxonomy obtained by

the Blast2go annotation was used in the diversity analysis and RDA using R-Vegan.

The data were further analyzed by R packages DeSeq2/Alx2 and the online server

GMine (http://cgenome.net/gmine) (Quinn et al., 2018). Since both RNA-seq data

sets were analyzed according to the pool of identified sequences, we treated them as

compositional data, considering the centered log-ratio of the values of each

gene/protein for each sample (Gloor et al., 2017). We report the log-ratio fold change

of the gene/proteins and Welch's t-test with Benjamini-Hochberg corrected statistical

probabilities, calculated by the R-packages DeSeq2 and AIx2 or the online server

Gmine.

3. Results

We evaluated the effects of *R. irregularis* inoculation compared to control soil.

Plants were harvested at 105 days, but no difference was observed in the shoot plant

biomass (Figure 1 A-B and SI S1).

We further evaluated the soil compartments that were divided in three

treatments with five replicates each: 1) control with static cores (C), 2) control with

rotated cores (rC) and 3) cores of soil inoculated with the AMF *R. irregularis* (A).

First, we detected that the C and A cores were colonized by AMF, while very few

AMF hyphae were observed in the rC (Figure 1.C). We also detected colonization by

non-AMF hyphae in all the cores, but there was no difference (P=0.356) among

treatments (Figure 1 D). Secondly, we detected increases in soil aggregation. At the

beginning of the experiment, all soil was sieved to 2 mm and we observed at the

harvest time 57 ± 4 % new formed aggregates, however no strong differences were

observed across treatments (P=0.121). We also observed increases in WSA during

the experiment, from 46 % ± 3 at the beginning, increased to 56-61 % after 105 days

(Figure 1.E.). At 105 days we found higher soil aggregation in the cores inoculated

with *R. irregularis* in relation to the rotated cores (P=0.036) (Figure 1.D-E). We also

detected a correlation between AMF hyphal length and WSA across the treatments

(p=0.67, P<0.01), as also between non-AMF hyphae and WSA (p=0.60, P<0.01).

Effects of AMF on soil community metabolism

The samples were further analyzed in terms of metatranscriptomics (Figures

2-4). The HiSeq sequencing run yielded 180 million reads (SI Figure S2). From that,

for the rRNA depletion libraries we recovered an average of 6 million reads for the C

treatment, 2.5 million for rC and 6.5 million for A. For the poly-A libraries we

recovered an average of 21.5 million reads for the C treatment, 30 million for rC and

25 million for A. For the Poly-A enrichment library, around of 50% of the transcripts

were still from bacteria (Figure 4). This indicates that Poly-A selection from soil is

challenging, since most of the transcripts in soils are generally from bacteria (Mendes

et al., 2018a; Schlüter et al., 2019). The method could be improved in future by, for

example, by repeating the poly-A selection for several times. Based on the Blast2go

annotation, from the poly-A library, we were able to recover around 0.47% of reads of

AMF (*Glomeromycota*) for C, 0.36% for rC and 0.46% for A (SI Figure S2). For the

rRNA depletion library we were able to recover 0.45% AMF reads for C, 0.32% reads

for rC and 0.42% for A.

The transcript analyses indicated several differences in gene expression

related to C, N, P and respiration processes clustered by the MG-RAST SEED terms

(Figure 2-3). From the 616,863 genes assembled by Trinity, 161,702 could be

annotated to the 29 SEED categories "level 1". We further segregated the

"GroEL/HSP60" category (SEED level 2) from the category "protein" in SEED level 1,

to better test our third hypothesis about HSP60. The first exploratory analysis, the

heat map and cluster analysis, indicated that the samples were distinctly clustered

between the rotated core and the other treatments with high abundance of AMF

hyphae (Figure 2.A). Some clustering was observed differentiating the static core

control and the *R. irregularis* treatment (Figure 2.A). Notice that the heat map scale is

not about negative or positive gene expression, but it is based on the scaled

centered log-ratio (CLR) transformation recommended for compositional data and

reflects the relative abundance of each metabolic category in each sample. From the

29 functional categories, 15 categories were associated by a network analysis to the

presence of AMF hyphae in soil (Figure 3). The analyses indicated that *R. irregularis*

inoculation had increased expression of genes related to RNA metabolism,

respiration and GroEL/HSP60. The control sample, that also had AMF hyphae, had

intermediate values between the inoculated and the rotated core treatments and was

related with C, N, P and S metabolism. The rotated control was related mostly with

cell division, mobility, fatty acids, and secondary metabolism (Figure 3). We found a

correlation between AMF hyphal length and transcripts related to respiration (r=0.66,

P=0.028), but no strong evidence for a correlation between hyphal length and total

GroEL/HSP60 transcripts (r=0.47, *P*=0.14).

For the proteomic data, we identified 1336 proteins using the RNA-seq as

reference, and the data normalized by ANPELA indicated a strong separation of the

samples by treatment (SI Figure S3), and a volcano plot identified 103 proteins that

were more abundant in C and A than in the rotated cores (Supp. Figure 3). From all

the proteins, 398 were identified at MG-RAST SEED categories (Figure 2.B), and,

again, there was a differentiation between rotate cores to the control and soil

inoculated with *R. irregularis*. Considering these 398 proteins, there was a decrease

in 19 SEED categories in the rotated cores (Figure 3.B) and thus the network

analyses were strongly influenced by the non-rotated treatments, that can be

observed by the color of the network nodes, indicating mainly the non-rotated

treatments (Figure 3.D). GroEL/HSP60 was more abundant in the soil cores

inoculated with *R. irregularis*. Using the *R. irregularis* genome as reference, we

identified 51 proteins and most of them were proteins related to protein metabolism,

including heat shock proteins (we report on this separately in the section about HSP).

Effects of AMF on soil community structure

The metatranscriptomic data were then analyzed taxonomically. From the

616,863 genes assembled by Trinity, 199,566 were annotated by Blast2go. All the

samples were dominated by Bacteria (Figure 4). The rRNA depletion library had

approx. 45% of Bacteria for the A and C treatments and approx. 65% in the rC

(Figure 4.A). For the poly-A libraries, it was also approx. 45% of Bacteria for the A

and C treatments, but approx. 50% of *Bacteria* in the rC. The microbial community

structure was further investigated by RDA (Figures 4. C1 and D1). For both, *Bacteria*

and *Eukaryota*, the samples were better clustered by the libraries made from the

mRNA enrichment by rRNA depletion libraries ("m" code in the figures). For the

depletion libraries, the rotated cores had a distinct community structure when

compared to the static-control soil and the soil inoculated with *R. irregularis*

(perMANOVA, *P*<0.001 for both, *Bacteria* and *Eukaryota*). However, clustering of the

samples from the poly-A libraries was not strongly supported (perMANOVA, *P*<0.12

for *Bacteria* and P<0.20 for *Eukaryota*).

The samples were further analyzed in detail in terms of the composition of

Bacteria and *Eukaryota* sequences. The RNA-seq data indicated that all the samples

were dominated by *Actinobacteria* and *Proteobacteria* (no difference among

treatments was observed; *P*=0.2), but a reduction of *Firmicutes* (Aldex2, *P*=0.052)

was observed in the rotated cores (Figure 4.C2). Considering *Eukaryota*, the

samples were dominated by *Longamoebia*, *Protosteliales* and *Streptophyta* (Figure

4. D2). *Fungi* accounted for around 15% of the *Eukaryota* reads, and *Glomeromycota*

for around 0.4% of the reads (SI Figure S2).

Effects of AMF on transcripts/proteins potentially linked to soil aggregation

We screened for genes/proteins potentially related to soil aggregation. We

explored a list of genes/proteins including genes related to EPS metabolism (Cania

et al., 2019), such as polysaccharide export outer membrane (Wza), capsular

polysaccharide export system (Kps) and lipopolysaccharide export proteins (Lpt). We

did not find differences for most of the genes/proteins, but found in the

metatranscriptome that Lpt (Figure 5A) was more abundant in rotated core samples

(z-score 0.6-0.9), when compared to the control and *R. irregularis* inoculated

treatments ($P<0.001$).

We further evaluated HSP60/GroEL, which was very abundant across all

treatments, especially in the cores with higher abundance of AMF hyphae (Figure 2).

In all treatments, at the RNA level, HSP60/GroEL was predominantly of bacterial

origin, particularly from *Acidobacteria* and *Proteobacteria* (SI Figure S5 A1-2). Of the

few reads of eukaryotic origin, the great majority was from *Basidiomycota*, with the

exception of the rotated cores, which had a high abundance of reads from

Longamoebia (SI Figure S5 A3). In the proteome, the vast majority of identified

HSP60/GroEL proteins came from *Bacteria* (app. 99% across all treatments), and

they were predominantly from *Acidobacteria* and *Proteobacteria* (SI Figure S5 B).

We found a correlation between the total HSP60/GroEL with the AMF hyphal

length for the total metaproteome (r=0.57, *P*=0.026), but little evidence for the total

metatranscriptome (r=0.47, *P*=0.14). However, as described before, most of these

HSP60/GroEL came from *Bacteria* (SI Figure S5 A1). We observed no correlation

between total HSP60/GroEL and WSA (all had $P>0.2$).

We further looked at the proteins identified from *R. irregularis* in the

metaproteome and there were no differences in HSP60/GroEL ($P=0.21$) across the

treatments (Figure 6). We compared this result with tubulins (Alfa+Beta) ($P=0.08$),

which are essential proteins related to fungal growth, and observed a very similar MS

signal intensity to the HSP60/GroEL protein. We screened the other HSPs from *R.*

irregularis and detected also little evidence for differences across treatments for two

other HSP, the HSP70 (Uniprot A0A2H5SZG6, $P=0.14$) and HSP70 (Uniprot

U9SYW5, $P=0.07$). None of these proteins were correlated with AMF hyphal length

or WSA (all had $P>0.2$). These results also indicated that *R. irregularis* transcripts

and proteins were a small proportion of the full microbial community (we estimated

AMF being between 0.3-0.5% of the active community, SI Figure S1).

4. Discussion

In this study, we investigated the effects of AMF on soil microbial communities

using soil RNA-seq and metaproteomics. We detected that AMF drove the active

microbial community as well as transcripts and proteins related to metabolic

functions. First, we showed that the presence of AMF induced changes in transcripts

and proteins in soil related to several metabolic processes, including protein

metabolism (N cycle) and respiration (C cycle). Secondly, we showed that *R.*

irregularis induces changes in the soil microbial community, which was largely

dominated by *Bacteria*. And thirdly, we detected transcripts and proteins that could

be related to soil aggregation.

Previous knowledge about AMF affecting soil functions was obtained mostly

by inoculation studies, or by quantifying AMF hyphae in soils and correlating them

with specific functions, such as nitrogen uptake or soil aggregation (Powell and Rillig,

2018; Hestrin et al., 2019; Rillig et al., 2019). To the best of our knowledge, this was

the first attempt to experimentally assess metatranscriptomes and metaproteomes to

estimate the contribution of AMF to the metabolism of the entire soil community. We

find that only 0.3-0.5% of the transcripts/proteins were of AMF origin and our data

indicate that AMF induced significant shifts in soil community metabolism when

compared to the rotated core control (Figures 2 and 3).

Our results bridge between microbial ecology and ecosystems functioning.

The data suggest that the AMF hyphosphere drive an active community related to

soil respiration and nutrient cycling, potentially improving nutrient mineralization from

soil organic matter and nutrient supply to plants. Previous studies also presented

several lines of evidence that AMF influence the microbial community and that these

fungi are important players in the carbon cycle, P and N dynamics and soil

respiration (Nottingham et al., 2010; Zhang et al., 2016; Powell and Rillig, 2018;

Hestrin et al., 2019). Interestingly, previous studies have suggested that AMF are

associated with specific bacterial communities and that these communities have

complementary functional roles to the AMF (Battini et al., 2016; Turrini et al., 2018).

Our data corroborate these findings in a general way, since the community had

increased expression of genes/proteins related to general soil processes in the

presence of the AMF hyphae (Figures 2 and 3).

Thus, we present evidence that arbuscular mycorrhizal fungi (AMF) select for

beneficial microbial communities in their hyphosphere. We suggest two possible

mechanisms to be tested in future: 1) the AMF hyphae produce exudates that select

a beneficial microbial community, analogous to reports for plant roots (Baltrus, 2017);

or, 2) AMF hyphae compete with the microbial community for available nutrients and

consequently induce the community to mineralize nutrients from soil organic matter

(Hestrin et al., 2019).

Moreover, our study showed that the soil communities were dominated by

transcripts and proteins of bacterial origin (Figure 4). We observed that the samples

were dominated by *Proteobacteria* and had an increased relative abundance of

Firmicutes in the hyphal compartments, and both groups were previously reported to

be predominantly associated with AMF (Battini et al., 2016; Turrini et al., 2018).

Interestingly in the rotated cores there was an increased expression of bacterial *lpt*

genes (lipopolysaccharide export proteins). These genes can be related to soil

aggregation and we speculate that these could be related to soil aggregation in the

rotated cores, since soil aggregates increased from the beginning of the experiment

to the harvest (Cania et al., 2019).

Lastly, our data showed a high abundance and diversity of HSP60 (GroEL)

across the soil samples detected by RNA-seq and by proteomics. This can be

explained in two ways; first, this chaperon occurs across all living organisms and it

has essential functions in protein folding and cell metabolism, in addition to moonlight

functions (Henderson et al., 2013). Second, for the proteome samples, our method

included the boiling of samples and this can likely select for thermostable proteins,

such as HSPs (Rosier et al., 2006; Gillespie et al., 2011). Furthermore, HSP60

abundance may not only reflect cell abundance, but also the metabolic status of cells

(Henderson et al., 2013). Moreover, our data clearly showed the high complexity of a

soil community sample and that HSP60 is highly abundant and diverse in soil. We

also observed that at nucleotide level the HSP from *R. irregularis* are divergent

enough to be differentiated from the community sequences (Magurno et al., 2019),

but due codon redundancy they are highly conserved at the amino acid level. This

implies that during processing for proteomics the HSP60 could potentially be cleaved

in positions to overlap MS peptides spectra of HSP60/GroEL from other members of

the microbial community, and theoretically causing noise in the quantification of

HSP60 from AMF. Previous studies using the monoclonal antibody MAb32B11 for *R.*

irregularis already showed that HSP60 quantifications in soil are also difficult due to

the complex composition of organic material in soil and could make quantifications

imprecise (Rosier et al., 2006). Our new findings also bring additional evidence to

avoid quantifying GRSPs as a proxy to quantify AMF proteins in soil (Rosier et al.,

2006; Gillespie et al., 2011). Nevertheless, we are optimistic that future works using

high-resolution proteomics, perhaps in simpler soil communities, will be able to

identify specific proteins from AMF in soil and potentially correlating them with

ecosystems functions, such as soil aggregation.

5. Conclusions

We provide evidence that the AMF hyphosphere drives an active community

related to soil respiration and nutrient cycling, potentially improving nutrient

mineralization from soil organic matter and nutrient supply to plants. Overall, our

results show that AMF have an intense effect on the soil microbial community and on

the metabolic pathways related to soil functions and nutrient dynamics. These

findings contribute to a better understanding of the role of AMF in soil and how they

drive soil metabolism and microbial community structure.

6. Acknowledgments

We acknowledge Alexander-von-Humboldt Foundation and CAPES Foundation for a postdoctoral grant awarded to DRL, and Freie Universität Berlin and the Max Planck Society for funding this work.

7. References

Baltrus, D. A. (2017). Adaptation, specialization, and coevolution within phytobiomes. *Current Opinion in Plant Biology* 38, 109–116. doi:10.1016/j.pbi.2017.04.023.

Battini, F., Cristani, C., Giovannetti, M., and Agnolucci, M. (2016). Multifunctionality and diversity of culturable bacterial communities strictly associated with spores of the plant beneficial symbiont Rhizophagus intraradices. *Microbiological Research* 183, 68–79. doi:10.1016/j.micres.2015.11.012.

Begum, N., Qin, C., Ahanger, M. A., Raza, S., Khan, M. I., Ashraf, M., et al. (2019). Role of Arbuscular Mycorrhizal Fungi in Plant Growth Regulation: Implications in Abiotic Stress Tolerance. *Front. Plant Sci.* 10. doi:10.3389/fpls.2019.01068.

Bhatt, P., Joshi, D., Kumar, N., and Kumar, N. (2019). "Recent Trends to Study the Functional Analysis of Mycorrhizosphere," in *Mycorrhizosphere and Pedogenesis*, eds. A. Varma and D. K. Choudhary (Singapore: Springer Singapore), 181–190. doi:10.1007/978-981-13-6480-8_11.

Cania, B., Vestergaard, G., Kublik, S., Köhne, J. M., Fischer, T., Albert, A., et al. (2019). Biological Soil Crusts from Different Soil Substrates Harbor Distinct Bacterial Groups with the Potential to Produce Exopolysaccharides and Lipopolysaccharides. *Microb Ecol*. doi:10.1007/s00248-019-01415-6.

Cavagnaro, T. R., Bender, S. F., Asghari, H. R., and Heijden, M. G. A. van der (2015). The role of arbuscular mycorrhizas in reducing soil nutrient loss. *Trends in Plant Science* 20, 283–290. doi:10.1016/j.tplants.2015.03.004.

Changey, F., Meglouli, H., Fontaine, J., Magnin-Robert, M., Tisserant, B., Lerch, T. Z., et al. (2019). Initial microbial status modulates mycorrhizal inoculation effect on rhizosphere microbial communities. *Mycorrhiza*. doi:10.1007/s00572-019-00914-1.

Delavaux, C. S., Smith-Ramesh, L. M., and Kuebbing, S. E. (2017). Beyond nutrients: a meta-analysis of the diverse effects of arbuscular mycorrhizal fungi on plants and soils. *Ecology* 98, 2111–2119. doi:10.1002/ecy.1892.

Gadkar, V., and Rillig, M. C. (2006). The arbuscular mycorrhizal fungal protein glomalin is a
putative homolog of heat shock protein 60. *FEMS Microbiol Lett* 263, 93–101.
doi:10.1111/j.1574-6968.2006.00412.x.

Gillespie, A. W., Farrell, R. E., Walley, F. L., Ross, A. R. S., Leinweber, P., Eckhardt, K.-U.,
et al. (2011). Glomalin-related soil protein contains non-mycorrhizal-related heat-
stable proteins, lipids and humic materials. *Soil Biology and Biochemistry* 43, 766–
777. doi:10.1016/j.soilbio.2010.12.010.

Gloor, G. B., Macklaim, J. M., Pawlowsky-Glahn, V., and Egozcue, J. J. (2017). Microbiome
Datasets Are Compositional: And This Is Not Optional. *Front. Microbiol.* 8.
doi:10.3389/fmicb.2017.02224.

Götz, S., García-Gómez, J. M., Terol, J., Williams, T. D., Nagaraj, S. H., Nueda, M. J., et al.
(2008). High-throughput functional annotation and data mining with the Blast2GO
suite. *Nucleic Acids Res* 36, 3420–3435. doi:10.1093/nar/gkn176.

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., et al.
(2011). Trinity: reconstructing a full-length transcriptome without a genome from
RNA-Seq data. *Nat Biotechnol* 29, 644–652. doi:10.1038/nbt.1883.

Henderson, B., Fares, M. A., and Lund, P. A. (2013). Chaperonin 60: a paradoxical,
evolutionarily conserved protein family with multiple moonlighting functions.
Biological Reviews 88, 955–987. doi:10.1111/brv.12037.

Hestrin, R., Hammer, E. C., Mueller, C. W., and Lehmann, J. (2019). Synergies between
mycorrhizal fungi and soil microbial communities increase plant nitrogen acquisition.
Commun Biol 2, 1–9. doi:10.1038/s42003-019-0481-8.

Kameoka, H., Maeda, T., Okuma, N., and Kawaguchi, M. Structure-Specific Regulation of
Nutrient Transport and Metabolism in Arbuscular Mycorrhizal Fungi. *Plant Cell*
Physiol. doi:10.1093/pcp/pcz122.

Lammel, D. R., Barth, G., Ovaskainen, O., Cruz, L. M., Zanatta, J. A., Ryo, M., et al. (2018).
Direct and indirect effects of a pH gradient bring insights into the mechanisms driving
prokaryotic community structures. *Microbiome* 6. doi:10.1186/s40168-018-0482-8.

Leifheit, E. F., Verbruggen, E., and Rillig, M. C. (2014). Rotation of hyphal in-growth cores
has no confounding effects on soil abiotic properties. *Soil Biology and Biochemistry*
79, 78–80. doi:10.1016/j.soilbio.2014.09.006.

Magurno, F., Malicka, M., Posta, K., Wozniak, G., Lumini, E., and Piotrowska-Seget, Z.
(2019). Glomalin gene as molecular marker for functional diversity of arbuscular
mycorrhizal fungi in soil. *Biol Fertil Soils* 55, 411–417. doi:10.1007/s00374-019-
01354-x.

Mateus ID, Masclaux FG, Aletti C, Rojas EC, Savary R, Dupuis C, et al. Dual RNA-seq
reveals large-scale non-conserved genotype × genotype-specific genetic reprograming
and molecular crosstalk in the mycorrhizal symbiosis. *ISME J* 2019; **13**: 1226–1238.

Mendes, L. W., Kuramae, E. E., Navarrete, A. A., van Veen, J. A., and Tsai, S. M. (2014).
Taxonomical and functional microbial community selection in soybean rhizosphere.
The ISME Journal 8, 1577–1587. doi:10.1038/ismej.2014.17.

Mendes, L. W., Mendes, R., Raaijmakers, J. M., and Tsai, S. M. (2018a). Breeding for soil-
borne pathogen resistance impacts active rhizosphere microbiome of common bean.
ISME J 12, 3038–3042. doi:10.1038/s41396-018-0234-6.

Mendes, L. W., Raaijmakers, J. M., de Hollander, M., Mendes, R., and Tsai, S. M. (2018b).
Influence of resistance breeding in common bean on rhizosphere microbiome
composition and function. *ISME J* 12, 212–224. doi:10.1038/ismej.2017.158.

Morin, E., Miyauchi, S., Clemente, H. S., Chen, E. C. H., Pelin, A., Providencia, I. de la, et al.
(2019). Comparative genomics of Rhizophagus irregularis, R. cerebriforme, R.
diaphanus and Gigaspora rosea highlights specific genetic features in
Glomeromycotina. *New Phytologist* 222, 1584–1598. doi:10.1111/nph.15687.

Nottingham, A. T., Turner, B. L., Winter, K., Heijden, M. G. A. van der, and Tanner, E. V. J.
(2010). Arbuscular mycorrhizal mycelial respiration in a moist tropical forest. *New
Phytologist* 186, 957–967. doi:10.1111/j.1469-8137.2010.03226.x.

Pepe, A., Sbrana, C., Ferrol, N., and Giovannetti, M. (2017). An in vivo whole-plant
experimental system for the analysis of gene expression in extraradical mycorrhizal
mycelium. *Mycorrhiza* 27, 659–668. doi:10.1007/s00572-017-0779-7.

Powell, J. R., and Rillig, M. C. (2018). Biodiversity of arbuscular mycorrhizal fungi and
ecosystem function. *New Phytologist* 220, 1059–1075. doi:10.1111/nph.15119.

Qian, C., and Hettich, R. L. (2017). Optimized Extraction Method To Remove Humic Acid
Interferences from Soil Samples Prior to Microbial Proteome Measurements. *J.
Proteome Res.* 16, 2537–2546. doi:10.1021/acs.jproteome.7b00103.

Quinn, T. P., Crowley, T. M., and Richardson, M. F. (2018). Benchmarking differential
expression analysis tools for RNA-Seq: normalization-based vs. log-ratio
transformation-based methods. *BMC Bioinformatics* 19, 274. doi:10.1186/s12859-
018-2261-8.

Rillig, M. C., Aguilar☐Trigueros, C. A., Camenzind, T., Cavagnaro, T. R., Degrune, F.,
Hohmann, P., et al. (2019). Why farmers should manage the arbuscular mycorrhizal
symbiosis. *New Phytologist* 222, 1171–1175. doi:10.1111/nph.15602.

Rillig, M. C., Mardatin, N. F., Leifheit, E. F., and Antunes, P. M. (2010). Mycelium of
arbuscular mycorrhizal fungi increases soil water repellency and is sufficient to
maintain water-stable soil aggregates. *Soil Biology and Biochemistry* 42, 1189–1191.
doi:10.1016/j.soilbio.2010.03.027.

Rillig, M. C., Muller, L. A., and Lehmann, A. (2017). Soil aggregates as massively concurrent
evolutionary incubators. *ISME J* 11, 1943–1948. doi:10.1038/ismej.2017.56.

Rillig, M. C., and Mummey, D. L. (2006). Mycorrhizas and soil structure. *New Phytologist*
171, 41–53. doi:10.1111/j.1469-8137.2006.01750.x.

Rosier, C. L., Hoye, A. T., and Rillig, M. C. (2006). Glomalin-related soil protein:
Assessment of current detection and quantification tools. *Soil Biology and
Biochemistry* 38, 2205–2211. doi:10.1016/j.soilbio.2006.01.021.

Schlüter, S., Eickhorst, T., and Mueller, C. W. (2019). Correlative Imaging Reveals Holistic
View of Soil Microenvironments. *Environmental Science & Technology* 53, 829–837.
doi:10.1021/acs.est.8b05245.

Schmidt, J. E., Kent, A. D., Brisson, V. L., and Gaudin, A. C. M. (2019). Agricultural
management and plant selection interactively affect rhizosphere microbial community
structure and nitrogen cycling. *Microbiome* 7, 146. doi:10.1186/s40168-019-0756-9.

Turrini, A., Avio, L., Giovannetti, M., and Agnolucci, M. (2018). Functional
Complementarity of Arbuscular Mycorrhizal Fungi and Associated Microbiota: The
Challenge of Translational Research. *Front. Plant Sci.* 9.
doi:10.3389/fpls.2018.01407.

Upton, R. N., Bach, E. M., and Hofmockel, K. S. (2019). Spatio-temporal microbial
community dynamics within soil aggregates. *Soil Biology and Biochemistry* 132, 58–
68. doi:10.1016/j.soilbio.2019.01.016.

Wilke, A., Bischof, J., Gerlach, W., Glass, E., Harrison, T., Keegan, K. P., et al. (2016). The
MG-RAST metagenomics database and portal in 2015. *Nucleic Acids Res* 44, D590–
D594. doi:10.1093/nar/gkv1322.

World reference base for soil resources Available at:
http://www.fao.org/3/w8594e/w8594e00.htm [Accessed August 3, 2019].

Wright, S. F., Franke-Snyder, M., Morton, J. B., and Upadhyaya, A. (1996). Time-course
study and partial characterization of a protein on hyphae of arbuscular mycorrhizal
fungi during active colonization of roots. *Plant Soil* 181, 193–203.
doi:10.1007/BF00012053.

Xu, J., Liu, S., Song, S., Guo, H., Tang, J., Yong, J. W. H., et al. (2018). Arbuscular
mycorrhizal fungi influence decomposition and the associated soil microbial
community under different soil phosphorus availability. *Soil Biology and Biochemistry*
120, 181–190. doi:10.1016/j.soilbio.2018.02.010.

Yan, Y., Kuramae, E. E., Hollander, M. de, Klinkhamer, P. G. L., and Veen, J. A. van (2017).
Functional traits dominate the diversity-related selection of bacterial communities in
the rhizosphere. *ISME J* 11, 56–66. doi:10.1038/ismej.2016.108.

Yao, Q., Li, Z., Song, Y., Wright, S. J., Guo, X., Tringe, S. G., et al. (2018). Community
proteogenomics reveals the systemic impact of phosphorus availability on microbial
functions in tropical soil. *Nat Ecol Evol* 2, 499–509. doi:10.1038/s41559-017-0463-5.

Zhang, B., Li, S., Chen, S., Ren, T., Yang, Z., Zhao, H., et al. (2016). Arbuscular mycorrhizal
fungi regulate soil respiration and its response to precipitation change in a semiarid
steppe. *Scientific Reports* 6, 19990. doi:10.1038/srep19990.

Zhang, Y., Cui, M., Duan, J., Zhuang, X., Zhuang, G., and Ma, A. (2019). Abundance, rather
than composition, of methane-cycling microbes mainly affects methane emissions

from different vegetation soils in the Zoige alpine wetland. *MicrobiologyOpen* 8,
e00699. doi:10.1002/mbo3.699.

Figures Captions

Figure 1 A. Dry shoot biomass of clover inoculated with the AMF *R. irregularis* (A)

and the non-inoculated controls (C) (n=10). B. Shoot biomass for the main

treatments A, C and including the rotated core control (rC) (n=5). The full

comparisons with all replicates are available in Supp. Material 1. C. AMF

hyphal length (in m g^{-1} of soil). D. Hyphal length of other fungi (in m g^{-1} of

soil). E. New aggregates > 2mm (%) F. Water stable aggregates (%). The

ANOVA *P*-value results are reported on the bottom of each graph and

small letters below the treatment codes, when different, indicate statistical

differences by the Tukey posthoc test (*P*<0.05). The gray dots on the box-

plots indicate the individual value of each replicate. The dark dot indicates

the mean and error bars the standard error.

Figure 2 Metabolic functions in soil clustered by the MG-RAST SEED classification

accessed by soil metatranscriptomics **(A)** and metaproteomics **(B)**. The

sample codes are: static soil core inoculated with the AMF *R. irregularis*

(A), non-inoculated static soil core (C), and non-inoculated rotated soil

core (rC). For the RNA samples, the prefix **m** indicates the mRNA libraries

enriched by rRNA depletion, the other samples without prefix indicate the

poly-A selection libraries.

Figure 3 The network connectivity of soil metabolic functions (SEED level 1) across

treatments for the metatranscriptome (**C.**) and the metaproteome (**D**.). In

blue the treatment inoculated with the AMF *R. irregularis* (A), in green the

control soil (C) and in salmon the control with rotated soil cores (rC).

Figure 4 Changes in the soil microbial community structures based on the taxonomic

annotation of the sequences. On top are the general high order taxonomy

for the RNA-seq data (**A.**) and for the proteome data (**B**.) (for details on

the proteome see Supp. Figure 4). In the middle are the microbial

community structures accessed by the redundancy analysis (RDA) based

on the RNA-seq for *Bacteria* (**C1.**) and for *Eukaryota* (**D1.**). Below are the

phyla relative abundance of *Bacteria* (**C2**) and *Eukaryota* (**D2**) across the

treatments. The sample codes are: static soil core inoculated with the AMF

R. irregularis (A), non-inoculated static soil core (C), and non-inoculated

rotated soil core (rC). For the RNA samples, the prefix **m** indicate the

mixed libraries made using the rRNA depletion.

Figure 5 The genes and proteins potentially related with soil aggregation. The panel

on top show the z-score of Lpt genes across the samples (**A.**). Below is

the quantification of proteins identified in relation to the *R. irregularis*

proteome database. The MS intensity signal for Tubulin (Alfa+Beta) (**B1.**),

GroeEL/HSP60 (**B2.**) and HSP70 (uniprot U9SYW5) (**B3.**). The sample

codes are: static soil core inoculated with the AMF *R. irregularis* (A), non-

inoculated static soil core (C), and non-inoculated rotated soil core (rC).

Supporting Information

Figure S1 Cumming plots for the classical variables evaluated in the experiment. **A.** Clover dry shoot biomass. **B.** AMF hyphae length (in m.g^{-1} of soil). **C.** Hyphae length of other fungi (in m.g^{-1} of soil). **D.** AMF spores observed during the hyphae length measurements **E.** New aggregates > 2mm (%) **F.** Water stable aggregates (%). The codes are: A, soil inoculated with the AMF *R. irregularis*, C, control soil (static core), and rC, the rotated core soil.

Figure S2 Summary of the RNA-seq results. In panel **A.** the read counts across the treatments; in **B.** Effect size and Volcano plot generated by Aldex2 showing all the transcripts as dots and in light colors are the transcripts that had statistical significant changes in relation to the rotated core ($P<0.05$).

Figure S3 Proteome data after the ANPELA server processing. A. Heatmap of all the identified proteins according to treatments (ANPELA normalized values). The sample codes are: static soil core inoculated with the AMF *R. irregularis* (A), non-inoculated static soil core (C), and non-inoculated rotated soil core (rC). B. A volcano plot showing significantly regulated proteins between the AMF treatments and the rotated core control ($P<0.01$).

Figure S4 The relative abundance of the all proteins identified by the taxonomic
annotation for *Bacteria* (**A.**) and *Eukaryota* (**B.**) across the treatments.
The sample codes are: static soil core inoculated with the AMF *R.*
irregularis (A), non-inoculated static soil core (C), and non-inoculated
rotated soil core (rC).

Figure S5 The relative abundance of the GroeL/HSP60 reads of the RNAseq based
on the high order taxonomy (**A1.**) and according to *Bacteria* (**A2.**) and
Eukaryote (**A3.**) phyla. On the bottom right is the relative abundance of
the GroeL/HSP60 proteins across the proteomes (**B**). The sample codes
are: static soil core inoculated with the AMF *R. irregularis* (A), non-
inoculated static soil core (C), and non-inoculated rotated soil core (rC).
For the RNA samples, the prefix m indicate the mixed libraries made
using the rRNA depletion.

A.

rC5 rC3 m**rC** rC3 m**rC** mA A5 A7 C2 C4 C3 C5 A3 A10 mC mA mC

B.

A.

	rC	A	C	mrC	mA	mC
LptC (Bacteria)	0.6	0.3	0.0	0.8	0.1	0.1
LptF (Bacteria)	0.7	0.1	0.2	0.9	0.1	0.0
LptG (Bacteria)	0.7	0.1	0.1	0.8	0.1	0.1

B1.

B2.

B3.