Title

Comparative population genetic structure of two Ixodidae ticks (*Ixodes ovatus* and *Haemaphysalis*

flava) in Niigata Prefecture, Japan

Authors

Maria Angenica Fulo Regilme[1], Megumi Sato[2], Tsutomu Tamura[3], Reiko Arai[3], Marcello Otake Sato[4],

Sumire Ikeda[5], Masaya Doi[1], Kohki Tanaka[1], Maribet Gamboa[1], Michael T. Monaghan[6, 7], Kozo

Watanabe[1]

[1] Department of Civil and Environmental Engineering, Ehime University, Matsuyama, Japan

[2] Graduate School of Health Sciences, Niigata University, Niigata, Japan

[3] Niigata Prefectural Institute of Public Health and Environmental Sciences, Niigata, Japan

[4] Department of Tropical Medicine and Parasitology, Dokkyo Medical University, 880 Kitakobayashi,

Mibu-machi, Shimotsuga-gun, Tochigi 321-0293, Japan

[5] Research Laboratories, Research and Development Headquarters, Earth Corporation

[6] Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin, Germany

[7] Institut für Biologie, Freie Universität Berlin, Germany

Corresponding author

watanabe_kozo@cee.ehime-u.ac.jp

Abstract

Ixodid tick species function as important vectors of tick-borne diseases in Japan. In this study, we

used genetic structure at two mitochondrial loci (*cox1*, 16S rRNA gene) to infer gene flow patterns

of *Ixodes ovatus* and *Haemaphysalis flava* from Niigata Prefecture, Japan. We hypothesized that *I.*

ovatus and *H. flava* have different population genetic structure because of their host mobility in

different tick life stages despite sharing of hosts. Samples (n = 1 to 77) were collected in 29 (*I.*

ovatus) and 17 (*H. flava*) sampling locations across Niigata. For *I. ovatus*, pairwise F_{ST} and analysis

of molecular variance (AMOVA) analyses of *cox1* sequences indicated significant among-

population differentiation. This was in contrast to *H. flava*, for which there were only two cases of

significant pairwise differentiation and no overall structure. A Mantel test revealed isolation by

distance and there was positive spatial autocorrelation of haplotypes in *I. ovatus cox1* and 16S

sequences, but non-significant results were observed in *H. flava* in both markers. Newly sampled *I.*

ovatus grouped together with a published *I. ovatus* sequence from northern Japan and were

distinct from two other *I. ovatus* groups that were reported from southern China. The three

genetic groups (China 1, China 2, and Japan) in our data set suggest the potential for cryptic

species within the lineage. While many factors can potentially account for the observed

differences in genetic structure, including population persistence and large-scale patterns of range

expansion, we propose that differences in the mobility of hosts of tick immature stages (small

mammals in *I. ovatus*; birds in *H. flava*) may be driving the observed patterns.

Keywords: gene flow, host mobility, Mantel test, Spatial autocorrelation, cryptic species

Background

Tick-borne diseases are a public health concern and their control is often challenging due to the fact that it involves a complex transmission chain of vertebrate hosts and ticks interacting in a changing environment [1]. In order to understand the vector-host relationship, studies on the spatial distribution and population genetic structure of ticks are needed. Population genetic studies have helped in understanding the dispersal patterns and potential pathogen transmission in ticks [2]. The direction, distance and gene flow patterns between tick populations can be inferred from population genetic studies. For example, if high gene flow is observed there might be a greater chance of colonizing new areas or of re-colonizing areas following successful vector control programs [3]. Therefore, estimates of genetic structure and gene flow in tick populations may give insights regarding the spread of tick-borne disease.

Host mobility can affect the genetic patterns of tick populations, although its effects are not consistent. Several studies have reported low levels of gene flow in ticks with less mobile hosts (e.g. smaller mammals) and high levels of gene flow in ticks with highly mobile hosts [2]. For example, *Amblyomma americanum* and *A. triste* (Ixodidae, Acari) exhibited high gene flow across various spatial scales (137, 000 km^2 to 2.78 million km^2) and this was attributed to their hosts' dispersal capabilities (large mammals and birds) [4-6], while low gene flow was observed in *Amblyomma dissimile* populations and this was attributed to the low mobility of its hosts (small mammals, reptiles and salamanders) [7]. In contrast, some studies have reported low gene flow despite the high mobility of the host, for example in *Ixodes scapularis* [8] and *Ornithodoros coriaceus* [9] which can be attributed to the host mobility.

In Japan, tick-borne diseases are an increasing public health concern, affecting humans and animals [10]. A total of 8 genera of ticks have been recorded in Japan, composed of 47 species: 43 Ixodidae species and 4 Argasidae species [11]. Out of 47 species, 21 species are known to

parasitize humans [12]. Among these 21 species, *Ixodes ovatus*, the main vector of Lyme disease

[13] and *Haemaphysalis flava,* a vector of severe fever with thrombocytopenia syndrome (SFTS)

and Japanese spotted fever (JSF), were reported in Japan [10,14]. A previous study found that the

hosts of adult *I. ovatus* were mainly hares and can also be large mammals (e.g. cows and horses),

and the hosts of immature forms were small rodents [15]. *H. flava* adult host preference are cows,

dogs, horses, wild boar, bear, and deer, while immature forms only parasitized birds [15]. Despite

the similarity of *I. ovatus* and *H. flava* in which blood meal from different vertebrate host is taken

in each life stage (larva, nymph and adult) [16], comparative population genetic studies on two

Ixodid tick species such as *I. ovatus* and *H. flava* from the same sampling location remain

nonexistent. Thus, our study is important because there are no previous studies about dispersal

movement of two important tick species: *I. ovatus* and *H. flava*, we conclude that low gene flow

might be observed in *I. ovatus* due to the lower mobility of hares in the adult stage and small

rodents in its immature stage while *H. flava* will exhibit high gene flow because of its avian

mediated dispersal in its immature stage.

In addition to tick population structure studies, genetic studies may also reveal the presence of

cryptic species, where morphologically identified individuals might represent more than one

species [17]. Previous studies on *Rhipecephalus appendiculatus* [18], *I. holocylus* [19] and *I. ovatus*

[20] have revealed the presence of cryptic species based on the clustering of haplotypes in a

phylogenetic tree, and concluded that morphological criteria for species differentiation alone are

equivocal and that genetic analysis is important. A putative *I. ovatus* species complex was

identified based on the presence of 3 distinct haplotype clusters in both *cox1* and 16S rRNA genes;

2 groups in China and one in Japan that included haplotypes from North America [20].

Here, we studied the population genetic structure of *I. ovatus* and *H. flava* and also tested for the

presence of cryptic species using mitochondrial DNA sequences of *cox1* and the 16S rRNA gene.

We hypothesized that *I. ovatus* and *H. flava* may have differences in their population genetic

structure despite some overlap in their adult host preference because of the differences in

mobility of the immature tick hosts. We predicted that *H. flava* exhibits high gene flow patterns

due to avian-mediated dispersal of immature ticks, in contrast to *I. ovatus* which uses small

mammals as hosts during its immature stages.

Methods

Study site, collection, sampling and identification

From April 2016 until November 2017, ticks were collected by the standard flagging methods [21]

across Niigata Prefecture in Japan and a total of 29 sampling locations were surveyed for *H. flava*

and *I. ovatus* ticks (Additional File 1. Table S1). Altitude per location ranged from 8 -1402 m/a.s.l.

and the geographic distance between sites ranged from 4.28 - 247.65 km. Collected ticks were

stored in microcentrifuge tubes with ethanol at 4^0C. A total of 2,103 individual tick samples were

collected. Identification of sex, developmental stage and species was performed using a

compound microscope and identification keys of [15].

DNA extraction, PCR amplification and sequencing

Genomic DNA (*I. ovatus* n=320*; H. flava* n=220) from each identified tick was extracted using the

Isogenome DNA extraction kits (NIPPON GENE Co. Ltd., Tokyo, Japan) following the manufacturer's

recommended protocol. Prior to DNA extraction, each tick was washed with alcohol and PBS

solution. DNA quality was checked using a NanoDrop™ 2000 Spectrophotometer (Thermo

Scientific™). Two mitochondrial genes were amplified by polymerase chain reaction (PCR):

Cytochrome Oxidase 1 (*cox1*) (658 base pairs) using the primer pairs LCO-1490 (5'-

GGTCAACAAATCATAAAGATATTGG -3') and HCO1-2198 (5'- AAACTTCAGGGTGACCAAAAAATCA-3')

[22] and the 16S rRNA gene (16S) (407 base pairs) with the following primer pairs 16S+1

(5'CTGCTCAATGAATATTTAAATTGC-3') and 16S − 1 (5' -CGGTCTAAACTCAGATCATGTAGG-3') [23].

All PCR amplifications of both *cox1* and 16S were performed with a final volume of 10 µl with 1µl

of genomic DNA. The PCR reaction for both markers were composed of the following: 10x Ex Taq

buffer, 25mM MgCl$_2$, 2.5mM dNTP, 10µm of forward and reverse primers and 5 units/µl of TaKaRa

Ex Taq™ (Takara Bio Inc.). The *cox1* PCR amplification were as follows: an initial denaturation of

94^{0}C for 2 minutes, denaturation of 94^{0}C for 30 seconds, annealing of 38^{0}C for 30 seconds,

extension of 72^{0}C for 1 minute for 30 cycles and final extension of 72^{0}C for 10 minutes. While the

16S PCR amplification followed the protocol of [23] with some modifications, initial denaturation

of 94^{0}C for 3 minutes, denaturation of 94^{0}C for 30 seconds, annealing of 50^{0}C for 40 seconds,

extension of 72^{0}C for 40 seconds for 30 cycles and final extension of 72^{0}C for 5 minutes. PCR

products were purified using the QIAquick 96 PCR Purification Kit (Qiagen) in accordance with the

manufacturer's instructions, and sequenced by Eurofin Genomics, Inc., Tokyo, Japan.

Sequence data analysis

We assembled forward and reverse reads for each individual using CodonCode Aligner ver 1.2.4

software (https://www.codoncode.com/aligner/). No ambiguous bases were observed and the

low quality bases were removed in the start and end of the reads.   Multiple alignment of

sequences was performed using the MAFFT alignment online program with default settings

(https://mafft.cbrc.jp/alignment/server/). To ensure sequence quality and correct species

identification, we checked the similarity of the sequences against reference sequences from

GenBank   using   BLASTN   (Basic   Local   Alignment   Search   Tool   −   Nucleotide,

https://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM=blastn&PAGE_TYPE=BlastSearch&LINK_LOC=bl

asthome). The quality of the final aligned sequences was checked in Mesquite ver 3.5 [24] wherein

the protein coding genes were translated into amino acids to confirm the absence of stop codons.

Population genetic analysis

We analyzed the final sequences of both *H. flava* and *I. ovatus* per genetic marker separately using

DNASp ver 6.12.03 [25] and Arlequin ver 3.5.2.2 [26] for the following parameters: number of

haplotypes, average number of polymorphic sites, average number of nucleotide differences,

gene/haplotype diversity and nucleotide diversity. The population genetic structure among and

within sampling locations in Niigata Prefecture was calculated by analysis of molecular variance

(AMOVA) performed in Arlequin with 1023 permutations. The degree of molecular genetic

differentiation between sampling locations defined as populations was assessed by calculation of

the pairwise F_{ST} values in Arlequin.

To determine if the genetic differentiation was influenced by geographical distance or altitudinal

differences among populations, we performed Mantel Test in GenAlEx ver 6.51b2 [27]. Two tests

per species and marker were conducted. First we compared pairwise genetic (pairwise F_{ST} values)

and geographical distances (km) and second we compared genetic distance (F_{ST} values) with

altitudinal differences (m/a.s.l.). The geographic distances were obtained from the GPS

coordinates (latitude and longitude) recorded during the sampling. All Mantel tests were assessed

using 9999 permutations for the significance of the correlation.

The spatial component of genetic variation was further assessed by spatial autocorrelation [28]

using GenAlEx [27]. The autocorrelation coefficient (*r*) was computed from the geographic

distance between sampling locations and the genetic distance (pairwise F_{ST} values). This measures

the genetic similarity between individual pairs within an identified distance class. The size of the

distance class influences the estimation of the spatial autocorrelation. We identified the

appropriate distance class based on the observed distribution of pairwise geographic distance

between the sampling locations (data not shown). The distance class sizes used were the

following: 20, 40, 60, 80, 100, 120, 140, 160, 180, 200, 220, 240, 260, 280, 300, and 320 kilometers.

Bootstrap estimates of *r* and random permutations were set at 9999 each for the test for

significance. The upper and lower error bars in the correlogram bound the 95% confidence interval

of *r* value as identified by bootstrap resampling. The upper and lower confidence limits bound the

95% confidence interval of the null hypothesis of no spatial structure in the data set.

The genetic relationships among the sampling locations (i.e., populations) were calculated by

unweighted pair group method with the arithmetic mean (UPGMA) cluster method using the APE

package [29] and R studio [30] to create dendrogram using the genetic distance matrix (pairwise

F_{ST} values) generated from GenAlEx.

Phylogenetic analysis

We constructed maximum likelihood (ML) gene trees for *cox1* and 16S haplotype sequences of *I.*

ovatus and *H. flava* using PhyML ver 3.1 [31] under the default settings. We applied HKY and GTR

nucleotide substitution models for *cox1* and 16S, respectively, as suggested by jModelTest ver 2

[32]. Additional sequences from China (MH208506, MH208512, MH208514, MH208522,

MH208515-19, MH208524, MH208531, MH208574, MH208577, MH208579, MH208681-87,

MH208689-93, MH208706, KU664519 [20]), Japan (Hokkaido, Yamanashi and Aomori) (AB231670,

AB819241, AB819243, AB819244 [33-34]) and the USA (U95900; [35]) were included. We used

Ixodes canisuga as an outgroup (KY962023, KY962074; [36]).

Results

The number of successfully sequenced and analyzed *H. flava* ranged from 1 to 36 individuals per

study site, with more successfully sequenced for *cox1* (220/223; 98.7%) than for 16S (172/223;

77.1%). The number of *I. ovatus* ranged from 1 to 36 individuals per study site and, as with *H. flava*,

more were successfully sequenced for *cox1* (307/320; 95.9%) than for 16S (284/320; 88.8%)

(Additional File 1. Table S1). There were 63 and 60 *cox1* haplotypes in *H. flava* and *I. ovatus*,

respectively, and 40 and 24 16S haplotypes (Table 1). Haplotype diversity ranged from 0.442 to

0.852 and nucleotide diversity ranged from 0.001 to 0.006 for both markers and species (Table 1).

AMOVA results revealed a high among-population divergence (41.54%) in *I. ovatus cox1* sequences,

whereas both *cox1* and 16S markers of *H. flava* indicated no significant genetic variation among

populations (Table 2). Global F_{ST} values were highest in *I. ovatus* with values 0.415 in *cox1* and

0.074 in 16S sequences. Pairwise F_{ST} values (0.500 to 1.00) of *I. ovatus* based on *cox1* showed

significant genetic differences between most pairs of populations such as between locations 18

and 25 and between locations 28 and 25 (Additional File 2 Table S2). No significant differences

were observed in 16S (Additional File 3 Table S3). Pairwise F_{ST} values from the *cox1* marker of *H.*

flava showed no significant genetic differences except between locations 18 and 15 and 18 and 25

(Additional File 4 Table S4). No significant genetic differentiation was also observed in 16S marker

of H. flava except in between locations 3 and 20 and 4 and 20 (Additional File 5 Table S5).

The Mantel test of *I. ovatus* showed evidence of isolation by geographic distance in both markers:

(*cox1 r* = 0.197, *p* = 0.008; 16S *r* = 0.293, *p* = 0.011) (Figures 1a, b). In contrast, there was no

evidence for isolation by altitudinal difference (*cox1* r = -0.066, *p*=0.225; 16S r = -0.023, p = 0.577).

Both Mantel tests were not significant for *H. flava* (distance: *cox1* r= -0.145, p= 0.23; 16S r = -0.049,

p= 0.33; altitude: r = 0.092, p = 0.30; 16S r = 0.217, p= 0.06). The spatial autocorrelation test in *I.*

ovatus revealed positive and significant correlation between genetic and geographic distance from

20 – 40 km, and 60 – 80 km, and 20 – 50 km and 120 – 160 km from *cox1* and 16S, respectively

(Figure 1c, d).

The UPGMA cluster dendrogram from the pairwise F_{ST} values from the *cox1*, *I. ovatus* (Figure 2)

revealed 2 genetic clusters across the 29 sampling locations, where the cluster 2 included

populations from the more western sites including the island (location 24 and 25) while cluster 1

mostly included populations in the northern and southern part of Niigata Prefecture (Figure 3). No

evident genetic clustering on the dendrogram was observed from the *I. ovatus* 16S sequences and

in both *cox1* and 16S *H. flava* sequences (Additional File 6 Figure S1; Additional File 7 Figure S2;

Additional File 8 Figure S3).

The *cox1* gene tree of 60 *I. ovatus* haplotypes indicated three groups: group 1 of the published

sequences from western China, one large group 2 containing the 58 haplotypes and 2 divergent

haplotypes (Hap 60 and Hap 59), and group 3 from western China (Figure 4). The 2 divergent

haplotypes were singletons. One was found in location 29 and the other in location 7. The

published haplotype from Hokkaido [33] occurred within our large group and the published

sequences from western China did not. The *I. ovatus* 16S gene tree (24 newly sequenced

haplotypes) (Additional File 9 Figure S4) also recovered published haplotypes from Japan

(Yamanashi and Aomori; [34]) as well as from North America [35] within the same cluster as our

new sequences. The *cox1* haplotype sequences of *H. flava* were similar to reference sequences

from China (Additional File 10 Figure S5) [unpublished manuscript] while the 16S haplotype

sequences (Additional File 11 Figure S6) showed similarities with Japan (Kagoshima, Fukui Aomori,

Fukui, Yamanashi, Kagawa and Ehime [34]) and China [unpublished manuscript] published

sequences.

Discussion

Contrasting population genetic structures between *H. flava* and *I. ovatus*

Our results supported our hypothesis that *H. flava* and *I. ovatus* display contrasting population

genetic structure. The noticeable high gene flow in *H. flava* may be a result of the higher level of

mobility of its hosts such as birds in tick's immature stage and the more pronounced genetic

structure and lower gene flow in *I. ovatus* may be due to the restricted ability of its small

mammalian hosts at the immature stage to disperse further. 28 species of birds were previously

reported as hosts of *H. flava,* mostly from the order Passeriformes known as migratory birds [37].

The host for adults of *I. ovatus* included large mammals such as hares while immature forms are

found on small mammals such as small rodents and rats [15]. The movement of small mammals is

limited because of environmental and ecological factors such as difficulty in crossing roads and

higher possibility of road kill due to vehicle collision and or traffic [38], which may explain the

genetic structuring in *I. ovatus*. The host preference of the adults *H. flava* are large mammals (e.g.

cows, horses, wild boar, deer, bear) and hares, dogs, rodents while the immature forms parasitize

birds [15]. Mammals and birds may have wide habitats ranges that may allow maintaining the

gene flow of *H. flava* between the locations in Niigata, as previously observed in *Amblyomma*

americanum populations [5,6,39], and *Ixodes ricinus* [40]. Birds present the greatest potential for

farther dispersal of *H. flava* that could reduce its genetic structuring [41] that supports the results

of our study.

Several alternative factors such as tick behavior, biology and ecology can also affect the tick

dispersal movement. For example, long movement patterns*,* resistance against unfavorable

weather conditions causing high survival in their hosts and low host specificity [4] are possible

reasons for the low genetic structuring in *H. flava* populations. It could also be influenced by

ecological factors such as changes in land use and cover, wildlife and forestry management as

observed in the previous study of [42] in the *Ixodes ricinus* tick populations from Europe. Previous

study on two tick species: *Hyalomma rufipes* and *Amblyomma hebraum* also displayed contrasting

genetic pattern despite the overlap sharing of highly mobiles hosts such as birds [43] due to the

capacity of the immature stages of ticks to survive off the hosts in various habitat conditions [44-

46]. Population structure can also be influenced by assortative mating (e.g. *Ixodes ricinus*) wherein

genetically similar individuals tend to mate, rather than random mating resulting in increased

genetic divergence [47]. Our study does not have supporting data to test these alternative factors

thus it is suggested in future studies to further analyze these factors that could play a role in the

tick's dispersal movement.

Based on previous studies [48-49], isolation by distance suggests that the populations are in

equilibrium of migration and genetic drift at the spatial scale of Niigata Prefecture. Our results

were congruent with a previous study of *Amblyomma ovale* that showed positive relationship of

geographic and genetic distance at the spatial scale of the whole of Brazil whose main hosts for

the immature stage are small rodents while tapir and jaguar in its adult stage [50]. In contrast, *H.*

flava populations showed an absence of isolation by distance. In circumstances when geographic

distance can't explain the genetic structure of tick populations, other factor such as ecological

variables (e.g. land use) maybe affecting the genetic structuring within tick populations [7, 40, 51].

A significant and positive spatial autocorrelation of *I. ovatus* in both *cox1* and 16S sequences was

observed within the geographic range of 20 – 40 km and 60- 80 km in *cox1* sequences and 20 – 50

km and 120 – 160 km range in the 16S sequences. We can infer that the positive *r* values at the

maximum geographic rages of 60-80 km in *cox1* and 120 - 160 km in 16S *I. ovatus* sequences

indicate that gene flow is occurring in the wide geographic range. The lack of significance at 40-60

km in *cox1* may suggest patch structure indicating genetic similarity or variation [28] as observed

on limpet species, *Siphonaria raoulensis* from New Zealand that also displayed recovery of positive

r values at greater distances [52]. However, no autocorrelation was observed in *H. flava* at any

distance class, suggesting that gene flow occurring has no specific pattern favoring certain

distances [53-55]. Since the sampling geographic range of our study was within Niigata Prefecture

and distances >220 km were not included in our sampling, we suggest that future studies collect

samples at a wider geographic scale to more precisely determine the geographic distance at which

autocorrelation exists.

16S *I. ovatus* sequences showed genetic differentiation (pairwise F_{ST} = 0.5-0.6) in few pairs of

sampling locations as compared to the *cox1* sequences which may be due to lower variability of

the marker thus providing inadequate genetic variation for population genetic analysis, as

previously observed in *Amblyomma ovale* [50] and *Rhipicephalus microplus* [56-57]. The low

variability of the 16S in *I. ovatus* may not be sufficient to infer intraspecific relationship and is

further supported by its low (*nd* = 0.001) nucleotide diversity as compared in *cox1 I. ovatus* (*nd* =

0.004)*. Thus, it may also explain the grouping observed in the *cox1 I. ovatus* tree which is absent in

the 16S *I. ovatus* sequences. Despite of these, we used 16S mitochondrial gene because this gene

mutates at a rate that is informative for species-level phylogenetic and broad biogeographic

inferences [2]

The negative pairwise F_{ST} values obtained in *H. flava* sequences do not have any biological meaning.

These results from the unbiased estimator allowing a negative F_{ST} when the sample size is small.

[58-60]. It can also be noted that we found significant pairwise F_{ST} values in 2 sampling location in

H. flava cox1 sequences probably due to the fact that both locations have only one tick sample.

Although, some sampling sites in this study have small sample size (n < 10), the *cox1* and 16S

sequences provided valuable insight into the genetic structure and gene flow patterns of the

present-day tick populations. Thus, for future research, we recommend to increase the sample

size per sampling site.

Species complex formation in *I. ovatus cox1* sequences

Our *cox1 I. ovatus* gene tree showed Japanese individuals to form a group that was distinct from

haplotypes from southern China [20]. Despite the low gene flow we found in *cox1 I. ovatus*,

haplotypes found from Niigata were closely related to the published sequence from Hokkaido,

Japan which may indicate that these ticks originated from a diverse set of geographical locations in

Japan which might be transported by its hosts or is undergoing recent population expansion from

northern Japan (Hokkaido) to south (Niigata) or *vice versa*. We found three groups (China 1, Japan

and China 2) and 2 slightly divergent *cox1* haplotypes (Hap 60 and 59) in the Japan group of *I.*

ovatus in Niigata. Considering two or more cryptic species can be concealed in one

morphologically described species [59], the occurrence of the 3 groups and the divergent

haplotypes suggests that *I. ovatus* may be a species complex. Previous studies have also observed

species complexes in *Ixodes* and *Rhipicephalus* [19-20,57;62] suggesting that morphological

criteria for species differentiation alone are equivocal and that genetic analysis is important.

Conclusion

Two Ixodid ticks, *H. flava* and *I. ovatus* exhibited contrasting patterns of genetic structure in

Niigata Prefecture, Japan. We suggest that these differences are due to differences in host

mobility of the immature life stages of the ticks. The more pronounced population genetic

structure of *I. ovatus* might be influenced by the restricted movement of its small mammalian host

in its immature stage, whereas the lack of structure in *H. flava* may be due to its avian-mediated

movement in its immature stage. Even though *I. ovatus* populations were genetically structured

within Niigata, a published haplotype from Hokkaido was also found, indicating that widespread

dispersal is possible. The occurrence of three groups and the divergent cox1 haplotypes in *I.*

ovatus emphasizes the need of additional methods in determining the existence of species

complex in *I. ovatus* populations in Japan.

List of Abbreviations

H. flava – Haemaphysalis flava

I. ovatus – Ixodes ovatus

AMOVA – Analysis of molecular variance

UPGMA - unweighted pair group method with arithmetic mean

ML – maximum likelihood

F_{ST} – fixation index

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All data generated or analyzed during the study are included in this published article and its

supplementary additional files. All the newly generated sequences are available in the Genbank

database under the accession numbers _______.

Competing interests

The authors declare that they have no competing interests.

Funding

This work is funded by the Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for

Scientific Research (16K00569, 19K21996, 19H02276), the Sumitomo Electric Industries Group

Corporate Social Responsibility Foundation and the Research Unit Program of Ehime University.

The research was partially supported by the German Academic Exchange Service (DAAD,

Programm Projektbezogener Personenaustausch Japan 2018, project 57402018).

Authors' contributions

MAFR, MG, MM, KW conceptualized and the designed the experiment. MS, TT, RA, SI and MOS designed the sampling collection, collected and identified the tick samples of this study. MD and KT conducted the molecular analyses. MAFR, MG and KW performed the data analysis and wrote the manuscript. All authors read and approve the manuscript.

Acknowledgements

The authors would like to thank Niigata Prefectural Office for their valuable help during the tick sampling collection. Thank you to Dr. Thaddeus Carvajal and Joeselle Serrana for their valuable suggestions in the population genetic analyses in this study and Micanaldo Francisco for the technical assistance in constructing the sampling map of this study.

References

1. Dantas-Torres F, Chomel BB, Otranto D. Ticks and tick-borne diseases: a One Health perspective. Trends in parasitology. 2012 Oct 1;28(10):437-46.

2. Araya-Anchetta A, Busch JD, Scoles GA, Wagner DM. Thirty years of tick population genetics: a comprehensive review. Infection, Genetics and Evolution. 2015 Jan 1;29:164-79.

3. Mccoy K. The population genetic structure of vectors and our understanding of disease epidemiology. Parasite. 2008Aug;15(3):444–8.

4. Guglielmone AA, Nava S, Mastropaolo M, Mangold AJ. Distribution and genetic variation of *Amblyomma triste* (Acari: Ixodidae) in Argentina. Ticks and tick-borne diseases. 2013 Sep 1;4(5):386-90.

5. Mixson TR, Lydy SL, Dasch GA, Real LA. Inferring the population structure and demographic history of the tick, *Amblyomma americanum* Linnaeus. Journal of Vector Ecology. 2006 Jun;31(1):181-93.

6. Trout RT, Steelman CD, Szalanski AL. Population genetics of *Amblyomma americanum* (Acari: Ixodidae) collected from Arkansas. Journal of medical entomology. 2014 Dec 1;47(2):152-61.

7. Lampo M, Rangel Y, Mata A. Population genetic structure of a three-host tick, *Amblyomma dissimile*, in eastern Venezuela. Journal of Parasitology. 1998 Dec 1;84:1137-42.

8. Qiu WG, Dykhuizen DE, Acosta MS, Luft BJ. Geographic uniformity of the Lyme disease spirochete (*Borrelia burgdorferi*) and its shared history with tick vector (*Ixodes scapularis*) in the northeastern United States. Genetics. 2002 Mar 1;160(3):833-49.

9. Teglas MB, May B, Crosbie PR, Stephens MR, Boyce WM. Genetic structure of the tick *Ornithodoros coriaceus* (Acari: Argasidae) in California, Nevada, and Oregon. Journal of medical entomology. 2005 May 1;42(3):247-53.

10. Yamaji K, Aonuma H, Kanuka H. Distribution of tick-borne diseases in Japan: past patterns and implications for the future. Journal of infection and chemotherapy. 2018 Jul 1;24(7):499-504.

11. Fujita H. Identification of immature ticks in Japan. Acari and Emerging/Reemerging Disease. 2007:53-68.

12. Okino T, Ushirogawa H, Matoba K, Hatsushika R. Bibliographical studies on human cases of hard tick (Acarina: Ixodidae) bites in Japan (7) cases of unidentified tick infestation. Kawasaki Medical Journal. 2010;36(2):127-41.

13. Miyamoto K, Nakao M, Fujita H, Sato F. The ixodid ticks on migratory birds in Japan and the isolation of Lyme disease spirochetes from bird-feeding ticks. Medical entomology and zoology. 1993 Dec 15;44(4):315-26.

14. Yu XJ, Liang MF, Zhang SY, Liu Y, Li JD, Sun YL, Zhang L, Zhang QF, Popov VL, Li C, Qu J. Fever with thrombocytopenia associated with a novel bunyavirus in China. New England Journal of Medicine. 2011 Apr 21;364(16):1523-32.

15. Yamaguti N, Tipton VJ, Keegan HL, Toshioka S. Ticks of Japan, Korea, and the Ryukyu islands. Brigham Young University Science Bulletin, Biological Series. 1971;15(1):1.

16. Herrmann C, Gern L. Search for blood or water is influenced by *Borrelia burgdorferi* in Ixodes ricinus. Parasites & vectors. 2015 Dec;8(1):6.

17. Fegan M, Prior P. How complex is the *Ralstonia solanacearum* species complex. Bacterial wilt disease and the *Ralstonia solanacearum* species complex. 2005 Jan 1;1:449-61.

18. Kanduma EG, Mwacharo JM, Githaka NW, Kinyanjui PW, Njuguna JN, Kamau LM, Kariuki E, Mwaura S, Skilton RA, Bishop RP. Analyses of mitochondrial genes reveal two sympatric but genetically divergent lineages of *Rhipicephalus appendiculatus* in Kenya. Parasites & vectors. 2016 Dec;9(1):353.

19. Song S, Shao R, Atwell R, Barker S, Vankan D. Phylogenetic and phylogeographic relationships in *Ixodes holocyclus* and *Ixodes cornuatus* (Acari: Ixodidae) inferred from COX1 and ITS2 sequences. International journal for parasitology. 2011 Jul 1;41(8):871-80.

20. Li LH, Zhang Y, Wang JZ, Li XS, Yin SQ, Zhu D, Xue JB, Li SG. High genetic diversity in hard ticks from a China-Myanmar border county. Parasites & vectors. 2018 Dec;11(1):469.

21. Ginsberg HS, Ewing CP. Comparison of flagging, walking, trapping, and collecting from hosts as sampling methods for northern deer ticks, *Ixodes dammini*, and lone-star ticks,

Amblyomma americanum (Acari: Ixodidae). Experimental & applied acarology. 1989 Sep 1;7(4):313-22.

22. Folmer O, Black, M, Hoeh W, Lutz R and Vrijenhoek R. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Molecular Marine Biology and Biotechnology. 1994;(3):294-9.

23. Tian Z, Liu G, Xie J, Yin H, Luo J, Zhang L, Zhang P, Luo J. Discrimination between *Haemaphysalis longicornis* and *H. qinghaiensis* based on the partial 16S rDNA and the second internal transcribed spacer (ITS-2). Experimental and applied acarology. 2011 Jun 1;54(2):165-72.

24. Maddison WP, Maddison DR. Mesquite: a modular system for evolutionary analysis. 2011. See http://mesquiteproject. org. 2011.

25. Rozas J, Ferrer-Mata A, Sánchez-DelBarrio JC, Guirao-Rico S, Librado P, Ramos-Onsins SE, Sánchez-Gracia A. DnaSP 6: DNA sequence polymorphism analysis of large data sets. Molecular biology and evolution. 2017 Sep 18;34(12):3299-302.

26. Excoffier L, Laval G, Schneider S. Arlequin (version 3.0): an integrated software package for population genetics data analysis. Evolutionary bioinformatics. 2005 Jan;1:117693430500100003.

27. Peakall RO, Smouse PE. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. Molecular ecology notes. 2006 Mar;6(1):288-95.

28. Smouse PE, Peakall RO. Spatial autocorrelation analysis of individual multiallele and multilocus genetic structure. Heredity. 1999 May;82(5):561-73.

29. Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and evolutionary analyses in R. Bioinformatics. 2018 Jul 17;35(3):526-8.

30. Team RC. R: A language and environment for statistical computing.

31. Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large phylogenies by maximum likelihood. Systematic biology. 2003 Oct 1;52(5):696-704.

32. Darriba D, Taboada GL, Doallo R, Posada D. jModelTest 2: more models, new heuristics and parallel computing. Nature methods. 2012 Aug;9(8):772.

33. Mitani H, Takahashi M, Masuyama M, Fukunaga M. *Ixodes philipi* (Acari: Ixodidae): phylogenetic status inferred from mitochondrial cytochrome oxidase subunit I gene sequence comparison. The Journal of parasitology. 2007 Jun 1:719-22.

34. Takano A, Fujita H, Kadosaka T, Takahashi M, Yamauchi T, Ishiguro F, Takada N, Yano Y, Oikawa Y, Honda T, Gokuden M. Construction of a DNA database for ticks collected in Japan: application of molecular identification based on the mitochondrial 16S rDNA gene. Medical Entomology and Zoology. 2014 Mar 25;65(1):13-21.

35. Norris DE, Klompen JS, Black WC. Comparison of the mitochondrial 12S and 16S ribosomal DNA genes in resolving phylogenetic relationships among hard ticks (Acari: Ixodidae). Annals of the Entomological Society of America. 1999 Jan 1;92(1):117-29.

36. Hornok S, Sándor AD, Beck R, Farkas R, Beati L, Kontschán J, Takács N, Földvári G, Silaghi C, Meyer-Kayser E, Hodžić A. Contributions to the phylogeny of *Ixodes* (Pholeoixodes) *canisuga*, I.(Ph.) kaiseri, I.(Ph.) hexagonus and a simple pictorial key for the identification of their females. Parasites & vectors. 2017 Dec;10(1):545.

37. Yamauchi T, Takeno K. Record of *Haemaphysalis flava* Neumann (Acari, Ixodidae) from *Zoothera dauma* (Latham)(Aves) in Kyushu, Japan. Esakia. 2000 Mar 31;40:99-101.

38. Haemig PD, Waldenström J, Olsen B. Roadside ecology and epidemiology of tick-borne diseases. Scandinavian journal of infectious diseases. 2008 Jan 1;40(11-12):853-8.

39. Reichard MV, Kocan AA, Van Den Bussche RA, Barker RW, Wyckoff J, Ewing SA. Sequence variation of the ribosomal DNA second internal transcribed spacer region in two spatially

distinct populations of *Amblyomma americanum* (L.)(Acari: Ixodidae). Journal of Parasitology. 2005 Apr;91(2):260-4.

40. Casati S, Bernasconi MV, Gern L, Piffaretti JC. Assessment of intraspecific mtDNA variability of European *Ixodes ricinus* sensu stricto (Acari: Ixodidae). Infection, Genetics and Evolution. 2008 Mar 1;8(2):152-8.

41. Mclain DK, Wesson DM, Oliver Jr JH, Collins FH. Variation in ribosomal DNA internal transcribed spacers 1 among eastern populations of *Ixodes scapularis* (Acari: Ixodidae). Journal of Medical Entomology. 1995 May 1;32(3):353-60.

42. Medlock JM, Hansford KM, Bormane A, Derdakova M, Estrada-Peña A, George JC, Golovljova I, Jaenson TG, Jensen JK, Jensen PM, Kazimirova M. Driving forces for changes in geographical distribution of *Ixodes ricinus* ticks in Europe. Parasites & vectors. 2013 Dec 1;6(1):1.

43. Cangi N, Horak IG, Apanaskevich DA, Matthee S, Das Neves LC, Estrada-Peña A, Matthee CA. The influence of interspecific competition and host preference on the phylogeography of two African Ixodid tick species. PLoS One. 2013 Oct 9;8(10):e76930.

44. Needham GR, Teel PD. Water balance by ticks between bloodmeals. Morphology, physiology, and behavioral biology of ticks/editors, John R. Sauer and J. Alexander Hair. 1986.

45. Cumming GS. Host distributions do not limit the species ranges of most African ticks (Acari: Ixodida). Bulletin of Entomological Research. 1999 Apr;89(4):303-27.

46. Estrada-Peña A. Ticks as vectors: taxonomy, biology and ecology. Revue scientifique et technique. 2015 Apr 1;34(1):53-65.

47. Kempf F, De Meeûs T, Arnathau C, Degeilh B, Mccoy KD. Assortative pairing in *Ixodes ricinus* (Acari: Ixodidae), the European vector of Lyme borreliosis. Journal of medical entomology. 2014 Oct 29;46(3):471-4.

48. Yaegashi S, Watanabe K, Monaghan MT, Omura T. Fine-scale dispersal in a stream caddisfly inferred from spatial autocorrelation of microsatellite markers. Freshwater Science. 2014 Feb 4;33(1):172-80.

49. Watanabe K, Monaghan MT, Takemon Y, Omura T. 1775201. Dispersal ability determines the genetic effects of habitat fragmentation in three species of aquatic insect. Aquatic conservation. 2010;20(5):574-9.

50. Bitencourth K, Amorim M, de Oliveira SV, Voloch CM, Gazêta GS. Genetic diversity, population structure and rickettsias in *Amblyomma ovale* in areas of epidemiological interest for spotted fever in Brazil. Medical and veterinary entomology. 2019 Jun;33(2):256-68.

51. Krakowetz CN, Dergousoff SJ, Chilton NB. Genetic variation in the mitochondrial 16S rRNA gene of the American dog tick, *Dermacentor variabilis* (Acari: Ixodidae). Journal of Vector Ecology. 2010 Jun;35(1):163-73.

52. Wood AR, Gardner JP. Small spatial scale population genetic structure in two limpets (*Scutellastra kermadecensis* and *Siphonaria raoulensis*) endemic to the Kermadec Islands, New Zealand. Marine Ecology Progress Series. 2007;349:159-70.

53. Pardanani N, Mather TN. Lack of spatial autocorrelation in fine-scale distributions of *Ixodes scapularis* (Acari: Ixodidae). Journal of medical entomology. 2004 Sep 1;41(5):861-4.

54. Kitron U, Kazmierczak JJ. Spatial analysis of the distribution of Lyme disease in Wisconsin. American journal of Epidemiology. 1997 Mar 15;145(6):558-66.

55. Nicholson MC, Mather TN. Methods for evaluating Lyme disease risks using geographic information systems and geospatial analysis. Journal of Medical Entomology. 2014 Sep 11;33(5):711-20.

56. Low VL, Tay ST, Kho KL, Koh FX, Tan TK, Lim YA, Ong BL, Panchadcharam C, Norma-Rashid Y, Sofian-Azirun M. Molecular characterisation of the tick *Rhipicephalus microplus* in Malaysia: new insights into the cryptic diversity and distinct genetic assemblages throughout the world. Parasites & vectors. 2015 Dec;8(1):341.

57. Burger TD, Shao R, Barker SC. Phylogenetic analysis of mitochondrial genome sequences indicates that the cattle tick, *Rhipicephalus* (Boophilus) *microplus*, contains a cryptic species. Molecular Phylogenetics and Evolution. 2014 Jul 1;76:241-53.

58. Weir BS, Cockerham CC. Estimating F-statistics for the analysis of population structure. evolution. 1984 Nov;38(6):1358-70.

59. Bitencourth K, Voloch CM, Serra-Freire NM, Machado-Ferreira E, Amorim M, Gazêta GS. Analysis of *Amblyomma sculptum* haplotypes in an area endemic for Brazilian spotted fever. Medical and veterinary entomology. 2016 Sep;30(3):342-50.

60. Vial L, Durand P, Arnathau C, Halos L, Diatta G, Trape JF, Renaud F. Molecular divergences of the *Ornithodoros sonrai* soft tick species, a vector of human relapsing fever in West Africa. Microbes and infection. 2006 Sep 1;8(11):2605-11.

61. Kwon-Chung KJ, Bennett JE, Wickes BL, Meyer W, Cuomo CA, Wollenburg KR, Bicanic TA. The Case for Adopting the" Species Complex" Nomenclature for the Etiologic Agents of *Cryptococcosis*. Sphere 2017; 2 (1): e00357-16. DOI: https://doi. org/10.1128/mSphere.:00357-16.

62. Xu G, Fang QQ, Keirans JE, Durden LA. Molecular phylogenetic analyses indicate that the *Ixodes ricinus* complex is a paraphyletic group. The Journal of parasitology. 2003 Jun 1:452-7.

Additional Files

Additional File 1.xls Table S1. Summary of *Haemaphysalis flava* and *Ixodes ovatus* collected from the different locations of Niigata Prefecture and its corresponding sample number

Additional File 2.xls Table S2. Pairwise comparison of genetic differentiation (F_{ST}) calculated for all *I. ovatus* using the *cox1* gene

Additional File 3.xls Table S3. Pairwise comparison of genetic differentiation (F_{ST}) calculated for all *I. ovatus* using the 16S gene

Additional File 4.xls Table S4. Pairwise comparison of genetic differentiation (F_{ST}) calculated for all *H. flava* using the *cox1* gene

Additional File 5.xls Table S5. Pairwise comparison of genetic differentiation (F_{ST}) calculated for all *H. flava* using the 16S gene

Additional File 6.docx Figure S1. An unweighted pair group method with the arithmetic mean (UPGMA) dendrogram of *I. ovatus* based on the pairwise genetic distance (F_{ST}) of 16S among 23 sampling locations across Niigata Prefecture, Japan.

Additional File 7.docx Figure S2 An unweighted pair group method with the arithmetic mean (UPGMA) dendrogram of *H. flava* based on the pairwise genetic distance (F_{ST}) of *cox1* among 17 sampling locations across Niigata Prefecture, Japan.

Additional File 8.docx Figure S3 An unweighted pair group method with the arithmetic mean (UPGMA) dendrogram of *H. flava* based on the pairwise genetic distance (F_{ST}) of 16S among 17 sampling locations across Niigata Prefecture, Japan.

Additional File 9.docx Figure S4 Phylogenetic tree of 16s haplotype sequences of I. ovatus inferred by maximum likelihood analysis with evolutionary model GTR with sequences from China (MH208506, MH208512, MH208514 – MH208519, MH208522, MH208524, MH208531, MH208574, MH208577, MH208579, KU664519), Japan (AB819241, AB819242, AB819243, AB819244) and US (U95900). An outgroup (Ixodes canisuga) was also included.

Additional File 10.docx Figure S5 Phylogenetic tree of *cox1* haplotype sequences of *H. flava* inferred by maximum likelihood analysis with evolutionary model HKY with sequences from China (KY021800- KY021807; KY021810-KY021819, KY003181, JQ62588 - JQ625889; JQ737097; JF758632). An outgroup (*Ixodes canisuga)* was also included.

Additional File 11.docx Figure S6 Phylogenetic tree of 16S haplotype sequences of *H. flava* inferred by maximum likelihood analysis with evolutionary model GTR with sequences from China (KC844858 -KC844867; KC844880-KC844882, M696720, KP324926, KX450279) and Japan (AB19177- AB819192). An outgroup (*Ixodes canisuga)* was also included.

Tables

Table 1. Summary of *cox1* and 16S haplotype distribution, variability, genetic diversity of *Ixodes ovatus* and *Haemaphysalis flava* populations in Niigata Prefecture

Marker	Species	n	h	s	k	hd±SD	nd±SD
cox1	*I. ovatus*	307	60	65	2.728	0.852 ± 0.012	0.004 ± 0.000
	H. flava	220	63	60	1.472	0.789 ± 0.026	0.002 ± 0.000
16S	*I. ovatus*	284	24	22	0.699	0.442 ± 0.037	0.001 ± 0.000
	H. flava	172	40	49	2.447	0.835 ± 0.024	0.006 ± 0.000

Abbreviations: *n* sample size; *h* number of haplotypes; *s* number of polymorphic sites; *k* mean number of nucleotide differences; *hd±SD* haplotype diversity ± Standard Deviation; *nd±SD* nucleotide diversity ± Standard Deviation

Table 2. Analysis of molecular variance (AMOVA) using *cox1* and 16S of *Ixodes ovatus* and *Haemaphysalis flava* populations

Marker	Species		df	ss	vc	pv	F_{ST}
cox1	*I. ovatus*	Among	28	190.38	0.5803 Va	41.54	0.4154**
		Within	278	227.04	0.8167 Vb	58.46	
	H. flava	Among	16	12.42	0.0039 Va	0.53	0.0053
		Within	203	148.73	0.7327 Vb	99.47	
16S	*I. ovatus*	Among	22	14.04	0.0260 Va	7.39	0.0739**
		Within	261	84.87	0.3252 Vb	92.61	
	H. flava	Among	12	23.6	0.0642 Va	5.21	0.0521**
		Within	159	185.58	1.1672 Vb	94.79	

Abbreviations: *df* degrees of freedom; *ss* sum of squares; *Va, Vb* and *Vc* are associate covariance components; *pv* percentage variation; *P<0.05 ** P<0.01

Figures

Figure 1. Plots (a) and (b) show the relationship between the pairwise F_{ST} values and the geographic distances for *I. ovatus* in both *cox1* and 16S markers from the Mantel test analysis. Figures (c) and (d) illustrate the influence of geographic distance in the spatial autocorrelation analysis of *cox1* and 16S *I. ovatus* sequences. The upper and lower 95% confidence intervals are depicted as the red dotted lines (U and L) and the correlation coefficient *r* is plotted. The values above and below the dashed lines indicate spatial structure. The error bars result from the 1000 permutations of individuals within each distance class. Both analyses were done in Genalex software.

Figure 2. An unweighted pair group method with the arithmetic mean (UPGMA) dendrogram of *I. ovatus* based on the pairwise genetic distance (F_{ST}) of *cox1* among 29 sampling locations across Niigata Prefecture, Japan.

Figure 3. The distribution of the two genetic clusters as observed in the UPGMA cluster dendrogram (Figure 3) of *I. ovatus cox1* sequences from Niigata Prefecture.

Figure 4. Maximum likelihood gene tree of *cox1* haplotype sequences of *I. ovatus* with sequences from China (MH208681-MH208687, MH208689-MH2086393, MH208706), Japan (AB231670) and an outgroup of *Ixodes canisuga.*

Yellow reference sequences from China (Li et al., 2018); *red* reference sequence from Hokkaido, Japan (Mitani et al., 2007); *blue I. ovatus* divergent *cox1* haplotypes; *green I. ovatus cox1* remaining haplotypes; *purple I. canisuga* outgroup