Comparative population genetic structure of two ixodid tick species (*Ixodes ovatus* and

Haemaphysalis flava) in Niigata Prefecture, Japan

Maria Angenica F. Regilme[a,b], Megumi Sato[c], Tsutomu Tamura[d], Reiko Arai[d], Marcello Otake Sato[e],

Sumire Ikeda[f], Maribet Gamboa[a,b], Michael T. Monaghan[g,h], Kozo Watanabe[a, b]

[a] Center for Marine Environmental Studies (CMES), Ehime University, Matsuyama, Ehime, 790-8577,

Japan

[b] Graduate School of Science and Engineering, Ehime University, Matsuyama, Ehime, 790-8577,

Japan

[c] Graduate School of Health Sciences, Niigata University, Niigata, 951-8518, Japan

[d] Niigata Prefectural Institute of Public Health and Environmental Sciences, Niigata, 950-2144, Japan

[e] Department of Tropical Medicine and Parasitology, Dokkyo Medical University, 880 Kitakobayashi,

Mibu-machi, Shimotsuga-gun, Tochigi 321-0293, Japan

[f] Research Laboratories, Research and Development Headquarters, Earth Corporation, Hyogo 678-

0192, Japan

[g] Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin, 12587, Germany

[h] Institut für Biologie, Freie Universität Berlin, 14195, Germany

Corresponding author

watanabe.kozo.mj@ehime-u.ac.jp

Abstract

Ixodid tick species such as *Ixodes ovatus* and *Haemaphysalis flava* are essential vectors of

tick-borne diseases in Japan. In this study, we investigated the population genetic structures and

gene flow of *I. ovatus* and *H. flava* as affected by the tick host mobility. We hypothesized that *I.*

ovatus and *H. flava* may have differences in their genetic structure due to the low mobility of small

rodent hosts of *I. ovatus* at the immature stage in contrast to the mediated dispersal of avian hosts

for immature *H. flava.* We collected 307 adult *I. ovatus* and 220 adult *H. flava* from 29 and 17

locations across Niigata Prefecture, Japan. We investigated the genetic structure at two

mitochondrial loci (*cox1*, 16S rRNA gene). For *I. ovatus*, pairwise F_{ST} and analysis of molecular

variance (AMOVA) analyses of *cox1* sequences indicated significant genetic variation among

populations. Both *cox1* and 16S rRNA markers showed non-significant genetic variation among

locations for *H. flava*. The Bayesian tree and haplotype network of *cox1* marker for *I. ovatus* samples

in Niigata Prefecture found 3 genetic groups wherein most haplotypes in group 2 were distributed

in low altitudinal areas. When we added *cox1* sequences of *I. ovatus* from China to the phylogenetic

analysis, three genetic groups (China 1, China 2, and Niigata and Hokkaido, Japan) were formed in

the tree suggesting the potential for cryptic species in the genetic group in Japan. Our results

support our hypothesis and suggest that the host preference of ticks at the immature stage may

influence the genetic structure and gene flow of the ticks. This information is vital in understanding

the tick-host interactions in the field to better understand the tick-borne disease transmission and

in designing an effective tick control program.

Keywords: genetic structure, gene flow, host mobility, cryptic species, haplotypes

Introduction

Tick-borne diseases are a public health concern, and their control is often challenging because of the complex transmission chain of vertebrate hosts and ticks interacting in a changing environment (Dantas-Torres et al., 2012). Population genetic studies helped understand the dispersal patterns and potential pathogen transmission in ticks (Araya-Anchetta et al., 2015). Population genetic studies can elucidate the direction, distance, and gene flow patterns between tick populations (McCoy, 2008). For example, if a high gene flow is observed, there might be a greater chance of colonizing new areas or re-colonizing areas following successful vector control programs.

Due to the small size of ticks and their vulnerability to harsh environments during off-host, tick dispersal is complex and is linked to its host movement (Falco and Fish, 1991). Host mobility can affect the genetic patterns of tick populations, although its effects are not consistent. Several studies have reported low gene flow in ticks with less mobile hosts (e.g., smaller mammals) and a high gene flow in ticks with highly mobile hosts (Araya-Anchetta et al., 2015). For example, *Amblyomma americanum* and *Amblyomma triste* (Acari, Ixodidae) exhibited a high gene flow across various spatial scales (137,000 km^2 to 2.78 million km^2) due to their hosts' dispersal capabilities (large mammals and birds) (Guglielmone et al., 2013; Mixson et al., 2006; Trout et al., 2010). Lampo et al., (1998) observed low gene flow in *Amblyomma dissimile* populations due to its hosts' low mobility (small mammals, reptiles, and salamanders).

In Japan, tick-borne diseases are an increasing public health concern, affecting humans and animals (Yamaji et al., 2018). A total of 8 genera of ticks have been recorded in Japan, composed of 47 species: 43 ixodid tick species and four argasid species (Fujita et al., 2006). Out of these 47 species, 21 species parasitize humans (Okino et al., 2010). Among these 21 species, *Ixodes ovatus*, the

primary vector of the causative agents of Lyme borreliosis (Miyamoto et al., 1993), and

Haemaphysalis flava, a vector of the causative agents of severe fever with thrombocytopenia

syndrome (SFTS) and Japanese spotted fever (JSF), was reported in Japan (Yamaji et al., 2018; Yu et

al., 2014). Yamaguti et al., (1971) observed that the hosts of adult *I. ovatus* were mainly hares but

could also be large mammals (e.g., cows and horses) and that the hosts of immature forms were

small rodents. They also found that the host preferences of adult *H. flava* were cows, dogs, horses,

wild boar, bear, and deer, while the immatures parasitized birds. Comparative population genetic

studies on two essential ixodid ticks: *I. ovatus* and *H. flava* are not existing.

In addition to population genetic structure, genetic analysis may also reveal the

phylogenetic pattern and the presence of cryptic species, where morphologically identified

individuals might represent more than one species (Fegan et al., 2005). Previous studies on

Rhipicephalus appendiculatus (Kanduma et al., 2016), *Ixodes holocyclus* (Song et al., 2011)*, and I.*

ovatus (Li et al., 2018) have revealed the presence of cryptic species based on the clustering of

haplotypes in a phylogenetic tree. Based on these previous studies, morphological criteria for

species differentiation alone are equivocal and that genetic analysis is essential.

Here, we studied the population genetic structure of *I. ovatus* and *H. flava* and also tested

for the presence of cryptic species using mitochondrial DNA sequences of *cox1* and the 16S rRNA

markers. We hypothesized that *I. ovatus* and *H. flava* would display contrasting population genetic

structures despite the overlap sharing of large mammalian hosts at the ticks' adult stage. The low

mobility of *I. ovatus'* host*,* mainly hares at the adult tick stage and small mammals during the

immature stage may contribute to the ticks' high genetic divergence. On the other hand, the

combined high mobility of large mammalian host at *H. flava* adult stage and avian mediated

dispersal at the immature stage may be affecting the ticks' homogenized population genetic

structure.

Material and methods

Study site, collection, sampling, and identification

From April 2016 until November 2017, ticks were collected by the standard flagging

methods (Ginsberg et al., 1989) in 29 sites (Additional File 1. Table S1) across Niigata Prefecture in

Japan. Ticks were collected 2 to 14 times in 6 core sites among the 29 sites (Additional File 2. Table

S2). Altitude per location ranged from 8 from 1402 m/a.s.l (mean=350). The geographic distance

between sites ranged from 4.28 to 247.65 km (mean=77.36). Collected ticks were stored in

microcentrifuge tubes with 70% ethanol at 4^0C. We identified the sex, developmental stage, and

species using a compound microscope and identification keys of Yamaguti et al., (1971).

DNA extraction, PCR amplification, and sequencing

Genomic DNA (*I. ovatus* n=320*; H. flava* n=220) from each identified adult tick individual

was extracted using the Isogenome DNA extraction kits (NIPPON GENE Co. Ltd., Tokyo, Japan)

following the manufacturer's recommended protocol. The total number of analyzed samples

(n=540) was chosen based on its species identity, while the other identified tick species were

excluded from this study. Before DNA extraction, each tick was washed with alcohol and a PBS

solution. DNA   concentration   and   quality   were   checked   using   a   NanoDrop™   2000

Spectrophotometer (Thermo Scientific™). Two mitochondrial genes were amplified by polymerase

chain reaction (PCR): Cytochrome Oxidase 1 (*cox1*) (658 base pairs) using the primer pairs LCO-1490

(5'- GGTCAACAAATCATAAAGATATTGG -3') and HCO1-2198 (5'- AAACTTCAGGGTGACCAAAAAATCA-

3') (Folmer et al., 1994) and the 16S rRNA gene (16S) (407 base pairs) with the following primer pairs

16S+1 (5'CTGCTCAATGAATATTTAAATTGC-3') and 16S − 1 (5' -CGGTCTAAACTCAGATCATGTAGG-3')

(Tian et al., 2011). All PCR amplifications of both *cox1* and 16S were performed with a final volume

of 10 µl with 1 µl of genomic DNA. The PCR reaction for both markers was composed of the

following: 10x Ex Taq buffer, 25 mM MgCl$_2$, 2.5 mM dNTP, 10 µm of forward and reverse primers,

and five units/µl of TaKaRa Ex Taq™ (Takara Bio Inc.). The *cox1* PCR amplification was as follows: an

initial denaturation of 94^{0}C for 2 min, denaturation of 94^{0}C for 30 s, annealing of 38^{0}C for 30 s, an

extension of 72^{0}C for 1 min for 30 cycles, and a final extension of 72^{0}C for 10 min. In contrast, the

16S PCR amplification followed the protocol of (Tian et al., 2011) with some modifications, initial

denaturation of 94^{0}C for 3 min, denaturation of 94^{0}C for 30 s, annealing of 50^{0}C for 40 s, an extension

of 72^{0}C for 40 s for 30 cycles and final extension of 72^{0}C for 5 min. PCR products were purified using

the QIAquick 96 PCR Purification Kit (Qiagen) following the manufacturer's instructions and

sequenced by Eurofin Genomics, Inc., Tokyo, Japan.

Sequence data analysis

We assembled forward and reverse reads for each individual using CodonCode Aligner

version 1.2.4 software (https://www.codoncode.com/aligner/). No ambiguous bases were observed,

and the low-quality bases were removed at the start and end of the reads. Multiple alignments of

sequences were performed using the MAFFT alignment online program with default settings

(https://mafft.cbrc.jp/alignment/server/). To ensure sequence quality and correct species

identification, we checked the sequences' similarity against reference sequences from GenBank

using BLASTN (Basic Local Alignment Search Tool – Nucleotide,

https://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM=blastn&PAGE_TYPE=BlastSearch&LINK_LOC=bl

asthome). We checked the quality of the final aligned sequences (*cox*1 = 658bp; 16S = 407bp) in

Mesquite version 3.5 (Maddison et al., 2011) protein-coding genes were translated into amino acids

to confirm the absence of stop codons.

Population genetic analysis

We combined the nearby sites with a geographic distance range of 8.83 km to 79.81

(mean=44.00) with less than 8 individuals for the population genetic analysis because the accurate

estimation of allele frequencies is difficult for small size population. A total of 8 populations (A to H)

(Additional File 1 Table S1) were formed. Some populations were not included in the population

genetic analysis because of the small sample size (< 8 individuals) and can't be combined with any

nearby sites.

We analyzed the sequences of both *H. flava* and *I. ovatus* per genetic marker separately

using DNASp version 6.12.03 (Rozas et al., 2017) and Arlequin version 3.5.2.2 (Excoffier and Lischer,

2010) for the following parameters: number of haplotypes (nh), the average number of polymorphic

sites (s), and average number of nucleotide differences (k). The haplotype diversity (h) and

nucleotide diversity (π) were calculated in Arlequin version 3.5.2.2. The population genetic structure

among and within populations in Niigata Prefecture was assessed by analysis of molecular variance

(AMOVA) performed in Arlequin with 9999 permutations. The degree of molecular genetic

differentiation between populations was assessed by calculating the pairwise F_{ST} values using

Arlequin.

To determine if the genetic differentiation was influenced by geographical distance or

altitudinal differences among populations, we performed Mantel Test in GenAlEx version 6.51b2

(Peakall and Smouse, 2006). Two tests per species and marker were conducted. First, we compared

pairwise genetic (pairwise F_{ST} values) and geographical distances (km), and second, we analyzed the

genetic distance (F_{ST} values) with altitudinal differences (m/a.s.l.) calculated from GenAlEx version

6.51b2. The geographic distances were obtained from the geographic midpoint of the populations

using the GPS coordinates (latitude and longitude) of each site recorded during the sampling. The

altitudinal differences are the calculated mean value per population. All Mantel tests were assessed

using 9999 permutations for the significance of the correlation.

The genetic relationships among the populations were calculated by the unweighted pair

group method with the arithmetic mean (UPGMA) cluster method using the APE package (Paradis

and Schliep, 2018) and R program (R Development Core team., 2016). To create a dendrogram, we

used the genetic distance matrix (pairwise F_{ST} values) generated from GenAlEx.

Haplotype network and Phylogenetic analysis

To evaluate the haplotypes' relationship, we constructed a haplotype network on the

PopART program version 1.7 per marker (*cox1* and 16S) and species (*I. ovatus* and *H. flava*

(http://popart.otago.ac.nz/index.shtml) using the median-joining (MJ) network algorithm (Bandelth

et al., 1999).

We performed a Bayesian phylogenetic analysis using BEAST v1.10.4 (Drummond and

Rambaut, 2007) to determine the phylogenetic structure of *I. ovatus* haplotypes within Niigata

Prefecture using the *cox1* marker. Additional sequences from China were not included in the

Bayesian analysis. We used the HKY substitution model with the estimated base frequencies. The

strict clock model was employed and the speciation yule process as the tree prior. A maximum clade

credibility tree was acquired using the TreeAnnotator v1.10.4 from the many trees obtained from

BEAUti v1.10.4 with 10% of trees used during burn-in. The maximum clade credibility tree was

viewed using FigTree v1.4.4.

We constructed maximum likelihood (ML) gene trees for *cox1* and 16S haplotype sequences

of *I. ovatus* and *H. flava* using PhyML version 3.1 (Guindon and Gascuel, 2003) default settings. We

applied HKY and GTR nucleotide substitution models for *cox1* and 16S, respectively, as suggested by

jModelTest version 2 (Darriba et al., 2012). Additional sequences from China (MH208506,

MH208512, MH208514, MH208522, MH208515-19, MH208524, MH208531, MH208574,

MH208577, MH208579, MH208681-87, MH208689-93, MH208706, KU664519 (Li et al., 2018),

Japan (Hokkaido AB231670, U95900; Yamanashi AB819241, AB819243 and Aomori AB819244)

(Mitani et al., 2007; Norris et al., 1999; Takano et al., 2014) were included to know the presence of

cryptic species. We used *Ixodes canisuga* as an outgroup because it is closely related to *I. ovatus* and

H. flava (KY962023 and KY962074; Hornok et al., 2015).

Results

A total of 2,374 individual ticks was collected. Adult and immature *Ixodes nipponensis*,

Dermacantor taiwanensis, *Ixodes persulcatus*, and *Ixodes monospinus* were also identified and used

for another research study. In this study, adult *I. ovatus* and *H. flava* were only analyzed because

these two species were the most abundant. Larval ticks were used in another study. The number

of *I. ovatus* ranged from 1 to 36 adult individuals per site and were more successfully sequenced for

cox1 (307/320; 95.9%) than for 16S (284/320; 88.8%). The analyzed *H. flava* ranged from 1 to 77

adult individuals per site and were also more successfully sequenced for *cox1* (220/223; 98.7%) than

for 16S (172/223; 77.1%). On the other hand, for the population genetic analysis, each population

(i.e., combined nearby sites) had the following tick individual range and mean per markers: *cox1 I.*

ovatus (28 to 62; mean=43), 16S *I. ovatus* (24 to 66; mean=40), *cox1 H. flava* (8 to 81; mean=43) and

16S *H.flava* (8 to 76; mean=34). There were 60 and 63 *cox1* haplotypes and 24 and 40 16S haplotypes

in *I. ovatus* and *H. flava,* respectively (Table 1). Haplotype diversity (h) ranged from 0.442 to 0.852,

and nucleotide diversity (π) ranged from 0.001 to 0.006 for both markers and species.

AMOVA results revealed a high among-population divergence (61.99%) in *I. ovatus cox1*

sequences, whereas both *cox1* and 16S markers of *H. flava* indicated no significant genetic variation

among populations (Table 2). Significant global F_{ST} values (P<0.01) were observed for *I. ovatus* with

significant values 0.3801 in *cox1* and 0.0378 in 16S markers. Pairwise F_{ST} values (0.0963 to 0.6808)

of *cox1* marker for *I. ovatus* showed significant genetic differences between most pairs of

populations such as between populations B and D and populations F and D (Additional File 3 Table

S3). We also observed significant genetic differences in 16S *I. ovatus* sequences ranged from 0.0514

to 0.0949 (Additional File 4 Table S4). Pairwise F_{ST} values from the *cox1* marker of *H. flava* showed

no significant genetic differences except in few population pairs such as A and C, C and E and D and

E (Additional File 5 Table S5). No significant genetic differentiation was also observed in 16S marker

of *H. flava* except between A and C (Additional File 6 Table S6).

The Mantel test of *I. ovatus* showed no significant isolation by geographic distance in both

markers: (*cox1 r* = 0.108, *p* = 0.269; 16S *r* = 0.518, *p* = 0.065) (Additional File 7 Figure S1, a and b)

and no evidence for isolation by altitudinal difference (*cox1* r = -0.066, *p*=0.225; 1S r = -0.023, p =

0.577). Both Mantel tests were also not significant for *H. flava* (distance: *cox1* r= 0.444, p= 0.130;

16S r = 0.355, p= 0.189; altitude: r = 0.092, p = 0.30; 16S r = 0.217, p= 0.06) (Additional File 7 Figure

S1, c and d).

The UPGMA cluster dendrogram constructed from the pairwise F_{ST} values of *I. ovatus* for the

cox1 marker (Figure 2) revealed two genetic clusters among the 7 populations, where cluster 2

included populations from the more western sites. In contrast, cluster 1 (A, B, and F) mostly had

populations in the northern and southern parts of Niigata Prefecture (Figure 3) and are distributed

in mountainous areas with high elevation. We observed no evident genetic clustering on the

dendrogram of *I. ovatus* for 16S marker and those of *H. flava* for *cox1* and 16S makers, respectively

(Additional File 8 Figure S2; Additional File 9 Figure S3; Additional File 10 Figure S4).

The haplotype network with the reference sequences from China and the Bayesian tree of

the *I. ovatus cox1* haplotypes showed similar patterns (Figure 5 and 6) of 4 genetic groups. The

haplotype network of the *cox1* marker of *I. ovatus* also displayed the two distinct haplotypes (Hap

60 and Hap 59) (Figure 6; Additional File 11 Figure S5). These two haplotypes were found from

sampling site 6 (Pop A) and sampling site 26 (Pop 6). The Bayesian tree of *I. ovatus cox1* haplotype

sequences with reference sequences from China also displayed the 4 genetic groups (Additional File

12 Figure S6). The haplotype network of *I. ovatus* from 16S marker (Additional File 13 Figure S7), *H.*

flava in both cox1 (Additional File 14 Figure S8) and 16S markers (Additional File 15 Figure S9) did

not display any distinct haplotypes and genetic groups.

A putative *I. ovatus* species complex was identified in the phylogenetic tree based on three

distinct haplotype groups found with both *cox1* and 16S rRNA markers; 2 groups in China and one

in Japan. The *cox1* phylogenetic tree of 60 *I. ovatus* haplotypes indicated three groups: group 1 of

the published sequences from western China, one large group 2 containing the 58 haplotypes and

two divergent haplotypes (Hap 60 and Hap 59), and group 3 from west China (Figure 4). The

published haplotype from Hokkaido, Northern Japan (Mitani et al., 2007) occurred within our large

group, and the published sequences from western China were in a different group.

The *I. ovatus* haplotypes in the 16S phylogenetic tree (Additional File 16 Figure S10) were

grouped with the published haplotypes from Japan (Yamanashi and Aomori Prefectures; Takano et

al., 2014) and Hokkaido (Norris et al., 1999). The *cox1* haplotype sequences of *H. flava* phylogenetic

tree (Additional File 17 Figure S11) were similar to reference sequences from China KY021800 –

KY021807, KY021810 – KY021819 and KY003181 (Li, Z., Cheng, T. and Liu, G.; Unpublished results

from NCBI); JQ625688 – JQ625689, JF758632 and JQ737097 (Lu et al. 2013) and JG737097 (Gou, H.,

Guan, G., Yin, H. and Luo, J.; unpublished results from NCBI). The *H. flava* 16S haplotype sequences

(Additional File 18 Figure S12) showed high similarities with reference sequences from Japan

(Kagoshima, Fukui Aomori, Fukui, Yamanashi, Kagawa, and Ehime Prefectures (Takano et al., 2014)

and China KC844858 –KC844867 (Cheng et al., 2013); KX450280 –KX450282 (Zhang, Y., Cui, Y., Peng,

Y., Yan, Y., Wang, X. and Ning, C. Liu, Q., Zhang, Y. and Zhu, D.; unpublished results from NCBI);

MG696720 (Zheng, W., Chen, S. and Chen, H.; unpublished results from NCBI) and KP324926 (Liu,

Q., Zhang, Y. and Zhu, D.; unpublished results from NCBI) sequences from GenBank.

Discussion

Contrasting population genetic structures between *I. ovatus* and *H. flava*

Our results supported our hypothesis that *I. ovatus* may display high genetic divergence

because of its low host mobility in contrast to the homogenized population genetic structure in *H.*

flava due to the combined large mammalian hosts at the ticks' adult stage and avian mediated

dispersal during the ticks' immature stage. The significant F_{ST} estimates in *I. ovatus cox1* (0.3801)

and 16S (0.0378) revealed population differentiation as supported by the high among populations

percentage variation (61.99) in the AMOVA analysis. The high genetic divergence in *I. ovatus* can be

due to the low mobility of its hosts, for example during the ticks' adult stage wherein they utilized

mostly hares while small rodents at the immature stage of *I. ovatus.* The host preference of *I. ovatus*

may also contribute to the observed 4 genetic groups in the Bayesian tree and the haplotype

network analysis. The haplotypes from groups 2 and 3 were mostly from low altitudinal areas as

compared to the haplotypes from group 1 with ticks from the high altitudinal area such as in

mountain areas. We assume that the large mammalian hosts such as cows and horses of *I. ovatus*

have enabled the ticks to reach high mountain ranges from group 1. The distribution of the type of

hosts may also affect altitude affecting also the formation of groups 1, 2, and 3 in the *cox1 I.ovatus*

dendrogram.

On the other hand, the homogenized population genetic structure observed in *H. flava*

might be because of the combined high host mobility of large mammals at the ticks' adult stage and

the avian mediated dispersal at the immature stage. Twenty-eight species of birds were previously

reported as hosts of immature *H. flava* from Japan, mostly from the order Passeriformes (Yamauchi

and Takeno, 2000). Large mammals and birds may have expansive habitats ranges that may allow

high gene flow of *H. flava* between the locations in Niigata, as previously observed in *Amblyomma*

americanum populations (Mixson et al., 2006; Reichard et al., 2005; Trout et al., 2010) and *I. ricinus*

(Casati et al.,2008).

Several alternative factors such as tick behavior, biology, and ecology can also affect tick's

dispersal patterns. A previous study on two tick species: *Hyalomma rufipes* and *Amblyomma*

hebraeum, also displayed contrasting genetic patterns despite the overlap sharing of highly mobile

hosts such as birds (Cangi et al., 2013). This might be due to the species specific capacity of the

immature stages of ticks to survive even after host detachment in various habitat conditions

(Cumming, 1999; Estrada-Peña, 2015; Needham et al., 1986). Population genetic structure can also

be influenced by assortative mating (e.g., *I. ricinus*) wherein genetically similar individuals tend to

mate rather than random mating resulting in decreased genetic divergence (Kempf et al., 2009). Our

study does not have supporting data to test these alternative factors; thus, it is suggested in future

studies to analyze further these factors that could play a role in the tick's dispersal movement.

16S marker showed significant genetic differentiation of *I. ovatus* (pairwise F_{ST} =0.0514 to

0.0949) in few pairs of sampling locations as compared to the *cox1* marker, which may be due to

lower variability of the marker thus providing inadequate genetic variation for population genetic

analysis, as previously observed in *Amblyomma ovale* (Bitencourth et al., 2019) and *Rhipicephalus*

microplus (Burger et al., 2014; Low et al., 2015). The low variability of 16S marker in *I. ovatus* may

not be sufficient to infer an intraspecific relationship and is further supported by its low (*nd* = 0.001)

nucleotide diversity as compared to *cox1* marker (*nd* = 0.004)*.* Thus, it may also explain the grouping

observed in the *cox1* phylogenetic tree of *I. ovatus* was absent in the 16S tree. Despite these, we

used the 16S marker because this gene mutates at a rate that is informative for inter-species level

phylogenetic analysis (Araya-Anchetta et al., 2015). The negative pairwise F_{ST} values observed in *H.*

flava do not have any biological meaning. These results from the unbiased estimator allowing a

negative F_{ST} when the sample size is small. (Bitencourth et al., 2016; Vial et al., 2006; Weir and

Cockerham, 1984) .

Species complex formation in *I. ovatus cox1* sequences

Our *cox1 I. ovatus* gene tree showed Japanese individuals to form a distinct group from

haplotypes from southern China (Li et al., 2018). Despite the low gene flow of *I. ovatus* in *cox1*

marker, we found that Niigata's haplotypes were near related to the published sequence from

Hokkaido in northern Japan. The result indicates that these ticks originated from a diverse set of

geographical locations in Japan, which might be transported by its hosts or are undergoing recent

population expansion from northern Japan (Hokkaido) to central Japan (Niigata) *vice versa*. We

found three groups (China 1, Japan, and China 2) and two slightly divergent *cox1* haplotypes (Hap

60 and 59) in the Japan group of *I. ovatus* in Niigata. Considering two or more cryptic species can be

concealed in one morphologically described species (Bitencourth et al., 2016), the occurrence of the

three groups and the divergent haplotypes suggests that *I. ovatus* may be a species complex. It can

be inferred that China 1 and China 2 might have a longer evolutionary time as compared to Japan.

The co-existence of a species in the same geographic area may explain the occurrence of the species

complex. One of the limitations of this study is that both Chinese and Japanese individuals were

collected in a limited geographical area. We suggest that future studies sample and sequence more

individuals from other locations. Extensive geographical sampling can explain better the taxonomy

and the genetic structure of species as suggested in the study of Liu et al., (2013) on *Rhipicephalus*

sanguineus. Previous studies have also observed species complexes in *Ixodes* and *Rhipicephalus*

(Burger et al., 2014; Li et al., 2018; Song et al., 2011; Xu et al., 2003), suggesting that morphological

criteria for species differentiation alone are equivocal and that genetic analysis is essential. Future

studies must tackle an integrative approach of the tick species concept that includes the morphology,

genetics, biology, and ecological trails of ticks (Dantas-Torres et al., 2012).

Conclusions

In summary, our findings revealed the contrasting patterns of the genetic structure of *I.*

ovatus and *H. flava* from Niigata Prefecture, Japan. The high genetic divergence in *I. ovatus* might

be influenced by the restricted movement of its small mammalian hosts during the ticks' immature

stage and hares at the adult stage. On the other hand, the homogenized genetic structure in *H. flava*

might be due to the avian mediated movement during the ticks' immature stage coupled with the

expansive movement of the large mammalian host at the ticks' adult stage. Our results suggest that

the host preference of ticks during its immature stage may strongly influence the population genetic

structure of ticks due to its higher ability to survive in an environment. Understanding the

population genetic structure of ticks such as *I. ovatus* and *H. flava* can give information about the

distribution and control of tick-borne diseases. Even though *I. ovatus* populations were genetically

structured within Niigata, a published haplotype from Hokkaido was also found, indicating that

widespread dispersal is possible. The occurrence of three groups and the divergent *cox1* haplotypes

in *I. ovatus* emphasizes the need for additional methods in determining the existence of a species

complex in *I. ovatus* populations in Japan.

List of Abbreviations

H. flava – Haemaphysalis flava

I. ovatus – Ixodes ovatus

AMOVA – analysis of molecular variance

UPGMA - unweighted pair group method with arithmetic mean

ML – maximum likelihood

F_{ST} – fixation index

PBS – phosphate-buffered saline

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All data generated or analyzed during the study are included in this published article and its

additional supplementary files. All the newly generated sequences are available in the GenBank

database under the accession numbers MW063669-MW064124 and MW065821 - MW066347.

Declarations of interests

The authors declare that they have no competing interests.

Authors' contributions

MAFR, MS, and KW conceptualized and designed the experiment. MS, TT, RA, SI, and MOS designed

the sampling collection, collected and identified the tick samples. MAFR and MG conducted the

molecular analyses. MAFR, MM, and KW performed the data analysis. MAFR and KW wrote the

manuscript. All authors read and approve the manuscript.

Acknowledgments

The authors would like to thank the Niigata Prefectural Office for their valuable help during the tick

sampling collection. We are also thankful to Masaya Doi, Kohki Tanaka, and Mizuki Ueda for their

technical assistance in the molecular analyses. Thank you to Dr. Thaddeus M. Carvajal and Dr.

Joeselle M. Serrana for their useful suggestions in the population genetic analyses in this study and

Micanaldo Francisco for technical assistance in constructing this study's sampling map.

Funding

This work is funded by the Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for Scientific Research (16K00569, 19K21996, 19H02276), and the Sumitomo Electric Industries Group Corporate Social Responsibility Foundation. The research was partially supported by the German Academic Exchange Service (DAAD, Programm Projektbezogener Personenaustausch Japan 2018, project 57402018).

References

Araya-Anchetta, A., Busch, J.D., Scoles, G.A., Wagner, D.M., 2015. Thirty years of tick population genetics: a comprehensive review. Infect. Genet. Evol. 29, 164–179. https://doi.org/10.1016/j.meegid.2014.11.008.

Bandelt, H.J., Forster, P., and Röhl, A., 1999. Median-joining networks for inferring intraspecific phylogenies. Mol. bio. and evo. 16, 37-48. https://doi.org/10.1093/oxfordjournals.molbev.a026036.

Bitencourth, K., Voloch, C.M., Serra-Freire, N.M., Machado-Ferreira, E., Amorim, M., Gazêta, G.S., 2016. Analysis of *Amblyomma sculptum* haplotypes in an area endemic for Brazilian spotted fever. Med. Vet. Entomol. 30, 342–350. https://doi.org/10.1111/mve.12174.

Bitencourth, K., Amorim, M., de Oliveira, S. V., Voloch, C.M., Gazêta, G.S., 2019. Genetic diversity, population structure and rickettsias in *Amblyomma ovale* in areas of epidemiological interest for spotted fever in Brazil. Med. Vet. Entomol. 33, 256–268. https://doi.org/10.1111/mve.12363.

Burger, T.D., Shao, R., Barker, S.C., 2014. Phylogenetic analysis of mitochondrial genome sequences indicates that the cattle tick, *Rhipicephalus* (*Boophilus*) *microplus*, contains a cryptic species. Mol. Phylogenet. Evol. 76, 241-253. https://doi.org/10.1016/j.ympev.2014.03.017.

Cangi, N., Horak, I.G., Apanaskevich, D.A., Matthee, S., das Neves, L.C.B.G., Estrada-Peña, A.,

Matthee, C.A., 2013. The influence of interspecific competition and host reference on the

phylogeography of two African ixodid tick species. PLoS One 8, 1–12.

https://doi.org/10.1371/journal.pone.0076930.

Casati, S., Bernasconi, M. V., Gern, L., Piffaretti, J.C., 2008. Assessment of intraspecific mtDNA

variability of European *Ixodes ricinus* sensu stricto (Acari: Ixodidae). Infect. Genet. Evol. 8,

152–158. https://doi.org/10.1016/j.meegid.2007.11.007.

Cheng, W.Y., Zhao, G.H., Jia, Y.Q., Bian, Q.Q., Du, S.Z., Fang, Y.Q., Qi, M.Z. and Yu, S.K., 2013.

Characterization of *Haemaphysalis flava* (Acari: Ixodidae) from Qingling subspecies of giant

panda (*Ailuropoda melanoleuca qinlingensis*) in Qinling Mountains (Central China) by

morphology and molecular markers. PLoS One. 8,7.

https://doi.org/10.1371/journal.pone.0069793.

Cumming, G.S., 1999. Host distributions do not limit the species ranges of most African ticks (Acari:

Ixodida). Bull. Entomol. Res. 89, 303–327. https://doi.org/10.1017/s0007485399000450.

Dantas-Torres, F., Chomel, B.B., Otranto, D., 2012. Ticks and tick-borne diseases: a one health

perspective. Trends Parasitol. 28, 437–446. https://doi.org/10.1016/j.pt.2012.07.003.

Darriba, D., Taboada, G.L., Doallo, R., Posada, D., 2012. JModelTest 2: more models, new heuristics

and parallel computing. Nat. Methods 9, 772. https://doi.org/10.1038/nmeth.2109.

Drummond AJ, Rambaut A. BEAST: Bayesian evolutionary analysis by sampling trees. BMC

evolutionary biology. 2007 Dec;7(1):1-8.

Estrada-Peña, A., 2015. Ticks as vectors: taxonomy, biology, and ecology. Rev. Sci. Tech. 34, 53–65.

https://doi.org/10.20506/rst.34.1.2345.

Excoffier, L., Lischer, H.E.L., 2010. Arlequin suite ver 3.5: a new series of programs to perform

population genetics analyses under linux and windows. Mol. Ecol. Resour. 10, 564–567.

https://doi.org/10.1111/j.1755-0998.2010.02847.

Falco RC, Fish D. Horizontal movement of adult Ixodes dammini (Acari: Ixodidae) attracted to CO2-

baited traps. Journal of Medical Entomology. 1991 Sep 1;28(5):726-9.

Fegan, M., Prior, P., 2005. How complex is the *Ralstonia solanacearum* species complex. Bacterial

wilt disease and the *Ralstonia solanacearum* species complex. APS Press., Minnesota, pp.

449-461.

Folmer, O., Black, M., Hoeh, W, Lutz, R., and Vrijenhoek, R.,1994. DNA primers for amplification of

mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Mol.

Mar. Bio. and Biotech.. 3, 294-299.

Fujita, H., Fournier, P.E., Takada, N., Saito, T., Raoult, D., 2006. *Rickettsia asiatica* sp. nov., isolated

in Japan. Int. J. Syst. Evol. Microbiol. 56, 2365–2368. https://doi.org/10.1099/ijs.0.64177-0.

Ginsberg, H.S., Ewing, C.P,.1989. Comparison of flagging, walking, trapping, and collecting from

hosts as sampling methods for northern deer ticks, *Ixodes dammini*, and lone-star ticks,

Amblyomma americanum (Acari: Ixodidae). Exp. Appl. Acarol.7, 313-322.

https://doi.org/10.1007/bf01197925.

Guglielmone, A.A., Nava, S., Mastropaolo, M., Mangold, A.J., 2013. Distribution and genetic

variation of *Amblyomma triste* (Acari: Ixodidae) in Argentina. Ticks Tick. Borne. Dis. 4, 386–

390. https://doi.org/10.1016/j.ttbdis.2013.01.009.

Guindon, S., Gascuel, O., 2003. A simple, fast, and accurate algorithm to estimate large

phylogenies by maximum likelihood. Syst. Biol. 52, 696–704.

https://doi.org/10.1080/10635150390235520.

Hornok, S., Kontschán, J., Estrada-Peña, A., De Mera, I.G.F., Tomanović, S., De La Fuente, J., 2015.

Contributions to the morphology and phylogeny of the newly discovered bat tick species,

Ixodes ariadnae in comparison with *I. vespertilionis* and *I. simplex.* Parasites Vectors. 8, 1–7.

https://doi.org/10.1186/s13071-015-0665-0.

Kanduma, E.G., Mwacharo, J.M., Githaka, N.W., Kinyanjui, P.W., Njuguna, J.N., Kamau, L.M.,

Kariuki, E., Mwaura, S., Skilton, R.A., Bishop, R.P., 2016. Analyses of mitochondrial genes

reveal two sympatric but genetically divergent lineages of *Rhipicephalus appendiculatus* in

Kenya. Parasites Vectors.9,353. https://doi.org/10.1186/s13071-016-1631-1.

Kempf, F., de Meeûs, T., Arnathau, C., Degeilh, B., McCoy, K.D., 2009. Assortative pairing in *Ixodes*

ricinus (Acari: Ixodidae), the European vector of Lyme borreliosis. J. Med. Entomol. 46, 471–

474. https://doi.org/10.1603/033.046.0309.

Lampo, M., Rangel, Y., Matat, A., 2015. Population genetic structure of a three-host population

genetic in *Amblyommma dissimile*. J. Parasitol. .84, 1137–1142.

https://doi.org/10.2307/3284662.

Li, L.H., Zhang, Y., Wang, J.Z., Li, X.S., Yin, S.Q., Zhu, D., Xue, J.B., Li, S.G., 2018. High genetic

diversity in hard ticks from a China-Myanmar border county. Parasites Vectors. 11, 1–11.

https://doi.org/10.1186/s13071-018-3048-5.

Liu, G.H., Chen, F., Chen, Y.Z., Song, H.Q., Lin, R.Q., Zhou, D.H., and Zhu, X.Q., 2013. Complete

mitochondrial genome sequence data provides genetic evidence that the brown dog tick

Rhipicephalus sanguineus (Acari: Ixodidae) represents a species complex. Int J Biol Sci. 9,361-

367. https://doi.org/10.7150/ijbs.6081.

Low, V.L., Tay, S.T., Kho, K.L., Koh, F.X., Tan, T.K., Lim, Y.A., Ong, B.L., Panchadcharam, C., Norma-

Rashid, Y., Sofian-Azirun.M., 2015. Molecular characterisation of the tick *Rhipicephalus*

microplus in Malaysia: new insights into the cryptic diversity and distinct genetic assemblages

throughout the world. Parasites Vectors. 8, 341. https://doi.org/10.1186/s13071-015-0956-5.

Lu, X., Lin, X.D., Wang, J.B., Qin, X.C., Tian, J.H., Guo, W.P., Fan, F.N., Shao, R., Xu, J. and Zhang,

Y.Z., 2013. Molecular survey of hard ticks in endemic areas of tick-borne diseases in China.

Ticks Tick Borne Dis. 4,288-296. https://doi.org/10.1016/j.ttbdis.2013.01.003.

Maddison, W.P., Maddison, D.R.,2011. Mesquite: a modular system for evolutionary analysis.

2011. See http://mesquiteproject. org.

McCoy, K., 2008. The population genetic structure of vectors and our understanding of disease

epidemiology. Parasite. 15, 444–8. https://doi.org /10.1051/parasite/2008153444.

Mitani, H., Takahashi, M., Masuyama, M., Fukunaga, M., 2007. *Ixodes philipi* (Acari: Ixodidae):

phylogenetic status inferred from mitochondrial cytochrome oxidase subunit I gene

sequence comparison. J. Parasitol. 1, 719-722. https://doi.org /10.1645/GE-953R.1.

Mixson, T.R., Lydy, S.L., Dasch, G.A., Real, L.A., 2006. Inferring the population structure and

demographic history of the tick, *Amblyomma americanum* Linnaeus. J. Vector Ecol. 31, 181–

192. https://doi.org/10.3376/1081-1710(2006)31[181:itpsad]2.0.co;2.

Miyamoto, K., Nakao, M., Fujita, H., Sato, F., 1993.The ixodid ticks on migratory birds in Japan and

the isolation of Lyme disease spirochetes from bird-feeding ticks. Med. entomol. and zoo. 44,

315-326. https://doi.org/10.7601/mez.44.315.

Needham, GR, Teel, PD., 1986. Water balance by ticks between bloodmeals. Morphology,

physiology, and behavioral biology of ticks/editors, Ellis Horwood Limited, Chichester, UK. pp.

100-151.

Norris, D.E., Klompen, J.S.H., Black IV, W.C., 1999. Comparison of the mitochondrial 12S and 16S

ribosomal DNA genes in resolving phylogenetic relationships among hard ticks (Acari:

Ixodidae). Ann. Entomol. Soc. Am. 92, 117–129. https://doi.org/10.1093/aesa/92.1.117.

Okino, T., Ushirogawa, H., Matoba, K., Hatsushika, R., 2010. Bibliographical studies on human

cases of hard tick (Acarina: Ixodidae) bites in Japan (7) cases of unidentified tick infestation.

Kawasaki Medical Journal. 36, 127-141.

Paradis, E., Schliep, K.,2018. APE 5.0: an environment for modern phylogenetics and evolutionary

analyses in R. Bioinformatics. 35, 526-528. https://doi.org/10.1093/bioinformatics/bty633

Peakall, R., Smouse, P.E., 2006. GENALEX 6: genetic analysis in excel. population genetic software

for teaching and research. Mol. Ecol. Notes 6, 288–295. https://doi.org/10.1111/j.1471-

8286.2005.01155.x.

R Development Core Team., 2016. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

Reichard, M.V., Kocan, A.A., Van Den Bussche, A., Barker, R.W., Wyckoff, J., Ewing, S.A., 2005.

Sequence variation of the ribosomal DNA second internal transcribed spacer region in two

spatially distinct populations of *Amblyomma americanum* (L.)(Acari: Ixodidae). J. Parasitol.91,

260-264. https://doi.org/10.1645/GE-292R1.

Rozas,J., Ferrer-Mata,A., Sánchez-DelBarrio,J.C., Guirao-Rico,S., Librado,P., Ramos-Onsins,S.E.,

Sánchez-Gracia,A.,2017. DnaSP 6: DNA sequence polymorphism analysis of large data sets.

Mol. bio.and evo.34, 3299-3302. https://doi.org/10.1093/molbev/msx248.

Song, S., Shao, R., Atwell, R., Barker, S., Vankan, D., 2011. Phylogenetic and phylogeographic

relationships in *Ixodes holocyclus* and *Ixodes cornuatus* (Acari: Ixodidae) inferred from *cox1*

and ITS2 sequences. Int. J. Parasitol. 41, 871–880.

https://doi.org/10.1016/j.ijpara.2011.03.008.

Takano, A., Fujita, H., Kadosaka, T., Takahashi, M., Yamauchi, T., Ishiguro, F., Takada, N., Yano, Y.,

Oikawa, Y., Honda, T., Gokuden, M., Tsunoda, T., Tsurumi, M., Ando, S., Andoh, M., Sato, K.,

Kawabata, H., 2014. Construction of a DNA database for ticks collected in Japan: application

of molecular identification based on the mitochondrial 16S rDNA gene. Med. Entomol. Zool.

65, 13–21. https://doi.org/10.7601/mez.65.13.

Teglas, M.B., Drazenovich, N.L., Stott, J., Foley, J.E., 2006. The geographic distribution of the

putative agent of epizootic bovine abortion in the tick vector, *Ornithodoros coriaceus*. Vet.

Parasitol. 140, 327–333. https://doi.org/10.1016/j.vetpar.2006.03.027.

Tian, Z., Liu, G., Xie, J., Yin, H., Luo, Jianxun, Zhang, L., Zhang, P., Luo, Jin, 2011. Discrimination

between *Haemaphysalis longicornis* and *H. qinghaiensis* based on the partial 16S rDNA and

the second internal transcribed spacer (ITS-2). Exp. Appl. Acarol. 54, 165–172.

https://doi.org/10.1007/s10493-010-9423-3.

Trout, R.T., Steelman, C.D., Szalanski, A.L., 2010. Population genetics of *Amblyomma americanum*

(Acari: Ixodidae) collected from Arkansas. J. Med. Entomol. 47, 152–161.

https://doi.org/10.1603/me09106.

Vial, L., Durand, P., Arnathau, C., Halos, L., Diatta, G., Trape, J.F., Renaud, F., 2006. Molecular

divergences of the *Ornithodoros sonrai* soft tick species, a vector of human relapsing fever in

West Africa. Microbes Infect. 8, 2605–2611. https://doi.org/10.1016/j.micinf.2006.07.012.

Weir, B.S., Cockerham, C.C., 1984. Estimating F-statistics for the analysis of population structure.

Evolution. 38, 1358–1370. https://doi.org/10.2307/2408641.

525 Xu, G., Fang, Q.Q., Keirans, J.E., Durden, L.A., 2003. Molecular phylogenetic analyses indicate that

526 the *Ixodes ricinus* complex is a paraphyletic group. J. Parasitol. 1, 452-457.

527 https://doi.org/10.1645/0022-3395(2003)089[0452:MPAITT]2.0.CO;2.

528 Yamaguti, N., Tipton, V.J., Keegan, H.L., Toshioka, S., 1971. Ticks for Japan, Korea, and the Ryukyu

529 islands. Brigham Young Univ. Sci. Bull. XV, 226.

530 Yamaji, K., Aonuma, H., Kanuka, H., 2018. Distribution of tick-borne diseases in Japan: past

531 patterns and implications for the future. J. Infect. Chemother. 24, 499–504.

532 https://doi.org/10.1016/j.jiac.2018.03.012.

533 Yamauchi, T., Takeno, K., 2000. Record of *Haemaphysalis flava* Neumann (Acari, Ixodidae) from

534 *Zoothera dauma* (Latham) (Aves) in Kyushu, Japan. Esakia 40, 99–101.

535 Yu, X.J., Liang, M.F., Zhang, S.Y., Liu, Y., Li, J.D., Sun, Y.L., Zhang, L., Zhang, Q.F., Popov, V.L., Li, C.,

536 Qu, J., 2014. Fever with thrombocytopenia associated with a novel bunyavirus in China. New

537 Eng. Jour. of Med. 364.1523-1532. https://doi.org/10.1056/NEJMoa1010095.

 Tables

542 **Table 1.** Summary of *cox1* and 16S haplotype, variability, genetic diversity of adult *I. ovatus* and

543 adult *H. flava* populations in Niigata Prefecture, Japan

Marker	Species	n	nh	s	k	h	π
cox1	*I. ovatus*	307	60	65	2.728	0.852	0.004
	H. flava	220	63	60	1.472	0.789	0.002
16S	*I. ovatus*	284	24	22	0.699	0.442	0.001
	H. flava	172	40	49	2.447	0.835	0.006

545 Abbreviations: n sample size; nh number of haplotypes; s number of polymorphic sites; k mean

546 number of nucleotide differences; h haplotype diversity; π nucleotide diversity

547 **Table 2.** Analysis of molecular variance (AMOVA) using *cox1* and 16S of adult *I. ovatus* and adult *H.*

548 *flava* populations

Marker	Species		df	ss	vc	pv	F_{ST}
cox1	*I. ovatus*	Among populations	6	144.45	0.5562 Va	38.01	0.3801*
		Within populations	229	262.17	0.9071 Vb	61.99	
	H. flava	Among populations	4	4.9	0.0135 Va	1.81	0.0181
		Within populations	213	156.12	0.7321 Vb	98.19	
16S	*I. ovatus*	Among populations	6	5.12	0.0133 Va	3.78	0.0378*
		Within populations	272	91.83	0.3376 Vb	96.22	

	Among populations	4	5.74	0.0092 Va	0.78	
H. flava						0.0079
	Within populations	164	191.31	1.1665 Vb	99.22	

549 Abbreviations: df degrees of freedom; ss sum of squares; Va, Vb and Vc are associate covariance

550 components; pv percentage variation; * P<0.01

551 **Figure 1.** Map of the 29 sampling sites used for this study. The populations labeled A to H are

552 composed of the nearby sites labeled 1 to 29 used for the population genetic analysis.

563 **Figure 2.** An unweighted pair group method with the arithmetic mean (UPGMA) dendrogram of

564 adult *I. ovatus* based on the pairwise genetic distance (F_{ST}) of *cox1* among the 7 populations across

565 Niigata Prefecture, Japan.

Figure 3. The distribution of the two genetic clusters as observed in the UPGMA cluster dendrogram

(Figure 2) of *I. ovatus cox1* sequences from Niigata Prefecture.

Figure 4. Maximum likelihood gene tree of *cox1* haplotype sequences of *I. ovatus* with sequences

from China (MH208681-MH208687, MH208689-MH2086393, MH208706), Japan (AB231670), and

an outgroup of *I. canisuga.*

Yellow reference sequences from China (Li et al., 2018); *red* reference sequence from Hokkaido,

Japan (Mitani et al., 2007); *blue I. ovatus* divergent *cox1* haplotypes; *green I. ovatus cox1*

remaining haplotypes; *purple I. canisuga* outgroup

Figure 5. Phylogenetic tree from Bayesian analysis of the *cox1 I. ovatus* haplotype sequences

586   **Figure 6.** Median-joining haplotype network of *I. ovatus* haplotypes with reference sequences

587   from China using the *cox1* marker. Each circle represents a unique haplotype and the lines

588   correspond to mutations.

590   For easier visualization, the mutation steps (n=70) were marked as ⭐ .

Additional Files

Additional File 1.xls Table S1. Summary of adult *Haemaphysalis flava* and *Ixodes ovatus* collected from the different locations of Niigata Prefecture and its corresponding sample number

Additional File 2.xls Table S2. Summary of tick collection per species, life stage, and locations

Additional File 3.xls Table S3. Pairwise comparison of genetic differentiation (F_{ST}) calculated for all *I. ovatus* using the *cox1* gene

Additional File 4.xls Table S4. Pairwise comparison of genetic differentiation (F_{ST}) calculated for all *I. ovatus* using the 16S gene

Additional File 5.xls Table S5. Pairwise comparison of genetic differentiation (F_{ST}) calculated for all *H. flava* using the *cox1* gene

Additional File 6.xls Table S6. Pairwise comparison of genetic differentiation (F_{ST}) calculated for all *H. flava* using the 16S gene

Additional File 7.docx Figure S1. Plots a to d show the relationship between the pairwise F_{ST} values and the geographic distances for *I. ovatus* and *H. flava* in both *cox1* and 16S markers from the Mantel test analysis.

Additional File 8.docx Figure S2. An unweighted pair group method with the arithmetic mean (UPGMA) dendrogram of *I. ovatus* based on the pairwise genetic distance (F_{ST}) of 16S among the 7 populations across Niigata Prefecture, Japan.

Additional File 9.docx Figure S3 An unweighted pair group method with the arithmetic mean (UPGMA) dendrogram of *H. flava* based on the pairwise genetic distance (F_{ST}) of *cox1* among the 5 populations across Niigata Prefecture, Japan.

Additional File 10.docx Figure S4 An unweighted pair group method with the arithmetic mean (UPGMA) dendrogram of *H. flava* based on the pairwise genetic distance (F_{ST}) of 16S among the 5 populations across Niigata Prefecture, Japan.

Additional File 11.docx Figure S5 Median-joining haplotype network of *I. ovatus* from *cox1*

marker. Each circle represents a unique haplotype and the lines correspond to mutations.

Additional File 12.docx Figure S6 Phylogenetic tree from Bayesian analysis of the *cox1 I. ovatus*

haplotype sequences with reference sequences from China

Additional File 13.docx Figure S7 Median-joining haplotype network of *I. ovatus* from 16S marker.

Each circle represents a unique haplotype and the lines correspond to mutations.

Additional File 14.docx Figure S8 Median-joining haplotype network of *H. flava* from *cox1* marker.

Each circle represents a unique haplotype and the lines correspond to mutations.

Additional File 15.docx Figure S9 Median-joining haplotype network of *H. flava* from 16S marker.

Each circle represents a unique haplotype, and the lines correspond to mutations.

Additional File 16.docx Figure S10 Phylogenetic tree of 16S haplotype sequences of *I. ovatus*

inferred by maximum likelihood analysis with evolutionary model GTR with sequences from China

(MH208506, MH208512, MH208514 – MH208519, MH208522, MH208524, MH208531,

MH208574, MH208577, MH208579, KU664519), Japan (AB819241, AB819242, AB819243,

AB819244, and U95900). An outgroup (*Ixodes canisuga*) was also included.

Additional File 17.docx Figure S11 Phylogenetic tree of *cox1* haplotype sequences of *H. flava*

inferred by maximum likelihood analysis with evolutionary model HKY with sequences from China

(KY021800- KY021807; KY021810-KY021819, KY003181, JQ62588 - JQ625889; JQ737097;

JF758632). An outgroup (*Ixodes canisuga)* was also included.

Additional File 18.docx Figure S12 Phylogenetic tree of 16S haplotype sequences of *H. flava*

inferred by maximum likelihood analysis with evolutionary model GTR with sequences from China

(KC844858 -KC844867; KC844880-KC844882, M696720, KP324926, KX450279) and Japan

(AB19177- AB819192). An outgroup (*Ixodes canisuga)* was also included.