Title

Comparative population genetic structure of two Ixodidae ticks (*Ixodes ovatus* and *Haemaphysalis*

flava) in Niigata Prefecture, Japan

Author names and affiliations

Maria Angenica Fulo Regilme[a], Megumi Sato[b], Tsutomu Tamura[c], Reiko Arai[c], Marcello Otake Sato[d],

Sumire Ikeda[e], Masaya Doi[a], Kohki Tanaka[a], Maribet Gamboa[a], Michael T. Monaghan[f,g], Kozo

Watanabe[a, h]

[a] Department of Civil and Environmental Engineering, Ehime University, Matsuyama, Ehime, 790-

8577, Japan

[b] Graduate School of Health Sciences, Niigata University, Niigata, 951-8518, Japan

[c] Niigata Prefectural Institute of Public Health and Environmental Sciences, Niigata, 950-2144, Japan

[d] Department of Tropical Medicine and Parasitology, Dokkyo Medical University, 880 Kitakobayashi,

Mibu-machi, Shimotsuga-gun, Tochigi 321-0293, Japan

[e] Research Laboratories, Research and Development Headquarters, Earth Corporation, Hyogo 678-

0192, Japan

[f] Leibniz-Institute of Freshwater Ecology and Inland Fisheries (IGB), Berlin, 12587, Germany

[g] Institut für Biologie, Freie Universität Berlin, 14195, Germany

[h] Center for Marine Environmental Studies (CMES), Ehime University, Matsuyama, Ehime, 790-8577,

Japan

Corresponding author

watanabe.kozo.mj@ehime-u.ac.jp

Abstract

Ixodid tick species such as *Ixodes ovatus* and *Haemaphysalis flava* are important vector of tick-borne

diseases in Japan. In this study, we used genetic structure at two mitochondrial loci (*cox1*, 16S rRNA

gene) to infer gene flow patterns of *I. ovatus* and *H. flava* from Niigata Prefecture, Japan. Samples

were collected in 29 (*I. ovatus*) and 17 (*H. flava*) sampling locations across Niigata Prefecture

(12,584.18 km²). For *I. ovatus*, pairwise F_{ST} and analysis of molecular variance (AMOVA) analyses of

cox1 sequences indicated significant among-population differentiation. This was in contrast to *H.*

flava, for which there were few cases of low significant pairwise differentiation. A Mantel test

revealed isolation by distance and there was positive spatial autocorrelation of haplotypes in *I.*

ovatus cox1 and 16S sequences, but non-significant results were observed in *H. flava* in both

markers. We found three genetic groups (China 1, China 2 and Japan) in the *cox1 I. ovatus* tree.

Newly sampled *I. ovatus* grouped together with a published *I. ovatus* sequence from northern Japan

and were distinct from two other *I. ovatus* groups that were reported from southern China. The

three genetic groups in our data set suggest the potential for cryptic species among the groups.

While many factors can potentially account for the observed differences in genetic structure

between the two species, including population persistence and large-scale patterns of range

expansion, the differences in the mobility of hosts of tick immature stages (small mammals in *I.*

ovatus; birds in *H. flava*) is possibly driving the observed patterns.

Keywords: gene flow, host mobility, Mantel test, Spatial autocorrelation, cryptic species

Introduction

Tick-borne diseases are a public health concern and their control is often challenging due to the fact

that it involves a complex transmission chain of vertebrate hosts and ticks interacting in a changing

environment (Dantas-Torres et al., 2012). Population genetic studies have helped in understanding

the dispersal patterns and potential pathogen transmission in ticks (Araya-Anchetta et al., 2015).

The direction, distance and gene flow patterns between tick populations can be inferred from

population genetic studies (Mccoy 2008). For example, if high gene flow is observed there might be

a greater chance of colonizing new areas or of re-colonizing areas following successful vector control

programs.

Host mobility can affect the genetic patterns of tick populations, although its effects are not

consistent. Several studies have reported low levels of gene flow in ticks with less mobile hosts (e.g.

smaller mammals) and high levels of gene flow in ticks with highly mobile hosts (Araya-Anchetta et

al., 2015). For example, *Amblyomma americanum* and *A. triste* (Ixodidae, Acari) exhibited high gene

flow across various spatial scales (137, 000 km^2 to 2.78 million km^2) and this was attributed to their

hosts' dispersal capabilities (large mammals and birds) (Guglielmone et al., 2013; Mixson et al.,

2006; Trout et al., 2010), while low gene flow was observed in *Amblyomma dissimile* populations

and this was attributed to the low mobility of its hosts (small mammals, reptiles and salamanders)

(Lampo et al., 1998). In contrast, some studies have reported low gene flow despite the high mobility

of the host, for example in *Ixodes scapularis* (Qiu et al., 2002) and *Ornithodoros coriaceus* (Teglas et

al., 2006) which can be attributed to the host mobility.

In Japan, tick-borne diseases are an increasing public health concern, affecting humans and animals

(Yamaji et al., 2018). A total of 8 genera of ticks have been recorded in Japan, composed of 47

species: 43 Ixodidae species and 4 Argasidae species (Fujita et al., 2006). Out of 47 species, 21

species are known to parasitize humans (Okino et al., 2010). Among these 21 species, *Ixodes ovatus*,

the main vector of Lyme disease (Miyamoto et al., 1993) and *Haemaphysalis flava,* a vector of severe

fever with thrombocytopenia syndrome (SFTS) and Japanese spotted fever (JSF), were reported in

Japan (Yamaji et al., 2018; Yu et al., 2011). A previous study found that the hosts of adult *I. ovatus*

were mainly hares and can also be large mammals (e.g. cows and horses), and the hosts of immature

forms were small rodents (Yamaguti et al., 1971). *H. flava* adult host preference are cows, dogs,

horses, wild boar, bear, and deer, while immature forms only parasitized birds (Yamaguti et al.,

1971). Despite the similarity of *I. ovatus* and *H. flava* in which blood meal from different vertebrate

host is taken in each life stage (larva, nymph and adult) (Herrmann and Gern, 2015), comparative

population genetic studies on two Ixodid tick species such as *I. ovatus* and *H. flava* from the same

sampling location remain nonexistent. Thus, our study field the gaps about the dispersal movement

of two important tick species: *I. ovatus* and *H. flava.* The low gene flow observed in *I. ovatus* might

be due to the lower mobility of hares in the adult stage and small rodents in its immature stage

while *H. flava* will exhibit high gene flow because of its avian mediated dispersal in its immature

stage.

In addition to tick population structure studies, genetic studies may also reveal the presence of

cryptic species, where morphologically identified individuals might represent more than one species

(Fegan et al., 2005). Previous studies on *Rhipecephalus appendiculatus* (Kanduma et al., 2016) , *I.*

holocylus (Song et al., 2011) and *I. ovatus* (Li et al., 2018) have revealed the presence of cryptic

species based on the clustering of haplotypes in a phylogenetic tree, and concluded that

morphological criteria for species differentiation alone are equivocal and that genetic analysis is

important.

Here, we studied the population genetic structure of *I. ovatus* and *H*. *flava* and also tested for the

presence of cryptic species using mitochondrial DNA sequences of *cox1* and the 16S rRNA gene. We

hypothesized that *I. ovatus* and *H. flava* may have differences in their population genetic structure

despite some overlap in their adult host preference can be attributed to the differences in mobility

of the immature tick hosts.

Material and methods

Study site, collection, sampling and identification

From April 2016 until November 2017, ticks were collected by the standard flagging methods (Ginsberg et al., 1989) across Niigata Prefecture in Japan and a total of 29 sampling locations were surveyed for *H. flava* and *I. ovatus* ticks (Additional File 1. Table S1). Altitude per location ranged from 8 -1402 m/a.s.l. and the geographic distance between sites ranged from 4.28 - 247.65 km. Collected ticks were stored in microcentrifuge tubes with ethanol at 4^0C. A total of 2,103 individual tick samples were collected. Identification of sex, developmental stage and species was performed using a compound microscope and identification keys of (Yamaguti et al., 1971) .

DNA extraction, PCR amplification and sequencing

Genomic DNA (*I. ovatus* n=320*; H. flava* n=220) from each identified tick was extracted using the Isogenome DNA extraction kits (NIPPON GENE Co. Ltd., Tokyo, Japan) following the manufacturer's recommended protocol. The total number of analyzed samples (n=540) was chosen based on its species identity while the other identified tick species were excluded for this study. Prior to DNA extraction, each tick was washed with alcohol and PBS solution. DNA quality was checked using a NanoDrop™ 2000 Spectrophotometer (Thermo Scientific™). Two mitochondrial genes were amplified by polymerase chain reaction (PCR): Cytochrome Oxidase 1 (*cox1*) (658 base pairs) using the primer pairs LCO-1490 (5'- GGTCAACAAATCATAAAGATATTGG -3') and HCO1-2198 (5'-AAACTTCAGGGTGACCAAAAAATCA-3') (Folmer et al., 1994) and the 16S rRNA gene (16S) (407 base pairs) with the following primer pairs 16S+1 (5'CTGCTCAATGAATATTTAAATTGC-3') and 16S – 1 (5' -CGGTCTAAACTCAGATCATGTAGG-3')(Tian et al., 2011). All PCR amplifications of both *cox1* and 16S were performed with a final volume of 10 μl with 1μl of genomic DNA. The PCR reaction for both markers were composed of the following: 10x Ex Taq buffer, 25mM MgCl$_2$, 2.5mM dNTP, 10μm of forward and reverse primers and 5 units/μl of TaKaRa Ex Taq™ (Takara Bio Inc.). The *cox1* PCR amplification were as follows: an initial denaturation of 94^0C for 2 minutes, denaturation of 94^0C for

30 seconds, annealing of 38^0C for 30 seconds, extension of 72^0C for 1 minute for 30 cycles and final

extension of 72^0C for 10 minutes. While the 16S PCR amplification followed the protocol of (Tian et

al., 2011) with some modifications, initial denaturation of 94^0C for 3 minutes, denaturation of 94^0C

for 30 seconds, annealing of 50^0C for 40 seconds, extension of 72^0C for 40 seconds for 30 cycles and

final extension of 72^0C for 5 minutes. PCR products were purified using the QIAquick 96 PCR

Purification Kit (Qiagen) in accordance with the manufacturer's instructions, and sequenced by

Eurofin Genomics, Inc., Tokyo, Japan.

Sequence data analysis

We assembled forward and reverse reads for each individual using CodonCode Aligner version 1.2.4

software (https://www.codoncode.com/aligner/). No ambiguous bases were observed and the low

quality bases were removed in the start and end of the reads. Multiple alignment of sequences was

performed using the MAFFT alignment online program with default settings

(https://mafft.cbrc.jp/alignment/server/). To ensure sequence quality and correct species

identification, we checked the similarity of the sequences against reference sequences from

GenBank using BLASTN (Basic Local Alignment Search Tool – Nucleotide,

https://blast.ncbi.nlm.nih.gov/Blast.cgi?PROGRAM=blastn&PAGE_TYPE=BlastSearch&LINK_LOC=bl

asthome). The quality of the final aligned sequences (*cox*1 = 658bp; 16S = 407bp) was checked in

Mesquite version 3.5 (Maddison et al., 2011) wherein the protein coding genes were translated into

amino acids to confirm the absence of stop codons.

Population genetic analysis

The group of sequences obtained from ticks collected in the same geographic location was

considered as belonging to the same population. We analyzed the final sequences of both *H. flava*

and *I. ovatus* per genetic marker separately using DNASp version 6.12.03 (Rozas et al., 2017) and

Arlequin version 3.5.2.2 (Excoffier and Lischer, 2010) for the following parameters: number of

haplotypes (nh), average number of polymorphic sites (s), and average number of nucleotide

differences (k). The haplotype diversity (h) and nucleotide diversity (π) were calculated in Arlequin

3.5.2.2. The population genetic structure among and within sampling locations in Niigata Prefecture

was calculated by analysis of molecular variance (AMOVA) performed in Arlequin with 1023

permutations. The degree of molecular genetic differentiation between sampling locations defined

as populations was assessed by calculation of the pairwise F_{ST} values in Arlequin. To determine if the

genetic differentiation was influenced by geographical distance or altitudinal differences among

populations, we performed Mantel Test in GenAlEx version 6.51b2 (Peakall and Smouse, 2006). Two

tests per species and marker were conducted. First we compared pairwise genetic (pairwise F_{ST}

values) and geographical distances (km) and second we compared genetic distance (F_{ST} values) with

altitudinal differences (m/a.s.l.). The geographic distances were obtained from the GPS coordinates

(latitude and longitude) recorded during the sampling. All Mantel tests were assessed using 9999

permutations for the significance of the correlation.

The spatial component of genetic variation was further assessed by spatial autocorrelation (Smouse

and Peakall, 1999) using GenAlEx (Peakall and Smouse, 2006). The autocorrelation coefficient (r)

was computed from the geographic distance between sampling locations and the genetic distance

(pairwise F_{ST} values). This measures the genetic similarity between individual pairs within an

identified distance class. The size of the distance class influences the estimation of the spatial

autocorrelation. We identified the appropriate distance class based on the observed distribution of

pairwise geographic distance between the sampling locations (data not shown). The distance class

ranged into 10 classes of 20 kilometers from 20 to 200 kilometers. Bootstrap estimates of r and

random permutations were set at 9999 each for the test for significance. The upper and lower error

bars in the correlogram bound the 95% confidence interval of r value as identified by bootstrap

resampling. The upper and lower confidence limits bound the 95% confidence interval of the null

hypothesis of no spatial structure in the distance class. The genetic relationships among the

sampling locations (i.e., populations) were calculated by unweighted pair group method with the

arithmetic mean (UPGMA) cluster method using the APE package (Paradis and Schliep., 2018) and

R program (R Development core team., 2016) to create dendrogram using the genetic distance

matrix (pairwise F_{ST} values) generated from GenAlEx.

To evaluate the relationship between the haplotypes, we constructed haplotype network on

PopART program version 1.7 per marker (cox1, 16S rRNA gene) and species (*I. ovatus* and *H. flava*

(http://popart.otago.ac.nz/index.shtml) using the median-joining (MJ) network algorithm (Bandelth

et al., 1999).

Phylogenetic analysis

We constructed maximum likelihood (ML) gene trees for *cox1* and 16S haplotype sequences of *I.*

ovatus and *H. flava* using PhyML version 3.1 (Guindon and Gascuel, 2003) under the default settings.

We applied HKY and GTR nucleotide substitution models for *cox1* and 16S, respectively, as suggested

by jModelTest ver 2 (Darriba et al., 2012). Additional sequences from China (MH208506, MH208512,

MH208514, MH208522, MH208515-19, MH208524, MH208531, MH208574, MH208577,

MH208579, MH208681-87, MH208689-93, MH208706, KU664519 (Li et al., 2018)), Japan (Hokkaido

AB231670, U95900; Yamanashi AB819241, AB819243 and Aomori AB819244) (Mitani et al., 2007;

(Norris et al., 1999), Takano et al., 2014)) were included. We used *Ixodes canisuga* an outgroup

because it is a closely related species (KY962023, KY962074; (Hornok et al., 2015)).

Results

The number of *I. ovatus* ranged from 1 to 36 individuals per study site and, as with *H. flava*, more

were successfully sequenced for *cox1* (307/320; 95.9%) than for 16S (284/320; 88.8%) (Additional

File 1. Table S1). The number of successfully sequenced and analyzed *H. flava* ranged from 1 to 36

individuals per study site, with more successfully sequenced for *cox1* (220/223; 98.7%) than for 16S

(172/223; 77.1%). There were 60 and 63 *cox1* haplotypes in *I. ovatus* and *H. flava* respectively, and

24 and 40 16S haplotypes (Table 1). Haplotype diversity (h) ranged from 0.442 to 0.852 and

nucleotide diversity (π) ranged from 0.001 to 0.006 for both markers and species (Table 1).

Table 1. Summary of *cox1* and 16S haplotype, variability, genetic diversity of *Ixodes ovatus* and

Haemaphysalis flava populations in Niigata Prefecture

Marker	Species	n	nh	s	k	h	π
cox1	*I. ovatus*	307	60	65	2.728	0.852	0.004
	H. flava	220	63	60	1.472	0.789	0.002
16S	*I. ovatus*	284	24	22	0.699	0.442	0.001
	H. flava	172	40	49	2.447	0.835	0.006

Abbreviations: n sample size; nh number of haplotypes; s number of polymorphic sites; k mean

number of nucleotide differences; h haplotype diversity; π nucleotide diversity

AMOVA results revealed a high among-population divergence (41.54%) in *I. ovatus cox1* sequences,

whereas both *cox1* and 16S markers of *H. flava* indicated no significant genetic variation among

populations (Table 2). Global F_{ST} values were highest in *I. ovatus* with values 0.415 in *cox1* and 0.074

in 16S sequences. Pairwise F_{ST} values (0.500 to 1.00) of *I. ovatus* based on *cox1* showed significant

genetic differences between most pairs of populations such as between populations18 and 25 and

between populations 28 and 25 (Additional File 2 Table S2). We also observed significant differences

in 16S ranged from (0.0225 to 0.6319) (Additional File 3 Table S3). Pairwise F_{ST} values from the *cox1*

marker of *H. flava* showed no significant genetic differences except in few population pairs such as

in 5 and 6 and 6 and 13 (Additional File 4 Table S4). No significant genetic differentiation was also

observed in 16S marker of *H. flava* except in few population pairs such as between 3 and 20 and 4

and 20 (Additional File 5 Table S5).

Table 2. Analysis of molecular variance (AMOVA) using *cox1* and 16S of *Ixodes ovatus* and

Haemaphysalis flava populations

Marker	Species		df	ss	vc	pv	F_{ST}
cox1	I. ovatus	Among populations	28	190.38	0.5803 Va	41.54	0.4154**
		Within populations	278	227.04	0.8167 Vb	58.46	
	H. flava	Among populations	16	12.42	0.0039 Va	0.53	0.0053
		Within populations	203	148.73	0.7327 Vb	99.47	
16S	I. ovatus	Among populations	22	14.04	0.0260 Va	7.39	0.0739**
		Within populations	261	84.87	0.3252 Vb	92.61	
	H. flava	Among populations	12	23.6	0.0642 Va	5.21	0.0521**
		Within populations	159	185.58	1.1672 Vb	94.79	

Abbreviations: df degrees of freedom; ss sum of squares; Va, Vb and Vc are associate covariance

components; pv percentage variation; *P<0.05 ** P<0.01

The Mantel test of *I. ovatus* showed evidence of isolation by geographic distance in both markers:

(*cox1 r* = 0.197, *p* = 0.008; 16S *r* = 0.293, *p* = 0.011) (Figures 1a, b). In contrast, there was no evidence

for isolation by altitudinal difference (*cox1* r = -0.066, *p*=0.225; 16S r = -0.023, p = 0.577). Both

Mantel tests were not significant for *H. flava* (distance: *cox1* r= -0.145, p= 0.23; 16S r = -0.049, p=

0.33; altitude: r = 0.092, p = 0.30; 16S r = 0.217, p= 0.06). The spatial autocorrelation test in *I. ovatus*

revealed positive and significant correlation between genetic and geographic distance from 20 – 40

km, and 60 – 80 km, and 20 – 50 km and 120 – 160 km from *cox1* and 16S, respectively (Figure 1c,

d).

The UPGMA cluster dendrogram from the pairwise F_{ST} values from the *cox1*, *I. ovatus* (Figure 2)

revealed 2 genetic clusters across the 29 populations, where the cluster 2 included populations from

the more western sites including the island (location 24 and 25) while cluster 1 mostly included

populations in the northern and southern part of Niigata Prefecture (Figure 3). No evident genetic

clustering on the dendrogram was observed from the *I. ovatus* 16S sequences and in both *cox1* and

16S *H. flava* sequences (Additional File 6 Figure S1; Additional File 7 Figure S2; Additional File 8

Figure S3).

A putative I. ovatus species complex was identified based on the presence of 3 distinct haplotype

clusters in both cox1 and 16S rRNA genes; 2 groups in China and one in Japan. The *cox1* gene tree

of 60 *I. ovatus* haplotypes indicated three groups: group 1 of the published sequences from western

China, one large group 2 containing the 58 haplotypes and 2 divergent haplotypes (Hap 60 and Hap

59), and group 3 from western China (Figure 4). The haplotype network of the *cox1* gene of *I. ovatus*

also displayed the 2 divergent haplotypes (Hap 60 and Hap 59) (Additional File 12 Supplementary

Figure S7). The 2 divergent haplotypes were only found once in the whole data set. One was found

in population 29 and the other in population 7. The published haplotype from Hokkaido (Mitani et

al., 2007) occurred within our large group and the published sequences from western China were in

a different clade. The *I. ovatus* 16S gene tree (24 newly sequenced haplotypes) (Additional File 9

Figure S4) also recovered published haplotypes from Japan (Yamanashi and Aomori; (Takano et al.,

2014) and Hokkaido (Norris et al., 1999) within the same cluster as our new sequences. The *cox1*

haplotype sequences of *H. flava* were similar to reference sequences from China (Additional File 10

Figure S5) [unpublished manuscript] while the 16S haplotype sequences (Additional File 11 Figure

S6) showed similarities with Japan (Kagoshima, Fukui Aomori, Fukui, Yamanashi, Kagawa and Ehime

(Takano et al., 2014) and China [unpublished manuscript] reference sequences from Genbank.

Discussion

Contrasting population genetic structures between *H. flava* and *I. ovatus*

Our results supported our hypothesis that *H. flava* and *I. ovatus* display contrasting population

genetic structure. The noticeable high gene flow in *H. flava* may be a result of the higher level of

mobility of its bird hosts in the tick's immature stage and the more pronounced genetic structure

and lower gene flow in *I. ovatus* may be due to the restricted ability of its small mammalian hosts

at the immature stage to disperse further. 28 species of birds were previously reported as hosts of

H. flava from Japan*,* mostly from the order Passeriformes known as migratory birds (Yamauchi and

Takeno, 2000). The host for adults of *I. ovatus* included large mammals such as hares while

immature forms are found on small mammals such as small rodents and rats (Yamaguti et al., 1971).

The movement of small mammals is limited because of environmental and anthropogenic factors

such as difficulty in crossing roads and higher possibility of road kill due to vehicle collision and or

traffic (Haemig et al., 2008), which may explain the genetic structuring in *I. ovatus*. The host

preference of the adults *H. flava* are large mammals (e.g. cows, horses, wild boar, deer, bear) and

hares, dogs, rodents while the immature forms parasitize birds (Yamaguti et al., 1971). Mammals

and birds may have wide habitats ranges that may allow maintaining the gene flow of *H. flava*

between the locations in Niigata, as previously observed in *Amblyomma americanum* populations

(Mixson et al., 2006; Trout et al., 2010; Reichard et al., 2005) , and *Ixodes ricinus* (Casati et al., 2008).

Birds present the greatest potential for farther dispersal of *H. flava* that could reduce its genetic

structuring (Mcclain et al., 1995) that supports the results of our study.

Several alternative factors such as tick behavior, biology and ecology can also affect the tick dispersal

movement. For example, long movement patterns, resistance against unfavorable weather

conditions causing high survival in their hosts and low host specificity (Guglielmone et al., 2013) are

possible reasons for the low genetic structuring in *H. flava* populations. Previous study on two tick

species: *Hyalomma rufipes* and *Amblyomma hebraum* also displayed contrasting genetic patterns

despite the overlap sharing of highly mobiles hosts such as birds (Cangi et al., 2013) due to the

capacity of the immature stages of ticks to survive off the hosts in various habitat conditions

(Needham et al., 1986; Cumming, 1999; Estrada-Peña, 2015). Population structure can also be

influenced by assortative mating (e.g. *Ixodes ricinus*) wherein genetically similar individuals tend to

mate, rather than random mating resulting in increased genetic divergence (Kempf et al., 2009). Our

study does not have supporting data to test these alternative factors thus it is suggested in future

studies to further analyze these factors that could play a role in the tick's dispersal movement.

Based on previous studies using microsatellites and Random Amplification of Polymorphic DNA

(RAPD) markers (Yaegashi et al., 2014; Watanabe et al., 2010), isolation by distance suggests that

the populations are in equilibrium of migration and genetic drift at the spatial scale of Niigata

Prefecture. Our results were congruent with a previous study of *Amblyomma ovale* that showed

positive relationship of geographic and genetic distance at the spatial scale of the whole of Brazil

whose main hosts for the immature stage are small rodents while tapir and jaguar in its adult stage

(Bitencourth et al., 2019). In contrast, *H. flava* populations showed an absence of isolation by

distance. In circumstances when geographic distance can't explain the genetic structure of tick

populations, other factor such as ecological variables (e.g. land use) maybe affecting the genetic

structuring within tick populations (Casati et al., 2008; Krakowetz et al., 2010; Lampo et al., 2015)

which can be inferred when using molecular markers with higher mutation rate such as

microsatellites.

A significant and positive spatial autocorrelation of *I. ovatus* in both *cox1* and 16S sequences was

observed within the geographic range of 20 – 40 km and 60- 80 km in *cox1* sequences and 20 – 50

km and 120 – 160 km range in the 16S sequences. We can infer that the positive *r* values at the

maximum geographic rages of 60-80 km in *cox1* and 120 - 160 km in 16S *I. ovatus* sequences indicate

that gene flow is occurring in the wide geographic range. The lack of significance at 40-60 km in *cox1*

may suggest patch structure indicating genetic similarity or variation (Smouse and Peakall, 1999) as

observed on limpet species, *Siphonaria raoulensis* from New Zealand that also displayed recovery

of positive *r* values at greater distances (Wood and Gardner, 2007). However, no autocorrelation

was observed in *H. flava* at any distance class, suggesting that gene flow occurring has no specific

pattern favoring certain distances (Pardanani and Mather, 2009; Kitron and Kazmierczak.,

1997;Nicholson et al., 2014). Since the sampling geographic range of our study was within Niigata

Prefecture and distances >220 km were not included in our sampling, we suggest that future studies

collect samples at a wider geographic scale to more precisely determine the geographic distance at

which autocorrelation exists.

16S *I. ovatus* sequences showed genetic differentiation (pairwise F_{ST} = 0.0532-0.5652) in few pairs

of sampling locations as compared to the *cox1* sequences which may be due to lower variability of

the marker thus providing inadequate genetic variation for population genetic analysis, as

previously observed in *Amblyomma ovale* (Bitencourth et al., 2019) and *Rhipicephalus microplus*

(Low et al., 2015;Burger et al., 2014). The low variability of the 16S in *I. ovatus* may not be sufficient

to infer intraspecific relationship and is further supported by its low (*nd* = 0.001) nucleotide diversity

as compared in *cox1 I. ovatus* (*nd* = 0.004). Thus, it may also explain the grouping observed in the

cox1 I. ovatus tree which is absent in the 16S *I. ovatus* sequences. Despite of these, we used 16S

mitochondrial gene because this gene mutates at a rate that is informative for species-level

phylogenetic and broad biogeographic inferences(Araya-Anchetta et al., 2015)

The negative pairwise F_{ST} values obtained in *H. flava* sequences do not have any biological meaning.

These results from the unbiased estimator allowing a negative F_{ST} when the sample size is small.

(Bitencourth et al., 2016; Vial et al., 2006; Weir and Cockerham, 1984) . It can also be noted that

we found significant pairwise F_{ST} values in 2 sampling location in *H. flava cox1* sequences probably

due to the fact that both locations have only one tick sample. Although, some sampling sites in this

study have small sample size (n < 10), the *cox1* and 16S sequences provided valuable insight into the

genetic structure and gene flow patterns of the present-day tick populations. Thus, for future

research, we recommend to increase the sample size per sampling site.

Species complex formation in *I. ovatus cox1* sequences

Our *cox1 I. ovatus* gene tree showed Japanese individuals to form a group that was distinct from

haplotypes from southern China (Li et al., 2018). Despite the low gene flow we found in *cox1 I.*

ovatus, haplotypes found from Niigata were closely related to the published sequence from

Hokkaido, Japan which may indicate that these ticks originated from a diverse set of geographical

locations in Japan which might be transported by its hosts or is undergoing recent population

expansion from northern Japan (Hokkaido) to south (Niigata) or *vice versa*. We found three groups

(China 1, Japan and China 2) and 2 slightly divergent *cox1* haplotypes (Hap 60 and 59) in the Japan

group of *I. ovatus* in Niigata. Considering two or more cryptic species can be concealed in one

morphologically described species (Bitencourth et al., 2016), the occurrence of the 3 groups and the

divergent haplotypes suggests that *I. ovatus* may be a species complex. Previous studies have also

observed species complexes in *Ixodes* and *Rhipicephalus* (Li et al., 2018; Song et al., 2011;Burger et

al., 2014;Xu et al., 2003) suggesting that morphological criteria for species differentiation alone are

equivocal and that genetic analysis is important. An integrative approach of tick species concept

that includes the morphology, genetics, biology and ecological trails of ticks must be tackled (Dantas

et al., 2019).

Conclusions

Two Ixodid ticks, *H. flava* and *I. ovatus* exhibited contrasting patterns of genetic structure in Niigata

Prefecture, Japan. We suggest that these differences are due to differences in host mobility of the

immature life stages of the ticks. The more pronounced population genetic structure of *I. ovatus*

might be influenced by the restricted movement of its small mammalian host in its immature stage,

whereas the lack of structure in *H. flava* may be due to its avian-mediated movement in its immature

stage. Even though *I. ovatus* populations were genetically structured within Niigata, a published

haplotype from Hokkaido was also found, indicating that widespread dispersal is possible. The

occurrence of three groups and the divergent cox1 haplotypes in *I. ovatus* emphasizes the need of

additional methods in determining the existence of species complex in *I. ovatus* populations in Japan.

List of Abbreviations

H. flava – Haemaphysalis flava

I. ovatus – Ixodes ovatus

AMOVA – Analysis of molecular variance

UPGMA - unweighted pair group method with arithmetic mean

ML – maximum likelihood

F_{ST} – fixation index

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

All data generated or analyzed during the study are included in this published article and its

supplementary additional files. All the newly generated sequences are available in the Genbank

database under the accession numbers _______.

Declarations of interests

The authors declare that they have no competing interests.

Funding

This work is funded by the Japan Society for the Promotion of Science (JSPS) Grant-in-Aid for

Scientific Research (16K00569, 19K21996, 19H02276), the Sumitomo Electric Industries Group

Corporate Social Responsibility Foundation and the Research Unit Program of Ehime University. The

research was partially supported by the German Academic Exchange Service (DAAD, Programm

Projektbezogener Personenaustausch Japan 2018, project 57402018).

Authors' contributions

MAFR, MG, MM, KW conceptualized and the designed the experiment. MS, TT, RA, SI and MOS

designed the sampling collection, collected and identified the tick samples of this study. MD and KT

conducted the molecular analyses. MAFR, MG and KW performed the data analysis and wrote the

manuscript. All authors read and approve the manuscript.

Acknowledgements

The authors would like to thank Niigata Prefectural Office for their valuable help during the tick

sampling collection. Thank you to Dr. Thaddeus Carvajal and Joeselle Serrana for their valuable

suggestions in the population genetic analyses in this study and Micanaldo Francisco for the

technical assistance in constructing the sampling map of this study.

References

Araya-Anchetta, A., Busch, J.D., Scoles, G.A., Wagner, D.M., 2015. Thirty years of tick population

genetics: A comprehensive review. Infect. Genet. Evol. 29, 164–179.

https://doi.org/10.1016/j.meegid.2014.11.008

Bandelt, H.J., Forster, P. and Röhl, A., 1999. Median-joining networks for inferring intraspecific

phylogenies. Mol. bio. and evo. 16, 37-48.

https://doi.org/10.1093/oxfordjournals.molbev.a026036

Bitencourth, K., Amorim, M., de Oliveira, S. V., Voloch, C.M., Gazêta, G.S., 2019. Genetic diversity,

population structure and rickettsias in *Amblyomma ovale* in areas of epidemiological interest

for spotted fever in Brazil. Med. Vet. Entomol. 33, 256–268.

https://doi.org/10.1111/mve.12363

Bitencourth, K., Voloch, C.M., Serra-Freire, N.M., Machado-Ferreira, E., Amorim, M., Gazêta, G.S.,

2016. Analysis of *Amblyomma sculptum* haplotypes in an area endemic for Brazilian spotted

fever. Med. Vet. Entomol. 30, 342–350. https://doi.org/10.1111/mve.12174

Burger,TD., Shao,R., Barker,SC., 2014. Phylogenetic analysis of mitochondrial genome sequences

indicates that the cattle tick, *Rhipicephalus* (Boophilus) *microplus*, contains a cryptic species.

Molecular Phylogenetics and Evolution. 76, 241-253.

https://doi.org/10.1016/j.ympev.2014.03.017

Cangi, N., Horak, I.G., Apanaskevich, D.A., Matthee, S., das Neves, L.C.B.G., Estrada-Peña, A.,

Matthee, C.A., 2013. The Influence of Interspecific Competition and Host Preference on the

Phylogeography of Two African Ixodid Tick Species. PLoS One 8, 1–12.

https://doi.org/10.1371/journal.pone.0076930

Casati, S., Bernasconi, M. V., Gern, L., Piffaretti, J.C., 2008. Assessment of intraspecific mtDNA

variability of European *Ixodes ricinus* sensu stricto (Acari: Ixodidae). Infect. Genet. Evol. 8,

152–158. https://doi.org/10.1016/j.meegid.2007.11.007

Cumming, G.S., 1999. Host distributions do not limit the species ranges of most African ticks (Acari:

Ixodida). Bull. Entomol. Res. 89, 303–327. https://doi.org/10.1017/s0007485399000450

Dantas-Torres, F., Chomel, B.B., Otranto, D., 2012. Ticks and tick-borne diseases: A One Health

perspective. Trends Parasitol. 28, 437–446. https://doi.org/10.1016/j.pt.2012.07.003

Darriba, D., Taboada, G.L., Doallo, R., Posada, D., 2012. JModelTest 2: More models, new heuristics

and parallel computing. Nat. Methods 9, 772. https://doi.org/10.1038/nmeth.2109

Estrada-Peña, A., 2015. Ticks as vectors: Taxonomy, biology and ecology. OIE Rev. Sci. Tech. 34,

53–65. https://doi.org/10.20506/rst.34.1.2345

Excoffier, L., Lischer, H.E.L., 2010. Arlequin suite ver 3.5: A new series of programs to perform

population genetics analyses under Linux and Windows. Mol. Ecol. Resour. 10, 564–567.

https://doi.org/10.1111/j.1755-0998.2010.02847.

Fegan,M., Prior,P., 2005. How complex is the *Ralstonia solanacearum* species complex. Bacterial

wilt disease and the *Ralstonia solanacearum* species complex. APS Press., Minnesota, pp.

449-461.

Folmer,O., Black, M., Hoeh,W, Lutz,R. and Vrijenhoek,R.,1994. DNA primers for amplification of

mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. Mol.

Mar. Bio. and Biotech.. 3, 294-299.

Fujita, H., Fournier, P.E., Takada, N., Saito, T., Raoult, D., 2006. *Rickettsia asiatica* sp. nov., isolated

in Japan. Int. J. Syst. Evol. Microbiol. 56, 2365–2368. https://doi.org/10.1099/ijs.0.64177-0

Ginsberg,H.S., Ewing,C.P,.1989. Comparison of flagging, walking, trapping, and collecting from

hosts as sampling methods for northern deer ticks, *Ixodes dammini*, and lone-star ticks,

Amblyomma americanum (Acari: Ixodidae). Exp. Appl. Acarol.7, 313-322.

https://doi.org/10.1007/bf01197925

Guglielmone, A.A., Nava, S., Mastropaolo, M., Mangold, A.J., 2013. Distribution and genetic

variation of *Amblyomma triste* (Acari: Ixodidae) in Argentina. Ticks Tick. Borne. Dis. 4, 386–

390. https://doi.org/10.1016/j.ttbdis.2013.01.009

Guindon, S., Gascuel, O., 2003. A Simple, Fast, and Accurate Algorithm to Estimate Large

Phylogenies by Maximum Likelihood. Syst. Biol. 52, 696–704.

https://doi.org/10.1080/10635150390235520

Haemig, P.D., Waldenström, J., Olsen, B., 2008. Roadside ecology and epidemiology of tick-borne

diseases. Scand. J. Infect. Dis. 40, 853–858. https://doi.org/10.1080/00365540802270003

Herrmann, C., Gern, L., 2015. Search for blood or water is influenced by Borrelia burgdorferi in

Ixodes ricinus. Parasites & Vectors. 8, 4–11. https://doi.org/10.1186/s13071-014-0526-2

Hornok, S., Kontschán, J., Estrada-Peña, A., De Mera, I.G.F., Tomanović, S., De La Fuente, J., 2015.

Contributions to the morphology and phylogeny of the newly discovered bat tick species,

Ixodes ariadnae in comparison with I. vespertilionis and I. simplex. Parasites & Vectors. 8, 1–

7. https://doi.org/10.1186/s13071-015-0665-0

Kanduma, E.G., Mwacharo, J.M., Githaka, N.W., Kinyanjui, P.W., Njuguna, J.N., Kamau, L.M.,

Kariuki, E., Mwaura, S., Skilton, R.A., Bishop, R.P., 2016. Analyses of mitochondrial genes

reveal two sympatric but genetically divergent lineages of Rhipicephalus appendiculatus in

Kenya. Parasites & Vectors.9,353. https://doi.org/10.1186/s13071-016-1631-1

Kempf, F., de Meeûs, T., Arnathau, C., Degeilh, B., McCoy, K.D., 2009. Assortative Pairing in *Ixodes*

ricinus (Acari: Ixodidae), the European Vector of Lyme Borreliosis. J. Med. Entomol. 46, 471–

474. https://doi.org/10.1603/033.046.0309

Kitron,U., Kazmierczak, J.J., 1997. Spatial analysis of the distribution of lyme disease in Wisconsin.

Am. J. Epidemiol. 145, 558–566. https://doi.org/10.1093/oxfordjournals.aje.a009145

Krakowetz, C.N., Dergousoff, S.J., Chilton, N.B., 2010. Genetic variation in the mitochondrial 16S

rRNA gene of the American dog tick, *Dermacentor variabilis* (Acari: Ixodidae). J. Vector Ecol.

35, 163–173. https://doi.org/10.1111/j.1948-7134.2010.00073.x

Lampo, M., Rangel, Y., Matat, A., 2015. Population genetic structure of a three-host population

genetic in *Amblyommma dissimile*. J. Parasitol. .84, 1137–1142.

https://doi.org/10.2307/3284662

Li, L.H., Zhang, Y., Wang, J.Z., Li, X.S., Yin, S.Q., Zhu, D., Xue, J.B., Li, S.G., 2018. High genetic

diversity in hard ticks from a China-Myanmar border county. Parasites & Vectors. 11, 1–11.

https://doi.org/10.1186/s13071-018-3048-5

Low,VL., Tay,ST., Kho,KL., Koh,FX., Tan,TK., Lim,YA., Ong,BL., Panchadcharam,C., Norma-Rashid,Y.,

Sofian-Azirun.M., 2015. Molecular characterisation of the tick Rhipicephalus microplus in

Malaysia: new insights into the cryptic diversity and distinct genetic assemblages throughout

the world. Parasites & Vectors. 8, 341. https://doi.org/10.1186/s13071-015-0956-5

Maddison,W.P., Maddison,D.R.,2011. Mesquite: a modular system for evolutionary analysis. 2011.

See http://mesquiteproject. org.

Mccoy ,K., 2008. The population genetic structure of vectors and our understanding of disease

epidemiology. Parasite. 15, 444–8. https://doi.org /10.1051/parasite/2008153444

Mclain,DK., Wesson,DM., Oliver,Jr,JH., Collins,FH., 1995. Variation in ribosomal DNA internal

transcribed spacers 1 among eastern populations of *Ixodes scapularis* (Acari: Ixodidae). J.

Med. Entomol. 32, 353-360. https://doi.org /10.1093/jmedent/32.3.353

Mitani, H., Takahashi,M., Masuyama,M., Fukunaga,M., 2007. *Ixodes philipi* (Acari: Ixodidae):

phylogenetic status inferred from mitochondrial cytochrome oxidase subunit I gene

sequence comparison. J. Parasitol. 1, 719-722. https://doi.org /10.1645/GE-953R.1

Mixson, T.R., Lydy, S.L., Dasch, G.A., Real, L.A., 2006. Inferring the population structure and

demographic history of the tick, *Amblyomma americanum* Linnaeus. J. Vector Ecol. 31, 181–

192. https://doi.org/10.3376/1081-1710(2006)31[181:itpsad]2.0.co;2

Miyamoto, K., Nakao,M., Fujita,H., Sato,F., 1993.The ixodid ticks on migratory birds in Japan and

the isolation of Lyme disease spirochetes from bird-feeding ticks. Med. entomol. and zoo. 44,

315-326. https://doi.org/10.7601/mez.44.315

Needham,GR, Teel,PD., 1986. Water balance by ticks between bloodmeals. Morphology,

physiology, and behavioral biology of ticks/editors, Ellis Horwood Limited,Chichester, UK. pp.

100-151.

Nicholson, MC, and Mather, TN., 2014. Methods for evaluating Lyme disease risks using

geographic information systems and geospatial analysis. J. Med. Entomol. 33, 711-20.

https://doi.org/10.1093/jmedent/33.5.711

Norris, D.E., Klompen, J.S.H., Black IV, W.C., 1999. Comparison of the mitochondrial 12S and 16S

ribosomal DNA genes in resolving phylogenetic relationships among hard ticks (Acari:

Ixodidae). Ann. Entomol. Soc. Am. 92, 117–129. https://doi.org/10.1093/aesa/92.1.117

Okino,T., Ushirogawa,H., Matoba,K., Hatsushika,R., 2010. Bibliographical studies on human cases

of hard tick (Acarina: Ixodidae) bites in Japan (7) cases of unidentified tick infestation.

Kawasaki Medical Journal. 36, 127-141.

Paradis, E., and Schliep, K.,2018. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics. 35, 526-528.

https://doi.org/10.1093/bioinformatics/bty633

Pardanani, N., Mather, T.N., 2009. Lack of Spatial Autocorrelation in Fine-Scale Distributions of

*Ixodes scapularis(*Acari: Ixodidae). J. Med. Entomol. 41, 861–864.

https://doi.org/10.1603/0022-2585-41.5.861

Peakall, R., Smouse, P.E., 2006. GENALEX 6: Genetic analysis in Excel. Population genetic software

for teaching and research. Mol. Ecol. Notes 6, 288–295. https://doi.org/10.1111/j.1471-

8286.2005.01155.x

Qiu, W.G., Dykhuizen, D.E., Acosta, M.S., Luft, B.J., 2002. Geographic uniformity of the lyme

disease spirochete (*Borrelia burgdorferi*) and its shared history with tick vector (I*xodes*

scapularis) in the Northeastern United States. Genetics 160, 833–849.

R Development Core Team., 2016. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

Reichard, M.V., Kocan,A.A., Van Den Bussche ,A., Barker,R.W., Wyckoff,J., Ewing,S.A., 2005.

Sequence variation of the ribosomal DNA second internal transcribed spacer region in two

spatially distinct populations of *Amblyomma americanum* (L.)(Acari: Ixodidae). J.

Parasitol..91, 260-264. https://doi.org/10.1645/GE-292R1

Rozas,J., Ferrer-Mata,A., Sánchez-DelBarrio,J.C., Guirao-Rico,S., Librado,P., Ramos-Onsins,S.E.,

Sánchez-Gracia,A.,2017. DnaSP 6: DNA sequence polymorphism analysis of large data sets.

Mol. bio.and evo.34, 3299-3302. https://doi.org/10.1093/molbev/msx248

Smouse, P.E., Peakall, R., 1999. Spatial autocorrelation analysis of multi-allele and multi-locus

genetic microstructure. Heredity (Edinb). 82, 561–573.

https://doi.org/10.1038/sj.hdy.6885180

Song, S., Shao, R., Atwell, R., Barker, S., Vankan, D., 2011. Phylogenetic and phylogeographic

relationships in *Ixodes holocyclus* and *Ixodes cornuatus* (Acari: Ixodidae) inferred from COX1

and ITS2 sequences. Int. J. Parasitol. 41, 871–880.

https://doi.org/10.1016/j.ijpara.2011.03.008

Takano, A., Fujita, H., Kadosaka, T., Takahashi, M., Yamauchi, T., Ishiguro, F., Takada, N., Yano, Y.,

Oikawa, Y., Honda, T., Gokuden, M., Tsunoda, T., Tsurumi, M., Ando, S., Andoh, M., Sato, K.,

Kawabata, H., 2014. Construction of a DNA database for ticks collected in Japan: application

of molecular identification based on the mitochondrial 16S rDNA gene. Med. Entomol. Zool.

65, 13–21. https://doi.org/10.7601/mez.65.13

Teglas, M.B., Drazenovich, N.L., Stott, J., Foley, J.E., 2006. The geographic distribution of the

putative agent of epizootic bovine abortion in the tick vector, *Ornithodoros coriaceus*. Vet.

Parasitol. 140, 327–333. https://doi.org/10.1016/j.vetpar.2006.03.027

Tian, Z., Liu, G., Xie, J., Yin, H., Luo, Jianxun, Zhang, L., Zhang, P., Luo, Jin, 2011. Discrimination

between *Haemaphysalis longicornis* and *H. qinghaiensis* based on the partial 16S rDNA and

the second internal transcribed spacer (ITS-2). Exp. Appl. Acarol. 54, 165–172.

https://doi.org/10.1007/s10493-010-9423-3

Trout, R.T., Steelman, C.D., Szalanski, A.L., 2010. Population Genetics of *Amblyomma americanum*

(Acari: Ixodidae) Collected From Arkansas. J. Med. Entomol. 47, 152–161.

https://doi.org/10.1603/me09106

Vial, L., Durand, P., Arnathau, C., Halos, L., Diatta, G., Trape, J.F., Renaud, F., 2006. Molecular

divergences of the Ornithodoros sonrai soft tick species, a vector of human relapsing fever in

West Africa. Microbes Infect. 8, 2605–2611. https://doi.org/10.1016/j.micinf.2006.07.012

Watanabe,K., Monaghan,M.T., Takemon,Y., Omura,T., 2010. Dispersal ability determines the

genetic effects of habitat fragmentation in three species of aquatic insect. Aquatic

conservation. 20, 574-579. https://doi.org/10.1002/aqc.1124

Weir, B.S., Cockerham, C.C., 1984. Estimating F-Statistics for the Analysis of Population Structure

Author (s): B . S . Weir and C . Clark Cockerham Published by : Society for the Study of

Evolution Stable URL : http://www.jstor.org/stable/2408641. Evolution (N. Y). 38, 1358–

1370. https://doi.org/10.2307/2408641

Wood, A.R., Gardner, J.P.A., 2007. Small spatial scale population genetic structure in two limpet

species endemic to the Kermadec Islands, New Zealand. Mar. Ecol. Prog. Ser. 349, 159–170.

https://doi.org/10.3354/meps07110

Xu,G., Fang,Q.Q., Keirans,J.E., Durden,L.A., 2003. Molecular phylogenetic analyses indicate that the

Ixodes ricinus complex is a paraphyletic group. J. Parasitol. 1, 452-457.

https://doi.org/10.1645/0022-3395(2003)089[0452:MPAITT]2.0.CO;2

Yamaguti, N., Tipton, V.J., Keegan, H.L., Toshioka, S., 1971. Ticks for Japan, Korea, and the Ryukyu

Islands. Brigham Young Univ. Sci. Bull. XV, 226.

Yaegashi,S., Watanabe,K., Monaghan,M.T., Omura,T., 2014. Fine-scale dispersal in a stream

caddisfly inferred from spatial autocorrelation of microsatellite markers. Freshwater Science.

33, 172-180. https://doi.org/10.1086/675076

Yamaji, K., Aonuma, H., Kanuka, H., 2018. Distribution of tick-borne diseases in Japan: Past

patterns and implications for the future. J. Infect. Chemother. 24, 499–504.

https://doi.org/10.1016/j.jiac.2018.03.012

Yamauchi, T., Takeno, K., 2000. Record of Haemaphysalis flava Neumann (Acari, Ixodidae) from

Zoothera dauma (Latham) (Aves) in Kyushu, Japan. Esakia 40, 99–101.

Yu,X.J., Liang,M.F., Zhang,S.Y., Liu,Y., Li,J.D., Sun,Y.L., Zhang,L., Zhang,Q.F., Popov,V.L., Li,C., Qu,J.,

2014. Fever with thrombocytopenia associated with a novel bunyavirus in China. New Eng.

Jour. of Med. 364.1523-1532. https://doi.org/10.1056/NEJMoa1010095

Additional Files

Additional File 1.xls Table S1. Summary of *Haemaphysalis flava* and *Ixodes ovatus* collected from

the different locations of Niigata Prefecture and its corresponding sample number

Additional File 2.xls Table S2. Pairwise comparison of genetic differentiation (F_{ST}) calculated for all

I. ovatus using the *cox1* gene

Additional File 3.xls Table S3. Pairwise comparison of genetic differentiation (F_{ST}) calculated for all

I. ovatus using the 16S gene

Additional File 4.xls Table S4. Pairwise comparison of genetic differentiation (F_{ST}) calculated for all

H. flava using the *cox1* gene

Additional File 5.xls Table S5. Pairwise comparison of genetic differentiation (F_{ST}) calculated for all

H. flava using the 16S gene

Additional File 6.docx Figure S1. An unweighted pair group method with the arithmetic mean

(UPGMA) dendrogram of *I. ovatus* based on the pairwise genetic distance (F_{ST}) of 16S among 23

sampling locations across Niigata Prefecture, Japan.

Additional File 7.docx Figure S2 An unweighted pair group method with the arithmetic mean

(UPGMA) dendrogram of *H. flava* based on the pairwise genetic distance (F_{ST}) of *cox1* among 17

sampling locations across Niigata Prefecture, Japan.

Additional File 8.docx Figure S3 An unweighted pair group method with the arithmetic mean

(UPGMA) dendrogram of *H. flava* based on the pairwise genetic distance (F_{ST}) of 16S among 17

sampling locations across Niigata Prefecture, Japan.

Additional File 9.docx Figure S4 Phylogenetic tree of 16s haplotype sequences of *I. ovatus* inferred

by maximum likelihood analysis with evolutionary model GTR with sequences from China

(MH208506, MH208512, MH208514 – MH208519, MH208522, MH208524, MH208531,

MH208574, MH208577, MH208579, KU664519), Japan (AB819241, AB819242, AB819243,

AB819244, and U95900). An outgroup (*Ixodes canisuga*) was also included.

Additional File 10.docx Figure S5 Phylogenetic tree of *cox1* haplotype sequences of *H. flava*

inferred by maximum likelihood analysis with evolutionary model HKY with sequences from China

(KY021800- KY021807; KY021810-KY021819, KY003181, JQ62588 - JQ625889; JQ737097;

JF758632). An outgroup (*Ixodes canisuga)* was also included.

Additional File 11.docx Figure S6 Phylogenetic tree of 16S haplotype sequences of *H. flava*

inferred by maximum likelihood analysis with evolutionary model GTR with sequences from China

(KC844858 -KC844867; KC844880-KC844882, M696720, KP324926, KX450279) and Japan

(AB19177- AB819192). An outgroup (*Ixodes canisuga)* was also included.

Additional File 12 Figure S7 Median-joining haplotype network of *I. ovatus* from *cox1* marker.

Each circle represents a unique haplotype and the lines correspond to mutations.

Additional File 13 Figure S8 Median-joining haplotype network of *I. ovatus* from 16S marker. Each

circle represents a unique haplotype and the lines correspond to mutations.

Additional File 14 Figure S9 Median-joining haplotype network of *H. flava* from *cox1* marker. Each

circle represents a unique haplotype and the lines correspond to mutations.

Additional File 15 Figure S10 Median-joining haplotype network of *H. flava* from 16S marker. Each

circle represents a unique haplotype and the lines correspond to mutations.

Figures

Figure 1. Plots (a) and (b) show the relationship between the pairwise F_{ST} values and the geographic

distances for *I. ovatus* in both *cox1* and 16S markers from the Mantel test analysis. Figures (c) and

(d) illustrate the influence of geographic distance in the spatial autocorrelation analysis of *cox1* and

16S *I. ovatus* sequences. The upper and lower 95% confidence intervals are depicted as the red

dotted lines (U and L) and the correlation coefficient *r* is plotted. The values above and below the

dashed lines indicate spatial structure. The error bars result from the 1000 permutations of

individuals within each distance class. Both analyses were done in Genalex software.

Figure 2. An unweighted pair group method with the arithmetic mean (UPGMA) dendrogram of *I.*

ovatus based on the pairwise genetic distance (F_{ST}) of *cox1* among 29 sampling locations across

Niigata Prefecture, Japan.

Figure 3. The distribution of the two genetic clusters as observed in the UPGMA cluster dendrogram

(Figure 3) of *I. ovatus cox1* sequences from Niigata Prefecture.

Figure 4. Maximum likelihood gene tree of *cox1* haplotype sequences of *I. ovatus* with sequences

from China (MH208681-MH208687, MH208689-MH2086393, MH208706), Japan (AB231670) and

an outgroup of *Ixodes canisuga.*

Yellow reference sequences from China (Li et al., 2018); *red* reference sequence from Hokkaido,

Japan (Mitani et al., 2007); *blue I. ovatus* divergent *cox1* haplotypes; *green I. ovatus cox1*

remaining haplotypes; *purple I. canisuga* outgroup