Direct ETTIN-auxin interaction controls chromatin state in gynoecium development

André Kuhn[1], Sigurd Ramans Harborough[2], Heather M. McLaughlin[1], Stefan Kepinski[2] and

Lars Østergaard[1*]

[1]Department of Crop Genetics, John Innes Centre, Norwich Research Park, Norwich NR4

7UH, United Kingdom.

[2]Centre for Plant Sciences, Faculty of Biological Sciences, University of Leeds, Leeds LS2

9JT, United Kingdom.

*For correspondence

Impact statement: Auxin binds to the ETTIN transcription factor to disrupt the interaction

between ETT and a TPL/TPR co-repressor and subsequently affecting chromatin dynamics

to ensure proper gynoecium development.

Mailing address of corresponding author
Lars Østergaard
John Innes Centre
Norwich Research Park
Norwich
NR4 7UH
United Kingdom
Phone +44 1603 450572
Email: lars.ostergaard@jic.ac.uk
Skype: lars.oest

Abstract

Hormonal signalling in animals often involves direct transcription factor-hormone interactions that modulate gene expression[1,2]. In contrast, plant hormone signalling is most commonly based on de-repression via the degradation of transcriptional repressors[3]. Recently, we uncovered a non-canonical signalling mechanism for the plant hormone auxin in organ development with strong similarity to animal hormonal pathways. In this mechanism, auxin directly affects the activity of the auxin response factor ETTIN (ETT) towards regulation of target genes without the requirement for protein degradation[4,5]. Here we show that auxin binds ETT to modulate gene expression and that this ETT-auxin interaction leads to the dissociation of ETT from co-repressor proteins of the TOPLESS/TOPLESS-RELATED family followed by histone acetylation and the induction of target gene expression. Whilst canonical ARFs are classified as activators or repressors[6], ETT is able to switch chromatin locally between repressive and de-repressive states in an instantly-reversible auxin-dependent manner.

Developmental programmes within multicellular organisms originate from a single cell (*i.e.* a

fertilised oocyte) that proliferates into numerous cells ultimately differentiating to make up

specialised tissues and organs. Tight temporal and spatial regulation of the genes involved

in these processes is essential for proper development of the organism. Changes in gene

expression are often controlled by mobile signals that translate positional information into

cell-type specific transcriptional outputs[7]. In plants, this coordination can be facilitated by

phytohormones such as auxin, which controls processes throughout plant development[8]. In

canonical auxin signalling, auxin-responsive genes are repressed when auxin levels are low

by Aux/IAA transcriptional repressors that interact with DNA-bound Auxin Response Factors

(ARFs). As auxin levels increase, the auxin molecule binds to members of the TIR1/AFB

family of auxin co-receptors[9,10]. This facilitates interaction with Aux/IAA repressors, Aux/IAA

ubiquitinylation and subsequent degradation by the 26S proteasome, while relieving the

repression of ARF-targeted loci[11,12].

We recently identified an alternative auxin-signalling mechanism whereby auxin directly

affects the activity of a transcription factor (TF) complex towards its downstream targets[4,5].

This mechanism mediates precise polarity switches during organ initiation and patterning

and includes the ARF, ETTIN (ETT/ARF3) as a pivotal component. However, ETT is an

unusual ARF lacking the Aux/IAA-interacting Phox/Bem1 (PB1) domain[4,13] and it is therefore

likely that ETT would mediate auxin signalling via an alternative pathway.

ETT can interact with a diverse set of TFs and these interactions are sensitive to the

naturally occurring auxin, indole 3-acetic acid (IAA). The region responsible for IAA-

sensitivity is situated within the C-terminal part of ETT, known as the ETT-Specific (ES)

domain[6]. A protein fragment containing 207 amino acids of the ES domain, $ES^{388-594}$,

sufficient for mediating IAA-sensitivity in ETT-protein interactions, was produced

recombinantly and shown to be intrinsically disordered[14]. The sensitivity of ETT-TF

interactions to IAA suggests a direct effect of the IAA molecule on the ETT protein.

Therefore, to test whether ETT binds IAA, we carried out heteronuclear single quantum

coherence (HSQC) nuclear magnetic resonance (NMR) experiments using [15]N-labelled

ES[388-594] protein. The HSQC spectrum, recorded at 5˚C, shows a prominent signal-dense

region consistent with the ES domain being largely intrinsically disordered. Interestingly, the

spectrum also shows dispersed peaks flanking the signal-dense region indicating that there

is nevertheless some propensity to form secondary structure, particularly with a helical

character (Fig. 1a). In addition to this overview of ETT structure, the HSQC NMR probes

chemical shifts of protein amide-NH bonds in response to the presence of ligand[15]. We

found that a number of residues shifted their position in the spectrum in response to the

addition of IAA, whereas addition of the related Benzoic Acid (BA) had no effect (Fig. 1a-c).

These shifts show that certain residues are experiencing a changed chemical environment

as a consequence of IAA-binding and this may include the conformational change of a

structural motif within the ETT protein. The HSQC experiment therefore demonstrates that

ETT binds IAA directly. This experiment has not allowed us to assign signals to specific

amino acids and hence there is some uncertainty associated with tracking the chemical

shifts of some residues. However, a particularly large change is observed when IAA is

added to the ETT fragment for the tryptophan NH cross peak (~10ppm, rectangle I in Fig.

1a,c). Since there is only one tryptophan in the ETT fragment used here (W505), this shift

can be assigned to this residue.

We also used the recombinant ETT fragment in an Isothermal Titration Calorimetry (ITC)

assay, which characterises binding of ligands to proteins by determining thermodynamic

parameters of the interaction as heat exchange. This experiment revealed interaction

between ETT and IAA, while control experiments titrating IAA into buffer without protein and

titrating buffer without IAA into the ETT fragment showed no heat exchange (Fig. 1d-f).

Together, these two independent biochemical methods demonstrate that ETT binds IAA

directly thus revealing a key molecular aspect of the non-canonical auxin-signalling pathway.

Previously, *PINOID* (*PID*)[16] and *HECATE1* (*HEC1*)[17] were identified as ETT target genes[4,5]

and both genes are upregulated in gynoecium tissue from the *ett-3* mutant compared to wild

type (Figure 2-figure supplement 1). We also observed that expression of both genes is

induced by IAA, but did not observe any additional induction beyond the constitutive

upregulation in the *ett-3* mutant background (Figure 2-figure supplement 1). This ETT-

dependent regulation does not require a functional TIR1/AFB machinery, since IAA-induction

of *PID* and *HEC1* is still observed in TIR1/AFB mutant combinations, whereas the known

TIR1/AFB-mediated auxin induction of the *IAA19* gene is completely abolished in these

mutants (Fig. 2a-c).

To further assess the TIR1/AFB independence of the ETT-mediated auxin signalling

pathway, we exploited a recently-developed synthetic auxin-TIR1 pair[18]. In this system, the

auxin-binding pocket of TIR1 has been engineered (ccvTIR1) to accommodate an IAA

derivative bearing a bulky side chain (cvxIAA). By expressing the ccvTIR1 in a *tir1 afb2*

mutant background, the canonical pathway will only respond to the addition of cvxIAA and

not IAA[18]. We performed an expression analysis on *ccvTIR1* gynoecia treated ±cvxIAA and

±IAA as well as control plants with the same treatments. In this experiment, *IAA19* served as

a control gene whose expression is known to be regulated in a TIR1/AFB-dependent

manner. Indeed, *IAA19* was strongly upregulated by cvxIAA in the ccvTIR1 line, but not by

IAA (Fig. 2d). In contrast, *PID* and *HEC1* expression was not significantly affected by

cvxIAA, whilst still responding to IAA in the ccvTIR1 background (Fig. 2d). These data

demonstrate that ETT-mediated auxin signalling can occur independently of the canonical

TIR1/AFB signalling pathway.

In a phylogenetic analysis of ETT protein sequences across the angiosperm phylum, we

identified a number of regions that are highly conserved (Figure 3-figure supplement 1).

Unsurprisingly, the DNA-binding domain characteristic to B3-type TFs such as ARF proteins

was conserved across all ETT proteins. Towards the C terminus of the ES domain we

identified an EAR-like motif with a particularly high level of conservation (Fig. 3a, Figure 3-

figure supplement 1). Ethylene-responsive element binding factor-associated Amphiphilic

Repression (EAR) motifs are also found in Aux/IAA proteins. Interactions between Aux/IAA

and members of the TOPLESS and TOPLESS-RELATED (TPL/TPR) family of co-

repressors occur via this motif[19]. TPL/TPRs mediate their repressive effect by attracting

histone deacetylases (HDACs) to promote chromatin condensation[20]. Since ETT functions

independently of the canonical auxin pathway, it is possible that its role in chromatin

remodelling occurs via direct interaction with TPL/TPRs through the EAR-like motif. To test

this, we carried out Yeast 2-Hybrid (Y2H) assays in which ETT was found to interact with

TPL, TPR2 and TPR4 (Fig. 3b, Figure 3-figure supplement 1). Moreover, mutating residues

in the EAR-like motif abolished the interactions demonstrating its requirement for the ETT-

TPL/TPR interaction (Fig. 3b).

Given that several ETT-protein interactions are affected by IAA and that part of the ETT

transcriptome changes in response to IAA[4,5], we tested the IAA sensitivity of ETT-TPL/TPR

interactions. In both Y2H and in co-immunoprecipitation (Co-IP) experiments, we observed

that the interactions were reduced with increasing IAA concentrations (Fig. 3b,c and Figure

3-figure supplement 2). Moreover, as described previously for other ETT-protein

interactions, the sensitivity was specific to IAA as other auxinic compounds tested did not

show this effect (Figure 3-figure supplement 2). Henceforth, 'auxin' will refer to IAA unless

stated otherwise. These data suggest that in conditions with low auxin levels, ETT can

interact with TPL/TPR proteins to repress the expression of target genes. An increase in

cellular auxin causes ETT to bind auxin thereby undergoing a conformational change that

abolishes interaction with TPL/TPR co-repressors.

TPL was originally identified as a key factor involved in setting up the apical-basal growth

axis during embryo development[21,22]. Large-scale interaction studies suggest that the five

Arabidopsis TPL/TPRs have roles throughout plant development[20,23]. Whilst ETT has been

implicated in a wide array of developmental processes[24-27], the most dramatic phenotypes of

ett loss-of-function mutants are observed during gynoecium development[13,28,29]. In

accordance with this, *ETT* is highly expressed in the gynoecium (Fig. 4a)[4]. We produced

reporter lines of *TPL*, *TPR2* and *TPR4* promoters fused to the *GUS* gene to test if they

overlap with *ETT* expression in the gynoecium. Both *pTPL:GUS* and *pTPR2:GUS* exhibited

strong expression in the apical part of the gynoecium where *ETT* is also expressed, while no

pTPR4:GUS expression was observed (Fig. 4a-d). Single loss-of-function mutants in *TPL*

and *TPR2* do not show any abnormal phenotypes during gynoecium development. However,

the *tpl tpr2* double mutant has defects in the development of the apical gynoecium similar to

ett mutants (Fig. 4e-g) demonstrating that *TPL* and *TPR2* function redundantly in gynoecium

development. Together with the protein interaction data and the overlapping expression

patterns, these results suggest that ETT and TPL/TPR2 cooperate to regulate gynoecium

development.

TPL was shown previously to recruit histone deacetylase, HDA19, during early Arabidopsis

flower development to keep chromatin in a repressed state[20]. Moreover, HDA19 was also

recently shown to participate in repression of the meristem identity gene, *SHOOT*

MERISTEMLESS (*STM*)[30]. Here, our analysis of gynoecia from the *hda19-4* mutant

demonstrate that HDA19 is also required for gynoecium development as the *hda19-4* mutant

has strong style defects (Fig. 4h). In agreement with this, the *HDA19* gene was highly

expressed in gynoecium tissue, whereas another member of the *HDA* gene family, *HDA6*,

was not (Figure 4-figure supplement 1). Moreover, HDA19 recruitment likely involves *ETT*,

since expression of the ETT target genes, *PID* and *HEC1*, are increased in the *tpl tpr2* and

hda19-4 mutants compared to wild type. Similar to the *ett* mutant, auxin treatments failed to

further induce expression in these mutants (Fig. 4i,j). These observations suggest that ETT,

TPL/TPR2 and HDA19 function in conjunction to control gene expression during gynoecium

development.

To test the direct interaction of ETT, TPL and HDA19 on chromatin, we performed

Chromatin-Immunoprecipitation (ChIP) using reporter lines expressing GFP fusion protein.

Although only ETT is expected to bind DNA, ChIP followed by qPCR revealed that all three

proteins associate with DNA elements in the same regions of the promoters of *PID* and

HEC1 (Fig. 5a). This supports a model in which ETT recruits TPL/TPR2 and HDA19 to ETT

target loci to keep chromatin in a condensed state through histone deacetylation. When

auxin levels increase, the ETT-TPL/TPR2 interaction is broken, presumably preventing

HDA19 from deacetylating histones. To test this, we assayed for H3K27 acetylation, which is

a substrate for HDA19. H3K27 acetylation increased in the absence of ETT and upon

treatment with auxin. This occurred in the same regions of the *PID* and *HEC1* promoters

where the proteins were found to associate (Fig. 5b,c). In agreement with ETT mediating the

association of TPL/TPR and HDA19 with these regions, there was no further increase of

acetylation in the *ett-3* mutant upon treatment with auxin (Fig. 5b,c).

The data presented in this paper provide molecular insight as to how auxin levels are

translated into changes in gene expression of ETT target genes. Our data lead to a model in

which low levels of auxin maintain ETT associations with TPL/TPR2 to repress gene

expression via H3K27 deacetylation. As auxin levels increase, TPL/TPR2 (and hence

HDA19) disassociate from ETT, promoting H3K27 acetylation (Fig. 5d). This model

molecularly underpins the published association between auxin dynamics and *PID*

expression at the gynoecium apex where *PID* is repressed at early stages of development to

allow symmetry transition, but subsequently de-repressed as auxin levels rise to facilitate

polar auxin transport[4,31].

The direct binding of auxin allows ETT to switch the chromatin locally between repressive

and de-repressive states, whilst other ARFs have been categorised as either repressors or

activators[8]. The effect of auxin is therefore instantly reversible, making it possible to switch

between states immediately in response to changes in auxin levels. This feature, which is

reminiscent of animal hormonal signalling pathways such as the Thyroid Hormone and

Wnt/ß-catenin pathways[1,2], may be particularly important in controlling changes in tissue

polarity during plant organogenesis as observed in the Arabidopsis gynoecium[31].

The identification of a direct auxin-ETT interaction to control gene expression adds an

additional layer of complexity to auxin biology, which contributes towards explaining how

auxin imparts its effect on highly diverse processes throughout plant development. In a wider

context, this work also opens for the exciting possibility that direct transcription factor-ligand

interactions is a general feature in the control of gene expression in plants as found in

animals.

MATERIAL AND METHODS

Plant materials and treatments

Plants were grown in soil at 22 °C in long day conditions (16hrs day/8 hrs dark). All

mutations were in the *Col-0* background. Mutant alleles described before include e*tt-3*[4,13]*,*

hda19-4 (SALK_139443)[32], *pETT:GUS*[33], *pETT:ETT-GFP* in *ett-3*[4], *pTPL:TPL:GFP*[34],

p35S:HDA19:GFP[34], *pTIR1:ccvTIR1* in *tir1-1 afb2-3*[19] and *tir1-1 afb2-3 afb3-4*[35].The *tpl*

mutant (SALK_034518C) was obtained from the European Arabidopsis Stock Centre.

For both expression and ChIP analysis, auxin treatments were applied by spraying bolting

Col-0 and *ett-3* inflorescences with a solution containing 100 μM IAA (Sigma) or cvxIAA and

0.015% Silwet L-77 (De Sangosse Ltd.). Treated samples were returned to the growth room

and incubated for two hours.

Expression analysis

Quantitative Real time PCR (qRT-PCR) was used for expression analysis. RNA was

extracted from floral buds using the RNeasy mini kit (Qiagen). Using the SuperScript™ IV

First-Strand Synthesis kit (ThermoFisher), cDNA was synthesised from 1 μg of total RNA.

Subsequently, qRT-PCR was carried out using SYBR Green JumpStart Taq ReadyMix

(Sigma) using the appropriate primers (Figure 2-source data 1). Relative expression values

were determined using the $2^{-\Delta\Delta Ct}$ method[36]. Data were normalised to *POLYUBIQUITIN 10*

(*UBQ10/*AT4G05320) expression.

ETT protein analysis by alignment

Published *ETT* sequences of 22 Angiosperm species were retrieved from Phytozome

version 12[37]. Nucleotide sequences were translated and aligned using MUSCLE in

Geneious version 6.1.8[38]. The EAR domain was extracted as a sequence logo (Fig. 3a;

Figure 3-figure supplement 1).

Generation of the *tpl tpr2* CRISPR mutant

The *tpl tpr2[ge]* mutant was generated using CRISPR/Cas9 technology by a method previously

described[39]. Briefly, for the construction of the RNA-guided genome-editing plasmid, DNA

sequences encoding the gRNA adjacent to the PAM sequences were designed to target two

specific sites in *TPR2* (AT3G16830). DNA-oligonucleotides (Figure 2-source data 1)

containing the specific gRNA sequence were synthesised and used to amplify the full gRNA

from a template plasmid (AddGene #46966). Using Golden Gate cloning[40] each gRNA was

then recombined in a L1 vector downstream of U6 promoter[39]. Finally, the resulting gRNA

plasmids were then recombined with a L1 construct containing *pYAO:Cas9_3:E9t*[39] (kindly

provided by Jonathan Jones) and a L1 construct containing Fast-Red selection marker

(AddGene #117499) into a L2 binary vector (AddGene #112207).

The construct was transformed into *Agrobacterium tumefaciens* strain GV3101 by

electroporation, followed by plant transformation by floral dip into the *tpl* single mutant[41].

Transgenic T0 seeds appear red under UV light and were selected under a Leica M205FA

stereo microscope. T0 plants were genotyped using PCR and the *TPR2* locus sequenced

(Oligonucleotides in Figure 2-source data 1). Genome edited plants were selected and the

next generation grown (T1). Seeds of this generation were segregating in a 3:1 ratio for the

transgene. Transgene negative plants were selected and grown on soil. To find homozygous

mutations T1 plants were again genotyped. The T2 generation was again checked for the

absence of the transgene.

Protein interaction

For Yeast-two-Hybrid (Y2H) assays coding sequences were cloned into pDONR207 and

recombined into the pGDAT7 and pGBKT7 (Clontech). Using the co-transformation

techniques[41] these constructs were transformed into the AH109 strain (Clontech).

Transformations were selected on Yeast Selection Medium (YSD) lacking Tryptophan (W)

and Leucine (L) at 28°C for 3-4 days. Transformed yeast cells were serially diluted (10^0, 10^1,

10^2 and 10^4) and dotted on YSD medium lacking Tryptophan (W), Leucin (L), adenine (A)

and Histidine (H) to test for interaction. To examine interaction strength 3-amino-1,2,4-

triazole (3-AT) was supplemented to the YSD (-W-L-A-H) medium with different

concentrations (0, 5, 10 mM). To determine the effect of auxinic compounds on the protein-

protein interactions benzoic acid (BA), IAA, NAA and 2,4D (all Sigma) were dissolved in

ethanol and added directly to the medium at the desired concentrations. Pictures were taken

after 3 days of growth at 28°C.

For the β-Galactosidase assay transgenic yeast was grown in liquid YSD (-W-L) medium

supplemented with/-out 100 μM IAA or NAA, to an OD_{600} of 0.5. The cells were then

harvested and lysed using 150 μL Buffer Z with β-mercaptoethanol (100 mM Phosphate

buffer pH 7, 10 mM KCl, 1mM Mg_2SO_4, β-mercaptoethanol 50 mM), 50 μL chloroform and

20 μL of 0.1% SDS. After lysis, the sample was incubated with 700 μL pre-warmed ONPG

solution (1mg/mL ONPG (o-Nitrophenyl-β-D-Galactopyranoside, Sigma) prepared in Buffer Z

without β-mercaptoethanol at 28°C until a yellow colour developed in the samples without

auxin treatment. After stopping all reactions (using 500 μL Na_2CO_3) the supernatant was

collected and OD_{405} determined. The β-Galactosidase activity was calculated as follows:

(A405*1000)/(A600*min*mL)

For co-immunoprecipitation, ETT-FLAG was generated using Golden Gate cloning[39] by

recombining a previously described L0 clone for ETT with a 35S promoter (AddGene

#50266), a C-terminal 3xFLAG epitope (AddGene #50308) and a Nos-terminator (AddGene

#50266) into a L1 vector (AddGene #48000). The pGWB14 TPL-HA construct was provided

by Salomé Prat and has been used in previous studies[42]. The epitope-tagged proteins were

transiently expressed in four-week-old *N. benthamiana* leaves for two days. Co-

immunoprecipitation was performed as described previously[43]. After harvest, 1 g of fresh leaf

tissue was ground in liquid nitrogen. The powder was homogenised for 30 min in two

volumes of extraction buffer (10% glycerol, 25 mM Tris-HCl pH 7.5, 1 mM EDTA, 150 mM

NaCl, 0.15% NP-40, 1mM PMSF, 10 mM DTT, 2% Polyvinylporrolidone, 1x cOmplete Mini

tablets EDTA-free Protease Inhibitor Cocktail (Roche). The homogenised samples were

cleared by centrifugation at 14,000 xg for 10 min and cleared lysates were incubated for 2h

with 20 µl anti-FLAG M2 magnetic beads (SIGMA-ALDRICH, M8823; lot: SLB2419). The

beads were washed five times with IP buffer (10% glycerol, 25 mM Tris-HCl pH 7.5, 1 mM

EDTA, 150 mM NaCl, 0.15% NP-40, 1 mM PMSF, 1 mM DTT, 1x cOmplete Mini tablets

EDTA-free Protease Inhibitor Cocktail (Roche)) and proteins were eluted by adding 80 µl 2x

SDS loading buffer followed by an incubation at 95 °C for 10 min. To examine auxin

sensitivity 4 g of fresh leaf tissue was collected, ground in liquid nitrogen and protein was

extracted. The lysate was then divided according to the number of treatments. The desired

concentration of IAA or NAA was added to each of the cleared lysates before the anti-FLAG

M2 magnetic beads were added. IAA or NAA at the desired concentration was also

supplemented to the IP buffer during the washes. The eluates were analysed by western

blot using an anti-FLAG antibody (M2, Abcam, ab49763, Lot: GR3207401-3) or an anti-HA

antibody (Abcam, ab173826, Lot: GR3255539-1). Both antibodies were used as 1:10000

dilutions. The antibodies were validated by the manufacturer.

Scanning electron microscopy

Whole inflorescences of *Col-0*, *ett-3*, *tpl tpr2^{ge}* and *hda19-4* were fixed overnight in FAA

(3.7% formaldehyde, 5% glacial acetic acid, 50% ethanol) and dehydrated through an

ethanol series (70% to 100%) as described previously[31]. The samples were then critical

point-dried, gynoecia dissected and mounted. After gold coating samples were examined

with a Zeiss Supra 55VP Field Emission Scanning electron microscope using an

acceleration voltage of 3 kV.

***TPL, TPR2* and *TPR4* reporter lines**

For the construction of the promoter:GUS reporter plasmids of *TPL*, *TPR2* and *TPR4*, 2.5 kb

of promoter sequences were isolated from genomic DNA and inserted upstream of the ß-

glucoronidase gene of pCambia1301 vectors using the In-Fusion Cloning Recombinase kit

(Clontech). The constructs were transformed into *Agrobacterium tumefaciens* strain GV3101

by electroporation, followed by plant transformation by floral dip into *Col-0*[41].

The GUS histochemical assay was performed in at least three individual lines per construct.

Inflorescences of each GUS line were pre-treated with ice cold acetone for 1h at -20°C and

washed two times for 5 minutes with 100 mM sodium phosphate buffer followed by one

wash with sodium phosphate buffer containing 1 mM $K_3Fe(CN)_6$ and 1 mM $K_4Fe(CN)_6$ (both

Sigma) at room temperature. Subsequently, samples were vacuum infiltrated for 5 minutes

with X-Gluc solution (100 mM sodium phosphate buffer, 10 mM EDTA, 0.5 mM $K_3Fe(CN)_6$, 3

mM $K_4Fe(CN)_6$, 0.1% Triton X100) containing 1 mg/ml of ß-glucoronidase substrate X-gluc

(5-bromo-4-chloro-3-indolylglucuronide, Melford) and incubated at 37°C. *pTPL:GUS* were

incubated for 20 minutes and *pTPR2:GUS* lines for 45 minutes to prevent overstaining.

pTPR4:GUS lines were incubated for 16h. After staining, the samples were washed in 70%

ethanol until chlorophyll was completely removed. Gynoecia were dissected and mounted in

chloral hydrate (Sigma). Samples were analysed using a Leica DM6000 light microscope.

Chromatin Immunoprecipitation

Transcription factor ChIP was performed in triplicate using the *pETT:ETT:GFP*,

pTPL:TPL:GFP and *p35S:HDA19-GFP* lines and data analysed as described previously[44].

Additionally, a *WUS* promoter fragment was used as a negative control for ETT binding[45]. IP

was conducted using the anti-GFP antibody (Roche, 11814460001, Lot: 19958500) and

Pierce Protein G magnetic beads (ThermoFisher, 88847, Lot: SI253639) were used for IP.

Histone acetylation ChIP was carried out and data were analysed as described previously[46].

The experiment was carried out in triplicate using 3 g auxin-treated or untreated *Col-0* or *ett-*

3 inflorescent tissue. The antibodies used for IP were anti-H3K27ac antibodies (Abcam,

ab4729, Lot: GR3231937-1) and anti-H3 (Abcam, ab1791, Lot: GR310541-1). All antibodies

were validated by the manufacturers.

In all ChIP experiments, DNA enrichment was quantified using quantitative PCR (qPCR)

with the appropriate primers (Supplementary Data). In case of H3K27ac, *ACTIN* was used

as an internal control and the data represented as ratio of (H3K27ac at *HEC1* or *PID* divided

by H3 at *HEC1or PID*) to H3K27ac at *ACT* divided by H3 at *ACT*).

Statistical analyses and replication

In all graphs error bars represent the standard deviation of the mean for all numerical

values. QRT-PCR and ChIP experiments have been carried out at least in triplicate. The

data presented here show an average of three replicates. For qRT-PCR data were analysed

using one-way ANOVA with post-hoc Tukey multiple comparison test. ChIP$_{qPCR}$ data were

analysed using two-way ANOVA with post hoc Bonferroni multiple comparison test. All

output of statistical tests can be found in the source data files. All statistical tests were

carried out using GraphPad Prism Version 5.04 (La Jolla California USA,

www.graphpad.com).

Protein production

The ES domain, ES$^{388-594}$, protein was isotopically labelled in preparation for NMR analysis.

The ES domain was expressed for as a fusion protein with a 6x Histidine tag in minimal

media with ^{15}N ammonium chloride. The ^{15}N isotope labelling of the expressed protein

involved a 125-fold dilution of cell culture in enriched growth media into minimal media with

^{15}N ammonium chloride and grown for 16 hours (37 ˚C / 200 rpm); followed by a further 40-

fold dilution into minimal media for the final period of cell growth and protein expression

(induced with L-arabinose 0.2 % w/v / 18 ˚C / 200 rpm and grown for a further 12 hours).

The fusion protein was isolated from soluble cell lysate by Co-NTA affinity chromatography

with two His-Trap 1 mL TALON Crude columns (GE Healthcare Life Sciences, 28953766).

Chromatography buffers contained sodium phosphate 20 mM pH 8.0, NaCl 500 mM and

either no-imidazole or 500 mM imidazole for wash and elution buffers respectively. The

majority of the non-specifically bound protein was removed by passing 20 mL of the wash

buffer through the columns. The protein eluted on a gradient of increasing imidazole

concentration of up to 30% elution buffer over 20 mL.

HSQC NMR

The ES domain, ES$^{388\text{-}594}$, protein was analysed by NMR at 5°C under reducing conditions

(DTT 10 mM), buffered at pH 8.0 (Tris 20 mM). ^{1}H-^{15}N HSQC was performed at 950 MHz,

TCI probe, Bruker following the parameters described in Figure 1-figure supplement 1.

Isothermal titration calorimetry (ITC)

ITC was carried out on a MicroCal PEAQ-ITC (Malvern) at 25 °C in a Buffer A (sodium

phosphate 20 mM, pH 8.0; NaCl 500 mM). Ligand (2 mM IAA) was injected (19 × 4.0 μl) at

150-s intervals into the stirred (500 rpm) calorimeter cell (volume 270 μl) containing 50 μM

ES$^{388\text{-}594}$ protein. Titration of Buffer A into 50 μM ES$^{388\text{-}594}$ protein and IAA (2 mM) into Buffer A

served as negative controls. Measurements of the binding affinity of all the titration data were

analysed using the MicroCal Software (Malvern).

Accessions

ETT, AT2G33860; TPL, AT1G15750; TPR1, AT1G80490, TPR2, AT3G16830; TPR3,

AT5G27030; TPR4, AT3G15880; HDA6, AT5G63110; HDA19, AT4G38130; HEC1,

AT5G67060; PID, AT2G34650; WUS, AT2G17950.

Acknowledgements

We are grateful to Yuli Ding, Yang Dong, Emilie Knight, Bhavani Natarajan, Mikhaela Neequaye, Nicola Stacey, Sophia Stavnstrup, Billy Tasker-Brown for critical comments on the manuscript, to Keiko Torii and Shinya Hagihara for the ccvTIR1 line and cvxIAA ligand, to Rebecca Mosher for assistance with the phylogenetic analysis of ETT protein sequences, to Thomas Laux for *TPL::TPL-GFP* and *35S::HDA19-GFP* lines and to Salomé Prat for *35S::TPL-HA* construct. We acknowledge Norwich Research Park Bioimaging for skillful assistance and the NMR facility in the Astbury Centre, Faculty of Biological Sciences for access to the 950 MHz and 600 MHz spectrometers funded by the University of Leeds. We thank Arnout Kalverda for assistance with analysing the NMR data.

Author Contributions

A.K. and L.Ø. conceived the experiments. A.K., S.R.H. and H.M.M. performed the experiments. A.K., S.R.H., H.M.M., S.K. and L.Ø. analysed the data. A.K. and L.Ø. wrote the manuscript and S.R.H., H.M.M. and S.K. commented on it. All authors read and approved the manuscript.

Funding

This work was supported by grant BB/S002901/1 to L.Ø., BB/L010623/1 to S.K., the UKRI Biotechnology and Biological Sciences Research Council Norwich Research Park Biosciences Doctoral Training Partnership [grant number BB/M011216/1 to A.K.], rotation PhD studentship from the John Innes Foundation to H.M.M. and by the Institute Strategic Programme grant (BB/J004553/1) to the John Innes Centre all from the UKRI Biotechnological and Biological Sciences Research Council.

Competing Interests

No competing interests declared.

    FIGURE LEGENDS

    Figure 1. ETT directly binds auxin (IAA).

    a, HSQC-NMR performed with ES$^{388-594}$ protein either alone (black), with indole-3-acetic acid

    (IAA, orange) or benzoic acid (BA, blue). **b,** zoom-in of the indicated rectangular region in a.

    c, zoom-in of the specific shifts (labelled I-V) in the indicated dotted rectangles in a and b.

    Changes in chemical shifts are indicated by arrows from control to IAA treatment. **d-f,** ITC

    spectre showing heat exchange between ES$^{388-594}$ protein and IAA (**d**), but not in controls

    (**e,f**). See Figure 1-figure supplement 1 for parameters used in the HSQC-NMR experiment.

    Figure 2. ETT regulates target gene expression independently of TIR1/AFB auxin

    receptors.

    Expression of the canonical auxin responsive *IAA19* gene (**a**) and the ETT-target genes

    HEC1 (**b**) and *PID* (**c**) in control-treated or 100 µM IAA-treated gynoecia assayed using qRT-

    PCR. **a,** *IAA19* expression is up-regulated in response to auxin in wild-type gynoecia (*Col-0*)

    but not in *tir1/afb* double and triple mutants. The ETT-target genes *HEC1* and *PID* are up-

    regulated in response to auxin in both wild-type and auxin receptor mutants (**b, c**). This

    suggests a TIR1/AFB independent regulation of these genes. **d,** Expression of *IAA19*, *HEC1*

    and *PID* in response to treatment with 100 µM IAA and 100 µM cvxIAA in wild-type (*Col-0*)

    and *pTIR1:ccvTIR1* gynoecia in the *tir1 afb2* double mutant (*ccvTIR1*). The data confirm

    TIR1/AFB independent regulation of *HEC1* and *PID* in the gynoecium. ***p <0.0001; Shown

    are mean ± standard deviation of three biological replicates. See Figure 2-source data 1 for

    statistical analyses.

    Figure 3. ETT interacts with members of the TPL/TPR co-repressor family in an auxin-

    sensitive manner.

    a, Schematic representation of ETT protein highlighting an EAR-like motif in the C-terminal

    ETT-specific domain. **b,** Y2H showing that ETT interacts with TPL, TPR2 and TPR4. These

interactions depend on the identified C-terminal RLFGF motif and are auxin-sensitive. DBD,

DNA-binding domain. **c,** Co-IP revealing that ETT interacts with TPL in an auxin-sensitive

manner with increasing IAA concentrations weakening the interaction.

Figure 4. ETT, TPL/TPR2 and HDA19 co-operatively regulate gene expression to

facilitate gynoecium development.

a-d, Promoter GUS expression analysis of *pETT:GUS* (**a**), *pTPL:GUS* (**b**), *pTPR2:GUS* (**c**)

and *pTPR4:GUS* (**d**) revealed that *ETT*, *TPL* and *TPR2* but not *TPR4* are co-expressed in

the Arabidopsis style. Scale bar = 300 μm. **e-h,** Gynoecium phenotypes of wild-type (**e**), *ett-*

3 (**f**) *tpl tpr2^{ge}* (**g**) and *hda19-4* (**h**). Scale bar = 100 μm. **i, j,** *HEC1* (**i**) and *PID* (**j**) are

constitutively mis-regulated in *ett-3*, *tpl tpr2^{ge}* and *hda19-4* gynoecia. This misregulation is

unaffected by treatment with 100 μM IAA. ***p-Values<0.0001; Shown are mean ± standard

deviation of three biological replicates. Differences between untreated and IAA-treated

mutants are not significant. See Figure 4-source data 1 for statistical analyses.

Figure 5. ETT, TPL and HDA19 co-operatively regulate *HEC1* and *PID* by modulation of

chromatin acetylation.

A, Chromatin immunoprecipitation (ChIP) shows ETT, TPL and HDA19 binding to conserved

regions of *HEC1* and *PID* loci. *WUS* served as negative control. **b, c,** H3K27ac accumulation

(from ChIP analysis) along the *HEC1* (**b**) and *PID* (**c**) loci in wild-type (*Col-0*) and *ett-3* plants

± treatment with 100 μM IAA. Numbers on the x axes are distances to the Transcription Start

Site (TSS). The schematic of the loci is shown below each panel. Dashed boxes represent

ETT binding regions. **d,** Schematic model illustrating alternative TIR1/AFB independent

auxin signaling. Under low auxin conditions an ETT-TPL-HDA19 complex binds to ETT-

target genes keeping their chromatin environments repressed, through de-acetylation. High

nuclear auxin concentrations abolish the ETT-TPL-HDA19 complex through direct ETT-

auxin interaction. This leads to an accumulation of histone acetylation and up-regulation of

ETT-target genes.

Values in **a**, **b** and **c** are means ± standard deviation of three biological replicates. See

Figure 5-source data 1 for statistical analyses.

Figure 1-figure supplement 1. Parameters for HSQC NMR experiment.

Figure 2-figure supplement 1. Expression of *HEC1* and *PID* in Col-0 and *ett-3*.

In wild-type gynoecia *HEC1* and *PID* are up-regulated upon auxin treatment while both

genes are constitutively up-regulated in *ett-3*. Treatment with 100 µM IAA does not affect

HEC1 and *PID* expression in the *ett-3* mutant suggesting that ETT acts as a transcriptional

repressor. Asterisks indicate significant change upon auxin treatment compared to untreated

Col-0 (*** indicating $p < 0.0001$). Shown are mean ± standard deviation of three biological

replicates. See Figure 2-source data 1 for statistical analyses.

Figure 3-figure supplement 1. ETT can interact with several members of the TPL/TPR

co-repressor family through a conserved EAR-like motif.

a, Alignment of ETT protein sequences of 22 species identified a conserved repressive motif

(RLFGF) at its c-terminal domain. **b,** ETT interacts with several members of the TPL/TPR

co-repressor family in Y2H. Additionally, controls for Fig. 2 are shown.

Figure 3-figure supplement 2. Interaction between ETT and TPL, TPR2 and TPR4 is

auxin-sensitive and specific to IAA.

a, In Y2H increasing concentrations of IAA lead to reduction of yeast growth abolishing the

interaction between ETT and its partners. The interactions are, therefore, auxin-sensitive. **b,**

Y2H to test specificity of auxin-sensitivity using benzoic acid (BA), NAA, and 2,4D in a yeast

growth assay. The data suggest that the auxin-sensitivity observed in (**a**) is IAA-specific. **c,**

Y2H based ONPG assay measuring the β-galactosidase activity as a measure of interaction

strength. **d,** Co-IP experiments show that the interaction between ETT and TPL cannot be

disrupted by NAA. The data support that the ETT TPL/TPR interactions are sensitive to IAA

but not to NAA. ***p <0.0001; Shown are mean ± standard deviation of three biological

replicates. See Figure 3-source data 1 for statistical analyses.

Figure 3-figure supplement 3. Original western blot images.

The red boxes indicate the areas used in Figure 3b and Figure 3-figure supplement 2d.

Figure 4-figure supplement 1. Expression of *TPL, TPRs* and *HDAs* genes in the

gynoecium.

Expression analysis using qRT-PCR in wild-type gynoecia showed that *TPL* and *TPR2* are

more strongly expressed than *TPR1,3* and *4*. Likewise, *HDA19* exhibits higher expression

compared to *HDA6*. ***p-Values<0.0001; Shown are mean ± standard deviation of three

biological replicates. See Figure 4-source data 1 for statistical analyses.

Source data

Figure 1-source data 1. Parameters for HSQC NMR.

Figure 2-source data 1. Output of statistical tests

Figure 3-source data 1. Output of statistical tests

Figure 4-source data 1. Output of statistical tests

Figure 5-source data 1. Output of statistical tests

Supplementary Data. Oligonucleotides used in this study.

REFERENCES

1. Tsai, M. J. & O'Malley, B. W. Molecular mechanisms of action of steroid/thyroid receptor

superfamily members. *Ann. Rev. Biochem.* **63,** 451-486 (1994).

2. Gammons, M. & Bienz, M. Multiprotein complexes governing Wnt signal transduction.

Curr. Opin. Cell Biol. **51,** 42-49 (2018).

3. Kelley, D.R. & Estelle, M. Ubiquitin-mediated control of plant hormone signaling. *Plant*

Physiol. **160,** 47-55 (2012).

4. Simonini, S. et al. A noncanonical auxin-sensing mechanism is required for organ

morphogenesis in Arabidopsis. *Genes Dev.* **30,** 2286-2296 (2016).

5. Simonini, S., Bencivenga, S., Trick, M. & Østergaard, L. Auxin-induced modulation of ETTIN

activity orchestrates gene expression in Arabidopsis. *Plant Cell* **29,** 1864-1882 (2017).

6. Mutte, S. K., Kato, H., Rothfels, C., Melkonian, M., Wong, G. K. & Weijers, D. Origin and

evolution of the nuclear auxin response sytem. *eLife* doi: 10.7554/elife.33399 (2018).

7. Hironaka, K. & Morishita, Y. Encoding and decoding of positional information in

morphogen-dependent patterning. *Curr. Opin. Genet. Dev.* **22,** 553-561 (2012).

8. Vanneste, S. & Friml, J. Auxin: a trigger for change in plant development. *Cell* **136,** 1005-

1016 (2009).

9. Kepinski, S. & Leyser, H. M. O. The Arabidopsis F-box protein TIR1 is an auxin receptor.

Nature **435,** 446-451 (2005).

10. Dharmasiri, N., Dharmasiri, S. & Estelle, M. The F-box protein TIR1 is an auxin receptor.

Nature **435,** 441-445 (2005).

11. Leyser, O. Auxin signaling. *Plant Physiol.* **176,** 465-479 (2018).

12. Weijers, D. & Wagner, D. Transcriptional responses to the auxin hormone. *Ann. Rev. Plant*

Biol. **67,** 539-574 (2016).

13. Sessions, R. A. et al. ETTIN patterns the Arabidopsis floral meristem and reproductive

organs. *Development* **124,** 4481-4491 (1997).

14. Simonini, S., Mas, P. J., Mas, C. M. V. S., Østergaard, L. & Hart, D. J. Auxin sensing is a

property of an unstructured domain in the auxin response factor ETTIN of *Arabidopsis*

thaliana. *Scientific Reports* **8:**13563 (2018).

15. Meyer, B. & Peters, T. NMR spectroscopy techniques for screening and identifying

ligand binding to protein receptors. *Angew. Chem, Int.* Ed. 42, 864 (2003).

16. Benjamins, R., Quint, A., Weijers, D., Hooykaas, P. & Offringa, R. The PINOID protein

kinase regulates organ development in *Arabidopsis* by enhancing polar auxin transport.

Development **128**, 4057-67 (2001).

17. Gremski, K, Ditta, G. & Yanofsky, M. F. The *HECATE* genes regulate female

reproductive tract development in *Arabidopsis thaliana*. *Development* **134,** 3593-3601

(2007).

18. Uchida, N. et al. Chemical hijacking of auxin signalling with an engineered auxin-TIR1

pair. *Nat. Chem. Biol.* **14,** 299-305 (2018).

19. Szemenyei, H., Hannon, M. & Long. TOPLESS mediates auxin-dependent

transcriptional repression during Arabidopsis embryogenesis. *Science* 319, 1384-1386

(2008).

20. Krogan, N. T., Hogan, K. & Long, J. A. APETALA2 negatively regulates multiple floral

organ identity genes in Arabidopsis by recruiting the co-repressor TOPLESS and the histone

deacetylase HDA19. *Development* **139,** 4180-4190 (2012).

21. Long, J. A., Ohno, C., Smith, Z. R. & Meyerowitz, E. M. TOPLESS regulates embryonic

fate in Arabidopsis. *Science* **312,** 1520-1523 (2006).

22. Smith, Z. R. & Long, J. A. Control of Arabidopsis apical-basal embryo polarity by

antagonistic transcription factors. *Nature* **464,** 423-426 (2010).

23. Causier, B., Ashworth, M., Guo, W. & Davies, B. The TOPLESS interactome: a

framework for gene expression in Arabidopsis. *Plant Physiol.* **158,** 423-438 (2012).

24. Garcia, D., Collier, S. A., Byrne, M. E. & Martienssen, R. A. Specification of leaf polarity

in Arabidopsis via the trans-acting siRNA pathway. *Curr. Biol.* **16,** 933-938 (2006).

25. Marin, E. *et al.* miR390, Arabidopsis TAS3 tasiRNAs, and their AUXIN RESPONSE

FACTOR targets define an autoregulatory network quantitatively regulating lateral root

growth. *Plant Cell* **22,** 1104-1117 (2010).

26. Kelley, D.R. et al. ETTIN (ARF3) physically interacts with KANADI proteins to form a

functional complex essential for integument development and polarity determination in

Arabidopsis. *Development* **139,** 1105-1109 (2012).

27. Pekker, I., Alvarez, J. P. & Eshed, Y. Auxin response factors mediate Arabidopsis organ

asymmetry via modulation of KANADI activity. *Plant Cell* **17,** 2899-2910 (2005).

28. Sessions, R.A. & Zambryski, P. C. Arabidopsis gynoecium structure in the wild and in

ettin mutants. *Development* **121,** 1519-1532 (1995).

29. Nemhauser, J., Feldmann, L. J. & Zambryski, P. C. Auxin and ETTIN in *Arabidopsis*

gynoecium morphogenesis. *Development* **127,** 3877-3888 (2000).

30. Chung, Y. et al. Auxin response factors promote organogenesis by chromatin-mediated

repression of the pluripotency gene SHOOTMERISTEMLESS. *Nature Communications*

10:886 (2019).

31. Moubayidin, L. & Østergaard, L. Dynamic control of auxin distribution imposes a

bilateral-to-radial symmetry switch during gynoecium development. *Curr. Biol.* **24,** 2743-

2748 (2014).

32. Kim, K.-C., Lai, Z., Fan, B. & Chen, Z. Arabidopsis WRKY38 and WRKY62 Transcription

Factors Interact with Histone Deacetylase 19 in Basal Defense. *Plant Cell* **20,** 2357-2371

(2008).

33. Ng, K. H., Yu, H. & Ito, T. AGAMOUS controls GIANT KILLER, a multifunctional

chromatin modifier in reproductive organ patterning and differentiation. *PLoS Biol.*

7:e1000251 (2009).

34. Pi, L. et al. Organizer-Derived WOX5 Signal Maintains Root Columella Stem Cells

through Chromatin-Mediated Repression of CDF4 Expression. *Dev. Cell* **33,** 576-588 (2015).

35. Parry, G. et al. Complex regulation of the TIR1/AFB family of auxin receptors. *Proc. Natl.*

Acad. Sci. USA **106,** 22540-22545 (2009).

36. Livak, K. J. & Schmittgen, T. D. Analysis of Relative Gene Expression Data Using Real-

Time Quantitative PCR and the 2−ΔΔCT Method. *Methods* **25,** 402-408 (2001).

37. Rokhsar, D. S. et al. Phytozome: a comparative platform for green plant genomics.

Nucleic Acids Research **40,** D1178-D1186 (2011).

38. Kearse, M. et al., Geneious Basic: An integrated and extendable desktop software

platform for the organization and analysis of sequence data. *Bioinformatics* **28,** 1647-1649

(2012).

39. Castel, B., Tomlinson, L., Locci, F., Yang, Y. & Jones, J. D. G. Optimization of T-DNA

architecture for Cas9-mediated mutagenesis in Arabidopsis. *PLoS One* **14,** e0204778

(2019).

40. Engler, C. et al. A Golden Gate Modular Cloning Toolbox for Plants. *Acs Synthetic*

Biology **3,** 839-843 (2014).

41. Clough, S. J. & Bent, A. F. Floral dip: a simplified method for Agrobacterium -mediated

transformation of Arabidopsis thaliana. *Plant Journal* **16,** 735-743 (1998).

42. Espinosa-Ruiz, A., Martínez, C., de Lucas, M., Fàbregas, N., Bosch, N., Caño-Delgado,

A. & Prat, S. TOPLESS mediates brassinosteroid control of shoot boundaries and root

meristem development in *Arabidopsis thaliana*. *Development* **144,** 1619-1628 (2017).

43. Egea-Cortines, M., Saedler, H. & Sommer, H. Ternary complex formation between the

MADS-box proteins SQUAMOSA, DEFICIENS and GLOBOSA is involved in the control of

floral architecture in Antirrhinum majus. *EMBO Journal* **18,** 5370-5379 (1999).

44. Schiessl, K., Muiño, J. M. & Sablowski, R. Arabidopsis JAGGED links floral organ

patterning to tissue growth by repressing Kip-related cell cycle inhibitors. *Proc. Nat. Acad.*

Sci. USA **111,** 2830-2835 (2014).

45. Liu, X. et al. AUXIN RESPONSE FACTOR 3 integrates the functions of AGAMOUS and

APETALA2 in floral meristem determinacy. *Plant Journal* **80,** 629-641 (2014).

46. Questa, J. I., Song, J., Geraldo, N., An, H. L. & Dean, C. Arabidopsis transcriptional

repressor VAL1 triggers Polycomb silencing at FLC during vernalization. *Science* **353,** 485-

488 (2016).

Figure 1. ETT directly binds auxin (IAA).

Figure 1. ETT directly binds auxin (IAA).
a, HSQC-NMR performed with ES[388-594] protein either alone (black), with indole-3-acetic acid (IAA, orange) or benzoic acid (BA, blue). **b,** zoom-in of the indicated rectangular region in a. **c,** zoom-in of the specific shifts (labelled I-V) in the indicated dotted rectangles in a and b. Changes in chemical shifts are indicated by arrows from control to IAA treatment. **d-f,** ITC spectre showing heat exchange between ES[388-594] protein and IAA (**d**), but not in controls (**e,f**). See Figure 1-figure supplement 1 for parameters used in the HSQC-NMR experiment.

Figure 2. ETT regulates target gene expression independently of TIR1/AFB auxin receptors.

Figure 2. ETT regulates target gene expression independently of TIR1/AFB auxin receptors.
Expression of the canonical auxin responsive *IAA19* gene (**a**) and the ETT-target genes *HEC1* (**b**) and *PID* (**c**) in control-treated or 100 µM IAA-treated gynoecia assayed using qRT-PCR. **a,** *IAA19* expression is up-regulated in response to auxin in wild-type gynoecia (*Col-0*) but not in *tir1/afb* double and triple mutants. The ETT-target genes *HEC1* and *PID* are up-regulated in response to auxin in both wild-type and auxin receptor mutants (**b, c**). This suggests a TIR1/AFB independent regulation of these genes. **d,** Expression of *IAA19, HEC1* and *PID* in response to treatment with 100 µM IAA and 100 µM cvxIAA in wild-type (*Col-0*) and *pTIR1:ccvTIR1* gynoecia in the *tir1 afb2* double mutant (*ccvTIR1*). The data confirm TIR1/AFB independent regulation of *HEC1* and *PID* in the gynoecium. ***p <0.0001; Shown are mean ± standard deviation of three biological replicates. See Figure 2-source data 1 for statistical analyses.

Figure 3. ETT interacts with members of the TPL/TPR co-repressor family in an auxin-sensitive manner.

Figure 3. ETT interacts with members of the TPL/TPR co-repressor family in an auxin-sensitive manner.
a, Schematic representation of ETT protein highlighting an EAR-like motif in the C-terminal ETT-specific domain. **b,** Y2H showing that ETT interacts with TPL, TPR2 and TPR4. These interactions depend on the identified C-terminal RLFGF motif and are auxin-sensitive. DBD, DNA-binding domain. **c,** Co-IP revealing that ETT interacts with TPL in an auxin-sensitive manner with increasing IAA concentrations weakening the interaction.

Fig. 4. ETT, TPL/TPR2 and HDA19 cooperatively regulate gene expression to facilitate gynoecium development.

Figure 4. ETT, TPL/TPR2 and HDA19 co-operatively regulate gene expression to facilitate gynoecium development.
a-d, Promoter GUS expression analysis of *pETT:GUS* (**a**), *pTPL:GUS* (**b**), *pTPR2:GUS* (**c**) and *pTPR4:GUS* (**d**) revealed that *ETT, TPL* and *TPR2* but not *TPR4* are co-expressed in the Arabidopsis style. Scale bar = 300 µm. **e-h,** Gynoecium phenotypes of wild-type (**e**), *ett-3* (**f**) *tpl tpr2^ge* (**g**) and *hda19-4* (**h**). Scale bar = 100 µm. **i, j,** *HEC1* (**i**) and *PID* (**j**) are constitutively mis-regulated in *ett-3, tpl tpr2^ge* and *hda19-4* gynoecia. This misregulation is unaffected by treatment with 100 µM IAA. ***p-Values<0.0001; Shown are mean ± standard deviation of three biological replicates. Differences between untreated and IAA-treated mutants are not significant. See Figure 4-source data 1 for statistical analyses.

Fig. 5. ETT, TPL and HDA19 co-operatively regulate *HEC1* and *PID* by modulation of chromatin acetylation.

Figure 5. ETT, TPL and HDA19 co-operatively regulate *HEC1* and *PID* by modulation of chromatin acetylation.
A, Chromatin immunoprecipitation (ChIP) shows ETT, TPL and HDA19 binding to conserved regions of *HEC1* and *PID* loci. *WUS* served as negative control. **b, c,** H3K27ac accumulation (from ChIP analysis) along the *HEC1* (**b**) and *PID* (**c**) loci in wild-type (*Col-0*) and *ett-3* plants ± treatment with 100 µM IAA. Numbers on the x axes are distances to the Transcription Start Site (TSS). The schematic of the loci is shown below each panel. Dashed boxes represent ETT binding regions. **d,** Schematic model illustrating alternative TIR1/AFB independent auxin signaling. Under low auxin conditions an ETT-TPL-HDA19 complex binds to ETT-target genes keeping their chromatin environments repressed, through de-acetylation. High nuclear auxin concentrations abolish the ETT-TPL-HDA19 complex through direct ETT-auxin interaction. This leads to an accumulation of histone acetylation and up-regulation of ETT-target genes.
Values in **a**, **b** and **c** are means ± standard deviation of three biological replicates. See Figure 5-source data 1 for statistical analyses.

Figure 1-figure supplement 1. Parameters for HSQC NMR experiment.

Experiment	Recycling Delays (S)	Scans	Nuclei		Spectral width (Hz)		Number of complex points	
			t1	t2	t1	t2	t1	t2
HSQC	1	56	^{15}N	^{1}H	2888.6	15243.9	256	2048

Figure 2-figure supplement 1. Expression *HEC1* and *PID* in Col-0 and *ett-3*.

Figure 2-figure supplement 1. Expression of *HEC1* and *PID* in Col-0 and *ett-3*.
In wild-type gynoecia *HEC1* and *PID* are up-regulated upon auxin treatment while both genes are constitutively up-regulated in *ett-3*. Treatment with 100 μM IAA does not affect *HEC1* and *PID* expression in the *ett-3* mutant suggesting that ETT acts as a transcriptional repressor. Asterisks indicate significant change upon auxin treatment compared to untreated *Col-0* (*** indicating $p < 0.0001$). Shown are mean ± standard deviation of three biological replicates. See Figure 2-source data 1 for statistical analyses.

Figure 3-figure supplement 1. ETT can interact with several members of the TPL/TPR co-repressor family through a conserved EAR-like motif.

Figure 3-figure supplement 1. ETT can interact with several members of the TPL/TPR co-repressor family through a conserved EAR-like motif.
a, Alignment of ETT protein sequences of 22 species identified a conserved repressive motif (RLFGF) at its c-terminal domain. **b,** ETT interacts with several members of the TPL/TPR co-repressor family in Y2H. Additionally, controls for Fig. 2 are shown.

Figure 3-figure supplement 2. Interaction between ETT and TPL, TPR2 and TPR4 is auxin-sensitive and specific to IAA.

Figure 3-figure supplement 2. Interaction between ETT and TPL, TPR2 and TPR4 is auxin-sensitive and specific to IAA.

a, In Y2H increasing concentrations of IAA lead to reduction of yeast growth abolishing the interaction between ETT and its partners. The interactions are, therefore, auxin-sensitive. **b,** Y2H to test specificity of auxin-sensitivity using benzoic acid (BA), NAA, and 2,4D in a yeast growth assay. The data suggest that the auxin-sensitivity observed in (**a**) is IAA-specific. **c,** Y2H based ONPG assay measuring the β-galactosidase activity as a measure of interaction strength. **d,** Co-IP experiments show that the interaction between ETT and TPL cannot be disrupted by NAA. The data support that the ETT TPL/TPR interactions are sensitive to IAA but not to NAA. ***p <0.0001; Shown are mean ± standard deviation of three biological replicates. See Figure 3-source data 1 for statistical analyses.

Figure 3-figure supplement 3. Original western blot images.

Figure 3-figure supplement 3. Original western blot images.
The red boxes indicate the areas used in Fig. 3b and Figure 3-figure supplement 2.

Figure 4-figure supplement 1. Expression TPL TPRs and HDAs in the gynoecium.

Figure 4-figure supplement 1. Expression of *TPL, TPRs* and *HDAs* genes in the gynoecium.
Expression analysis using qRT-PCR in wild-type gynoecia showed that *TPL* and *TPR2* are stronger expressed then *TPR1,3* and *4*. Likewise, *HDA19* exhibits higher expression compared to *HDA6*. ***p-Values<0.0001; Shown are mean ± standard deviation of three biological replicates. See Figure 4-source data 1 for statistical analyses.