Computation-guided optimization of split protein systems

Taylor B. Dolberg[1,2,9], Anthony T. Meger[3,4,9], Jonathan D. Boucher[2,5], William K. Corcoran[2,5], Elizabeth E.

Schauer[1,2], Alexis N. Prybutok[1,2], Srivatsan Raman[3,4,8*], Joshua N. Leonard[1,2,5,6,7*]

[1]Department of Chemical and Biological Engineering, Northwestern University, Evanston, Illinois 60208,

United States

[2]Center for Synthetic Biology, Northwestern University, Evanston, Illinois 60208, United States

[3]Department of Biochemistry, University of Wisconsin-Madison, Madison, Wisconsin 53706, United States

[4]Great Lakes Bioenergy Research Center, University of Wisconsin-Madison, Madison, Wisconsin 53706,

United States

[5]Interdisciplinary Biological Sciences Program, Northwestern University, Evanston, Illinois 60208, United

States

[6]Chemistry of Life Processes Institute, Northwestern University, Evanston, Illinois 60208, United States

[7]Member, Robert H. Lurie Comprehensive Cancer Center, Northwestern University, Evanston, Illinois

60208, United States

[8]Department of Bacteriology, University of Wisconsin-Madison, Madison, Wisconsin 53706, United States

[9]These authors contributed equally to this work

Contact Information

*Corresponding authors: Srivatsan Raman (sraman4@wisc.edu) and Joshua N. Leonard (j-

leonard@northwestern.edu)

ABSTRACT

Splitting bioactive proteins, such as enzymes or fluorescent reporters, into conditionally reconstituting

fragments is a powerful strategy for building tools to study and control biochemical systems. However, split

proteins often exhibit a high propensity to reconstitute even in the absence of the conditional trigger, which

limits their utility. Current approaches for tuning reconstitution propensity are laborious, context-specific, or

often ineffective. Here, we report a computational design-driven strategy that is grounded in fundamental

protein biophysics and which guides the experimental evaluation of a focused, sparse set of mutants—

which vary in the degree of interfacial destabilization while preserving features such as stability and catalytic

activity—to identify an optimal functional window. We validate our method by solving two distinct split

protein design challenges, generating both broad insights and new technology platforms. This method will

streamline the generation and use of split protein systems for diverse applications.

KEYWORDS: synthetic biology, split proteins, computational protein design, protein engineering

INTRODUCTION

Split proteins and conditional reconstitution systems are powerful tools for interrogating biology and controlling cell behavior.[1-4] These systems work by splitting a protein into two fragments to disrupt the protein's function. Each fragment is then fused to a partner domain such that the split protein is reconstituted, and its function is restored only when the partner domains interact. This modular strategy may be applied to diverse functional proteins to control bioluminescence[5, 6], fluorescence[7], proteolytic cleavage[8-10] and transcription[11, 12]. As a result, conditionally-reconstituted split proteins have been employed in a variety of applications including probing and discovering new protein-protein interactions[13-16], studying post-translational modifications[17], imposing small molecule-regulated control over enzymatic activity[18, 19], and rewiring cellular signaling[9, 20].

Despite their utility in certain contexts, broader application of split protein systems is largely limited by the spontaneous reconstitution of fragments, resulting in high background activity (**Fig. 1a**). Splitting a protein tends to expose its hydrophobic core, creating highly unfavorable interactions between the core and solvent. Reconstitution is driven by a strong inherent preference to desolvate by recombining the fragments. Evaluating alternative splitting sites can vary reconstitution propensity, but this approach often only partially ameliorates the problem because changing splitting sites may not significantly affect underlying hydrophobic forces. Therefore, it is necessary to identify variants with a reconstitution propensity that precludes spontaneous reconstitution but enables reconstitution under desired conditions. Variants with a range of reconstitution propensities can be generated by random mutagenesis and screened for the desired property. However, high-throughput screening is not readily available for all split protein systems, and low-throughput clonal testing of variants can be laborious and suffer from inefficient exploration of sequence space. Even when screens generate improved variants, it may be difficult to interpret why only certain mutants were successful, and as a result, generalizable rules cannot be transferred to guide the tuning of new split protein systems. Furthermore, split protein systems tuned by mutagenesis exhibit performance characteristics determined by (and limited to) the conditions used in the initial screens, again posing a barrier to general applications.

Here, we report a general strategy based on fundamental principles of protein biophysics for optimizing

split protein systems which we term **S**plit **P**rotein **O**ptimization by **R**econstitution **T**uning, or SPORT. We

use computational mutagenesis with the Rosetta macromolecular modeling suite to guide limited

experimental screening and thereby discretely map the sequence-energy landscape of the split interface.

This allows us to determine optimal interaction energies that maximize the performance of the split protein

system. We demonstrate proof-of-concept by optimizing a split protease system for conditional

reconstitution in two different contexts: membrane-embedded and cytosolic. Our approach generates

simple design rules that may be extended to tune other split protein systems for distinct design goals and

can be implemented by most research laboratories. This work demonstrates a new method for efficiently

engineering split protein systems, which will streamline the generation and expand the use of split protein

systems for diverse applications.

RESULTS

Formulation of the design challenge and strategy

As a first step toward developing a strategy that addresses the challenge of designing split proteins, we

selected a model system based on the well-studied Tobacco Etch Virus protease (TEVp)—we sought to

tune the reconstitution propensity of split TEVp. For this purpose, we modified a synthetic receptor system

that we previously reported (Modular Extracellular Split Architecture, or MESA)[21] to serve as a reporter of

conditional split TEVp reconstitution. In this testbed, ligand binding-induced dimerization of a membrane

receptor reconstitutes an intracellular split TEVp, which then autolytically liberates a sequestered

transcription factor to drive reporter gene expression (**Fig. 1b**). Our initial evaluation demonstrated that the

canonical split TEVp (split between residues 118/119)[22] showed high propensity to reconstitute, resulting

in high background from ligand-independent signaling (**Supplementary Fig. 1**). As the original screens

used to identify this split were performed in a soluble rather than membrane-bound context, these data

suggest that tethering split TEVp to a membrane may promote reconstitution. Furthermore, we determined

that this problem is not limited to the canonical split site, as other TEVp partitioning also yielded poor

performance (**Supplementary Fig. 2**). Given these observations, we formulated a design goal: rationally

mutate split TEVp to optimize two key MESA performance characteristics—minimal reporter gene

expression in the absence of ligand and a substantial fold increase in reporter expression upon ligand

addition.

Biophysical principles underlying SPORT

We developed SPORT, a computation-guided workflow to rationally design split protein interfaces to

optimize reconstitution propensity (**Fig. 1c**). SPORT employs Rosetta, a state-of-the-art software package

for protein design.[23] Given a protein with a predetermined split site, our first step was to identify key

interfacial residues to target for mutagenesis. Residues with large differences in solvent-accessible surface

area (SASA)—when comparing intact protein and split fragments—were classified as buried residues.

These buried residues are ideal targets for mutagenesis as they likely contribute substantially to the driving

force for spontaneous reconstitution. For each buried residue, we performed a comprehensive, *in silico*

mutational scan to evaluate the energy perturbation of all possible single-point mutations on the interaction

energy across the split protein interface ($\Delta\Delta G_{Interfacial}$) and total stability of the mutated protein ($\Delta\Delta G_{Total}$)

relative to the parent. The degree of disruption is a critical design consideration. Insufficient disruption may

retain high background activity while excessive disruption may impair catalytic activity due to loss of overall

protein stability. Therefore, the interface must be carefully tuned so that the driving force provided by ligand

binding-induced dimerization promotes reconstitution. This "Goldilocks zone" likely differs for each

individual protein and perhaps depends upon context, and this zone is difficult to define *a priori*. Therefore,

our strategy was to identify the Goldilocks zone for a given protein by choosing mutations that span the

range of $\Delta\Delta G$ values. We hypothesized that a limited test set of mutants would direct subsequent

mutagenesis efforts by predicting desirable mutant combinations from a vast amount of sequence space.

Each of these propositions was tested using experimental case studies.

Validating SPORT by tuning a membrane-tethered split TEV protease

To investigate and validate SPORT, we applied our design workflow to the split TEVp MESA system. We

first assessed the per-residue change in solvent accessible surface area (ΔSASA) between the intact form

and split fragments (**Fig. 2a**). In total, 130 of the 218 residues showed increased SASA in the isolated

fragments. We excluded from this set the catalytic triad and 27 residues lying within a 6 Å coordination

sphere around the catalytic triad to avoid perturbing the catalytic function of reconstituted TEVp. Of the

remaining 100 positions, we chose the 15 positions with the largest ΔSASA (9 in N-terminal and 6 in C-

terminal halves of split TEVp) as candidates for mutagenesis. Next, we evaluated the energy perturbation

of all possible single-point mutations (285 in total) at these positions using Rosetta. As expected, few

mutations were predicted to increase stability, and a vast majority were destabilizing (**Fig. 2b,**

Supplementary Figure 3). Many positions exhibited a variety of stabilizing, benign, and destabilizing point

mutations that were consistent with structure. For instance, bulky sidechain substitutions (W, F, Y, R, K and

H) at position 103 resulted in many steric clashes with neighboring residues (**Fig. 2b** right panel) and

subsequently conferred large decreases in predicted stability.

Guided by these predictions, we next experimentally characterized 20 single-mutant split TEV variants that

span a wide range of $\Delta\Delta G_{Interfacial}$ (3.1 to 16.1 Rosetta Energy Units, or REU) and $\Delta\Delta G_{Total}$ (-1.9 to 30 REU)

energies (**Fig. 2c**). We observed high background signaling (i.e., reporter expression) for disruptions up to

$\Delta\Delta G_{Interfacial}$ ~6.6 REU, which suggested that destabilization was insufficient. However, four out of ten

variants with $\Delta\Delta G_{Interfacial}$ >10 REU exhibited reduced background signaling and substantial ligand-induced

activation (**Fig. 2c**). The remaining six were completely inactive (or "dead"); they induced no signaling under

any conditions. Energy-based partitioning across different phenotypes (inducible, not inducible, and dead)

became evident when comparing $\Delta\Delta G_{Interfacial}$ and $\Delta\Delta G_{Total}$ of all 20 single mutant split TEV variants (**Fig.**

2d). Variants with high background activity due to insufficient destabilization fell in the region where

$\Delta\Delta G_{Interfacial}$ < 10 and $\Delta\Delta G_{Total}$ < 10 REU. Variants with the dead phenotype had $\Delta\Delta G_{Interfacial}$ > 10 and

$\Delta\Delta G_{Total}$ > 10 REU; since these mutants were expressed, as confirmed by Western blot (**Supplementary**

Fig. 4), the lack of signaling suggested that these mutations directly preclude reconstitution. Most of the

inducible variants (three out of four) were observed in the energy window where $\Delta\Delta G_{Interfacial}$ > 10 and

$\Delta\Delta G_{Total}$ < 10 REU, which may represent the Goldilocks zone we hypothesized to exist. An additional region

contained a mixture of inducible and dead phenotypes. By inspection of these results, we then proposed a

model for broadly classifying experimental phenotypes based on energy partitions (**Fig. 2e**).

SPORT predicts outcomes of combining mutations

We next evaluated whether our proposed classifier model—developed based upon observations with single

mutants—could predict the phenotypes of combined NTEVp and CTEVp mutations, including both double

(two mutations on one chain) and paired (one mutation on each chain) mutants derived by combining the

initial 14 single non-dead mutations tested (**Fig. 3a**). Of the 67 possible double and paired mutants tested,

28 were predicted to be inducible. We experimentally tested 14 of these and found that 10 exhibited

inducible signaling as predicted, one was dead, and three were not inducible; this yields an observed

accuracy of 0.71 for inducible predictions (**Fig. 3b,c**). Interestingly, three of the prediction failures fell at the

low end of the range of predicted changes in interfacial energy, suggesting potential opportunities for

refining the classifier model. We also noted an interesting trend in our $\Delta\Delta G$ (total and interfacial)

calculations—for the sixty-seven mutants tested, the calculated $\Delta\Delta G$ for the double and paired mutants

were nearly identical to the sums of the $\Delta\Delta G$ calculated for the associated single mutants (**Supplementary**

Fig. 5). Thus, for subsequent analyses of combined mutants, we simply added the effects of single mutants

in our calculations.

We next investigated how variations in expression level might impact the inducibility of the mutants. We

used Western blot analysis to normalize and vary chain expression levels by adjusting DNA doses

(**Supplementary Fig. 6**). Notably, these constructs remained inducible across the entire range of

expression levels tested, suggesting that the biophysical mechanism of optimized split protein reconstitution

is robust to variations in the expression levels and ratio of the membrane-bound split TEVp fragments.

However, we observed that the performance of these constructs (i.e., fold induction of signaling upon ligand

addition) could be substantially improved through tuning expression such that protein levels of each

fragment are comparable (**Fig. 3d, Supplementary Fig. 7**). Taken together, these results suggest that a

classifier calibrated with a limited set of experimental observations spanning the full range of ΔΔG can

predict function of new mutants with high accuracy in a manner that is independent of expression level of

the construct.

SPORT predicts phenotypes of novel mutations and combinations

All mutations previously tested were derived from predictions based on our computational method. Next,

we wanted to test a broad set of mutants (outside the calibration set) to investigate the accuracy of our

classification scheme. Therefore, we experimentally characterized additional single mutants (not included

in the original calibration set) and all combinations of paired mutations derived from both the original and

this expanded mutation set (omitting dead constructs); mutants were selected to explore the boundaries of

the energy landscape classifier model and were expected to reflect a wide range of induced and uninduced

reporter expression levels and ratios (**Figs. 2e**, **3a**). We also sought to investigate whether the phenotypic

partitioning demonstrated for inducibility (**Fig. 3b,c**) is extensible to the other phenotype classes (i.e., dead

and not inducible). This expanded set paired 10 N-terminal mutations with 16 C-terminal mutations. In

general, we observed that variants with larger $\Delta\Delta G_{Interfacial}$ energies had lower reporter expression levels in

both the background (OFF) and ligand-induced (ON) states (**Fig. 4a**, left and middle panels). Thus, the cost

of lowering the OFF state is to also lower the ON state, but these reductions are not always proportional.

This is evident by the diversity of calculated fold inductions (**Fig. 4a**, right panel). Only one variant,

H75P/W198E, exhibited a significant decrease in the OFF state and an increase in the ON state relative to

wild type (WT). However, the variants with highest fold inductions, such as H75S/I163P (17.3 fold induction)

and H75T/I163P (9.92 fold induction), exhibited significantly lower OFF and ON states than did the WT,

reflecting a tradeoff between desirable performance characteristics. Overall, we observed moderate

agreement between the actual and predicted phenotypes for these novel variants and combinations (**Fig.**

4b). The classifier model was most accurate for predicting the not inducible phenotype (25 of 29, 86%).

Many inducible phenotype predictions were also confirmed (31 of 52, 60%). This success rate is impressive

given that phenotypic classification boundaries were set roughly based upon the sparse calibration set (**Fig.**

2e). Interestingly, this analysis also indicates that the energy landscape calculated by SPORT correlates

with each phenotype to differing degrees.

In order to gain additional insight into how choice of calibration set and sample size may impact the accuracy

of SPORT predictions, we performed retrospective bootstrapping analysis of the data presented in **Figure**

4 (see **Supplementary Note 2** for full details). Experimental data were stratified by the energy landscape

and partitioned randomly into calibration and prediction subsets. Logistic regression modeling was applied

to evaluate the accuracy of classification under various conditions (**Supplementary Fig. 8**). We observed

robust prediction accuracy using multiple unique calibration sets and using sample sizes ranging from 4 to

28. This outcome suggests that our ability to generate a general classification model was not dependent

upon the specific calibration data we used in our initial characterization experiment (**Fig. 2**), and that a

relatively small set of calibration data drawn from a distribution like that included in **Figure 4** would be

sufficient to generate a general classification model.

Extension of SPORT predictions to new design goals

A major limitation to current approaches for employing split proteins is that often a variant selected to

perform well in one context fails in a different context. To investigate whether the SPORT design method is

generalizable beyond our initial model system, we developed a distinct model system. This new system

employs split TEVp in a soluble form, where we hypothesized that a different reconstitution propensity

would be required compared to the membrane-bound model system. To generate such a soluble test

system (**Fig. 5a**), ligand binding domains were fused to split TEVp domains along with a soluble

transcription factor flanked by protease cleavage sites and nuclear export signals (NES); thus, TEVp-

mediated cleavage removes the NES from the transcription factor to enable nuclear localization and

reporter expression. We first developed and tested a panel of soluble transcription factors that could

implement this mechanism. This evaluation included varying the number of NES elements, their placement

at N and/or C terminus of the transcription factor, and the P1' residue of the TEVp cleavage sequence

which governs cleavage kinetics[24] (**Fig. 5b**). Several soluble transcription factor constructs exhibited the

desired phenotype of low signaling in the absence of TEVp and high signaling when co-expressed with full

TEVp; construct TF10 was selected for evaluating split TEVp variants.

Using our soluble split TEVp test system, we evaluated a panel of 10 single TEVp mutants and 10 paired

TEVp mutants spanning a range of interfacial energies **(Fig. 5c)**. The construct based upon WT split TEVp

exhibited a substantial fold induction, which is consistent with the fact that this split protein was identified

by screens performed in the soluble phase[22]. However, the WT construct also yielded high background

signaling in the absence of ligand, indicating an opportunity to improve performance. For modest increases

in $\Delta\Delta G_{Interfacial}$ (~0–3.6 REU), background signaling persisted. For intermediate increases in $\Delta\Delta G_{Interfacial}$ (~6-

10 REU), a mixture of phenotypes was observed, including dead constructs and those with both

substantially reduced background signaling and substantial fold inductions. For large increase in

$\Delta\Delta G_{Interfacial}$ > ~10 REU, constructs were generally weakly-inducible or dead. Thus, a focused evaluation of

20 design variants, guided by SPORT, yielded ~5 variants exhibiting improved performance compared to

the WT construct. Altogether, these observations demonstrate that the SPORT method and associated

energy landscape concept can be employed to efficiently solve distinct split protein optimization challenges.

DISCUSSION

In this study, we developed and validated what is—to our knowledge—the first computational strategy for

tuning split protein reconstitution propensity. Although the split TEVp MESA used as our first model system

would have been deemed infeasible using standard evaluations of split proteins (**Supplementary Figs. 1-**

2), application of SPORT to tune this system yielded multiple high-performing new synthetic receptor

scaffolds (**Fig. 3d**). We show that unlike the classical MESA receptors we have characterized in prior work[21,]

[25, 26], split TEVp MESA tuned by SPORT exhibit excellent performance characteristics (i.e., low-background

and high fold-induction) in a manner that is robust to variations in both biosensor expression level and the

ratio at which biosensor chains are expressed (**Supplementary Figure 7**). This property is of great practical

utility, as it precludes the need to carefully tune the implementation of each biosensor.

Several important insights emerged from this study. First, our approach demonstrated that testing a sparse

set of mutants along the energy landscape is an effective strategy to choose optimal interfacial energies to

promote conditional reconstitution. Second, multiple point mutations with similar energies exhibited similar

performance, which suggests reconstitution propensity depends on the energy of destabilization but is

agnostic to specific mutations. Third, the concept of a Goldilocks zone is likely generalizable to different

proteins and application contexts, but the optimal energy window may have to be adjusted on a case-by-

case basis. We find that membrane-bound split proteins must be destabilized to a greater degree than

soluble split proteins in order to avoid spontaneous reconstitution. Altogether, these results show that split

protein systems can be engineered based on fundamental principles of protein biophysics, which obviates

the need for exhaustive screening and generates rules applicable to new candidate proteins.

There are several interesting opportunities for extending and improving SPORT in future work. First,

although our analysis showed that SPORT can be used to identify mutations that confer specific energy

changes, it does not yet enable *a priori* prediction of where the Goldilocks zone will fall for new applications.

It is possible that subsequent analysis of many case studies could identify trends that enable such

predictions and thus harness SPORT to further focus experimental investigations. An additional opportunity

is pairing SPORT with a multiparameteric optimization framework for exploring pareto-optimal tradeoffs

between performance characteristics; for example, in our model system, there seems to be such a tradeoff

between low background in the ligand-free state and high signaling in the ligand-induced state. Finally, the

SPORT algorithm itself may be refined to better avoid false positives (e.g., dead mutants that share a region

of the energy landscape with inducible variants). Altogether, our findings suggest many opportunities for

expanding the utility of split proteins for many new applications and highlight the impact of SPORT-guided

development of novel biochemical and synthetic biology tools.

METHODS

General DNA assembly

Plasmid cloning was performed using standard molecular biology techniques of PCR and restriction

enzyme cloning with Phusion DNA Polymerase (NEB), restriction enzymes (NEB; Thermo Fisher), T4 DNA

Ligase (NEB), and Antarctic Phosphatase (NEB). Development of the tTA-responsive YFP reporter plasmid

was described previously[21]. Plasmids were transformed into chemically competent TOP10 *E. coli* (Thermo

Fisher) and grown at 37˚C. Plasmid maps are provided as GenBank files (**Supplementary Data 1**).

Plasmid preparation

Plasmid DNA used for transfection was prepared using the PEG precipitation method, which was previously

described in detail.[27]

Cell culture

HEK293FT cells (Life Technologies/Thermo) were maintained at 37°C incubator and 5% CO_2. Cells were

cultured in DMEM (Gibco 31600-091) with 10% FBS, 6 mM L-glutamine (2 mM from Gibco 31600-091 and

4 mM from additional Gibco 25030-081), penicillin (100 U/µL), and streptomycin (100 µg/mL) (Gibco

15140122).

Transfection

Transfections were performed in 24 well plates seeded at 1.5×10^5 cell in 0.5 mL of DMEM media. At 6-8

hours post-seeding, cells were transfected using calcium phosphate method with a total DNA content of 1-

2 ug DNA per mL of media, using DNA prepared by PEG precipitation. All experiments included blue

fluorescent protein (BFP) as a control to assess transfection efficiency. The exact DNA amounts added to

the mix per well are as follows, unless otherwise stated in figure captions: 25 ng of each TEVp chain, 200

ng of BFP control, 200 ng of YFP reporter plasmid, and 150 ng of pcDNA plasmid. This mixture was added

dropwise to an equal-volume solution of 2x HEPES-Buffered Saline (280 mM NaCl, 0.5 M HEPES, 1.5 mM

Na_2HPO_4) and gently pipetted up and down four times. After 2.5 minutes, the solution was mixed vigorously

by pipetting ten times and 100 µL of this mixture was added dropwise to each well of the plated cells, and

the plates were swirled gently. For functional experiments, 12 hours post-transfection, media containing 0.1

µM rapamycin analog (Takara AP21967) or 0.1% ethanol as a control was added to cells. At 24-30 hours

post-media change, cells were harvested for flow cytometry with Trypsin-EDTA, which was then quenched

with medium, and the resulting cell solution was added to at least 2 volumes of FACS buffer (PBS pH 7.4

with 2–5 mM EDTA and 0.1% BSA). Cells were spun at 150 x g for 5 min, FACS buffer was decanted, and

fresh FACS buffer was added. All experiments were performed in biological triplicate.

Flow Cytometry

Approximately 10^4 live cells from each transfected well of the 24-well plate were analyzed using a BD LSR

Fortessa Special Order Research Product (Robert H. Lurie Cancer Center Flow Cytometry Core) running

FACSDiva software. Samples were analyzed using FlowJo v10 software (FlowJo, LLC). The HEK293FT

cell population was identified by FSC-A vs. SSC-A gating, and singlets were identified by FSC-A vs. FSC-

H gating. A control sample of cells—generated by transfecting cells with a mass of pcDNA (empty vector)

equivalent to the mass of DNA used in other samples in the experiment—was used to distinguish

transfected and non-transfected cells. For the single-cell subpopulation of the pcDNA-only sample, a gate

was made to identify cells that were positive for the constitutively driven blue fluorescent protein (BFP) used

as a transfection control in other samples such that the gate included no more than 1% of the non-

fluorescent cells. The mean fluorescence intensity (MFI) of the single-cell transfected population was

calculated and exported for further analysis. BD LSR Fortessa settings used were as follows: BFP was

collected in the Pacific Blue channel (405 nm excitation, 450/50 nm filer) and EYFP was collected in the

FITC channel (488 nm excitation, 505 LP and 530/30 nm filter). To quantify reporter expression, the FITC

channel MFI was averaged across three biological replicates. Cell autofluorescence was subtracted and

MFI was converted to Mean Equivalents of Fluorescein (MEFLs) using the coefficient determined by the

calibration curve of UltraRainbow Calibration Particles (Spherotech URCP-100-2H) run in each individual

experiment. Standard error was propagated through all calculations.

Western Blotting

Western blots were performed to evaluate protein expression and normalize total expression of each TEVp

chain. A 3X-FLAG tagged NanoLuciferase as a normalization control, and images were analyzed using

ImageJ software. A detailed western blot protocol was previously described.[27]

Solvent-accessible Surface Area

The structure of TEVp was obtained from the Research Crystallography for Structural Bioinformatics

(RCSB) PDB (ID code: 1LVM). Per-residue solvent-accessible surface areas (SASA) were computed using

GROMACS v2018.1, which utilizes the double cubic lattice method (DCLM) described by Eisenhaber et

al.[28] The change in solvent accessible area was computed as

$$\Delta SASA = SASA_{fragments} - SASA_{reconstituted}$$

where structures of the N and C-terminal fragments were isolated from the crystal structure.

Computational Interface Scanning

All modeling calculations were performed using the *Rosetta* molecular modeling suite v3.9. Single-point

mutants were generated using the standard Relax application, which enables local conformational sampling

to minimize energy (**Supplementary Note 1** includes full details). The total energy (ΔG_{Total}) of each mutant

was computed as the average of 100 relaxed models. The energy perturbation to total energy was

computed as

$$\Delta\Delta G_{Total} = \Delta G_{Total}^{Mutant} - \Delta G_{Total}^{WT}$$

The Rosetta Scripts application with the InterfaceAnalyzeMover was applied to each relaxed model to

compute the average residue-residue interaction energies between the N- and C-terminal fragments

(**Supplementary Note 1** includes the full details). The interfacial energy was computed as the pair-wise

sum of all short-range interaction energies as shown by

$$\Delta G_{Interfacial} = \sum_{i} \sum_{j} Energy_{i-j}^{SR}$$

where i and j denote the sets of residues within each fragment. The energy perturbation of each mutation

to the interfacial energy was then computed as

$$\Delta\Delta G_{Interfacial} = \Delta G_{Interfacial}^{Mutant} - \Delta G_{Interfacial}^{WT}$$

Phenotype Classifier

Experimentally characterized variants were assigned class labels (not-inducible, inducible and dead) based

on reporter expression levels in the ligand-absent and ligand-induced states. Variants with ≥1.2 fold higher

reporter expression in the ligand-induced state relative to the ligand-absent state were labeled as inducible.

For variants with expression levels <5% of wild-type (WT) sequence in the ligand-induced state and <1.2

fold activation were classified as functionally dead. The remaining variants were assigned the not-inducible

class label.

STATISTICAL ANALYSIS

Statistical details for each experiment are included the figure legends. The data shown reflect the mean

across these biological replicates of the mean fluorescence intensity (MFI) of approximately 2,000–3,000

single, transfected cells. Error bars represent the SEM (standard error of the mean). For statistical

analyses, two-tailed Student's t-tests were used to evaluate whether a significant difference exists between

two groups of samples, and the reported comparisons meet the two requirements of this test: (1) the values

compared are expected to be derived from a normal distribution, and (2) the variance of each group is

expected to be comparable to that of the comparison group since the same transfection methodologies and

data collection methods were used for all samples that were compared. A p value of ≤ 0.05 was considered

to be statistically significant.

DATA AVAILABILITY

Data reported in composite figures (**Figs. 2a,b,d**, **3a, 4b, Supplementary Fig. 4**) are included as **Source**

Data.

CODE AVAILABILITY

Rosetta details and script are provided in **Supplementary Note 1**.

ACKNOWLEDGEMENTS

This work was supported in part by the National Institute of Biomedical Imaging and Bioengineering of the NIH under Award Number 1R01EB026510 (JNL); the Northwestern University Flow Cytometry Core Facility supported by Cancer Center Support Grant (NCI 5P30CA060553); T.B.D was supported by the Department of Defense (DoD) through the National Defense Science & Engineering Graduate Fellowship (NDSEG). This work is also supported in part by the Great Lakes Bioenergy Research Center, U. S. Department of Energy, Office of Science, Office of Biological and Environmental Research under Award Number DE-SC0018409 (S.R and A.T.M). The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH, Department of Defense, Department of Energy or other federal agencies.

AUTHOR CONTRIBUTIONS

T.B.D., A.T.M., S.R, and J.N.L conceptualized the project. T.B.D., J.D.B, W.K.C., E.E.S, created reagents, designed and performed experiments, and analyzed the data. A.N.P. assisted in analyzing and visualizing the data. A.T.M., developed the computational model and code. T.B.D., A.T.M., S.R., and J.N.L. drafted the manuscript, T.B.D., A.T.M., and A.N.P. created the figures. J.N.L. and S.R. supervised the work. All authors edited and approved the final manuscript.

COMPETING INTERESTS

J.N.L is a co-inventor on a patent that covers the MESA technology used in this manuscript (US Patent 9,732,392 B2).

REFERENCES

1. Romei, M.G. & Boxer, S.G. Split Green Fluorescent Proteins: Scope, Limitations, and Outlook.

Annual Review of Biophysics **48**, 19-44 (2019).

2. Shekhawat, S.S. & Ghosh, I. Split-protein systems: beyond binary protein-protein interactions. *Curr*

Opin Chem Biol **15**, 789-797 (2011).

3. Wehr, M.C. & Rossner, M.J. Split protein biosensor assays in molecular pharmacological studies.

Drug Discovery Today **21**, 415-429 (2016).

4. Muller, J. & Johnsson, N. Split-ubiquitin and the split-protein sensors: chessman for the endgame.

Chembiochem **9**, 2029-2038 (2008).

5. Paulmurugan, R. & Gambhir, S.S. Monitoring protein-protein interactions using split synthetic

renilla luciferase protein-fragment-assisted complementation. *Anal Chem* **75**, 1584-1589 (2003).

6. Dixon, A.S. et al. NanoLuc Complementation Reporter Optimized for Accurate Measurement of

Protein Interactions in Cells. *ACS chemical biology* **11**, 400-408 (2016).

7. Ozawa, T., Kaihara, A., Sato, M., Tachihara, K. & Umezawa, Y. Split luciferase as an optical probe

for detecting protein-protein interactions in mammalian cells based on protein splicing. *Analytical*

Chemistry **73**, 2516-2521 (2001).

8. Gray, D.C., Mahrus, S. & Wells, J.A. Activation of specific apoptotic caspases with an engineered

small-molecule-activated protease. *Cell* **142**, 637-646 (2010).

9. Gao, X.J., Chong, L.S., Kim, M.S. & Elowitz, M.B. Programmable protein circuits in living cells.

Science **361**, 1252-1258 (2018).

10. Fink, T. et al. Design of fast proteolysis-based signaling and logic circuits in mammalian cells.

Nature chemical biology **15**, 115-122 (2019).

11. Zetsche, B., Volz, S.E. & Zhang, F. A split-Cas9 architecture for inducible genome editing and

transcription modulation. *Nature biotechnology* **33**, 139-142 (2015).

12. Nihongaki, Y., Otabe, T., Ueda, Y. & Sato, M. A split CRISPR–Cpf1 platform for inducible genome

editing and gene activation. *Nature chemical biology* **15**, 882-888 (2019).

13.   Paulmurugan, R., Umezawa, Y. & Gambhir, S.S. Noninvasive imaging of protein-protein

interactions in living subjects by using reporter protein complementation and reconstitution

strategies. *Proc Natl Acad Sci U S A* **99**, 15608-15613 (2002).

14.   Fetchko, M. & Stagljar, I. Application of the split-ubiquitin membrane yeast two-hybrid system to

investigate membrane protein interactions. *Methods* **32**, 349-362 (2004).

15.   Pandey, N., Nobles, C.L., Zechiedrich, L., Maresso, A.W. & Silberg, J.J. Combining random gene

fission and rational gene fusion to discover near-infrared fluorescent protein fragments that report

on protein-protein interactions. *ACS Synth Biol* **4**, 615-624 (2015).

16.   Jones, K.A. et al. Development of a Split Esterase for Protein–Protein Interaction-Dependent Small-

Molecule Activation. *ACS Central Science* (2019).

17.   Wehr, M.C., Reinecke, L., Botvinnik, A. & Rossner, M.J. Analysis of transient phosphorylation-

dependent protein-protein interactions in living mammalian cells using split-TEV. *BMC Biotechnol*

8, 55 (2008).

18.   Camacho-Soto, K., Castillo-Montoya, J., Tye, B. & Ghosh, I. Ligand-gated split-kinases. *J Am*

Chem Soc **136**, 3995-4002 (2014).

19.   Camacho-Soto, K., Castillo-Montoya, J., Tye, B., Ogunleye, L.O. & Ghosh, I. Small molecule gated

split-tyrosine phosphatases and orthogonal split-tyrosine kinases. *J Am Chem Soc* **136**, 17078-

17086 (2014).

20.   Fink, T. et al. Design of fast proteolysis-based signaling and logic circuits in mammalian cells. *Nat*

Chem Biol **15**, 115-122 (2019).

21.   Daringer, N.M., Dudek, R.M., Schwarz, K.A. & Leonard, J.N. Modular extracellular sensor

architecture for engineering mammalian cell-based devices. *ACS Synth Biol* **3**, 892-902 (2014).

22.   Wehr, M.C. et al. Monitoring regulated protein-protein interactions using split TEV. *Nature Methods*

3, 985-993 (2006).

23.   Crescitelli, R. et al. Distinct RNA profiles in subpopulations of extracellular vesicles: apoptotic

bodies, microvesicles and exosomes. *J Extracell Vesicles* **2** (2013).

24.   Kapust, R.B., Tozser, J., Copeland, T.D. & Waugh, D.S. The P1' specificity of tobacco etch virus

protease. *Biochem Biophys Res Commun* **294**, 949-955 (2002).

25.   Hartfield, R.M., Schwarz, K.A., Muldoon, J.J., Bagheri, N. & Leonard, J.N. Multiplexing Engineered

481   Receptors for Multiparametric Evaluation of Environmental Ligands. *Acs Synthetic Biology* **6**, 2042-

482   2055 (2017).

483   26.   Schwarz, K.A., Daringer, N.M., Dolberg, T.B. & Leonard, J.N. Rewiring human cellular input-output

484   using modular extracellular sensors. *Nat Chem Biol* **13**, 202-209 (2017).

485   27.   Donahue, P.S., Draut, J.W., Muldoon, J.J., Edelstein, H.I., Bagheri, N., Leonard, J.N. COMET: A

486   toolkit for compsoing customizable genetic programs in mammalian cells. *bioRxiv* (2019).

487   28.   Eisenhaber, F., Lijnzaad, P., Argos, P., Sander, C. & Scharf, M. The double cubic lattice method:

488   Efficient approaches to numerical integration of surface area and volume and to dot surface

489   contouring of molecular assemblies. *Journal of Computational Chemistry* **16**, 273-284 (1995).

    FIGURES

Fig. 1 Design-driven strategy for tuning split protein systems. a, Current methods for optimizing split proteins are limited (left); an ideal tool would enable adapting split proteins for multiple applications, each of which may require distinct reconstitution propensities (right). **b,** This cartoon illustrates the experimental testbed used here; ligand binding-induced chain dimerization results in split TEVp reconstitution, trans-cleavage, and release of a previously sequestered transcription factor to drive reporter expression. **c,** Split Protein Optimization by Reconstitution Tuning (SPORT) workflow: identify important residues at the split interface which are mutable, identify mutations that alter the total and interfacial energy, and use an application-specific model, trained on limited experiments, to identify those mutations that are predicted to yield a desired functional phenotype.

Fig. 2 Computation guided method development and experimental analysis. a, Left, characterization

of the solvent accessible surface area (SASA) of each residue of 118/119 split TEVp. Right, 3D depiction

of 118/119 split TEVp, showing the catalytic triad (orange), coordination sphere (yellow), and ΔSASA

(greyscale). **b,** Mutational scanning of high ΔSASA residues (left) and example of all possible mutations of

residue 103 (right), with change in interfacial energy indicated by color. **c,** Experimental analysis of TEVp

mutations predicted to span a range of interfacial energies. Error bars depict S.E.M. (*$p \leq 0.05$, ***$p \leq$

0.001). **d,** Experimental phenotypes observed in **c** were plotted on an energy landscape and annotated as

indicated by color (reporter expression normalized to WT < 0.05 is "dead", fold induction < 1.2 is "not

inducible", fold induction ≥ 1.2 is "inducible"). **e,** Proposed model for predicting zones of functional

phenotypes based upon total and interfacial energy; the boundaries were proposed based upon

observations using the initial 20 mutants tested in **c**.

Fig. 3 Evaluation of model-predicted phenotypes for combined mutations. a, Computed energies and predicted phenotypes based on the classifier model—proposed in **Fig. 2e**—of all possible double and paired mutants constructed by combinatorial sampling of the initial single mutants tested (omitting dead mutations) in **Fig. 2c**. **b,** Experimental evaluation of selected mutants predicted to be inducible. **c,** Experimentally observed phenotypes for the fourteen mutants predicted to be inducible (from **b**), showing that the model predicts inducibility at a fairly high rate (10/14). **d,** Normalizing protein expression levels improves performance (fold induction) of selected mutants (from **b**), whereas WT function is not changed. Normalization was achieved using Western Blot analysis (**Supplementary Fig. L4**) to adjust DNA doses transfected (per well, N-terminal chains: 0.4 ng WT, 1 ng 75S, 1.4 ng 75E; C-terminal chains: 5 ng WT, 12 ng 190K). Error bars depict S.E.M. (*p ≤ 0.05, ***p ≤ 0.001).

Fig. 4 Evaluation of model-predicted phenotypes for novel mutations and combinations. a, For each

experimentally characterized construct, reporter output was quantified in the absence of ligand (OFF state)

and following ligand addition (ON state), and the fold induction was calculated. Calculated interfacial energy

for each construct is indicated by circle *color*, magnitude of reporter expression (or fold-induction) is

indicated by circle *size*, and constructs with a fold induction ≥ 1.2 are denoted with a black border. Single

mutants observed to be dead (**Fig. 2**) were not carried forward to this analysis. **b,** Left, experimentally

observed phenotypes (data point color) were mapped onto the proposed classifier model from **Fig. 2e**, with

observed frequency distributions shown as histograms. Right, evaluation of model prediction accuracy

compared to random assignment of phenotypes.

Fig. 5 Model guided design of a new split TEVp application in soluble context. a, This cartoon illustrates the soluble split TEVp testbed. Ligand-binding-induced dimerization mediates reconstitution of split TEVp, which then cleaves one or more nuclear export sequence (NES) elements from a soluble transcription factor, leading to nuclear import and reporter expression. **b,** Developing the testbed by evaluating engineered transcription factors (TF) for consistency with the mechanism proposed in **a**; shaded cleavage sequence (CS) domains indicate a G residue in the P1' position, unshaded CS domains indicate

549   a M residue in this position.[24] **c,** Experimental analysis of single and paired mutants sampling a range of

550   interfacial energies (indicated by color and labeled), employing TF10 from **b**. Error bars depict S.E.M. (*$p \leq$

551   0.05, ***$p \leq 0.001$).

a Tools for optimizing split protein systems

b Experimental testbed

1. Dimerization
2. TEVp reconstitution
3. Cleavage
4. TF release

c

Identify interfacial residues

Computational interface scanning

Calibration with experimental data

Predictions