Excess Centrosomes Disrupt Vascular Lumenization and Endothelial Cell Adherens Junctions

Danielle B Buglak[a], Erich J Kushner[b], Allison P Marvin[b], Katy L Davis[b], and Victoria L Bautch[a,b,c,#]

[a]Curriculum in Cell Biology and Physiology, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA.
[b]Department of Biology, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA.
[c]McAllister Heart Institute, The University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA.

[#]Corresponding author: Victoria L Bautch, PhD
    Professor of Biology
    Department of Biology, CB#3280
    University of North Carolina at Chapel Hill
    Chapel Hill, NC 27599 USA
    Phone: 919-966-6797
    E-mail: bautch@med.unc.edu

Running head: CENTROSOMES IN VASCULAR LUMENIZATION AND JUNCTIONS

WORKING TITLE: 10 words
Abstract: 145 words
Text: 5,181 words
Figures: 4 main, 3 supplemental

ABSTRACT

Proper blood vessel formation requires coordinated changes in endothelial cell polarity and rearrangement of cell-cell junctions to form a functional lumen. One important regulator of cell polarity is the centrosome, which acts as a microtubule organizing center. Excess centrosomes perturb aspects of endothelial cell polarity linked to migration, but whether centrosome number influences apical-basal polarity and cell-cell junctions is unknown. Here, we show that excess centrosomes alter the apical-basal polarity of endothelial cells in angiogenic sprouts and disrupt endothelial cell-cell adherens junctions. Endothelial cells with excess centrosomes had narrower lumens in a 3D sprouting angiogenesis model, and zebrafish intersegmental vessels had reduced perfusion following centrosome overduplication. These results indicate that endothelial cell centrosome number regulates proper lumenization downstream of effects on apical-basal polarity and cell-cell junctions. Endothelial cells with excess centrosomes are prevalent in tumor vessels, suggesting how centrosomes may contribute to tumor vessel dysfunction.

Keywords: Endothelial cell, centrosome, junctions, polarity, lumen, angiogenesis

INTRODUCTION

Angiogenesis is the sprouting of new blood vessels from pre-existing vessels and

is crucial during development, pregnancy, wound healing, and tumorigenesis [1–3].

Endothelial cells (EC) form new sprouts to provide conduits for blood flow and create a

functional vascular network. This process requires dynamic rearrangement of

endothelial cell-cell junctions and establishment of apical-basal polarity [4–8].

Lumenization closely follows sprout formation and extension temporally, and many

sprout tips have multiple EC that are polarized in the apical-basal axis [9]. However, it is

not known how centrosome number contributes to this process.

As tumors grow, hypoxia leads to elevated levels of pro-angiogenic factors that

promote neo-angiogenesis to vascularize the tumor. However, the blood vessels

surrounding these tumors are often leaky, tortuous, and not properly lumenized [10–

12], suggesting defects in EC junctions and polarity. Interestingly, tumor vessels are

also characterized by EC with excess centrosomes [13,14]. The centrosome acts as a

major microtubule organizing center and determinant of cell polarity [15,16]. High

VEGF-A signaling in EC results in centrosome overduplication, which affects interphase

cells by increasing invasiveness, elevating microtubule nucleations, changing aspects of

polarity and migration, and altering signaling dynamics [13,17–19]. However, it is

unclear how excess centrosomes impact cell-cell interactions and lumenogenesis.

Cadherins form cell-cell adherens junctions that link to the actin cytoskeleton and

restrict the apical vs. basolateral domains [20,21]. In EC, VE-cadherin is the primary

component of the vascular adherens junction and is required for localization of apical

markers during lumen formation in zebrafish [7]. Depletion of VE-cadherin is

embryonically lethal in zebrafish and mice due to severe vascular defects, including a

lack of vascular lumens [22,23]. However, the effects of centrosome number on

adherens junctions have not been investigated.

Here, we show for the first time that excess centrosomes prevent the proper

polarization of interacting EC. Excess centrosomes resulted in EC with destabilized

adherens junctions and narrow and closed vascular lumens both *in vitro* and *in vivo*.

These findings reveal a novel role for the centrosome in EC apical-basal polarity, and in

junction and lumen formation, and suggest how excess centrosomes in the vasculature

may contribute to poor perfusion in tumor vessels.

RESULTS

Excess centrosomes perturb polarization of junctionally linked EC

We previously showed that excess centrosomes interfere with repolarization along the forward-rearward axis in sprouting EC [17], and thus we hypothesized that supernumerary centrosomes disrupt multiple EC polarity axes. We examined the polarity between EC sharing junctions using an inducible system to overexpress polo-like kinase 4 (Plk4) in human umbilical vein endothelial cells (HUVEC) **(Fig. 1a)**. Plk4 regulates centriole duplication, and its overexpression downstream of a TRE (tet-responsive element) upon addition of doxycycline (DOX) results in supernumerary centrosomes [13,17,24,25]. TRE-Plk4 HUVEC were seeded onto large H-shaped micropatterns that allow for polarity assessment between two EC that form a cell-cell junction. Polarity was defined based on centrosome position relative to the nucleus and cell-cell junction, with a "proximal" position being near the cell-cell junction, a "central" position in the nuclear region, and a "distal" position between the nucleus and the cell periphery **(Fig. 1b)**. Two EC with a normal centrosome number (1-2) typically had a distal centrosome polarity **(Fig. 1c-d)**. In contrast, two EC with excess centrosomes (>2) showed a significant increase in central/proximal polarity, indicating that centrosome number affects EC polarization relative to cell-cell junctions **(Fig. 1c-d)**. Interestingly, when a normal EC and an EC with excess centrosomes were linked on the same pattern (N:O), normal EC had a higher frequency of aberrant centrosome localization than N:N counterparts, while EC with excess centrosomes had a higher frequency of normal centrosome localization relative to O:O counterparts, suggesting that EC polarity influences polarity in neighboring cells **(Fig. 1d)**. Furthermore, when two EC with normal

centrosome number (N:N) formed a junction, both cells usually had centrosomes in the

distal position, which we termed "distal" polarity **(Fig. 1e)**. All other polarity combinations

were deemed "other". When both EC forming a junction had excess centrosomes (O:O),

polarity was randomized between distal and other **(Fig. 1e)**. Additionally, N:N EC

combinations were more likely to have cells with matching polarity, while EC with an

O:O combination were less likely to have the same polarity as their neighbor **(Fig. 1f)**.

Taken together, these data indicate that excess centrosomes prevent proper EC

polarization relative to cell-cell junctions.

EC with excess centrosomes form disorganized adherens junctions

Because interacting EC with excess centrosomes had perturbed polarity in the

junction axis, we asked whether EC cell-cell interactions were influenced by centrosome

number. Adherens junctions are required for proper apical-basal polarity in epithelial

cells [7,26], and the distribution of VE-cadherin at EC adherens junctions is indicative of

junction maturity; a linear VE-cadherin pattern is associated with mature stabilized

junctions, while a serrated and punctate VE-cadherin pattern is associated with

activated and/or disrupted junctions [27]. To determine how excess centrosomes

influence adherens junctions, we seeded induced tet-Plk4 HUVEC onto H-shaped

micropatterns. Due to the heterogeneous nature of Plk4-induced centrosome over-

duplication, a given cell-cell junction could have contributions from two EC with a normal

number of centrosomes (N:N), two EC with excess centrosomes (O:O), or one of each

(N:O) **(Fig. 2a-a")**. We found that junctions with contributions from EC with normal

centrosome numbers had more linear and stable junctions, while junctions with

contributions from EC with excess centrosomes formed more chaotic and disrupted

junctions **(Fig. 2b-b", c-c")**. These chaotic junctions had increased total VE-cadherin

area **(Fig. 2d)**, and line scans revealed a more dispersed VE-cadherin junction

distribution when one or both EC had excess centrosomes **(Fig. 2e)**. We also assessed

the overall width of the junction and found that junctional VE-cadherin distribution was

wider in junctions formed when one or both EC had excess centrosomes **(Fig. 2f)**.

Together, these data suggest that both the junction shape and VE-cadherin distribution

are perturbed in EC with supernumerary centrosomes. Because tight junctions are

maintained along with ectopic patches of VE-cadherin in migrating and anastomosing

EC [28], we examined tight junctions via ZO-1 staining and found that both N:O and

O:O patterns had missing or abnormal tight junctions downstream of disrupted

adherens junctions, with O:O patterns having a more severe disruption **(Fig. S1)**. In

these cases, there was no organized ZO-1 staining in the large patches of VE-cadherin

beyond the junction site. Overall, these findings indicate that EC with excess

centrosomes do not maintain proper endothelial cell-cell adherens or tight junctions.

EC with excess centrosomes have abnormal lumens

We hypothesized that the disrupted polarity seen between EC with

supernumerary centrosomes had physiological consequences. To further examine EC

polarity, we analyzed lumenized sprouts that form and polarize in the apical-basal axis

in a 3D sprouting angiogenesis assay [9,29]. DOX-treated tet-Plk4 HUVEC expressing

centrin::eGFP were stained for Moesin1 as an apical marker, since Moesin1 localizes to

F-actin found at apical EC cell-cell contacts [6] **(Fig. 3a-b)**. We found that EC with

normal centrosome numbers (1-2) typically had apically positioned centrosomes, while

EC with excess centrosomes (>2) had randomized centrosome polarity in the apical-

basal axis **(Fig. 3b-c)**. In addition, EC with excess centrosomes were more likely to

have punctate Moesin1, basal Moesin1, or unpolarized abnormal Moesin1 staining

patterns compared to normal linear and apical patterns **(Fig. 3b, d)**, indicative of polarity

defects. Furthermore, VE-cadherin around EC with excess centrosomes was more

dispersed in 3D sprouts compared to EC with normal centrosome number **(Fig. S2a)**.

Since both apical-basal polarity and proper junction formation are important for

lumen formation [4–8], we hypothesized that excess centrosomes in EC perturb vessel

lumenogenesis. After normalizing for overall vessel width, vessel lumen widths were

narrower near EC with excess centrosomes (excluding closed lumens) compared to EC

with normal centrosome numbers **(Fig. 3b, e)**, and raw lumen widths were also

decreased **(Fig. S2b)**. Interestingly, some lumens were completely closed around EC

with supernumerary centrosomes, and the incidence of closed and narrowed lumens

was elevated near sprout EC with excess centrosomes **(Fig. 3f)**. These findings indicate

that EC centrosome number regulates proper vessel lumen diameter, likely through

effects on apical-basal polarity.

Overexpression of Plk4 in zebrafish results in fewer lumenized vessels

To determine if centrosome number affects lumen formation *in vivo*, we utilized a

zebrafish model with vascular-specific Cre-mediated overexpression of *Plk4* **(Fig. S3a)**,

and crossed to *Tg(fli:centrin-GFP)* fish to visualize EC centrosomes. Fish

overexpressing *Plk4* had increased EC with >2 centrosomes **(Fig. S3b-c)**, and

microangiography at 72 hpf revealed that these fish also had fewer perfused

intersegmental vessels (ISVs) compared to wild type controls or fish lacking Cre **(Fig.**

4a-b). Lumens of perfused vessels in fish with *Plk4* overexpression were narrower

compared to controls, and more lumens were completely closed or narrowed **(Fig. 4c-**

d). To further test the effects of supernumerary centrosomes, we generated a second

zebrafish model by utilizing the *Gal4-UAS* system to transiently overexpress *Plk4* in

blood vessels **(Fig. S3d)**. ISVs containing EC with excess centrosomes were narrower

than neighboring ISVs consisting of EC with 1-2 centrosomes, and ISVs from fish

lacking *Tg(fli:Gal4)* **(Fig. 4e-f)**. These findings indicate that ISV narrowing is specific to

sprout regions with EC harboring excess centrosomes. Reduced vessel perfusion,

coupled with narrow or blocked lumens and vessels, suggests that excess centrosomes

in EC lead to disrupted lumenogenesis *in vivo*.

DISCUSSION

Angiogenesis involves migration and rearrangement of EC, and leads to new conduits through anastomosis and lumen formation between adjacent EC [1–3]. Cell-cell interactions and polarization in several axes are important for lumen formation, but the role of the centrosome in these processes is poorly understood. Here, we show that supernumerary centrosomes perturb EC polarity and cell adhesion and disturb lumen formation *in vitro* and *in vivo*, implying that the centrosome is involved in both junctional adhesion and the apical-basal polarization that precedes vascular lumen formation or stabilization.

Our data highlights important roles for the centrosome in maintaining junction integrity and lumen development in interphase cells. Besides overduplication, centrosome function is impacted by centriole loss or structural defects. Mis-localization of centrosomal proteins, structural deficits, and centrosome loss all result in G1 cell-cycle arrest [30], and compromise analysis of effects of centrosome loss on interphase cell behavior. Although supernumerary centrosomes affect cellular processes, they do not acutely block cell-cycle progression [17] and thus allow for analysis of centrosome perturbation on interphase behaviors.

Our previous work showed that EC centrosome overduplication is linked to microtubule defects, with increased nucleations and catastrophes accompanied by forward-rearward polarity defects [17]. Changes in microtubule dynamics are linked to regulation of adherens junctions in EC [31]. Our data indicate that adherens junctions

are significantly disrupted in EC with excess centrosomes, suggesting that junctions are

disrupted downstream of centrosome-induced microtubule defects.

Lumen structure is also disrupted downstream of excess centrosomes in

sprouting EC, suggesting an apical-basal polarity defect. It is likely that junctional

defects contribute to poor lumenization *in vitro* and *in vivo*. VE-cadherin at EC junctions

is linked to vascular lumen formation during anastomosis [32,33]. Moreover, genetic

loss of VE-cadherin in mice or zebrafish leads to lumen defects, vascular dysfunction,

and embryonic lethality [22,23], supporting a link between disrupted adherens junctions

and malformed lumens in EC sprouts. Proper adherens junctions are necessary to

stabilize tight junctions and form lumens [34,35] , and here we show that tight junction

formation is also compromised in EC with excess centrosomes, suggesting that tight

junction defects contribute to lumen defects.

We show that lumens are perturbed following centrosome overduplication in both

an *in vitro* angiogenesis model lacking physiological flow and *in vivo*, in zebrafish with

blood flow. These findings indicate that the lumenal defects resulting from excess

centrosomes do not depend on shear stress exerted by blood flow. While lumenization

requires blood flow during late stages of angiogenesis [36], lumens also form *in vitro*

and *in vivo* absent blood flow [7,37]. Disruption of the VE-cadherin junctional complex is

sufficient to prevent luminal expansion independent of blood flow through changes in

cell contractility [37]; thus, it is likely that the profound changes in VE-cadherin

localization at EC cell-cell junctions following centrosome over-duplication alter the

ability of EC to properly coordinate the cytoskeletal forces needed for polarization and

lumenogenesis.

Centrosome amplification is a common phenomenon in tumors and has been

linked to changes in the tumor microenvironment, including endothelial cells [13,14].

Local hypoxia in tumors increases VEGF-A signaling, leading to high CyclinE/cdk2 that

promotes EC centrosome overduplication [19], and elevated *Plk4* expression occurs in

human and mouse tumor vessels [38,39]. Although therapies targeting angiogenesis

have had mixed results [40,41], understanding tumor EC responses is necessary for

design of more complex multi-modal anti-tumor therapies [41–44]. In summary, by

revealing a unique link between centrosome perturbation and adherens junction stability

and lumenization, our data indicate a novel role for the centrosome in EC polarity,

junctions, and lumenization in blood vessels. These abnormalities may contribute to

tumor blood vessel dysfunction.

MATERIALS & METHODS

Cell culture

HUVEC (C2519A, lot 486264, Lonza) were cultured in EBM-2 (CC-3162, Lonza) supplemented with the Endothelial Growth Medium (EGM-2) bullet kit (CC-3162, Lonza) and 1x antibiotic-antimycotic (Gibco). Normal human lung fibroblasts (CC2512, Lonza) were cultured in DMEM (Gibco) with 10% fetal bovine serum (FBS) and 1x antibiotic-antimycotic (Gibco). All cells were maintained at 37°C and 5% CO_2. Tetracycline-inducible Plk4-overexpression HUVEC were generated as previously described [13,17,25]. Viruses were produced by the UNC Lenti-shRNA Core Facility. To induce centrosome overduplication, tet-Plk4 HUVEC were treated with 1µg/ml doxycycline (DOX, D9891, Sigma) for 24hr followed by washout. Experiments were performed 48hr following DOX addition.

3D sprouting angiogenesis assay

The 3D sprouting angiogenesis assay was performed as previously described with the following changes [17,45]. Tet-Plk4 HUVEC were treated with 1µg/ml DOX for 24hr at 37°C. 13µl/ml centrin:eGFP lentivirus and 1µg/ml Polybrene were added to HUVEC 16hr post DOX addition for the last 8hr of DOX treatment. Cells were washed with Dulbecco's phosphate-buffered saline (DPBS) (MT-21-031-CV, Corning) and fresh media was added. The following day, infected cells were coated onto cytodex 3 microcarrier beads (17048501, GE Healthcare Life Sciences) and embedded in a fibrin matrix as described. Z-stacks were acquired with an Olympus confocal microscope (FV1200 or FV3000) and processed in Image J (National Institutes of Health). Total

vessel width was measured in the YZ plane by drawing a line through the nucleus and

across the entire vessel. Lumen width was measured in the YZ plane using Moesin1

staining (1:1000, ab52490, Abcam) as a surrogate for the apical surface and normalized

to total vessel width. Lumens were considered "narrow" if the normalized lumen width

fell below the mean lumen width minus one standard deviation of the normal condition.

Lumens were considered "closed" if the lumen width was zero. DAPI (10236276001,

Sigma) was used to visualize the nucleus, and VE-cadherin was used to visualize

adherens junctions (1:1000, 2500S, Cell Signaling).

Micropatterns

Large H-micropatterns (1600 µm^2, CYTOO) were coated with 5µg/ml fibronectin

and seeded with HUVEC according to manufacturer's instructions. Cells were allowed

to spread for 24hr, then fixed with ice-cold methanol for 10 min at 4°C.

Immunofluorescence and quantification

Fixed HUVEC were blocked for 1hr at RT in blocking solution (5% FBS, 2X

antibiotic-antimycotic (Gibco), 0.1% sodium azide (s2002-100G, Sigma) in DPBS). Cells

were incubated in primary antibody overnight at 4°C, then washed 3X for 5 min in

DPBS. Secondary antibody and DRAQ7 (1:1000, ab109202, Abcam) were added for

1hr at RT followed by 3X washes for 10 min each in DPBS. Slides were mounted with

coverslips using Prolong Diamond Antifade mounting medium (P36961, Life

Technology) and sealed with nail polish. Primary and secondary antibodies were diluted

in blocking solution. Images were acquired with an Olympus confocal microscope

312 (FV1200 or FV3000) and analyzed using ImageJ. Centrosome overduplication was

313 quantified by counting centrosome number in at least 500 cells per replicate. Junction

314 area on micropatterns was quantified by thresholding VE-cadherin staining and

315 measuring total area within the junction in ImageJ. VE-cadherin distribution on doublet

316 micropatterns was measured in ImageJ by drawing a line across the junction (from

317 centrosome in first cell to centrosome in second cell) and acquiring a line scan of VE-

318 cadherin intensity. Peaks were counted for each pattern and normalized to the line

319 length. Only peaks above half of the maximum VE-cadherin intensity were counted.

320 Junction width was measured by creating a histogram of the VE-cadherin signal

321 integrated over the y-axis and measuring the width of the most prominent peak(s). The

322 following primary antibodies were used: anti-γ-tubulin (1:5000, T6557, Sigma-Aldrich),

323 anti-VE-cadherin (1:500, 2500S, Cell Signaling), anti-ZO-1 (1:500, MABT339, Millipore

324 Sigma), anti-Moesin-1 (1:1000, ab52490, Abcam). The following secondary antibodies

325 from Life Technologies were used: goat anti-mouse AlexaFluor 488 (1:500, A11029),

326 goat anti-mouse AlexaFluor 594 (1:500, A11005), goat anti-mouse AlexaFluor 647

327 (1:500, A21236) goat anti-rabbit AlexaFluor 488 (1:500, A11034), goat anti-rabbit

328 AlexaFluor 594 (1:500, A11037).

330 ***Zebrafish***

331  Zebrafish (*Danio rerio*) were housed in an institutional animal care and use

332 committee (IACUC)-approved facility and maintained as previously described (Wiley, *et*

333 *al*., 2011). *Tg(kdrl:mCherry)* was a gift from D. Stainier. *Tg(fli:Gal4)* was a gift from W.

334 Herzog. *Tg(fli:centrin-GFP)* was created by subcloning centrin:GFP into the

pCR8/GW/TOPO middle entry vector (Thermo Fisher) and then gateway cloning into

the pDestTol2CG2 plasmid (REF Tol2 kit). Tol2 transposase RNA was generated using

the sp6 mMessage mMachine synthesis kit (AM1340, Thermo Fisher). For

overexpression of *Plk4*, *p2a-Lox-mCherry-STOP-Lox-Plk4* was fused to the 3' end of

the endogenous *kdrl* gene. The assembled construct was subcloned into the pKHR4

backbone and linearized by I-SceI digest. This construct, 200ng/µl sgRNA (5'-

TCTGGTTTGGAAGGACACAG-3'), and 700ng/µl of Cas9 recombinant protein

(PNABio) were injected into one-cell stage *Tg(fli:centrin-GFP)* embryos to generate

double stranded breaks at the 3' end of the *kdrl* gene and induce homologous

recombination as previous described [46]. F1 fish were injected with Cre mRNA at the

1-2 cell stage. Cre mRNA was obtained by subcloning the *Cre* gene from the pME-

ERT2-Cre-ERT2 plasmid into the pCR8/GW/TOPO middle entry vector, gateway

cloning into a pCS Dest vector, then preparing RNA with the sp6 mMessageMachine

synthesis kit. pME-ERT2-Cre-ERT2 was a gift from Kryn Stankunas (Addgene plasmid

82587; http://n2t.net/addgene:82587; RRID:Addgene 82587). Embryos were collected

from desired crosses and grown in E3 at 28.5°C. At 72 hpf, embryos were injected with

Qdot605 (Q10103MP, Thermo Fisher). *Plk4* overexpression was determined by the loss

of mCherry signal. A second fish model was generated as follows: a *5X-UAS-Plk4*

construct was generated using Gibson cloning by fusing a *5X-UAS* tag upstream of

zebrafish *Plk4*. pUAS:Cas9T2AGFP;U6:sgRNA1;U6:sgRNA2 was a gift from Filippo Del

Bene (Addgene plasmid #74009; http://n2t.net/addgene:74009;RRID:Addgene_74009)

[47]. This construct and I-SceI meganuclease were injected into one-cell fish from

crosses between Tg(fli:centrin-GFP) and Tg(fli:Gal4;kdrl:mCherry), and fish were grown

in E3 at 28.5°C to 72 hpf. Fish were fixed by incubating dechorionated embryos in ice

cold 4% PFA at 4°C overnight. Embryos were rinsed in PBS and mounted using a fine

probe to de-yolk and a small blade to separate the trunk from the cephalic region, and

mounting with VECTASHIELD® Hardset™ Antifade Mounting Medium. The coverslip

was sealed with petroleum jelly before imaging on an Olympus FV1200 confocal

microscope. Centrosome overduplication was confirmed based on centrin-GFP centriole

labeling. ISV perfusion was measured by counting the number of perfused ISVs based

on Qdot605 signal as a percentage of total ISV number (visualized using

Tg(kdrl:mCherry)). Lumen width was measured by measuring the diameter of perfused

ISVs near the junction with the dorsal aorta. Lumens were considered "narrow" if the

lumen width fell below the mean lumen width minus one standard deviation of the

WT+Cre condition. Lumens were considered closed if the ISV perfusion was

discontinuous.

Statistics

Student's two-tailed *t* test was used to determine statistical significance in

experiments with 2 groups. One-way ANOVA with Tukey's multiple comparisons test

was used to determine statistical significance for experiments with 3 groups, and X^2 was

used for categorical data. Error bars represent the mean ± standard deviation. Statistical

tests and graphs were made using the Prism 7 software (GraphPad Software).

ACKNOWLEGMENTS

We thank Tony Perdue and the UNC Biology Microscopy Core, the UNC Lenti-

shRNA Core, Maryanna Parker and Kaitlyn Quigley for fish room support, and Bautch

lab members for critical discussions.

FUNDING

This work was supported by grants from the National Institutes of Health

(HL43174, HL116719, HL117256, HL139950 to VLB), a K99/R00 (1K99HL124311-

01A1 to EJK), the Integrated Vascular Biology Training Grant (5T32HL069768-17 to

DBB), and an American Heart Association Predoctoral Fellowship (19PRE34380887 to

DBB).

DECLARATION OF INTERESTS: None.

REFERENCES

1.   Carmeliet P, Jain RK. (2011) Molecular mechanisms and clinical applications of

angiogenesis. Nature. 473(7347):298–307.

2.   Bautch VL, Caron KM. (2015) Blood and lymphatic vessel formation. Cold Spring

Harb Perspect Biol. 7(3):a008268.

3.   Kushner E, Bautch V. (2013) Building blood vessels in development and disease.

Curr Opin Hematol. 20(3):231–6.

4.   Blum Y, Belting HG, Ellertsdottir E, Herwig L, Luders F, Affolter M. (2008)

Complex cell rearrangements during intersegmental vessel sprouting and vessel

fusion in the zebrafish embryo. Dev Biol. 316:312–22.

5.   Iruela-Arispe ML, Davis GE. (2009) Cellular and molecular mechanisms of

vascular lumen formation. Dev Cell. 16(2):222–31.

6.   Strilic B, Kucera T, Eglinger J, Hughes MR, McNagny KM, Tsukita S, et al. (2009)

The molecular basis of vascular lumen formation in the developing mouse aorta.

Dev Cell. 17(4):505–15.

7.   Wang Y, Kaiser MS, Larson JD, Nasevicius A, Clark KJ, Wadman SA, et al.

(2010) Moesin1 and VE-cadherin are required in endothelial cells during in vivo

tubulogenesis. Development. 137(18):3119–28.

8.   Lee CY, Bautch VL. (2011) Ups and downs of guided vessel sprouting: the role of

polarity. Physiology. 26(5):326–33.

9.   Pelton JC, Wright CE, Leitges M, Bautch VL. (2014) Multiple endothelial cells

constitute the tip of developing blood vessels and polarize to promote lumen

formation. Development. 141(21):4121–6.

10.   Aird WC. (2012) Endothelial cell heterogeneity. Cold Spring Harb Perspect Med.

2(1):a006429.

11.   Hashizume H, Baluk P, Morikawa S, McLean JW, Thurston G, Roberge S, et al.

(2000) Openings between defective endothelial cells explain tumor vessel

leakiness. Am J Pathol. 156(4):1363–80.

12.   Carmeliet P, Jain RK. (2011) Principles and mechanisms of vessel normalization

for cancer and other angiogenic diseases. Nat Rev Drug Discov. 10:417–27.

13.   Kushner EJ, Ferro LS, Liu J-Y, Durrant JR, Rogers SL, Dudley AC, et al. (2014)

Excess centrosomes disrupt endothelial migration via centrosome scattering. J

Cell Biol. 206(2):257–72.

14.   Hida K, Hida Y, Amin DN, Flint AF, Panigrahy D, Morton CC, et al. (2004) Tumor-

Associated Endothelial Cells with Cytogenetic Abnormalities. Cancer Res.

64(22):8249–55.

15.   Kollman JM, Merdes A, Mourey L, Agard DA. (2011) Microtubule nucleations by

gamma-tubulin complexes. Nat Rev Mol Cell Biol. 12:709–21.

16.   Bettencourt-Dias M, Glover DM. (2007) Centrosome biogenesis and function:

centrosomics brings new understanding. Nat Rev Mol Cell Biol. 8:451–63.

17.   Kushner EJ, Ferro LS, Yu Z, Bautch VL. (2016) Excess centrosomes perturb

dynamic endothelial cell repolarization during blood vessel formation. Mol Biol

Cell. 27(12):1911–20.

18.   Godinho SA, Picone R, Burute M, Dagher R, Su Y, Leung CT, et al. (2014)

Oncogene-like induction of cellular invasion from centrosome amplification.

Nature. 510(7503):167–71.

19. Taylor SM, Nevis KR, Park HL, Rogers GC, Rogers SL, Cook JG, et al. (2010)

Angiogenic factor signaling regulates centrosome duplication in endothelial cells

of developing blood vessels. Blood. 116(16):3108–17.

20. Spracklen AJ, Peifer M. (2015) Actin and apical constriction: some (re)-assembly

required. Dev Cell. 35(6):662–4.

21. Nowotarski SH, Peifer M. (2014) Cell biology: a tense but good day for actin at

cell-cell junctions. Curr Biol. 24(15):R688–90.

22. Carmeliet P, Lampugnani M-G, Moons L, Breviario F, Compernolle V, Bono F, et

al. (1999) Targeted deficiency or cytosolic truncation of the VE-cadherin gene in

mice impairs VEGF-mediated endothelial survival and angiogenesis. Cell.

98(2):147–57.

23. Montero-Balaguer M, Swirsding K, Orsenigo F, Cotelli F, Mione M, Dejana E.

(2009) Stable vascular connections and remodeling require full expression of VE-

cadherin in zebrafish embryos. PLoS One. 4(6):e5772.

24. Habedanck R, Stierhof YD, Wilkinson CJ, Nigg EA. (2005) The Polo kinase Plk4

functions in centriole duplication. Nat Cell Biol. 7(11):1140–6.

25. Yu Z, Ruter DL, Kushner EJ, Bautch VL. (2017) Excess centrosomes induce p53-

dependent senescence without DNA damage in endothelial cells. FASEB J.

10:4295–304.

26. Iden S, Rehder D, August B, Suzuki A, Wolburg-Buchholz K, Wolburg H, et al.

(2006) A distinct PAR complex associates physically with VE-cadherin in

vertebrate endothelial cells. EMBO Rep. 7(12):1239–46.

27. Bentley K, Franco CA, Philippides A, Blanco R, Dierkes M, Gebala V, et al. (2014)

The role of differential VE-cadherin dynamics in cell rearrangement during angiogenesis. Nat Cell Biol. 16(4):309–21.

28. Paatero I, Sauteur L, Lee M, Lagendijk AK, Heutschi D, Wiesner C, et al. (2018) Junction-based lamellipodia drive endothelial cell rearrangements in vivo via a VE-cadherin-F-actin based oscillatory cell-cell interaction. Nat Commun. 9(1):3545.

29. Lampugnani M-G, Orsenigo F, Rudini N, Maddaluno L, Boulday G, Chapon F, et al. (2010) CCM1 regulates vascular-lumen organization by inducing endothelial polarity. J Cell Sci. 123:1073–80.

30. Mikule K, Delaval B, Kaldis P, Jurcyzk A, Hergert P, Doxsey S. (2007) Loss of centrosome integrity induces p38-p53-p21-dependent G1-S arrest. Nat Cell Biol. 9:160–70.

31. Komarova YA, Huang F, Geyer M, Daneshjou N, Garcia A, Idalino L, et al. (2012) VE-cadherin signaling induces EB3 phosphorylation to suppress microtubule growth and assemble adherens junctions. Mol Cell. 48(6):914–25.

32. Lenard A, Ellertsdottir E, Herwig L, Krudewig A, Sauteur L, Belting H-G, et al. (2013) In vivo analysis reveals a highly stereotypic morphogenetic pathway of vascular anastomosis. Dev Cell. 25(5):492–506.

33. Herwig L, Blum Y, Krudewig A, Ellertsdottir E, Lenard A, Belting H-G, et al. (2011) Distinct cellular mechanisms of blood vessel fusion in the zebrafish embryo. Curr Biol. 21(22):1942–8.

34. Campbell HK, Maiers JL, DeMali KA. (2017) Interplay between tight junctions and adherens junctions. Exp Cell Res. 358(1):39–44.

486   35.   Odenwald MA, Choi W, Buckley A, Shashikanth N, Joseph NE, Wang Y, et al.

487   (2017) ZO-1 interactions with F-actin and occludin direct epithelial polarization

488   and single lumen specification in 3D culture. J Cell Sci. 130(1):243–59.

489   36.   Gebala V, Collins R, Geudens I, Phng L-K, Gerhardt H. (2016) Blood flow drives

490   lumen formation by inverse membrane blebbing during angiogenesis in vivo. Nat

491   Cell Biol. 18(4):443–50.

492   37.   Hultin S, Zheng Y, Mahdi M, Vertuani S, Gentili C, Balland M, et al. (2014)

493   AmotL2 links VE-cadherin to contractile actin fibres necessary for aortic lumen

494   expansion. Nat Commun. 5:3743.

495   38.   Roudnicky F, Poyet C, Wild P, Krampitz S, Negrini F, Huggenberger R, et al.

496   (2013) Endocan is upregulated on tumor vessels in invasive bladder cancer

497   where it mediates VEGF-A-induced angiogenesis. Cancer Res. 73(3):1097–106.

498   39.   Goveia J, Rohlenova K, Taverna F, Treps L, Conradi L, Pircher A, et al. (2020) An

499   integrated gene expression landscape profiling approach to identify lung tumor

500   endothelial cell heterogeneity and angiogenic candidates. Cancer Cell. 37(1):21–

501   36.

502   40.   Brossa A, Buono L, Fallo S, Pla AF, Munaron L, Bussolati B. (2019) Alternative

503   strategies to inhibit tumor vascularization. Int J Mol Sci. 20:6180.

504   41.   Tamura R, Tanaka T, Akasaki Y, Murayama Y, Yoshida K, Sasaki H. (2019) The

505   role of vascular endothelial growth factor in the hypoxic and immunosuppressive

506   tumor microenvironmnet: perspectives for therapeutic implications. Med Oncol.

507   37(2).

42.   Sakurai Y, Akita H, Harashima H. (2019) Targeting tumor endothelial cells with

nanoparticles. Internaltional J Mol Sci. 20(23):5819.

43.   Zhao Y, Wang X. (2019) Plk4: a promising target for cancer therapy. J Cancer

Res Clin Oncol. 145:2413–22.

44.   Rivera-Rivera Y, Saavedra HI. (2016) Centrosome - a promising anti-cancer

target. Biologics. 10:167–76.

45.   Nakatsu MN, Hughes CCW. (2008) An optimized three-dimensional in vitro model

for the analysis of angiogenesis. Methods Enzymol. 443:65–82.

46.   Hoshijima K, Jurynec MJ, Grunwald DJ. (2016) Precise editing of the zebrafish

genome made simple and efficient. Dev Cell. 36(6):654–67.

47.   Di Donato V, De Santis F, Auer TO, Testa N, Sánchez-Iranzo H, Mercader N, et

al. (2016) 2C-Cas9: a versatile tool for clonal analysis of gene function. Genome

Res. 26(5):681–92.

FIGURE LEGENDS

Figure 1. Excess centrosomes perturb polarity between EC sharing junctions.

(a) Schematic of *Plk4* overexpression in HUVEC. TRE, tet-responsive element; rtTA,

reverse tet transactivator. **(b)** Schematic of H-micropattern with possible centrosome

positions. **(c)** Representative images of EC on H-micropatterns with either 1-2

centrosomes (left) or >2 centrosomes (right). Yellow arrowheads, 1-2 centrosomes with

distal localization; white arrowheads, >2 centrosomes with central localization;

centrosome (γ-tubulin, green); DNA (blue, DRAQ7). Insets, γ-tubulin. Scale bar, 10μm.

(d) Quantification of EC centrosome position based on pattern orientation and EC with

1-2 centrosomes (N) or >2 centrosomes (O). **(e)** Quantification of polarity based on

pattern orientation. **(f)** Quantification of polarity mismatch. N=4 replicates. Statistics: χ^2

analysis. (N:N), both EC 1-2 centrosomes/cell; (O:O), both EC >2 centrosomes/cell;

(N:O) one EC 1-2 centrosomes, one EC >2 centrosomes.

Figure 2. EC with excess centrosomes form disorganized adherens junctions.

(a-a'') Schematics showing potential arrangement of EC on H-patterns. **(b-b'')**

Representative images of cell-cell junctions on H patterns with EC centrosome status

N:N **(b)**, N:O **(b')**, or O:O **(b'')**. EC were stained for γ-tubulin (green, centrosome), VE-

cadherin (red, junctions), and DRAQ7 (blue, DNA). Scale bar, 10μm. **(c-c'')**

Representative VE-cadherin line scans between centrosomes in N:N **(c)**, N:O **(c')**, or

O:O **(c'')** EC junctions. #, positive peak. **(d)** Quantification of VE-cadherin area in EC

junctions with the indicated centrosome complements. **(e)** Quantification of peaks/length

of VE-cadherin line scans of EC junctions with indicated centrosome complements. **(f)**

Quantification of junction width of EC with indicated centrosome complements. N=4

replicates. Statistics: one-way ANOVA with Tukey's correction. *, $p \leq 0.05$, **, $p \leq 0.01$, ***,

$p \leq 0.001$, ****, $p \leq 0.0001$.

Figure 3. Sprouts containing EC with excess centrosomes have abnormal

lumens.

(a) Schematic showing the XY plane of vessel sprout and centrosome position scoring

criteria. **(b)** Representative images showing lumen diameter in EC with indicated

centrosome number. Left panels, XY axis; right panels, YZ axis. Centrin::GFP (green,

centrosome), Moesin1 (red, apical), DNA (cyan, DAPI). Insets, centrin-GFP. Scale bar,

10µm. **(c)** Apical-basal centrosome position in sprouting EC with indicated centrosome

number. N=3 replicates. Statistics: χ^2 analysis. **(d)** Moesin1 staining pattern in EC with

indicated centrosome numbers. N=3 replicates. Statistics: χ^2 analysis. **(e)** Quantification

of lumen diameter in EC with indicated centrosome number. N=3 replicates. Statistics:

one-way ANOVA with Tukey's correction. **, $p \leq 0.01$. **(f)** Quantification of open, narrow,

or closed lumens in EC with indicated centrosome number. N=3 replicates. Statistics: χ^2

analysis.

Figure 4. *Plk4* overexpression in zebrafish results in fewer lumenized vessels.

(a-a'') Representative images of zebrafish tails at 72 hpf perfused with Qdot605 to

assess lumen formation in embryos of indicated genotype and indicated injection with

Cre. Blue arrowheads, open lumens; red arrowheads, narrow lumens; green

arrowheads, closed lumens. Insets, examples of different lumen categories. LSL, lox-

STOP-lox. **(b)** Quantification of ISV perfusion in 72 hpf fish with indicated

gentoypes/condition (WT + Cre, n=5 fish; LSL-Plk4 uninjected, n=31; LSL-Plk4 + Cre,

n=28). **(c)** Quantification of lumen width in 72 hpf fish with indicated genotypes/condition

(WT + Cre, n=86 lumens; LSL-Plk4 uninjected, n=628; LSL-Plk4 + Cre, n=450).

Statistics: one-way ANOVA with Tukey's correction. *, $p{\le}0.05$, ****, $p{\le}0.0001$. **(d)**

Quantification of open, narrow, or closed lumens in vessels of 72 hpf fish with indicated

genotypes/condition. N=5 fish/condition. Statistics: χ^2 analysis. **(e)** Representative

images of ISVs from 72 hpf embryos characterized by presence of *Tg(fli:Gal4)* and

centrosome number. *Tg(kdrl:mCherry)* (red), *Tg(fli:centrin-GFP)* (green). Insets, centrin-

GFP. Scale bar, 20µm. **(f)** Quantification of ISV width with indicated genotypes/condition

(Fli:Gal4 negative, n=4 fish; Fli:Gal4 positive, n=13). Statistics: one-way ANOVA with

Tukey's correction.

a

TRE — ┤×→ Plk4

↓ +Doxycycline

rtTA
TRE →Transcription Plk4

b

Distal
Central
Proximal

c

γ-tubulin
DRAQ7

1-2 centrosomes (N)   >2 centrosomes (O)

d

%EC with centrosome position

p=0.0009
p=0.038
p=0.027

■ Distal
■ Proximal/Central

EC: N   O   N   O
Pattern: N:N O:O N:O N:O

e

% EC polarity arrangement

p=0.014

■ Distal:Distal
■ Other

N:N   O:O

f

% EC polarity arrangement

p=0.026

■ Match
■ Mismatch

N:N   O:O

a N

a' O

a'' O

N

N

O

b γ-tubulin
VE-cadherin
DRAQ7

b'

b''

c
VE-cadherin intensity (AU)
#
Distance (μm)

c'
VE-cadherin intensity (AU)
#
Distance (μm)

c''
VE-cadherin intensity (AU)
#
Distance (μm)

d
VE-cadherin covered area (μm²)
****
*
N:N N:O O:O

e
Peaks/μm
**
***
N:N N:O O:O

f
Junction width (μm)
**
****
N:N N:O O:O

a Basal / Middle / Apical

b
1-2 centrosomes (N)
Centrin::GFP
Moesin1
DAPI

>2 centrosomes (O)

XY   YZ

c
% of EC with centrosome position

p=0.014

Apical
Middle
Basal

N   O

d
Moesin1 pattern % of EC

p=0.015

Normal
Abnormal

N   O

e
Lumen width/ Total vessel width

**

N   O

f
N   O

Open
Narrow
Closed

Total=27   Total=26

p=0.006

SUPPLEMENTARY MATERIAL

Supplementary Figure 1. EC with excess centrosomes do not form proper tight junctions.

Representative images of micropatterns with N:N, N:O, or O:O configurations. EC were labeled for DNA (cyan, DAPI), centrosomes (green, centrin::GFP), adherens junctions (VE-cadherin, red), and tight juntions (ZO-1, gray). Insets, centrin-GFP. Scale bar, 10µm.

Supplementary Figure 2. EC with excess centrosomes have disrupted junctions and lumens in 3D sprouts.

(a) Representative images of 3D sprouts showing junctions in EC with indicated centrosome number. EC are labeled for DNA (cyan, DAPI), centrosomes (green, centrin::GFP), and adherens junctions (VE-cadherin, red). Insets, centrin-GFP. Scale bar, 10 µm. **(b)** Quantification of raw lumen width (not normalized to vessel width). n=3 replicates. Statistics: one-way ANOVA with Tukey's correction.

Supplementary Figure 3. *Plk4* **overexpression in zebrafish leads to excess centrosomes.**

(a) Schematic for Cre-mediated *Plk4* overexpression in zebrafish. **(b)** Representative image of *Plk4* overexpression in 72 hpf zebrafish. Yellow arrowheads, EC centrosome number. **(c)** Quantification of EC with >2 centrosomes in zebrafish with indicated genotypes/conditions (WT + Cre, n=17 fish; LSL-Plk4 uninjected, n=15; LSL-Plk4 + Cre, n=14). Statistics: one-way ANOVA with Tukey's correction. ****, $p \leq 0.0001$. **(d)** Schematic for transient *Gal4-UAS* mediated overexpression of *Plk4* in zebrafish.