1
2
3
4
5 **Excess Centrosomes Disrupt Vascular Lumenization and Endothelial Cell**
6 **Adherens Junctions**

13 Danielle M Berlin[a], Erich J Kushner[b], Katy L Davis[b], and Victoria L Bautch[a,b,c,#]
14
[a]Curriculum in Cell Biology and Physiology, The University of North Carolina at Chapel Hill,
Chapel Hill, North Carolina 27599, USA.
[b]Department of Biology, The University of North Carolina at Chapel Hill, Chapel Hill, North
Carolina 27599, USA.
[c]McAllister Heart Institute, The University of North Carolina at Chapel Hill, Chapel Hill, North
Carolina 27599, USA.
21
22 [#]Corresponding author: Victoria L Bautch, PhD
23   Professor of Biology
24   Department of Biology, CB#3280
25   University of North Carolina at Chapel Hill
26   Chapel Hill, NC 27599 USA
27   Phone: 919-966-6797
28
29   E-mail: bautch@med.unc.edu

30
31 Running head: CENTROSOMES IN VASCULAR LUMENIZATION AND JUNCTIONS
41 WORKING TITLE: 10 words
42 Abstract: 146 words
43 Text: 4,397 words
44 Figures: 4
45
46

ABSTRACT

Proper blood vessel formation requires coordinated changes in endothelial cell polarity and rearrangement of cell-cell junctions to form a functional lumen. One important regulator of cell polarity is the centrosome, which acts as a microtubule organizing center. Excess centrosomes perturb aspects of endothelial cell polarity linked to migration, but whether centrosome number influences apical-basal polarity and cell-cell junctions is unknown. Here, we show that excess centrosomes alter the apical-basal polarity of endothelial cells in angiogenic sprouts and disrupt endothelial cell-cell adherens junctions. Endothelial cells with excess centrosomes had narrower lumens in a 3D sprouting angiogenesis model, and zebrafish intersegmental vessels had reduced perfusion following centrosome overduplication. These results indicate that endothelial cell centrosome number regulates proper lumenization downstream of effects on apical-basal polarity and cell-cell junctions. Endothelial cells with excess centrosomes are prevalent in tumor vessels, suggesting how centrosomes contribute to the dysfunction of tumor vasculature.

Keywords: Endothelial cell, centrosome, junctions, polarity, lumen, angiogenesis

INTRODUCTION

Angiogenesis is the sprouting of new blood vessels from pre-existing vasculature, and it is crucial during development, pregnancy, wound healing, and tumorigenesis [1–3]. Endothelial cells (EC) form new sprouts that then extend, lumenize, and connect with other sprouts and vessels to provide new conduits for blood flow and create a functional vascular network. This process requires dynamic rearrangement of endothelial cell-cell junctions and establishment of apical-basal polarity, such that the apical surface of EC faces the vessel lumen and the basal surface faces the tissue [4–8]. Lumen formation closely follows sprout formation and extension temporally, and many sprout tips have two or more EC that are polarized in the apical-basal axis [9]. However, it is not known how centrosome number contributes to this process.

As tumors grow, hypoxia leads to elevated levels of pro-angiogenic factors such as vascular endothelial growth factor (VEGF-A) that promote neo-angiogenesis to vascularize the tumor. However, the blood vessels surrounding these tumors are often leaky, tortuous, and not properly lumenized [10–12], suggesting defects in EC junctions and polarity. Interestingly, tumor vessels are also characterized by EC with excess centrosomes [13,14]. High VEGF-A signaling in EC results in centrosome overduplication, which affects interphase cells by increasing invasiveness, elevating microtubule (MT) nucleations, changing aspects of polarity and migration, and altering signaling dynamics [13,15–17]. However, it is unclear how excess centrosomes impact

cell-cell interactions and lumenogenesis, which require proper cytoskeletal dynamics

[18,19].

Cadherins form cell-cell adherens junctions that link to the actin cytoskeleton and

restrict the apical vs. basolateral domains in epithelial cells [20,21]. In EC, VE-cadherin

is the primary component of the vascular adherens junction and is required for

localization of apical markers during lumen formation in zebrafish intersegmental

vessels (ISVs) [7]. Depletion of VE-cadherin is embryonically lethal in zebrafish and

mice due to severe vascular defects, including a lack of vascular lumens [22,23]. The

centrosome acts as a major microtubule organizing center and determinant of cell

polarity [24,25]. As EC sprout and lumenize, they are not columnar, but flattened in the

apical-basal plane. Although more filamentous actin appears on the apical side of EC

[26–28], the topology of the centrosome is not well-defined, and the effects of

centrosome perturbation on apical-basal polarity have not been investigated.

Here, we show for the first time that excess centrosomes prevent the proper

polarization of interacting EC. Excess centrosomes resulted in EC with destabilized

adherens junctions and narrow and closed vascular lumens both *in vitro* and *in vivo*.

These findings reveal a novel role for the centrosome in EC apical-basal polarity, and in

junction and lumen formation, and suggest how excess centrosomes in the vasculature

may contribute to poor perfusion in tumor vessels.

RESULTS

Excess centrosomes perturb polarization of junctionally linked EC

We previously showed that excess centrosomes interfere with repolarization along the forward-rearward axis in sprouting EC [15], and thus we hypothesized that supernumerary centrosomes disrupt multiple EC polarity axes. We examined the polarity between EC sharing junctions using an inducible system to overexpress polo-like kinase 4 (Plk4) in human umbilical vein endothelial cells (HUVEC) **(Fig. 1a)**. Plk4 regulates centriole duplication, and its overexpression downstream of a TRE (tet-responsive element) upon addition of doxycycline (DOX) results in supernumerary centrosomes [13,15,29,30]. HUVEC were seeded onto H-shaped micropatterns that allow for polarity assessment between two EC that form a cell-cell junction. Polarity was defined based on centrosome position relative to the nucleus and cell-cell junction, with a "proximal" position being near the cell-cell junction, a "central" position in the nuclear region, and a "distal" position between the nucleus and the cell periphery **(Fig. 1b)**. EC with a normal centrosome number (1-2) typically had a distal centrosome polarity (**Fig. 1c-d**). In contrast, EC with excess centrosomes (>2) showed a significant increase in centrosome proximal polarity, indicating that centrosome number affects EC polarization relative to cell-cell junctions **(Fig. 1c-d)**. Interestingly, when two EC with normal centrosome number (N:N) formed a junction, both cells usually had centrosomes in the distal position, which we termed "normal" polarity **(Fig. 1e)**. All other polarity combinations were deemed abnormal. When both EC forming a junction had excess centrosomes (O:O), polarity was randomized between normal and abnormal **(Fig. 1e)**. Furthermore, N:N EC combinations were more likely to have cells with matching

polarity, while EC with an O:O combination were less likely to have the same polarity as

their neighbor **(Fig. 1f).** Taken together, these data indicate that excess centrosomes

prevent proper polarization of EC relative to cell-cell junctions.

EC with excess centrosomes form disorganized adherens junctions

Because interacting EC with excess centrosomes had perturbed polarity in the

junction axis, we asked whether EC cell-cell interactions were influenced by centrosome

number. Adherens junctions are required for proper apical-basal polarity in epithelial

cells [7,31], and the distribution of VE-cadherin at EC adherens junctions is indicative of

junction maturity; a linear VE-cadherin pattern is associated with mature stabilized

junctions, while a serrated and punctate VE-cadherin pattern is associated with

activated and/or disrupted junctions [32]. To determine how excess centrosomes

influence adherens junctions, we seeded induced tet-Plk4 HUVEC onto H-shaped

micropatterns. Due to the heterogeneous nature of Plk4-induced centrosome over-

duplication a given cell-cell junction could have contributions from two EC with a normal

number of centrosomes (N:N), two EC with excess centrosomes (O:O), or one of each

(N:O) **(Fig. 2a-a")**. We found that junctions with contributions from EC with normal

centrosome numbers had more linear and stable junctions, while junctions with

contributions from EC with excess centrosomes formed more chaotic and disrupted

junctions **(Fig. 2b-b", c-c")**. These chaotic junctions had an increase in the total area of

VE-cadherin **(Fig. 2d)**, and line scans across the junctions revealed a more dispersed

VE-cadherin distribution when one or both EC had excess centrosomes **(Fig. 2e)**.

Overall, these findings indicate that EC with excess centrosomes do not maintain proper

endothelial cell-cell adherens junctions.

EC with excess centrosomes have abnormal lumens

We hypothesized that the disrupted polarity seen between EC with

supernumerary centrosomes had physiological consequences. To examine apical-

basal polarity in EC, we used a 3D sprouting angiogenesis assay in which lumenized

sprouts form that appropriately polarize in the apical-basal axis [9,33]. We analyzed

sprouts formed from DOX-treated tet-Plk4 HUVEC expressing centrin::eGFP and

stained for Moesin1 as an apical marker **(Fig. 3a-b)** and found that EC with normal

centrosome numbers (1-2) typically had apically positioned centrosomes, while EC with

excess centrosomes (>2) had randomized centrosome polarity in the apical-basal axis

(Fig. 3b-c). In addition, EC with excess centrosomes were more likely to have punctate

Moesin1, basal Moesin1, or unpolarized abnormal Moesin1 staining patterns compared

to normal linear and apical patterns **(Fig. 3b, d)**, indicative of polarity defects.

Since both apical-basal polarity and proper junction formation are important for

lumen formation [4–8], we hypothesized that excess centrosomes in EC perturb vessel

lumenogenesis. To normalize for overall vessel structure, we measured the width of the

vessel lumen relative to the total vessel width. We found that EC with excess

centrosomes had narrower lumens compared to EC with normal centrosome numbers

in 3D sprouts **(Fig 3b, e)**. Interestingly, there were several examples where lumens

were completely closed around EC with supernumerary centrosomes. These findings

indicate that EC centrosome number regulates proper vessel lumen diameter, likely

through effects on apical-basal polarity.

Overexpression of Plk4 in zebrafish results in fewer lumenized vessels

To determine if centrosome number affects lumen formation *in vivo*, we fused a

p2a-loxP-mCherry-STOP-loxP-Plk4 transgene to the 3' end of the endogenous *kdrl*

gene for vascular-selective expression of Plk4 after loxP site excision **(Fig. 4a)**. Adult

fish were then crossed to a *Tg(fli:centrin-GFP)* line, which allowed for visualization of

EC centrosomes. F1 embryos were injected with Cre mRNA and *Plk4* overexpression

was determined by loss of the mCherry signal. Fish with *Plk4* overexpression had an

increase in the percentage of EC with >2 centrosomes **(Fig. 4b-c)**. To determine if

overexpression of *Plk4* affected lumenogenesis, we performed microangiography at 72

hpf by injecting fluorescent microcarrier beads into the fish vasculature and examining

the intersegmental vessels (ISVs) for patency. Embryos with *Plk4* overexpression had

fewer perfused ISVs compared to wild type controls or fish lacking Cre **(Fig. 4d-e)**.

Reduced perfusion is indicative of narrower or absent lumens, suggesting that excess

centrosomes induced by *Plk4* overexpression lead to disrupted lumenogenesis *in vivo*.

DISCUSSION

Angiogenesis involves migration and rearrangement of EC, and leads to new

conduits through anastomosis and lumen formation between adjacent EC [1–3]. It is

clear that cell-cell interactions and polarization in several axes are important for lumen

formation, but the role of the centrosome in these processes is poorly understood. Here,

we show that supernumerary centrosomes perturb EC polarity and cell adhesion and

disturb lumen formation *in vitro* and *in vivo*, implying that the centrosome is involved in

both junctional adhesion and the apical-basal polarization that precedes vascular lumen

formation or stabilization.

Our data highlights important roles for the centrosome in maintaining junction

integrity and lumen development in interphase cells. Besides overduplication,

centrosome function can be impacted by centriole loss or structural defects. While over-

duplication allows for progression through the cell-cycle, mis-localization of centrosomal

proteins, structural deficits, and centrosome loss all result in cell-cycle arrest during G1

[34]. The ability to examine effects of centrosome loss on interphase cell behavior is

therefore compromised, highlighting the utility of a centrosome overduplication model to

study centrosome function.

We show that EC cell-cell junctions are disrupted downstream of centrosome

perturbations. It is likely that these junctional defects contribute to poor lumenization *in*

vitro and *in vivo*. EC junctions are linked to vascular lumen formation during

anastomosis, where VE-cadherin facilitates tip cell interactions and subsequent junction

rearrangements necessary to form a continuous lumen [35,36]. Moreover, genetic loss

of VE-cadherin in mice or zebrafish leads to lumen defects, vascular dysfunction, and

embryonic lethality [22,23], supporting a link between disrupted adherens junctions and

malformed lumens in EC sprouts.

Our results reveal that lumens are perturbed following centrosome

overduplication in both an *in vitro* angiogenesis model lacking physiological flow and in

an *in vivo* zebrafish model with blood flow. These findings indicate that lumenal defects

resulting from excess centrosomes do not depend on shear stress exerted by blood

flow. During later stages of angiogenesis, high pressure from blood flow is necessary for

changes in cell morphology resulting in functional lumens [37]. However, lumens can

also form *in vitro* and *in vivo* absent blood flow [7,38]. Hultin and colleagues showed

that disruption of the VE-cadherin junctional complex via the scaffolding protein amotL2

was sufficient to prevent lumenal expansion independent of blood flow through changes

in cell contractility [38]. Thus, it is likely that the profound changes in VE-cadherin

localization at EC cell-cell junctions following centrosome over-duplication alter the

ability of EC to properly coordinate the cytoskeletal forces needed for polarization and

lumenogenesis, regardless of blood flow.

In summary, by providing the first link between centrosome perturbation and

adherens junction stability and lumenization, our data indicate a novel role for the

centrosome in EC polarity, junctions, and lumenization in blood vessels. These

abnormalities may contribute to tumor blood vessel dysfunction.

MATERIALS & METHODS

Cell culture

HUVEC (C2519A, Lonza) were cultured in EBM-2 (CC-3162, Lonza) supplemented with the Endothelial Growth Medium (EGM-2) bullet kit (CC-3162, Lonza) and 1x antibiotic-antimycotic (Gibco). Normal human lung fibroblasts (CC2512, Lonza) were cultured in DMEM (Gibco) with 10% fetal bovine serum (FBS) and 1x antibiotic-antimycotic (Gibco). All cells were maintained at 37°C and 5% CO_2. Tetracycline-inducible Plk4-overexpression HUVEC were generated as previously described [13,15,30]. Viruses were produced by the UNC Lenti-shRNA Core Facility. To induce centrosome overduplication, tet-Plk4 HUVEC were treated with 1µg/ml doxycycline (DOX, D9891, Sigma) for 24hr followed by washout. Experiments were performed 48hr following DOX addition.

3D sprouting angiogenesis assay

The 3D sprouting angiogenesis assay was performed as previously described with the following changes [15,39]. Tet-Plk4 HUVEC were treated with 1µg/ml DOX for 24hr at 37°C. 13µl/ml centrin:eGFP lentivirus and 1µg/ml Polybrene were added to HUVEC 16hr post DOX addition for the last 8hr of DOX treatment. Cells were washed with Dulbecco's phosphate-buffered saline (DPBS) (MT-21-031-CV, Corning) and fresh media was added to remove the virus and DOX. The following day, infected cells were coated onto cytodex 3 microcarrier beads (17048501, GE Healthcare Life Sciences) and embedded in a fibrin matrix as described. Z-stacks were acquired with an Olympus confocal microscope (FV1200 or FV3000) and processed in Image J (National Institutes

of Health). Total vessel width was measured in the YZ plane by drawing a line through

the nucleus and across the entire vessel. Lumen width was measured in the YZ plane

using Moesin1 staining (1:1000, ab52490, Abcam) as a surrogate for the apical surface

and normalized to total vessel width. DAPI (10236276001, Sigma) was used to visualize

the nucleus.

Micropatterns

Large H-micropatterns (CYTOO) were coated with 5µg/ml fibronectin and seeded

with HUVEC according to manufacturer's instructions. Cells were allowed to spread for

24hr, then fixed with ice-cold methanol for 10 min at 4°C.

Immunofluorescence and quantification

Fixed HUVEC were blocked for 1hr at RT in blocking solution (5% FBS, 2X

antibiotic-antimycotic (Gibco), 0.1% sodium azide (s2002-100G, Sigma) in DPBS). Cells

were incubated in primary antibody overnight at 4°C, then washed 3X for 5 min in

DPBS. Secondary antibody and DRAQ7 (1:1000, ab109202, Abcam) were added for

1hr at RT followed by 3X washes for 10 min each in DPBS. Slides were mounted with

coverslips using Prolong Diamond Antifade mounting medium (P36961, Life

Technology) and sealed with nail polish. Primary and secondary antibodies were diluted

in blocking solution. Images were acquired with an Olympus confocal microscope

(FV1200 or FV3000) and analyzed using ImageJ. Centrosome overduplication was

quantified by counting centrosome number in at least 500 cells per replicate. Junction

area on micropatterns was quantified by thresholding the VE-cadherin staining and

measuring total VE-cadherin area in ImageJ. VE-cadherin distribution on doublet

micropatterns was measured in ImageJ by drawing a line across the junction (from

centrosome in first cell to centrosome in second cell) and acquiring a line scan of VE-

cadherin intensity. Peaks were counted for each pattern and normalized to the line

length. Only peaks above half of the maximum VE-cadherin intensity were counted. The

following primary antibodies were used: anti-γ-tubulin (1:5000, T6557, Sigma-Aldrich),

anti-VE-cadherin (1:500, 2500S, Cell Signaling), anti-Moesin-1 (1:1000, ab52490,

Abcam). The following secondary antibodies from Thermo Fisher Scientific were used:

goat anti-mouse AlexaFluor 488 (1:500, A11029), goat anti-mouse AlexaFluor 594

(1:500, A11005), goat anti-rabbit AlexaFluor 488 (1:500, A11034), goat anti-rabbit

AlexaFluor 594 (1:500, A11037).

Zebrafish

Zebrafish (*Danio rerio*) were housed in an institutional animal care and use

committee (IACUC)-approved facility. Zebrafish were maintained as previously

described (Wiley, *et al*., 2011). *Tg(kdrl:mCherry)* was a gift from D. Stainier.

Tg(fli:centrin-GFP) was created by subcloning centrin:GFP into the pCR8/GW/TOPO

middle entry vector (Thermo Fisher) and then gateway cloning into the pDestTol2CG2

plasmid (REF Tol2 kit). Tol2 transposase RNA was generated using the sp6 mMessage

mMachine synthesis kit (AM1340, Thermo Fisher). For overexpression of *Plk4*, *p2a-*

Lox-mCherry-STOP-Lox-Plk4 was fused to the 3' end of the endogenous *kdrl* gene. The

assembled construct was subcloned into the pKHR4 backbone and linearized by IsceI

digest. 200ng/µl sgRNA (5'-*TCTGGTTTGGAAGGACACAG*-3') and 700ng/µl of Cas9

recombinant protein (PNABio) were injected into one-cell stage *tg(fli:centrin-GFP)*

embryos alongside the construct to generate double stranded breaks at the 3' end of

the *kdrl* gene and induce homologous recombination as previous described [40]. F1

were injected with Cre mRNA at the 1-2 cell stage. Cre mRNA was obtained by

subcloning the *Cre* gene from the pME-ERT2-Cre-ERT2 plasmid into the

pCR8/GW/TOPO middle entry vector, gateway cloning into a pCS Dest vector, then

preparing RNA with the sp6 mMessageMachine synthesis kit. pME-ERT2-Cre-ERT2

was a gift from Kryn Stankunas (Addgene plasmid # 82587;

http://n2t.net/addgene:82587; RRID:Addgene 82587). Embryos were collected from

desired crosses and grown in E3 at 28.5°C. At 72 hpf, embryos were injected with

Qdot605 (Q10103MP, Thermo Fisher). Fish were fixed by incubating dechorionated

embryos in ice cold 4% PFA at 4°C overnight. Embryos were rinsed in PBS and

mounted using a fine probe to de-yolk and a small blade to separate the trunk from the

cephalic region and using VECTASHIELD® Hardset™ Antifade Mounting Medium. The

coverslip was sealed with petroleum jelly before imaging on an Olympus FV1200

confocal microscope. Centrosome overduplication was confirmed in Plk4-

overexpressing embryos based on centrin-GFP centriole labeling. ISV perfusion was

measured by counting the number of perfused ISVs based on Qdot605 signal as a

percentage of total ISV number (visualized using *Tg(kdrl:mCherry)*).

Statistics

Student's two-tailed *t* test was used to determine statistical significance in

experiments with 2 groups. One-way ANOVA with Tukey's multiple comparisons test

was used to determine statistical significance for experiments with 3 groups, and X^2 or

two-way ANOVA with Sidak's correction were used for categorical data. Error bars

represent the mean ± standard deviation. Statistical tests and graphs were made using

the Prism 7 software (GraphPad Software).

ACKNOWLEGMENTS

We thank Tony Perdue and the UNC Biology Microscopy Core, the UNC Lenti-

shRNA Core, Maryanna Parker and Allison Marvin for fish room support, and Bautch lab

members for critical discussions.

FUNDING

This work was supported by grants from the National Institutes of Health

(HL43174, HL116719, HL117256, HL139950 to VLB), a K99/R00 (1K99HL124311-

01A1 to EJK), the Integrated Vascular Biology Training Grant (5T32HL069768-17 to

DMB), and an American Heart Association Predoctoral Fellowship (19PRE34380887 to

DMB).

DECLARATION OF INTERESTS: None.

REFERENCES

1. Carmeliet P, Jain RK. (2011) Molecular mechanisms and clinical applications of angiogenesis. Nature. 2011;473(7347):298–307.

2. Bautch VL, Caron KM. (2015) Blood and lymphatic vessel formation. Cold Spring Harb Perspect Biol. 2015;7(3):a008268.

3. Kushner E, Bautch V. (2013) Building blood vessels in development and disease. Curr Opin Hematol. 2013;20(3):231–6.

4. Blum Y, Belting HG, Ellertsdottir E, Herwig L, Luders F, Affolter M. (2008) Complex cell rearrangements during intersegmental vessel sprouting and vessel fusion in the zebrafish embryo. Dev Biol. 2008;316:312–22.

5. Iruela-Arispe ML, Davis GE. (2009) Cellular and molecular mechanisms of vascular lumen formation. Dev Cell. 2009;16(2):222–31.

6. Strilic B, Kucera T, Eglinger J, Hughes MR, McNagny KM, Tsukita S, et al. (2009) The molecular basis of vascular lumen formation in the developing mouse aorta. Dev Cell. 2009;17(4):505–15.

7. Wang Y, Kaiser MS, Larson JD, Nasevicius A, Clark KJ, Wadman SA, et al. (2010) Moesin1 and VE-cadherin are required in endothelial cells during in vivo tubulogenesis. Development. 2010;137(18):3119–28.

8. Lee CY, Bautch VL. (2011) Ups and downs of guided vessel sprouting: the role of polarity. Physiology. 2011;26(5):326–33.

9. Pelton JC, Wright CE, Leitges M, Bautch VL. (2014) Multiple endothelial cells constitute the tip of developing blood vessels and polarize to promote lumen formation. Development. 2014;141(21):4121–6.

10.   Aird WC. (2012) Endothelial cell heterogeneity. Cold Spring Harb Perspect Med.

2012;2(1):a006429.

11.   Hashizume H, Baluk P, Morikawa S, McLean JW, Thurston G, Roberge S, et al.

(2000) Openings between defective endothelial cells explain tumor vessel

leakiness. Am J Pathol. 2000;156(4):1363–80.

12.   Carmeliet P, Jain RK. (2011) Principles and mechanisms of vessel normalization

for cancer and other angiogenic diseases. Nat Rev Drug Discov. 2011;10:417–27.

13.   Kushner EJ, Ferro LS, Liu J-Y, Durrant JR, Rogers SL, Dudley AC, et al. (2014)

Excess centrosomes disrupt endothelial migration via centrosome scattering. J

Cell Biol. 2014;206(2):257–72.

14.   Hida K, Hida Y, Amin DN, Flint AF, Panigrahy D, Morton CC, et al. (2004) Tumor-

Associated Endothelial Cells with Cytogenetic Abnormalities. Cancer Res.

2004;64(22):8249–55.

15.   Kushner EJ, Ferro LS, Yu Z, Bautch VL. (2016) Excess centrosomes perturb

dynamic endothelial cell repolarization during blood vessel formation. Mol Biol

Cell. 2016;27(12):1911–20.

16.   Godinho SA, Picone R, Burute M, Dagher R, Su Y, Leung CT, et al. (2014)

Oncogene-like induction of cellular invasion from centrosome amplification.

Nature. 2014;510(7503):167–71.

17.   Taylor SM, Nevis KR, Park HL, Rogers GC, Rogers SL, Cook JG, et al. (2010)

Angiogenic factor signaling regulates centrosome duplication in endothelial cells

of developing blood vessels. Blood. 2010;116(16):3108–17.

18.   Komarova YA, Huang F, Geyer M, Daneshjou N, Garcia A, Idalino L, et al. (2012)

VE-cadherin signaling induces EB3 phosphorylation to suppress microtubule

growth and assemble adherens junctions. Mol Cell. 2012;48(6):914–25.

19.   Phng L-K, Gebala V, Bentley K, Philippides A, Wacker A, Mathivet T, et al. (2015)

Formin-mediated actin polymerization at endothelial junctions is required for

vessel lumen formation and stabilization. Dev Cell. 2015;32(1):123–32.

20.   Spracklen AJ, Peifer M. (2015) Actin and apical constriction: some (re)-assembly

required. Dev Cell. 2015;35(6):662–4.

21.   Nowotarski SH, Peifer M. (2014) Cell biology: a tense but good day for actin at

cell-cell junctions. Curr Biol. 2014;24(15):R688–90.

22.   Carmeliet P, Lampugnani M-G, Moons L, Breviario F, Compernolle V, Bono F, et

al. (1999) Targeted deficiency or cytosolic truncation of the VE-cadherin gene in

mice impairs VEGF-mediated endothelial survival and angiogenesis. Cell.

1999;98(2):147–57.

23.   Montero-Balaguer M, Swirsding K, Orsenigo F, Cotelli F, Mione M, Dejana E.

(2009) Stable vascular connections and remodeling require full expression of VE-

cadherin in zebrafish embryos. PLoS One. 2009;4(6):e5772.

24.   Kollman JM, Merdes A, Mourey L, Agard DA. (2011) Microtubule nucleations by

gamma-tubulin complexes. Nat Rev Mol Cell Biol. 2011;12:709–21.

25.   Bettencourt-Dias M, Glover DM. (2007) Centrosome biogenesis and function:

centrosomics brings new understanding. Nat Rev Mol Cell Biol. 2007;8:451–63.

26.   Pelaseyed T, Bretscher A. (2018) Regulation of actin-based apical structures on

epithelial cells. J Cell Sci. 2018;131:jcs221853.

27.   Wu S-Z, Bezanilla M. (2018) Actin and microtubule cross talk mediates persistent

421   polarized growth. J Cell Biol. 2018;217(10):3531.

422 28. St Johnston D, Sanson B. (2011) Epithelial polarity and morphogenesis. Curr

423   Opin Cell Biol. 2011;23(5):540–6.

424 29. Habedanck R, Stierhof YD, Wilkinson CJ, Nigg EA. (2005) The Polo kinase Plk4

425   functions in centriole duplication. Nat Cell Biol. 2005;7(11):1140–6.

426 30. Yu Z, Ruter DL, Kushner EJ, Bautch VL. (2017) Excess centrosomes induce p53-

427   dependent senescence without DNA damage in endothelial cells. FASEB J.

428   2017;10:4295–304.

429 31. Iden S, Rehder D, August B, Suzuki A, Wolburg-Buchholz K, Wolburg H, et al.

430   (2006) A distinct PAR complex associates physically with VE-cadherin in

431   vertebrate endothelial cells. EMBO Rep. 2006;7(12):1239–46.

432 32. Bentley K, Franco CA, Philippides A, Blanco R, Dierkes M, Gebala V, et al. (2014)

433   The role of differential VE-cadherin dynamics in cell rearrangement during

434   angiogenesis. Nat Cell Biol. 2014;16(4):309–21.

435 33. Lampugnani M-G, Orsenigo F, Rudini N, Maddaluno L, Boulday G, Chapon F, et

436   al. (2010) CCM1 regulates vascular-lumen organization by inducing endothelial

437   polarity. J Cell Sci. 2010;123:1073–80.

438 34. Mikule K, Delaval B, Kaldis P, Jurcyzk A, Hergert P, Doxsey S. (2007) Loss of

439   centrosome integrity induces p38-p53-p21-dependent G1-S arrest. Nat Cell Biol.

440   2007;9:160–70.

441 35. Lenard A, Ellertsdottir E, Herwig L, Krudewig A, Sauteur L, Belting H-G, et al.

442   (2013) In vivo analysis reveals a highly stereotypic morphogenetic pathway of

443   vascular anastomosis. Dev Cell. 2013;25(5):492–506.

36.   Herwig L, Blum Y, Krudewig A, Ellertsdottir E, Lenard A, Belting H-G, et al. (2011)

Distinct cellular mechanisms of blood vessel fusion in the zebrafish embryo. Curr

Biol. 2011;21(22):1942–8.

37.   Gebala V, Collins R, Geudens I, Phng L-K, Gerhardt H. (2016) Blood flow drives

lumen formation by inverse membrane blebbing during angiogenesis in vivo. Nat

Cell Biol. 2016;18(4):443–50.

38.   Hultin S, Zheng Y, Mahdi M, Vertuani S, Gentili C, Balland M, et al. (2014)

AmotL2 links VE-cadherin to contractile actin fibres necessary for aortic lumen

expansion. Nat Commun. 2014;5:3743.

39.   Nakatsu MN, Hughes CCW. (2008) An optimized three-dimensional in vitro model

for the analysis of angiogenesis. Methods Enzymol. 2008;443:65–82.

40.   Hoshijima K, Jurynec MJ, Grunwald DJ. (2016) Precise editing of the zebrafish

genome made simple and efficient. Dev Cell. 2016;36(6):654–67.

FIGURE LEGENDS

Figure 1. Excess centrosomes perturb polarity between EC sharing junctions.

(a) Schematic of *Plk4* overexpression in HUVEC. TRE, tet-responsive element; rtTA, reverse tet transactivator **(b)** Schematic of H-micropattern with possible centrosome positions. **(c)** Representative images of EC on H-micropatterns with either 1-2 centrosomes (left) or more than 2 centrosomes (right). Yellow arrowheads, normal centrosomes with distal localization. White arrowheads, over-duplicated centrosomes with proximal (top cell) and central (bottom cell) localization. Insets show centrosome number. Scale bar, 10μm. **(d)** Quantification of EC centrosome position. Significance of proximal location for 1-2 centrosomes vs. >2 centrosomes by two-way ANOVA with Sidak's correction. **, $p \leq 0.01$. **(e)** Quantification of polarity pattern. **(f)** Quantification of polarity mismatch. N=4 replicates. Statistics: χ^2 analysis unless otherwise stated. (N:N), both EC 1-2 centrosomes/cell; (O:O), both EC >2 centrosomes/cell.

Figure 2. EC with excess centrosomes form disorganized adherens junctions.

(a-a'') Schematics showing potential arrangements of EC on H-patterns. **(b-b'')** Representative images of cell-cell junctions on H patterns with EC centrosome status N:N **(b)**, N:O **(b')**, or O:O **(b'')**. EC were stained for γ-tubulin (green), VE-cadherin (red), and DNA (blue, DRAQ7). Scale bar, 10μm. **(c-c'')** Representative line scans of VE-cadherin signal between centrosomes in N:N **(c)**, N:O **(c')**, or O:O **(c'')** EC junctions. #, positive peak. **(d)** Quantification of VE-cadherin area in EC junctions with the indicated centrosome complements. **(e)** Quantification of peaks/length of VE-cadherin line scans

of EC junctions with indicated centrosome complements. N=4 replicates. Statistics: one-

way ANOVA with Tukey's correction. *, $p \leq 0.05$, **, $p \leq 0.01$, ***, $p \leq 0.001$, ****, $p \leq 0.0001$.

Figure 3. Sprouts containing EC with excess centrosomes have abnormal

lumens.

(a) Schematic showing the XY plane of a vessel sprout and indicating centrosome

position scoring criteria. **(b)** epresentative images showing lumen diameter in EC with

indicated centrosome number. Left panels, XY axis; right panels, YZ axis. Centrin::GFP

(green, centrosomes), Moesin1 (red, apical), DNA (cyan, DAPI). Insets show

centrosome number. Scale bar, 10μm. **(c)** Apical-basal centrosome position in sprouting

EC with indicated centrosome number. N=3 replicates. Statistics: χ^2 analysis. **(d)**

Moesin1 staining pattern in EC with indicated centrosome numbers. N=3 replicates.

Statistics: X^2 analysis. **(e)** Quantification of lumen diameter in EC with indicated

centrosome number. N=3 replicates. Statistics: one-way ANOVA with Tukey's

correction. **, $p \leq 0.01$.

Figure 4. *Plk4* overexpression in zebrafish results in fewer lumenized vessels.

(a) Schematic for Cre mediated overexpression of *Plk4* in zebrafish embryos. **(b)**

Representative image of *Plk4* overexpression in 72 hpf zebrafish embryo. tg(Fli:Centrin-

GFP) (green), Qdot605 (purple). Yellow arrowheads, centrosomes in EC. Scale bar, 20

μm. **(c)** Quantification of EC with >2 centrosomes in zebrafish with indicated

genotypes/conditions (WT + Cre, n=17; LSL-Plk4 uninjected, n=15; LSL-Plk4 + Cre,

n=14). Statistics: one-way ANOVA with Tukey's correction. ****, $p \leq 0.0001$. **(d-d'')**

representative images of zebrafish tails at 72 hpf perfused with Qdot605 to assess

lumen formation in embryos of indicated genotype and indicated injection with Cre.

DLAV, dorsal longitudinal anastomotic vessel; ISV, intersegmental vessel; CVP, caudal

vein plexus; LSL, lox-STOP-lox. **(e)** Quantification of ISV perfusion in 72 hpf embryos

with indicated gentoypes/condition (WT + Cre, n=5; LSL-Plk4 uninjected, n=31; LSL-

Plk4 + Cre, n=28). Statistics: one-way ANOVA with Tukey's correction. *, $p \leq 0.05$, ****,

$p \leq 0.0001$.

a

TRE → ✗ → Plk4

↓ +Doxycycline

rtTA
TRE → Transcription
Plk4

b

Distal
Central
Proximal

c

γ-tubulin
DRAQ7

1-2 centrosomes

>2 centrosomes

d

%EC with centrosome position

■ Distal
■ Central
■ Proximal

**

1-2   >2
Centrosome #

e

% EC arrangement

■ Normal
■ Abnormal

p=0.014

N:N   O:O
Centrosome #

f

% EC arrangement

■ Match
■ Mismatch

p=0.026

N:N   O:O
Centrosome #

a

Basal
Middle
Apical

b

Centrin::GFP
Moesin1
DAPI

1-2 centrosomes

>2 centrosomes

XY   YZ

c

$p=0.014$

■ Apical
■ Middle
■ Basal

% of EC with centrosome position

1-2   >2

Centrosome #

d

$p=0.015$

■ Normal
■ Abnormal

Moesin1 pattern % of EC

1-2   >2

Centrosome #

e

**

Lumen width/ Total vessel width

0.8

0.6

0.4

0.2

0.0

1-2   >2

Centrosome #

a

kdrl

Cas9
3'-kdrl sgRNA

DSB

kdrl

p2a | mCherry | STOP | Plk4
LoxP | | *LoxP*

kdrl | p2a | mCherry | STOP | Plk4
| *LoxP* | | *LoxP*

Cre

kdrl | p2a | Plk4

b

Tg(Fli:Centrin-GFP)   Qdot605

LSL-Plk4+Cre

72 hpf   DLAV

20μm

CentrinNeonGreen

c

% EC >2 centrosomes

****
****

WT   LSL   LSL
      Plk4   Plk4

Cre: +   -   +

d

WT+Cre

DLAV

ISV

CVP

d'

LSL-Plk4

DLAV

ISV

CVP

d''

LSL-Plk4+Cre

DLAV

ISV

CVP

e

% ISVs Perfused

*
****

WT   LSL   LSL
      Plk4   Plk4

Cre: +   -   +