Characterizing the tolerance of phage therapy candidate OMKO1 to
environmental stress

Michael Blazanin[1,⊗], Wai Tin Lam[1,⊗], Benjamin K. Chan[1], and Paul E. Turner[1,2,*]

[1]Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT 06511, USA.

[2]Program in Microbiology, Yale School of Medicine, New Haven, CT 06520, USA.

[⊗]These authors contributed equally to the work, and are listed in alphabetical order.

*Correspondence to: Paul E. Turner, Department of Ecology and Evolutionary Biology, Yale University,
165 Prospect St, New Haven, CT 06520, USA; paul.turner@yale.edu

Keywords: bacteriophage, thermotolerance, virus

Short title: Tolerance of OMKO1 to environmental stress

Abstract

Antibiotic resistant bacterial pathogens are increasingly prevalent, driving the need for alternative

approaches to chemical antibiotics when treating infections. One such approach is bacteriophage

therapy: the use of bacteria-specific viruses that lyse (kill) their host cells. However, while the effect of

environmental conditions (e.g. elevated temperature) on antibiotic efficacy is well-studied, the

generalized effects of environmental stressors on the potency of phage therapy candidates are seldom

studied. Therapeutic phage OMKO1 infects and kills the opportunistic human pathogen *Pseudomonas*

aeruginosa, while selecting for the evolution of antibiotic re-sensitivity in the target bacterial

population. Here, we used phage OMKO1 as a model to test how a therapeutic virus degrades in

different environments, and whether exposure to an environmental stressor affects subsequent growth

ability on host bacteria. We observed that OMKO1 particles were highly tolerant to different saline

concentrations, but rapidly deactivated at elevated temperatures and under high concentrations of

urea. We also found that exposure to elevated temperature reduced the growth ability of surviving

OMKO1 particles, suggesting a temperature-induced phenotypic shift. Our findings demonstrate that

OMKO1 is highly tolerant to a range of conditions that could be experienced in and outside the human

body, while also showing the need for careful characterization of therapeutic phages to ensure that

environmental exposure does not compromise their expected potency, dosing, and pharmacokinetics.

Introduction

Widespread use of antibiotics – particularly in human therapy and animal agriculture – has selected for the evolution of multi-drug resistant bacterial pathogens, commonly associated with poorer prognosis and higher morbidity in human infections [1, 2]. One alternative or complementary approach to treating bacterial infections with chemical antibiotics is bacteriophage therapy [3], where bacteria-specific viruses with lytic replication cycles are used to kill (lyse) target bacterial cells. With any therapeutic treatment, understanding the effect of environmental conditions on efficacy is paramount. The effects of conditions like high temperature on the stability and application of chemical antibiotics is classically shown [4-7]. In contrast, despite increasing development of phage therapy approaches, the effects of various environmental conditions on phage stability and subsequent reproduction on host cells often remains untested. Here, we use a therapeutic phage for a common human pathogen as a model of phage stability under a range of environmental conditions.

The need for novel approaches to treat bacterial pathogens varies greatly between infections: some strains of bacterial pathogens remain treatable with conventional antibiotics, while others show resistance across multiple drug classes, sometimes to all currently-approved antibiotics [8]. One of these multi-drug resistant bacterial pathogens is *Pseudomonas aeruginosa*, which has been identified by the World Health Organization as a high-priority threat to human health [9]. *P. aeruginosa* is a widespread gram-negative opportunistic pathogen that is commonly found in both natural habitats (e.g., soil, fresh water) and artificial environments (e.g., sewage, households, hospitals, and contaminated medical equipment) [10-13]. For these reasons, *P. aeruginosa* is frequently encountered by humans and causes urinary-tract and respiratory infections in immunocompromised individuals, as well as fouling surgically-implanted materials and devices [14, 15] . Individuals with cystic fibrosis (CF), non-CF bronchiectasis, and chronic-obstructive pulmonary disease (COPD) are especially vulnerable to lung infections caused by *P. aeruginosa*. Treatment of *P. aeruginosa* with chemical antibiotics is often ineffective because the bacteria readily form biofilms that limit the penetrance of antibiotic molecules and due to multi-drug efflux (Mex) systems: protein complexes that actively remove various types of antibiotics from the cell [16-18].

As an alternative or complementary approach to treating *P. aeruginosa*, phage therapy has many attractive advantages, such as the ability for the phage 'drug' to self-amplify within the infection site. However, phage therapy also has some disadvantages, most notably the rapid evolution of bacterial resistance to lytic phage infection [19]. We previously described a naturally-occurring phage that leverages this inevitability of phage-resistance evolution as a strength [20]. The dsDNA phage OMKO1 (virus family Myoviridae) attacks *P. aeruginosa*, while affecting the ability for the target bacteria to maintain resistance to various antibiotics. When bacterial strains evolve resistance to phage OMKO1, the mutants can show drug re-sensitivity, suggesting compromised ability for mechanisms such as MexAB and MexXY efflux pumps to remove antibiotics from the cell. Thus, phage OMKO1 can be doubly effective – the virus kills phage-susceptible bacterial cells, and also can drive evolution of phage resistance-associated loss of antibiotic resistance. This 'evolutionary tradeoff' is a mechanistic example of how phages can synergistically interact with chemical antibiotics, to beneficially extend the usefulness of currently-approved drugs. For example, phage OMKO1 and ceftazidime were used successfully in

emergency treatment of a 76-year-old patient, to resolve a chronic multi-drug resistant *P. aeruginosa*
mediastinal and aortic graft infection [21]. Phage OMKO1 is also currently being tested to resolve or
reduce *P. aeruginosa* infections in the lungs of CF, non-CF bronchiectasis and COPD patients when
administered via aerosol-delivery (nebulizer) treatment.

While phage OMKO1 already has been used successfully for patient treatment, its stability across
environmental conditions remains uncharacterized. Phages for therapeutic use can be manufactured in
large volumes then stored, and treatment can be administered through various routes, including
intravenous, oral suspension, nebulizer, and direct injection to the site of infection. However, it is well-
known that virus particles can degrade over time (i.e., become inactivated for cellular infection), and
that the rate of degradation can depend on the degree of environmental stress, including elevated
temperature, and exposure to high concentrations of salt and urea [22-30]. Because degradation alters
the concentration of active virus particles, it can create inaccuracies for treatment dosing and
administration. In addition, depending on the sensitivity of phage particles to conditions present in the
human body, environmentally-induced degradation could also play a role in the pharmacokinetics
during treatment. Thus, it is vital to test how exposure to environmental stressors affects the titer and
activity of phage-therapy candidates. Here, we use phage OMKO1 as a general model to test the adverse
biological effects of three environmental stressors (salt, urea, and heat) – within and beyond the
conditions experienced in the human body – on the stability and subsequent infectivity of a virus
successfully used in phage therapy.

89
Materials and Methods
Strains
P. aeruginosa strain PA01 was provided by B. Kazmierczak (Yale School of Medicine) and cultured on
1.5% agar plates and liquid media made from Lysogeny Broth (LB: 10 g/L tryptone, 10 g/L NaCl, 5 g/L
yeast extract). Bacteria were incubated at 37°C, and overnight batch cultures were grown with shaking
(100 rpm) in 10mL broth, by inoculating with a single randomly-chosen colony. Bacterial stocks were
stored in 40% glycerol at -80°C.

Phage OMKO1 was originally isolated from an aquatic sample (Dodge Pond, East Lyme, CT) enriched on
PAO1 (Chan et al., 2016). High-titer stocks (lysates) of the phage were obtained by mixing 10μl of the
original stock with 10ml of PAO1 culture in exponential phase. After 12 hours to allow phage population
growth, the mixture was centrifuged and filtered (0.22μm) to remove bacteria to obtain a cell free
lysate. Phage lysates were stored at 4°C. Five biological replicate OMKO1 stocks (S3, S8, S11, S16, R3)
used in some experiments were generated by adding 10μl of the original stock into 5 independent 10ml
PAO1 cultures in exponential phase. For simplicity, these were labeled stocks A, B, C, D, and E,
respectively.

Measuring Bacterial and Phage Densities

The density of a bacterial suspension was estimated using measurements of optical density at 600nm
(OD600), based on a pre-generated standard curve for conversions between OD600 and colony forming
units (CFU) per mL in a bacterial culture.

Estimates of phage titers were obtained via classic plaque-assay methods [31]. Lysate samples were
serially diluted in LB, and 100 µL of phage dilution were mixed with 100 µL overnight PAO1 culture and
added to 4 mL 'soft' (0.75%) agar LB held at 55°C. After gentle vortexing, the mixture was immediately
spread onto a 1.5% agar LB plate. When solidified, plates were inverted and incubated overnight at
37°C. Plates with countable plaques were used to calculate the number of plaque-forming-units (PFUs)
per mL in the lysate.

Analysis

All analysis was carried out in R (3.6.0) using dplyr (0.8.2), figures were made using ggplot2 (3.2.0) and
ggsignif (0.6.0) [32-35].

Urea & salt stress survival
To measure effects of urea (carbamide) and salt (NaCl) on phage stability, 10 µL of phage stock A (see
Strains) was added to 1 mL of either urea or salt solution at a defined concentration, and vortexed to
mix thoroughly. Only a single replicate (n=1) was exposed to each concentration. After a specified time,
50 µL were removed and immediately diluted 200-fold to terminate the stress condition. Then, the
stressed samples were titered in triplicate to calculate survival following stress (except the controls,
0.17M [i.e., LB medium] for saline and 0M for urea, which were titered singly). We then fit a generalized
linear model and carried out an ANCOVA. For urea, the model was of $\log_{10}$(percent survival) as a
function of urea concentration (as a factor) and the interaction between concentration and duration; for
salt, of $\log_{10}$(percent survival) as a function of saline concentration (as a factor) and the interaction
between concentration and duration. Values below the limit of detection and treatments that had fewer
than 3 points above the limit of detection were excluded from statistical analyses and plots.

Heat stress survival
To measure effects of heat on phage stability, for each temperature-duration treatment 100 µL of a
single OMKO1 phage stock (for Fig 3A, only Stock A) stored in LB media was split evenly into two
replicate 250 µL Eppendorf tubes and placed in a pre-heated MyCycler PCR block (BioRad Laboratories,
Inc., Hercules, CA) with an established temperature gradient (30-80°C, at 5°C intervals). Tubes were
removed from the heat block after specified times, and immediately cooled on ice to halt the stressful
condition. The two replicate tubes were mixed together, and then titered in triplicate to calculate
percent survival. We then fit a generalized linear model and carried out an ANCOVA. For Fig 3A, the
model was of $\log_{10}$(percent survival) as a function of temperature (as a factor) and the interaction
between temperature and duration; for Fig 3B, of $\log_{10}$(percent survival) as a function of phage stock
and the interaction of phage stock and duration. Values below the limit of detection and treatments that
had fewer than 3 points above the limit of detection were excluded from statistical analyses and plots.

Measuring phage fitness by bacterial growth curves
To measure phage fitness post heat stress, phages were subjected to heat stress as described above
then titered. These samples were then normalized by concentration to combine a defined number of
post-stress PFUs with bacteria in a 200 µL volume of LB in replicate wells (n = 3 or 6) of a 96-well plate
(Corning Inc., Corning, NY). For most populations, 200 PFUs were inoculated in each well. However,
because of low phage survival rates at longer heat shock times, some wells received fewer PFUs (Fig S2).

Across all wells, however, bacteria and phage were inoculated at a constant multiplicity of infection
(MOI; ratio of phage particles to bacterial cells) of 10^{-5}. Plates were incubated overnight at 37°C and an
automated spectrophotometer (INFINITE F500 microplate reader, TECAN Schweiz AG, Männedorf,
Switzerland) was used to monitor changes in the bacterial density (OD600), as described above. Growth
curves were smoothed and the first local maximum optical density of the bacterial population was used
as a reverse proxy for the fitness of phage [36]; see R script at https://github.com/mikeblazanin/tin-
omko1. We then fit a linear model and carried out an ANOVA for peak bacterial density as a function of
heat shock treatment (as a factor), phage stock, and treatment-stock interaction. Treatments which
were not initially inoculated with 200 PFUs were excluded from statistical analyses. Post-hoc pairwise
comparisons between all levels of the heat shock treatment were performed with Tukey's Honest
Significant Differences.

Results

To estimate the tolerance of phage OMKO1 to salt stress, we exposed samples of phage OMKO1 to
seven different NaCl concentrations ranging from 0.5M to 5M saline (n=1 per treatment), including
0.17M saline (i.e., LB medium) as a control. In each treatment environment, subsamples were obtained
at 5, 30, 60 and 90 minutes to create 28 test measures (7 salt environments x 4 time points). We then
measured the number of phage particles (PFU) that remained viable ('survived') for infection ability on
P. aeruginosa bacteria in each subsample. These measures were compared to the number of phage
particles measured at 5 minutes in the control environment to estimate changes in percent survival over
time. Results (Fig. 1) showed that saline concentration did not significantly alter the rate of particle
decay over time (slope of the lines in Fig 1; ANCOVA: $F_{(8, 16)} = 0.93$, $p = 0.52$). We concluded that phage
OMKO1 viability was unaffected by salt exposure under the conditions tested.

Figure 1. Phage OMKO1 stability is unaffected by saline concentration

To measure phage particle survival of saline stress, phage OMKO1 was exposed to a range of salt concentrations then titered over time. Mean percent survival was calculated and is plotted relative to the titer of the 5-minutes control (0.17M – LB media). The dashed line denotes 100% survival, while the dotted line denotes the limit of detection.

To estimate the tolerance of phage OMKO1 to urea stress, we exposed samples of phage OMKO1 to eleven different urea concentrations ranging from 1M to 10M urea (n=1 per treatment), including 0M urea as a control. In each treatment environment, subsamples were obtained at 5, 30, 60 and 90 minutes, to create 44 test measures (11 urea environments x 4 time points). We then measured the number (PFU) of surviving viable phage particles over time. These measures were compared to the particle number measured at 5 minutes in the 0M control to estimate changes in percent survival over time. Results (Fig 2) showed that urea concentration had a significant effect on the rate of decay of phage particles (slope of the lines in Fig 2; ANCOVA: $F_{(6, 12)} = 128.49$, $p < 0.001$). Specifically, we observed that particle survival was unaffected in 0M urea, whereas phage samples showed significantly negative rates of decay in the five lowest urea concentrations (1M through 5M urea), and that the remaining highest concentrations of urea (6M through 10M urea) caused such rapid particle decay that most measures in these environments were unobtainable (i.e., fell below the limit of detection) (Table 1). We concluded that urea concentration was an environmental factor capable of affecting phage OMKO1 survival, depending on the level and time duration of urea exposure.

Figure 2. Degradation of phage OMKO1 under different urea concentrations.

To measure phage particle survival of urea stress, phage OMKO1 was exposed to a range of urea
concentrations then titered over time. Mean percent survival was calculated and is plotted
relative to the titer of the 0-minute control (0M). The dashed line denotes 100% survival, the
dotted line denotes the limit of detection, and asterisks denote observations that fell below that
limit. Concentrations 7M through 10M were tested but caused such rapid particle decay that all
measures fell below the limit of detection.

**Table 1. Multiple regression shows significant phage OMKO1 decay at 1M and higher urea
concentrations.** Parameter estimates of the rate of decay over time (slope) depending on urea
concentration were evaluated to detect increased decay (decreased slopes) against the null
hypothesis of 0 slope using one-tailed t-tests (df = 12) and a Bonferroni correction (6 tests).

Urea Concentration (M)	Estimated Coefficient	t-value	Bonferroni-adjusted p-value
0	0.0014	2.57	1
1	-0.0023	-3.64	0.010
2	-0.0020	-3.08	0.028
3	-0.0052	-8.26	<0.001
4	-0.0087	-13.70	<0.001
5	-0.0228	-22.04	<0.001
6-10	NA	NA	NA

To estimate the tolerance of phage OMKO1 to heat stress, we exposed samples of phage OMKO1 to six different temperatures ranging from 55°C to 80°C for 5, 30, 60, or 90 minutes, (n = 1 per temperature-duration combination) creating 24 test combinations (6 temperature environments x 4 time points). We then measured the percent of surviving viable phage particles, relative to the number estimated at 5 minutes at 55°C, as described above. Results (Fig 3A) showed a significant effect of temperature shock on phage OMKO1 decay rate (ANCOVA, $F_{(4, 8)}$ = 125.01, p = <0.001). At the two lowest temperatures (55°C and 60°C), there was no significant evidence of OMKO1 decay over time (Table 2). However, at 65°C there was a significant moderate signal of decay, which became more pronounced at 70°C. 75°C and 80°C caused such rapid particle decay that most measures in these environments were unobtainable (i.e., fell below the limit of detection) (Table 1). Abnormal plaque morphology was also observed in heat-treated phages (data not shown), suggesting possible phenotypic alteration of phage OMKO1 following heat treatment, explored further below. We concluded that heat shock was an environmental factor capable of affecting phage OMKO1 survival, depending on the level and time duration of temperature exposure.

Table 2. Multiple regression shows significant phage OMKO1 decay at 65°C and higher temperatures. Parameter estimates of the rate of decay over time (slope) depending on temperature were evaluated to detect increased decay (decreased slopes) against the null hypothesis of 0 slope using one-tailed t-tests (df = 8) and a Bonferroni correction (4 tests).

Temperature (°C)	Estimated Coefficient	t-value	Bonferroni-adjusted p-value
55	0.0008	0.98	1
60	0.0005	0.58	1
65	-0.0028	-3.35	0.020
70	-0.0182	-22.08	<0.001
75-80	NA	NA	NA

Given the observed substantial decay of phage OMKO1 induced by elevated temperatures (Fig. 3A), we next sought to examine whether heat-induced decay continues exponentially over longer periods of time, as well as the degree of variation in heat-induced decay among different stocks of the phage. We generated five independent phage stocks (A through E) and exposed each to a 5-minute control and four prolonged periods (90, 180, 270 and 360 minutes) of 70°C heat stress (n=1 per stock-duration combination). We then measured the percentage of surviving viable phage particles. Results (Fig 3B) showed significant decay over time (ANCOVA effect of duration, $F_{(1, 20)}$ = 519.1, p < 0.001) and some weak variation among biological replicates in decay dynamics (ANCOVA effect of stock, $F_{(4, 20)}$ = 2.54, p = 0.07; ANCOVA interaction between stock and duration, $F_{(4, 20)}$ = 0.95, p = 0.45). Additionally, as expected, we found very similar fit estimates for the comparable decay rate of phage OMKO1 between the two experiments [Fig 3A slope of $\log_{10}$(percent) at 70°C = -0.0167, Fig 3B slope of $\log_{10}$(percent) of stock A = -0.0161]. We concluded that prolonged exposure of phage OMKO1 to a deleterious heat-shock

environment caused reduced percent survival over time, and that biological replicates of the phage
behaved similarly in the assayed conditions.

Finally, we sought to understand whether the effects of temperature might extend beyond the simple
viability of phage particles (as assessed by their ability to form plaques). In particular, we tested how the
duration of high temperature exposure affected the growth ability of surviving phage particles on
susceptible host bacteria. To do so, we took the phage particles that had been heat-shocked at 70°C for
different durations and used dilutions in LB to normalize their concentrations according to how many
plaques they formed. Because of low titers, some populations after 270 minutes of heat shock and all
populations after 300 minutes of heat shock were unable to be standardized to the necessary
concentration (see Fig S1 for data). Then, we inoculated these standardized suspensions with *P.*
aeruginosa host bacteria in assay wells of a 96-well plate. We grew these cultures for >12 hours,
measuring the density (OD600) of bacteria every five minutes using an automated spectrophotometer
(see Methods). As expected, the observations revealed that bacterial density increased initially, peaked
at a maximum value, and then declined over time as the phage in the mixture lysed the bacterial
population (data not shown). As a proxy for phage fitness (virus growth ability on the bacterial
population), we computationally determined the peak bacterial density in each well. Here, the peak
density is thus negatively related to the fitness of the test phage population, because more fit phage
would suppress bacterial density sooner and to a greater degree, producing lower peak densities. For
comparison, we included two controls where bacteria were cultured in the absence of phage for >12
hours as described above, in normal LB medium and in LB medium that was heat-shocked for 360
minutes.

As shown in Fig. 3C, we observed a significant effect of phage OMKO1 heat-shock exposure on
subsequent fitness of the viruses, as measured by peak density of the bacterial population (ANOVA, $F_{(6,}$
$_{49)}$ = 1478.5, $p < 0.001$); in addition, we observed significant variation among the phage stocks in both
peak density in the absence of heat shock, and how peak bacterial density responded to heat-shocked
viruses (ANOVA effect of stock, $F_{(4, 49)}$ = 13.73, $p =\, < 0.001$; ANOVA interaction between stock and
treatment, $F_{(11, 49)}$ = 27.95, $p < 0.001$). As expected, the presence of phage OMKO1 reduced peak
bacterial density relative to the uninfected bacterial controls, indicating that phage negatively affected
bacterial growth (Table 3). Additionally, any phage exposure to heat shock allowed the peak bacterial
density to measurably increase, demonstrating that heat shock generally decreased virus fitness (Tukey
Honest Significant Differences among all heat stress treatments showed significant differences between
0 minutes and all other treatments at $p < 0.01$, see Fig 3C). However, increasing durations of heat shock
exposure did not lead to further measurable increases in peak bacterial density, indicating that phage
fitness did not further decline with heat-shock duration (Tukey HSD finds no significant differences
among 5 minutes through 270 minutes treatments, $p \geq 0.69$). Surprisingly, this remained true even after
extremely long heat shocks, where the survival rate was very low (e.g. after 180 minutes survival was
~0.01%; Fig. 3B). Note that this increase in peak bacterial density could not be explained by the effect of
heat-shocked medium alone, as heat shocked media reduces peak bacterial density (+ Ctrl Shock vs +
Ctrl, two-sample t-test of unequal variance, t = -4.18, df = 8.86, $p = 0.002$). These results indicated that,
in addition to the negative effects of heat shock on phage survival (Figs 3A and B), any exposure to heat

stress reduced the subsequent phenotypic ability (fitness) of phage OMKO1, shown by decreased impact
of viruses on the growth of susceptible bacteria.

281   **Figure 3. Inactivation & fitness suppression of OMKO1 by thermal stress**

A. To measure phage particle survival of heat stress at different temperatures, OMKO1 was exposed to one of a range of temperatures for 5, 30, 60, or 90 minutes, then titered. Percent survival was calculated and is plotted relative to the titer of the 0-minute 55°C treatment. The dashed line denotes 100% survival, while asterisks denote the first observations that fell below the limit of detection (dotted line). 80°C was tested but caused such rapid particle decay that all measures fell below the limit of detection. **B.** To measure phage particle survival of heat stress over longer periods of time, and to measure variation in heat stress tolerance between different stocks of OMKO1, five stocks were exposed to 70°C for 5, 90, 180, 270, or 360 minutes, then titered. Percent survival was calculated and is plotted relative to the source stock. The dashed line denotes 100% survival, while the dotted line denotes the highest limit of detection across the five stocks. **C.** To determine whether phage growth was altered by exposure to heat stress among surviving phage particles, heat shocked phage particles (from Fig 3B) were inoculated with bacteria and grown overnight with repeated measurements of bacterial density. The height of the peak bacterial density was computationally determined and is plotted relative to the negative control wells with no bacteria (dotted line). Thus, this peak height is a reverse proxy for phage fitness. Error bars denote 95% confidence intervals among replicate wells (non-visible error bars are smaller than the points). Positive control bacteria were grown in the absence of phage in LB media ("+ Ctrl") or LB media that had been heat shocked for 360 mins at 70°C ("+ Ctrl Shock"). T-tests showed significant reduction in peak bacterial density between positive controls and unshocked phage (see Table 3), and Tukey Honest Significant Differences were used among all heat shock treatments (see main text).

Table 3. Phage OMKO1 suppression of peak bacterial density is robust to presence/absence of heat-shocked media in controls. Two-sample one-tailed unequal variance t-tests between both positive controls (+ Ctrl is bacteria alone, + Ctrl Shock is bacteria alone in media shocked at 70°C for 360 mins) and each of the 0 minutes heat shock populations show Bonferroni-corrected (10 tests) significant reductions in peak bacterial growth.

Contrast	Estimated Difference	df	t-value	Bonferroni-adjusted p-value
+ Ctrl – 0 mins Phage A	-1.24	5.35	-106.6	<0.001
+ Ctrl – 0 mins Phage B	-0.99	6.38	-65.2	<0.001
+ Ctrl – 0 mins Phage C	-1.26	5.77	-105.6	<0.001
+ Ctrl – 0 mins Phage D	-1.27	5.24	-109.2	<0.001
+ Ctrl – 0 mins Phage E	-1.25	6.29	-101.8	<0.001
+ Ctrl Shock – 0 mins Phage A	-1.16	5.17	-68.9	<0.001
+ Ctrl Shock – 0 mins Phage B	-0.90	6.98	-46.6	<0.001
+ Ctrl Shock – 0 mins Phage C	-1.18	5.38	-69.1	<0.001
+ Ctrl Shock – 0 mins Phage D	-1.18	5.11	-70.5	<0.001
+ Ctrl Shock – 0 mins Phage E	-1.17	5.68	-67.7	<0.001

Discussion

Due to increasing interest in phage therapy to complement or replace traditional chemical antibiotics, we sought to understand the stability and response of phage particles to environmental stressors. Here, we examined how the therapeutic phage OMKO1 responded to possible degradation caused by salt, urea, and heat stressors as a model of the environmental robustness of a phage therapy candidate. We observed no measurable adverse effect of salinity on phage OMKO1 survival (Fig 1), but strong effects of both increased urea concentrations (Fig 2) and elevated temperatures (Fig 3A, B) on virus survival. In addition, we found that heat stress generally induced additional reductions in virus fitness, as heat-shock surviving phage particles had reduced abilities to suppress a susceptible bacterial population when compared to non-stressed viruses (Fig 3C).

High temperatures induced both increased decay of phage particles and decreased phage lysis of their bacterial hosts. Phage OMKO1 particles decay approximately exponentially (Fig 3B, linear on log-linear axes), with higher temperatures increasing the rate of degradation, although we have limited resolution to detect a bi-phasic decay, as observed elsewhere [25]. Intriguingly, however, OMKO1 populations experience a drastic reduction in fitness after even a brief exposure to heat stress, while increased durations of heat stress have little additional effect (Fig 3C). Along with the observed abnormal plaque morphology after heat shock, these results indicate a strong adverse phenotypic response to environmental temperature conditions. We propose two possible explanations for this observation. The first is that the population of OMKO1 particles is heterogenous in resistance to heat shock, and that stability trades off with fitness. While this sort of stability-function tradeoff is thought to be common for proteins [37, 38], this explanation is incompatible with the observation of simultaneous high survival and large decreases in fitness for two of the replicate populations (A and E, see Fig 3B, C). The second explanation is that high temperatures alter viral particle conformations to a lower-fitness state. Thus, any exposure to heat shock alters the state and phenotype of all surviving viral particles, consistent with the observed changes in fitness following heat shock (Fig 3C). While this bistability is environmentally induced, phage particle bistability elsewhere has been previously reported to promote the evolution of novel host use [39] .

Our observation of deactivation of phage OMKO1 by elevated urea concentrations was consistent with previously-published findings on other phages. For example, in some of the earliest work published on this topic tested the survival of particles of coliphages (viruses that specifically infect *Escherichia coli*) over time in a 4.61M urea solution, observing decay anywhere from 0 to 7 orders of magnitude within 30 minutes [27]. Similarly, phage T4 experienced 90% decay after one minute of exposure to 2.5M urea [28]. By comparison, phage OMKO1 decayed more slowly (~80% after 30 minutes at 5M, Fig 2), suggesting that this virus has relatively greater tolerance for elevated urea concentrations compared with the limited evidence from other published studies on phage survival.

In contrast to our findings with urea, our failure to find an effect of saline stress diverged from previously-published results on other phages. Phages T2, T4, T6 and ps$_1$ have been reported to be deactivated when they are rapidly osmotically shocked from 4M to 0M NaCl, and several phages decay in 0M NaCl, all conditions where we observed no effect [26, 29, 30]. Experimental differences may have contributed to these divergent results, including differences in the rate of dilution and thus severity of

osmotic shock (in the former case) or in the shorter length of time our experiments lasted (in the latter case). Additionally, our findings likely reflect some degree of underlying differences in susceptibility to osmotic shock between different phages, where OMKO1 is more tolerant of osmotic shock and saline stress than other reported phages.

Our results are also relevant to the understanding of how environmental stressors might affect use of phage OMKO1 in human therapy. All of our experiments were designed to span (and exceed) the range of environmental conditions that phage particles could experience during storage or within therapy applications in the human body. For example, within the human body the concentrations of salt range from roughly 30mM in urine to 135-145 mM in blood [40], all conditions where we saw no increase in phage OMKO1 decay rate (Fig 1); urea concentrations range from 2.6-6.5mM in human blood up to 325mM in human urine [41], again conditions where we observed no increase in phage decay rate (Fig 2); and human body temperatures can reach a maximum of around 40°C while phage storage conditions could reach perhaps 50°C, again conditions where we expect no effect on phage OMKO1 decay rate based on our findings (Fig 3A). These indications of the general stability of phage OMKO1 despite possible environmental stressors are promising for its further potential uses in clinical applications.

Future studies on the stability of phage OMKO1 and other therapeutic phages should expand the types of examined stressors, consider interactions between multiple stressors, and begin to more deeply elucidate the nature of the observed phage particle bistability. Our findings were limited by the scope and nature of our experimental designs. Additionally, we carried out 14 statistical tests, so we would expect nearly one of those tests to have been significant by chance (we observed 11 to be significant at $p < 0.05$). Future work can expand on the scope of this study by considering the effects of pH and electromagnetic radiation, as well as interactions among any environmental stressors or between these stressors and storage conditions [e.g. freezing, lyophilizing (freeze drying)]. Additionally, biophysical and imaging approaches could be utilized to understand the mechanistic details behind phage particle stability and conformation, like those underlying the observed changes in phage fitness after any duration of heat stress exposure. While these experiments would deepen our understanding of phage biology and stability, our current work has revealed the limits of the stability of phage OMKO1 and highlighted intriguing nonlinear responses of phages to environmental stress, confirming the need for careful characterization and storage of therapeutic phages for widespread clinical use.

Acknowledgements

We thank Alita Burmeister, Caroline Turner, and Michael Wiser for helpful feedback on earlier versions of the manuscript, and the support for Wai Tin Lam from the Chinese University of Hong Kong.

Data Availability

All data and analysis are available at https://github.com/mikeblazanin/tin-omko1.

References

1.   de Kraker ME, Davey PG, Grundmann H. Mortality and hospital stay associated with resistant
Staphylococcus aureus and Escherichia coli bacteremia: estimating the burden of antibiotic resistance in
Europe. PLoS medicine. 2011;8(10):e1001104. Epub 2011/10/25. doi: 10.1371/journal.pmed.1001104.
PubMed PMID: 22022233; PubMed Central PMCID: PMCPMC3191157.
2.   Michael CA, Dominey-Howes D, Labbate M. The antimicrobial resistance crisis: causes,
consequences, and management. Frontiers in public health. 2014;2:145. Epub 2014/10/04. doi:
10.3389/fpubh.2014.00145. PubMed PMID: 25279369; PubMed Central PMCID: PMCPMC4165128.
3.   Kortright KE, Chan BK, Koff JL, Turner PE. Phage Therapy: A Renewed Approach to Combat
Antibiotic-Resistant Bacteria. Cell host & microbe. 2019;25(2):219-32. Epub 2019/02/15. doi:
10.1016/j.chom.2019.01.014. PubMed PMID: 30763536.
4.   Landerkin GB, Katznelson H. Stability of antibiotics in honey and sugar syrup as affected by
temperature. Applied microbiology. 1957;5(3):152-4. Epub 1957/05/01. PubMed PMID: 13435725;
PubMed Central PMCID: PMCPMC1057279.
5.   Wick WE. Influence of antibiotic stability on the results of in vitro testing procedures. Journal of
bacteriology. 1964;87(5):1162-70. Epub 1964/05/01. PubMed PMID: 5874538; PubMed Central PMCID:
PMCPMC277162.
6.   Samara E, Moriarty TF, Decosterd LA, Richards RG, Gautier E, Wahl P. Antibiotic stability over six
weeks in aqueous solution at body temperature with and without heat treatment that mimics the curing
of bone cement. Bone & joint research. 2017;6(5):296-306. Epub 2017/05/19. doi: 10.1302/2046-
3758.65.bjr-2017-0276.r1. PubMed PMID: 28515059; PubMed Central PMCID: PMCPMC5457644.
7.   Svahn O, Bjorklund E. Describing sorption of pharmaceuticals to lake and river sediments, and
sewage sludge from UNESCO Biosphere Reserve Kristianstads Vattenrike by chromatographic
asymmetry factors and recovery measurements. Journal of chromatography A. 2015;1415:73-82. Epub
2015/09/13. doi: 10.1016/j.chroma.2015.08.061. PubMed PMID: 26362805.
8.   Theuretzbacher U. Global antibacterial resistance: The never-ending story. Journal of global
antimicrobial resistance. 2013;1(2):63-9. Epub 2013/06/01. doi: 10.1016/j.jgar.2013.03.010. PubMed
PMID: 27873580.
9.   Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, et al. Discovery,
research, and development of new antibiotics: the WHO priority list of antibiotic-resistant bacteria and
tuberculosis. The Lancet Infectious Diseases. 2018;18(3):318-27. doi: https://doi.org/10.1016/S1473-
3099(17)30753-3.
10.   Fazeli H, Akbari R, Moghim S, Narimani T, Arabestani MR, Ghoddousi AR. Pseudomonas
aeruginosa infections in patients, hospital means, and personnel's specimens. J Res Med Sci.
2012;17(4):332-7. Epub 2012/12/26. PubMed PMID: 23267393; PubMed Central PMCID:
PMCPMC3526125.
11.   Mena KD, Gerba CP. Risk assessment of Pseudomonas aeruginosa in water. Reviews of
environmental contamination and toxicology. 2009;201:71-115. Epub 2009/06/02. doi: 10.1007/978-1-
4419-0032-6_3. PubMed PMID: 19484589.

12.   Remold SK, Brown CK, Farris JE, Hundley TC, Perpich JA, Purdy ME. Differential habitat use and
niche partitioning by Pseudomonas species in human homes. Microbial ecology. 2011;62(3):505-17.
Epub 2011/04/20. doi: 10.1007/s00248-011-9844-5. PubMed PMID: 21503776.
13.   Slekovec C, Plantin J, Cholley P, Thouverez M, Talon D, Bertrand X, et al. Tracking down
antibiotic-resistant Pseudomonas aeruginosa isolates in a wastewater network. PloS one.
2012;7(12):e49300-e. doi: 10.1371/journal.pone.0049300. PubMed PMID: 23284623.
14.   Garau J, Gomez L. Pseudomonas aeruginosa pneumonia. Current opinion in infectious diseases.
2003;16(2):135-43. Epub 2003/05/08. doi: 10.1097/00001432-200304000-00010. PubMed PMID:
12734446.
15.   Mittal R, Aggarwal S, Sharma S, Chhibber S, Harjai K. Urinary tract infections caused by
Pseudomonas aeruginosa: A minireview. Journal of Infection and Public Health. 2009;2(3):101-11. doi:
https://doi.org/10.1016/j.jiph.2009.08.003.
16.   Aeschlimann JR. The role of multidrug efflux pumps in the antibiotic resistance of Pseudomonas
aeruginosa and other gram-negative bacteria. Insights from the Society of Infectious Diseases
Pharmacists. Pharmacotherapy. 2003;23(7):916-24. Epub 2003/07/30. doi:
10.1592/phco.23.7.916.32722. PubMed PMID: 12885104.
17.   Mah TF, Pitts B, Pellock B, Walker GC, Stewart PS, O'Toole GA. A genetic basis for Pseudomonas
aeruginosa biofilm antibiotic resistance. Nature. 2003;426(6964):306-10. Epub 2003/11/25. doi:
10.1038/nature02122. PubMed PMID: 14628055.
18.   Walters MC, 3rd, Roe F, Bugnicourt A, Franklin MJ, Stewart PS. Contributions of antibiotic
penetration, oxygen limitation, and low metabolic activity to tolerance of Pseudomonas aeruginosa
biofilms to ciprofloxacin and tobramycin. Antimicrobial agents and chemotherapy. 2003;47(1):317-23.
Epub 2002/12/25. PubMed PMID: 12499208; PubMed Central PMCID: PMCPMC148957.
19.   Seed KD. Battling Phages: How Bacteria Defend against Viral Attack. PLoS pathogens.
2015;11(6):e1004847. Epub 2015/06/13. doi: 10.1371/journal.ppat.1004847. PubMed PMID: 26066799;
PubMed Central PMCID: PMCPMC4465916.
20.   Chan BK, Sistrom M, Wertz JE, Kortright KE, Narayan D, Turner PE. Phage selection restores
antibiotic sensitivity in MDR Pseudomonas aeruginosa. Sci Rep. 2016;6:26717. Epub 2016/05/27. doi:
10.1038/srep26717. PubMed PMID: 27225966; PubMed Central PMCID: PMCPMC4880932.
21.   Chan BK, Turner PE, Kim S, Mojibian HR, Elefteriades JA, Narayan D. Phage treatment of an
aortic graft infected with Pseudomonas aeruginosa. Evol Med Public Health. 2018;2018(1):60-6. Epub
2018/03/29. doi: 10.1093/emph/eoy005. PubMed PMID: 29588855; PubMed Central PMCID:
PMCPMC5842392.
22.   Pollard EC, Solosko W. The thermal inactivation of T4 and lambda bacteriophage. Biophysical
journal. 1971;11(1):66-74. doi: 10.1016/S0006-3495(71)86195-7. PubMed PMID: 5539000.
23.   Goldhill DH, Turner PE. The evolution of life history trade-offs in viruses. Current opinion in
virology. 2014;8:79-84. Epub 2014/08/05. doi: 10.1016/j.coviro.2014.07.005. PubMed PMID: 25087040.
24.   Jonczyk E, Klak M, Miedzybrodzki R, Gorski A. The influence of external factors on
bacteriophages--review. Folia microbiologica. 2011;56(3):191-200. Epub 2011/06/01. doi:
10.1007/s12223-011-0039-8. PubMed PMID: 21625877; PubMed Central PMCID: PMCPMC3131515.

25. Bleichrodt JF, Blok J, Berends-Van Abkoude ER. Thermal inactivation of bacteriophage phi X174
and two of its mutants. Virology. 1968;36(3):343-55. Epub 1968/11/01. doi: 10.1016/0042-
6822(68)90160-8. PubMed PMID: 5722182.
26. Anderson TF. The reactions of bacterial viruses with their host cells. The Botanical Review.
1949;15(7):464-505.
27. Burnet FM. The Classification of Dysentery-Coli Bacteriophages. III. A Correlation of the
Serological Classification with Certain Biochemical Tests. Journal of Pathology and Bacteriology.
1933;37:179-84.
28. Sato GH. Activation of bacteriophage by urea. Science (New York, NY). 1956;123(3203):891-2.
Epub 1956/05/18. doi: 10.1126/science.123.3203.891. PubMed PMID: 13324109.
29. Seaman PF, Day MJ. Isolation and characterization of a bacteriophage with an unusually large
genome from the Great Salt Plains National Wildlife Refuge, Oklahoma, USA. FEMS microbiology
ecology. 2007;60(1):1-13. Epub 2007/01/26. doi: 10.1111/j.1574-6941.2006.00277.x. PubMed PMID:
17250749.
30. Whitman PA, Marshall RT. Characterization of two psychrophilic Pseudomonas bacteriophages
isolated from ground beef. Applied microbiology. 1971;22(3):463-8. Epub 1971/09/01. PubMed PMID:
4107517; PubMed Central PMCID: PMCPMC376334.
31. Dulbecco R, Vogt M. Some problems of animal virology as studied by the plaque technique. Cold
Spring Harbor symposia on quantitative biology. 1953;18:273-9. Epub 1953/01/01. doi:
10.1101/sqb.1953.018.01.039. PubMed PMID: 13168995.
32. Team RC. R: A Language and Environment for Statistical Computing [Manual]. Vienna, Austria: R
Foundation for Statistical Computing; 2019. Available from: https://www.r-project.org/.
33. Wickham H, François R, Henry L, Müller K. dplyr: A Grammar of Data Manipulation [Manual].
2019. Available from: https://cran.r-project.org/package=dplyr.
34. Wickham H. ggplot2: Elegant Graphics for Data Analysis: Springer-Verlag New York; 2016.
35. Ahlmann-Eltze C. ggsignif: Significance Brackets for 'ggplot2' [Manual]. 2019. Available from:
https://cran.r-project.org/package=ggsignif.
36. Turner PE, Draghi JA, Wilpiszeski R. High-throughput analysis of growth differences among
phage strains. Journal of microbiological methods. 2012;88(1):117-21. Epub 2011/11/22. doi:
10.1016/j.mimet.2011.10.020. PubMed PMID: 22101310.
37. Beadle BM, Shoichet BK. Structural bases of stability-function tradeoffs in enzymes. Journal of
molecular biology. 2002;321(2):285-96. Epub 2002/07/30. doi: 10.1016/s0022-2836(02)00599-5.
PubMed PMID: 12144785.
38. Studer RA, Christin PA, Williams MA, Orengo CA. Stability-activity tradeoffs constrain the
adaptive evolution of RubisCO. Proceedings of the National Academy of Sciences of the United States of
America. 2014;111(6):2223-8. Epub 2014/01/29. doi: 10.1073/pnas.1310811111. PubMed PMID:
24469821; PubMed Central PMCID: PMCPMC3926066.
39. Petrie KL, Palmer ND, Johnson DT, Medina SJ, Yan SJ, Li V, et al. Destabilizing mutations encode
nongenetic variation that drives evolutionary innovation. Science (New York, NY). 2018;359(6383):1542-
5. Epub 2018/03/31. doi: 10.1126/science.aar1954. PubMed PMID: 29599247.
40. Reynolds RM, Padfield PL, Seckl JR. Disorders of sodium balance. BMJ (Clinical research ed).
2006;332(7543):702-5. doi: 10.1136/bmj.332.7543.702. PubMed PMID: 16565125.

41.   Liu L, Mo H, Wei S, Raftery D. Quantitative analysis of urea in human urine and serum by 1H
nuclear magnetic resonance. The Analyst. 2012;137(3):595-600. Epub 12/16. doi: 10.1039/c2an15780b.
PubMed PMID: 22179722.

Figure S1. Fitness suppression of OMKO1 by thermal stress across all tested conditions. To determine whether phage growth was altered by exposure to heat stress among surviving phage particles, heat shocked phage particles (from Fig 3B) were inoculated with bacteria and grown overnight with repeated measurements of bacterial density. The height of the peak bacterial density was computationally determined and is plotted. Thus, this peak height is a reverse proxy for phage fitness. Error bars denote 95% confidence intervals among replicate wells. Positive control bacteria were grown in the absence of phage in LB media ("+ Ctrl") or LB media that had been heat shocked for 360 mins at 70°C ("+ Ctrl Shock"), while negative control wells contained only media. Note that, due to low survival after long durations of heat shock, some growth curves were initially inoculated with reduced numbers of bacteria and phage (holding MOI constant), confounding observed differences in peak bacterial density.