1   Parkinson Disease

2   Flanker task-elicited Event Related Potential sources reflect human recombinant
3   Erythropoietin differential effects on Parkinson's patients

Maria L Bringas Vega[1,2*], Shengnan Liu[1], Min Zhang[1], Ivonne Pedroso Ibañez[2], Lilia M.
Morales Chacon[2], Lidice Galan Garcia[3] Vanessa Perez Bocourt[4], Marjan Jahanshahi[1,5], Pedro
A Valdes-Sosa[1,3*]

1.   The Clinical Hospital of Chengdu Brain Science Institute, MOE Key Lab for Neuroinformation,
University of Electronic Science and Technology of China, Chengdu, China;
2.   Centro Internacional de Restauracion Neurologica CIREN, La Habana, Cuba;
3.   Centro de Neurociencias de Cuba CNEURO La Habana Cuba
4.   Miami Dade College, Florida USA
5.   UCL Queen Square Institute of Neurology, London UK;

*These authors contributed equally to the paper.

Correspondence ML Bringas Vega and Pedro Valdes-Sosa maria.bringas@neuroinformatics-
collaboratory.org pedro.valdes@neuroinformatics-collaboratory.org

16   Abstract

We used EEG source analysis to identify which cortical areas were involved in the automatic
and controlled processes of inhibitory control on a flanker task and compared the potential
efficacy of recombinant-human erythropoietin (rHuEPO) on the performance of Parkinson' s
Disease patients.

The samples were 18 medicated PD patients (nine of them received rHuEPO in addition to
their usual anti-PD medication through random allocation and the other nine patients were
on their regular anti-PD medication only) and 9 age and education-matched healthy controls
(HCs) who completed the flanker task with simultaneous EEG recordings. N1 and N2 event-
related potential (ERP) components were identified and a low-resolution tomography
(LORETA) inverse solution was employed to localize the neural generators.

Reaction times and errors were increased for the incongruent flankers for PD patients
compared to controls. EEG source analysis identified an effect of rHuEPO on the lingual gyri
for the early N1 component. N2-related sources in middle cingulate and precuneus were
associated with the inhibition of automatic responses evoked by incongruent stimuli
differentiated PD and HCs.

From our results rHuEPO, seems to mediate an effect on N1 sources in lingual gyri but not on
behavioural performance. N2-related sources in middle cingulate and precuneus evoked by
incongruent stimuli differentiated PD and HCs.

Introduction

Discovering neuroprotective agents to slow down the progression of Parkinson's Disease (PD)
and, importantly, to improve cognitive deficits is an active area of research (Athauda &
Foltynie, 2015). The search for agents to supplement usual dopaminergic treatments directed
towards motor symptoms is not surprising since the characteristic motor impairment of
patients is usually accompanied by cognitive deficits (Kehagia, Barker, & Robbins, 2010). Since
cognitive dysfunction has a negative impact on the quality of life of patients(Schrag,
Jahanshahi, & Quinn, 2000); finding effective therapies that target cognition in PD is of
paramount importance. As an example, we found that human recombinant erythropoietin
(rHuEPO) (Pedroso et al., 2012) improved general measures of cognition in chronically
medicated PD patients, an additional benefit to that obtained on their usual medical
treatment. This result extends to PD the evidence for neuroprotective properties of rHuEPO
already described in other neurologic diseases (Brines & Cerami, 2005) and is supported by
the anti-apoptotic, anti-inflammatory and cytoprotective effects of EPO in PD animal models
(Sirén, Faßhauer, Bartels, & Ehrenreich, 2009)(Xue, Zhao, & Guo, 2007). This promising result
suggested the need to further study the effect of rHuEPO on cognition in PD.

We believe that to further understand the effect of rHuEPO on cognition in PD patients we
need to examine its effect on specific stages of information processing. This is because the
overt behavioural measures used in our previous study: a) do not have temporal sensitivity,
being the end outcome of many sequential processes, and b) do not reflect localized neural
activity. Consequently, and as a first objective, we zeroed in on very early automatic neural
processes involved in inhibitory control, the lack of which is so common in non-demented PD
patients. This early lack of inhibitory control is easily measured in a number of tasks such as
the Stop signal, go no-go, Stroop, Hayling Sentence Completion task and the Simon task
described in (Obeso et al., 2011) and (Seer, Lange, Georgiev, Jahanshahi, & Kopp, 2016).
However, we decided to use a very well-studied paradigm: Ericksen's Flanker Task (Eriksen &
Eriksen, 1974). It explores the lack of inhibition related to the difficulty in suppressing
interference by incongruent stimuli. It allows the evaluation of very short latency automatic
activation to incongruent flankers around 100 msec. and other, controlled processes, around
200 msec. These produce increased reaction times (RTs) and errors in incongruent trials
versus congruent trials in PD patients in comparison with normal (eg. (P Praamstra, Stegeman,
Cools, & Horstink, 1998; Peter Praamstra, Plat, Meyer, & Horstink, 1999; S A Wylie et al., 2009;
Scott A Wylie, Stout, & Bashore, 2005). It is, however, the early ERP responses that are of
interest here, not the overt behavioural response indexed by the RT which occurs later about
400 msec.

There is no clear way to study these early responses behaviourally. However, these
processes might be probed by direct measurements of fast neural responses such as those
provided by event-related responses (ERPs). In particular, the Flanker task elicits the N1, N2
and P3 ERP components, which are related to automatic and controlled process respectively
(Pires, Leitai, Guerrini, & Simoes, 2014). Here, we will focus only on the early components
N1 and N2. The N1 component has not been, to our knowledge sufficiently studied in the
Flanker task in PD. However, the fronto-central N2 on incongruent trials of flanker tasks in
patients with PD have received more attention (M Falkenstein, Willemssen, Hohnsbein, &
Hielscher, 2006; J. R. Folstein & Van Petten, 2008; Verleger et al., 2010; S A Wylie et al.,
2009; Scott A Wylie et al., 2005). The comparison of medicated PD patients and drug-naïve

de novo PD patients showed that neither the presence of PD (see also (Verleger et al., 2010)
nor dopaminergic medication modulates N2 amplitude variability on incongruent conditions
of flanker tasks (for a discussion see a review of ERP and cognition in PD by Seer et al.,
2017). It seems logical then to determine if the additional cognitive improvement produced
by rHuEPO with respect to dopaminergic treatment, is accompanied by changes in the early
components in the N1 and N2 ERP components, helping us to pinpoint one of the stages of
cognitive processing affected by this drug. Furthermore, in addition to finer grained timing
information, it is possible to leverage source localization methods to identify the neural
sources of any ERP component change.

Therefore, the aim of our study is to use a flanker task to identify if rHuEPO improves
automatic and controlled inhibitory control in PD patients and to locate the neural generators
of these processes. This could be a first step in identifying an ERP biomarker for this type of
cognitive process to be used in clinical trials.

Materials and Methods

Methods:

Description of the Sample and Clinical Trial

Eighteen PD patients (Hoehn and Yahr stages I to III, mean age 53.9, SD 3.2 years) were
recruited at the Clinic of Movement Disorders and Neurodegeneration, Centro International
de Restauracion Neurologica (CIREN) in La Habana, Cuba to participate in a safety clinical
assay of Erythropoietin (rHuEPO) in PD. The design of this investigation, results, scheme of
application and doses employed may be found in (Pedroso et al., 2012). Inclusion criteria were
a clinical diagnosis of idiopathic PD according to the UK Brain Bank criteria and a good
response to dopaminergic treatment and aged between 45-75 years (Hughes, Daniel, Kilford,
& Lees, 1992). Exclusion criteria were manifestation or indicative signs of major cognitive
impairment, psychotic symptoms, and/or presence of other chronic diseases. Nine of the PD
patients, through random allocation, received additionally to their usual anti-parkinsonism
medication, rHuEPO for five weeks and the other nine did not. rHuEPO approved and
registered for use in humans was obtained at the Centro de Inmunologia Molecular, La
Habana Cuba (ior® EPOCIM). There were no significant differences in age, years of education
or duration of illness between the two PD groups. To exclude dementia and major depression,
the Mini Mental State Examination and the Hamilton Depression Scale were respectively
administered (M. F. Folstein, Folstein, & McHugh, 1975; Hamilton, 1960). All patients were
assessed on the motor subscale of the Unified Parkinson´s Disease Rating Scale (UPDRS) both
during "on" (mean 6.3, SD1.1) and "off" medication (mean 21.7, SD 4.3) states.

For the purpose of comparisons, 9 healthy controls (HCs) matched in age (mean 51.2, SD 3.9
years) and educational level were recruited at the same clinic. The PD patients were tested
on their usual anti-parkinsonism medication. The patients and controls signed an informed
consent to participate in this study as a complement of the clinical trial following the CIREN
ethics committee regulations.

Eriksen´s Flanker Task

All participants completed the Eriksen´s Flanker task, while the EEG was simultaneously recorded. Each trial of the task consisted of the presentation of a set of 5 ordered letters (HHHHH or SSSSS) for the congruent condition and 5 letters with H or S at the centre and different laterals or flankers (SSHSS or HHSHH) for the incongruent condition. Participants were instructed to respond to the central letter, whether H or S, by pressing a key with the index finger of the right or left hand respectively. Participants were instructed to respond as fast and as accurately as possible. A total of 480 trials in two blocks, each lasting 8 minutes were completed. In each block 80 stimuli were shown for the congruent condition and 160 for the incongruent. Only the correct responses with reaction times (RTs) >150 and <800 msec. were selected for analysis.

The physical characteristics of the stimuli were black letters on a white frame with a height = 1.5 cm. and Lenght= 7 cm., under 6^0 a visual angle. The distance of the participant to the computer monitor was 60 cm. Each stimulus was presented at the centre of the screen for 190 msec., followed by a fixed interstimulus interval (ITI) of 1735 msec. A training block of 40 stimuli was designed to ensure task instructions were understood.

ERP measurement

The Electroencephalogram (EEG) was continuously recorded at a sampling rate of 512 Hz from 64 electrodes located at standard positions of the International 10/20 System using a Brain Vision system (*https://www.brainproducts.com/products_by_apps.php?aid=5*) (Jasper, 1958). Linked ears were used as on-line reference and the front as earth. To monitor eye movement artefacts, the electro-oculogram (EOG, horizontal and vertical) was recorded from electrodes placed 1 cm to the left and right of the external canthi, and from an electrode beneath the right eye.

Data were filtered using 1-30 Hz and a notch filter to eliminate the 60Hz powerline artefact. All data were referenced using an average reference to all the channels. The baseline was corrected between -200 to - 0 msec. Epochs with electric activity exceeding baseline activity by 100 μV were considered as artefacts and were automatically rejected from further processing (15% of epochs related to hits and 11% of the epochs related to errors). For the analysis, several electrodes were excluded (EOG, ECG, TP9 and TP10).

ERPs were obtained from the EEG recordings for each participant for all the electrodes within the two experimental conditions and averaged over the two groups using Analyzer software (*https://www.brainproducts.com/productdetails.php?id=17*). Epochs of 800 msec. (from -200 msec. (baseline) until 600 msec. post-stimulus onset) were analyzed locked to the stimulus. We selected two windows to examine the stimulus-locked ERPs, using only the correct response averages for the N1 (80-180 msec.) and N2 (200-300 msec.) components in the expected time-windows (see ERPs guidelines in (Picton et al., 2000). Henceforth we will refer to these averages simply as the amplitude of the N1 and N2 components. The average waveform for each participant and each condition was estimated in all the electrodes, but the averaged waveform for group are plotted below for the electrode with the higher statistics amplitudes.

160 In order to localize the generators of the ERP components, a lead field was constructed for
161 each participant to calculate the (volume-constrained) inverse solution, at the two selected
162 latencies using LORETA (Low Resolution Tomography) (*http://www.uzh.ch/keyinst/loreta*) .
163 (Pascual-Marqui et al., 1999)For LORETA, the intracerebral volume is partitioned into 6239
164 voxels at 5mm spatial resolution.

Statistical analysis

We now summarize the experimental design. Our sample is divided into 3 **Groups**: 9
Parkinson patients with the usual treatment (PD Control), 9 patients with the usual treatment
plus EPO (PD rHuEPO), and 9 healthy controls (HCs). Additionally, the ERPs for each
participant was recorded in two conditions: congruent and incongruent.

For each participant the following variables were used in this paper:

1. Reaction Time and errors to the Flanker task
2. Amplitude of the N1 and N2 ERP component at the 60 EEG scalp electrodes.
3. Power of the N1 and N2 sources component for the 6239 source voxels.

The statistical analyses performed were:

a) Reaction Times and errors were analysed using a two-way repeated measure ANOVA
with the with Group (HCs, PD Control and PD rHuEPO) as the between group factor
and the experimental condition (incongruent versus congruent) as the within-subject
repeated measures factor. We report the F statistic and the p value for tests of the
main effect and the interaction. The Greenhouse-Geisser adjustment was applied
since lack of sphericity was observed. These analyses were completed with STATISTICA
7.0.
b) An exploratory analysis of the differences in ERP amplitude topographies between the
HCs and PD Control +PD rHuEPO groups was carried out by means of a multivariate t
test that corrects for multiple comparisons by means of a permutation technique. The
permutation test has the following advantages: the tests are distribution free that
control the experiment-wise error for the simultaneous univariate comparisons, no
assumptions of an underlying correlation structure are required, and they provide
exact p-values valid for any number of subjects, timepoints and all 60 electrodes. The
overall significance level was selected to be 0.05. The method is described in (Galán,
Biscay, Rodríguez, Pérez-Abalo, & Rodriguez, 1997; Galan, Biscay, Valdes, Neira, &
Virues, 1994) as implemented in the software NEEST from Neuronic
http://www.neuronicsa.com/. This allowed the selection of a:
193   1- A subset of electrodes to be subjected to Multivariate Analysis of Variance
194   (MANOVA) to be described in c) below.
195   2- The selection of most representative electrodes to plot the N1 and N2 grand
196   average ERPs.
197   3- The analysis of time intervals to be further studied.
c) Examine for each ERP component, and for their selected group of electrodes, a
repeated measures Multivariate Analysis of Variance (r-MANOVA) for the design
Group by Condition with a significance level set at the 0.05 level. The different

contrasts for the interaction and main effects were tested by using the Wilk's lambda,
approximated by an F function and the p value reported. Note that this allows a
simultaneous confidence interval for contrasts on group differences and to examine
which electrode contribute to the effects. The MANOVA was that implemented in the
STATISTICA 7.0. package.

d) Further analysis for selected differences of the ERP component source images
between selected groups was carried out using the LORETA-built-in voxel-wise
randomization tests with 2000 permutations (Nichols & Holmes, 2001), based on
statistical nonparametric mapping. Voxels with significant differences ($p<0.01$,
corrected for multiple comparisons) between contrasted conditions were located with
the coordinates of the AAL (Automated Anatomical Labelling of Activations) 116
structures atlas of the Montreal Neurological Institute (MNI) (Tzourio-Mazoyer et al.,
2002).

Results

Behavioural results:

Reaction time. The differences between the three groups were significant for Factor Group:
($F_{(2,24)}=7.47$, $p=0.003$), the Condition was not significant as we predicted in the preliminary
analysis ($F_{(2,24)}=3.22$, $p=0.06$). The interaction of Group*Condition also was not significant
($p>0.8$). The contrast between the two groups of patients (PD Control and PD rHuEPO) didn't
show differences in the reaction time ($F_{(2,15)}=0.62$, $p=0.55$). Table 1 shows the performance
of the PD groups separately and Table 2 the fusion of PD patients versus HCs.

Errors. The differences between the errors in the three groups were significant for Factor
Group: ($F_{(2,15)}=10.49$, $p=0.0014$), and for Condition, ($F_{(2,24)}=11.6$, $p=0.0003$), but not for the
interaction Group*Condition ($p=0.1$). The comparison between the two PD groups were
significant only for Condition, incongruent ($F_{(1,16)}=55.3$, $p=0.00001$, and not for the
congruent condition ($F_{(1,16)}=1.88$, $p=0.18$).

	PD rHuEPO n=9		PD Control n=9	
	congruent	incongruent	congruent	incongruent
	Means (SD)	Means (SD)	Means (SD)	Means (SD)
Reaction times msec.	459.33 (71.76)	479.89 (49.43)	460.22 (72.10)	488.22 (63.76)
Percent errors	13.22 (7.76)	43.22 (21.37)	8.78 (6.76)	32.00 (15.79)

Table 1: The results of the reaction times and the percent errors for the congruent and incongruent trials
for the PD patients with and without rHuEPO. The values in the table are means with standard deviations
in parenthesis.

When using the contrast comparing all PD patients and HCs (Table 2), the results were
consistent with previous findings where the RTs increased with incongruent flankers
compared to congruent for both groups.

	(PD rHuEPO + PD Control) n=18		HCs n=9	
	congruent	incongruent	congruent	incongruent
	Means (SD)	Means (SD)	Means (SD)	Means (SD)
Reaction time msec.	459.78 (69.79)	484.06 (55.51)	411.22 (52.00)	431.33 (43.47)
Percent errors	9.00 (3.81)	37.61 (19.12)	3.33 (2.40)	11.00 (7.42)

Table 2: The results of the reaction times and percent errors for the congruent and incongruent trials for the Parkinson's. disease (PD) patients and healthy control (HCs) groups. The values in the table are means with standard deviations in parenthesis.

Exploratory results of ERPs

As mentioned in the Methods, the multivariate t tests corrected for multiple comparisons with permutation tests provides exact p-values, valid for any number of participants, timepoints and recording sites yielded as significant the ERP components in the midline at the 0.05 level. Within this group the most significant ERP was Oz for N1 and Cz for N2 as described in the literature. We will therefore concentrate on these electrode sets henceforth since they all are significant above the globally valid significance threshold.

The same procedure allows, additionally, to select the time windows and which factor (Condition or Group) to be further analyzed. Figure 1 illustrates, for one derivation, the statistics shown above the red line, the latencies with significance for each factor (Group or Condition) in all the time window for analysis. The interaction between them was not significant at any time. The exploratory analysis between experimental conditions did not reflect significant differences in the time range for the early ERP components N1 and N2 (around 100 and 200 msec. respectively).

Note that the significant differences for Condition are in the range of the P300 or later, not in the scope of our study. For that reason, we focus all the further analysis on the incongruent condition, which is the condition which elicits inhibitory control. Nevertheless, henceforth we continue to report the full two-way analysis (Group x Condition), though concentrating on the Group Factor analyses.

Figure 1. Left: t values for the tests of differences between Groups independently of Condition. Right: the t values tests for differences between Condition independently of Group. The read line indicates the statistical significant threshold (corrected for all electrodes and all times by a multivariate permutation test).

Analysis of the N1 component:

We tested the N1 amplitudes with the repeated measures rMANOVA (Group X Condition),
and examined the main effects and the interaction between them. The interaction and the
factor Condition were not significant (p=0.23). However, the main effect of Group was
significant with a Wilk's Lambda=0.40, F(8,42)=2.97, p=0.009. A contrast between the two
groups of patients was also significant with a Wilk's Lambda=0.47, F(4,13)=1.2, p=0.003.
Furthermore, with electrode-wise contrasts 13 electrode sites **F4, FC2, FC4, FC6, C2, C4, C6,**
CP2, O1, O2, Oz, PO3, PO4, PO7, PO8 retained significance. Note that the N1 at the O1
electrode followed the following pattern (See Figure 2): the amplitude of the PD rHuEPO
group (-4.2 μV) was not different statistically from that of the HCs. On the other hand, the
amplitude of the PD Control group (-1.2 μV) was significantly lower.

Figure 2: Left: the group average N1 waveform for each group in the window (80-180 msec.) in the electrode
site O1 with the highest amplitude. The N1 peak was at 152 msec. Right: The Lingual Gyri are the sources of
the N1 component according to AAL coordinates (X=92, Y=76, Z=172). The scale of statistical significance is
self generated using the real values of the original data. All the voxels plotted were significant at p< 0.01).

The localization of the differences between the two Parkinson groups of this component are
localized anatomically by means of the randomized nonparametric test for LORETA. This
showed that the PD rHuEPO had a larger N1 component than the PD Control group at the p<
0.01 level (corrected for multiple comparisons) at the lingual gyri (See right side of Figure 2).

Analysis of the N2 component:

We tested the N2 amplitudes with the repeated measures rMANOVA (Group X Condition),
and examined the interaction and the main effects. The interaction was not significant with a
Wilk's Lambda=0.43, F(6,44)=2.97 and the factor Condition was also not significant (p=0.323).

The main effect of Group (comparing **three groups**) was significant, F(2,24)=6.14, p=0.006, in
seven fronto-central electrodes: Cz(F(2,24)=6.50, p=0.005),CPz (F(2,24)=4.43, p=0.02), CP1
(F(2,24)=5.9, p=0.008), CP2 (F(2,24)=5.6966, p=0.00945), C1(F(2,24)=3.6125, p=0.04251), C2
(F(2,24)=4.6242, p=0.02).

A contrast between the **two groups** of patients was also significant in fronto-central areas,
the electrodes Cz (F(2,24)=4.43, p=0.002), CPz (F(2,24)=6.5, p=0.005) and FC1, FC2, C1, C2
(p<0.05). There were no significant differences between conditions or interaction between
factors.

Note that the N2 grand average at the Cz electrode followed an opposite pattern than N1
(See Figure 4): the amplitude of the PD Control group (-2.10 μV) and healthy controls (-2.46
μV) was not different statistically. On the other hand, the amplitude of the PD rHuEPO group
(-0.67 μV) was significantly lower than both of them. See Table 3 for details of amplitude and
latencies of N2 in Cz.

Figure 3: Left: the N2 waveform averaged by groups in the window (200-300 msec.) in the electrode site Cz
with the highest amplitude. Note for the HC group the early 195 msec. latency and for both PD patients a later
peak around 224 msec. Right: The N2 component showed maximal activation at middle cingulum and
precuneus bilaterally (left located at X=92, Y=108, Z=156). To the right the localization of the precuneus left.
The bicolour scale is showing all the signifcant values after Bonferroni correction and using permutations.

Grupos	Amplitude (μV) CONDITION		Latency (msec.) CONDITION	
	Cz-Cong	Cz-Incong	Cz-Cong	Cz-Incong
HCs	-2.4	- 2.45	195	199
PD Control	- 2.10	- 2.09	226	224
PD rHuEPO	- 0.56	- 0.67	224	223

Table 3: The measures of amplitude and latency of the N2 component for the two conditions Congruent and
Incongruent at the electrode Cz which exhibited the highest amplitude.

The source analysis of the differences (**comparing the three groups**), for the N2 component,
was localized anatomically by means of the LORETA randomized nonparametric test (p< 0.01
level corrected for multiple comparisons) at the the middle cingulum and precuneus
bilaterally. See Figure 3, right side.

In order to know if the errors were related to the N2 amplitude, we select a linear mixed
effect model, and carried out a repeated measures ANOVA log(errors) x Group x amplitudeN2.
But the results were not significant for the interaction of log(errors) with the N2 amplitude,

only   the   main   effect   for   Group   (p=0.001675).   See   Figure   4.

Figure 4: the plot of the N2 and log(errors) of the three groups. Note the variability of the data with 2 outliers of the HCs and 1 outlier of the PD Control group with positive amplitudes of N2.

Discussion

The current study was designed to examine if the novel rHuEPO neuroprotective compound, given to Parkinson patients in addition to their usual medication changed the amplitude of ERP components during an inhibitory control task.

The behavioural results were consisted with previous studies in PD patients in both the rHuEPO and PD Control groups. Both groups showed significantly increased reaction times and a higher number of errors to the incongruent stimuli during the performance of the flanker task as compared to age and education matched HCs. These higher error rates in PD controls are consistent with the proposal that the basal ganglia together with the anterior cingulate (Botvinick, Cohen, & Carter, 2004) participate in the monitoring of incongruence and error monitoring (Brázdil et al., 2002)(Michael Falkenstein, Christ, & Hohnsbein, 2000) which may be impaired in PD due to the dopamine deficiency (for a recent revision of how the progressive dopamine deficiency reduces striatal cholinergic interneuron activity see (McKinley et al., 2019).

It should be noted that we did not find the expected beneficial effect of rHuEPO on behavioural performance (RT and accuracy) in PD patients who received the neuroprotective agent as compared to those that only received the usual treatment. Rather, the differences between groups of patients were found in the ERP components. This is in accordance with our hypothesis that an overall behavioural response might be noisier than some of its time parsed substages. This suggests further studies to identify overt behavioural responses at similar short time scales as ERP components. On the other hand, as it sometimes happens with this type of clinical study the small sample size may lead to lack of power to detect subtle effects.

Regarding the N1 component. This component reflects selective attention, linked to the basic characteristics of a stimulus, and also to the recognition of a specific visual pattern (Luck, Woodman, & Vogel, 2000). N1 amplitude also has been hypothesized to reflect sustained covert visual attention (Di Russo, Martinez, & Hillyard, 2003) being associated with the intensity of covert attention to the central target in the flanker task. In terms of spatial localization, the N1 amplitude is greater in occipital regions (Luck et al., 2000; Mangun & Hillyard, 1990). The neural sources of the N1 in Flanker tasks were located at the brain visual areas of the occipital cortex (Herrmann & Knight, 2001; Hillyard & Anllo-Vento, 1998; Luck et

al., 2000). For example, Bokura (2001) using LORETA identified additional sources of the visual
N1 in the occipito-temporal lobe (Bokura, Yamaguchi, & Kobayashi, 2001) and Zhang (2017)
(Zhang, Brandt, Schrempf, Beste, & Stock, 2017) also localized N1 for Flanker source in extra-
striate visual cortex. We thus expected the differences between PD groups to be localized on
the scalp in the occipital electrodes and the sources to be in brain occipital areas.

This is what we found: the generators of N1, both in the scalp topography and using LORETA,
in the visual areas of the occipital lobe of both hemispheres. The activation of the source for
the PD patients who received rHuEPO was much larger that of the PD group who did not
receive it. In fact, the response of the rHuEPO group became statistically indistinguishable
from that of the HCs, suggestive of a possible neuroprotective effect of rHuEPO on the lingual
gyrus, a region associated with the early and automatic processing of visual stimuli. In
summary, our findings suggested an effect of rHuEPO on the visual attentional window in the
early information-processing stage, thus enhancing the automatic processing of flankers
regardless of their compatibility.

Regarding the N2 component. The second component N2 has been found in several studies
of inhibition using the Flanker task and its amplitude and latency was unaltered in medicated
PD patients (for a review see (Seer et al., 2016)). Van Veen and Carter (Veen & Carter, 2002)
used BESA source localization to study inhibition and response conflict in the Eriksen Flanker
Task, determining that the N2 amplitude associated with incongruent trials can be explained
by a dipole that is located in the ACC. Bokura et al. (Bokura et al., 2001) also conducted an
experiment to understand the anatomical structures that are involved in N2 using a visual
modality of the Flanker paradigm and LORETA which located the N2 generators at cingulate
and the right lateral orbitofrontal cortex.

In our study, we found that the amplitude of N2 component for the PD control and HC groups,
were statistically indistinguishable. But the N2 amplitude in the rHuEPO PD group was
diminished with respect to the other two groups. These effects were topographically located,
as expected, in the fronto-central areas, with neural generators of these differences localised
to the posteromedial portion of the parietal lobe, the precuneus, a structure involved in the
processing of perceptual ambiguities of stimuli (Cavanna & Trimble, 2006) and in the middle
cingulate cortex, probably related to monitoring of conflict in the Flanker task (Enriquez-
Geppert et al., 2013). In comparison with previous reports, we concur with Van Eimeren who
found dysfunction of the default mode network and particularly deactivation of the posterior
cingulate cortex and the precuneus (van Eimeren, Monchi, Ballanger, & Strafella, 2009) in PD
relative to healthy controls, considering these changes in PD closely related to higher errors
in executive tasks in PD compared with healthy controls.

However, in our study the striking decrease of the N2 produced by rHuEPO needs further
research to find an adequate explanation.

Behaviour versus ERPs

Contrary to our expectation, rHuEPO was not associated with a significant improvement in
behavioural performance and did not influence the neural generators of the N2.

The ERP allows neural activity tracking on a millisecond time scale and represents a
continuous measure of information processing, for that reason we selected the ERP to study
a more refined measure of the process of inhibitory control.

This apparent contradiction between behavioural and electrophysiological results could be
related to their different temporal course. Note that the inhibition is a complex process that
can be automatically initiated in the first 100 msec. post-stimulus and extend its action
through both automatic and controlled processes until 800 msec. Reaction time, on the other
hand, started much later >400 milliseconds after the stimulus presentation, with a strong
motor component to complete the response.

Therefore, the aim of our study is to use a flanker task to identify if rHuEPO improves
automatic and controlled inhibitory control in PD patients and to locate the neural generators
in these processes. This could be a first step in identifying an ERP biomarker for this type of
cognitive process to be used in clinical trials.

Limitations

Since this study was completed as part of a safety trial, the samples and the doses employed
were small. This might also explain the lack of clear correlations with behaviour, for example
reaction time with N2 amplitude. Thus, the results require confirmation with larger samples
in future studies. However, the results highlighted the role of EEG source analysis and
advantages of electrophysiology with its high temporal resolution and insensitivity to placebo
effects, in identifying brain changes after an intervention such as rHuEPO.

Conclusions

-We found that rHuEPO improved automatic inhibitory control in PD patients but did not
improve behavioural performance.

-The differences between PD rHuEPO and PD Control groups was in the N1 component at the
lingual gyrus. The differences between PD and healthy controls was on the N2 component in
the cingulate and precuneus.

-Electrophysiology is potentially a useful tool for identifying effects of neuroprotective
compounds on different stages of information processing.

-The components N1 and N2 as well as others like P3 should be further studied as possible
biomarkers for the evaluation of neuroprotective drugs in Parkinson's Disease.

Data Availability

The tables with the behavioural performance (reaction time, hits and errors) and the N2
amplitude for the averaged time window (200-300 milliseconds) of the samples was
submitted in the supplementary material 1. The raw and pre-processing EEG recordings in
BrainVision format with all the individual and grand average potentials for group and
condition can be available by request to maria.bringas@neuroinformatics-collaboratory.org

Conflicts of Interest

The authors declare that there is no conflict of interest regarding the publication of this
paper.

Funding Statement

This paper received support from the NSFC (China-Cuba-Canada) project (No. 81861128001) and the funds from National Nature and Science Foundation of China (NSFC) with funding No. 61871105, 61673090, and 81330032, and CNS Program of UESTC (No. Y0301902610100201).

Acknowledgments

The authors would like to thank to the Centro de Neurociencias de Cuba, specially to Valia Rodriguez and Indira Alvarez for the support during the recordings of the ERPs and the Centro Internacional de Restauracion Neurologica for the recruitment and neuropsychological evaluation of the patients. We are in debt with all the PD patients and their caretakers who volunteer to participated in our study.

Supplementary Materials

The supplementary material 1 consisted in one excel table with the behavioural performance of the subjects during the Flanker task. Tab "Answers": the hits, errors and non-answers in the congruent and incongruent condition. Tab "Reaction Time": the mean and standard deviation (SD) of the hits, errors of each subject in each group for congruent and incongruent trials.

References

Athauda, D., & Foltynie, T. (2015). The ongoing pursuit of neuroprotective therapies in Parkinson disease. *Nature Reviews Neurology*, *11*(1), 25–40. https://doi.org/10.1038/nrneurol.2014.226

Bokura, H., Yamaguchi, S., & Kobayashi, S. (2001). Electrophysiological correlates for response inhibition in a Go/NoGo task. *Clinical Neurophysiology*, *112*(12), 2224–2232.

Botvinick, M. M., Cohen, J. D., & Carter, C. S. (2004). Conflict monitoring and anterior cingulate cortex : an update. *Trends in Cognitive Sciences*, *8*(12), 8–12. https://doi.org/10.1016/j.tics.2004.10.003

Brázdil, M., Roman, R., Falkenstein, M., Daniel, P., Jurák, P., & Rektor, I. (2002). Error processing – evidence from intracerebral ERP recordings. *Experimental Brain Research*, *146*(4), 460–466. https://doi.org/10.1007/s00221-002-1201-y

Brines, M., & Cerami, A. (2005). Emerging biological roles for erythropoietin in the nervous system. *Nature Reviews Neuroscience*, *6*(6), 484–494. https://doi.org/10.1038/nrn1687

Cavanna, A. E., & Trimble, M. R. (2006). The precuneus: a review of its functional anatomy and behavioural correlates. *Brain*, *129*(3), 564–583. https://doi.org/10.1093/brain/awl004

Di Russo, F., Martinez, A., & Hillyard, S. A. (2003). Source analysis of event-related cortical activity during visuo-spatial attention. *Cerebral Cortex*, *13*(5), 486–499. https://doi.org/10.1093/cercor/13.5.486

Enriquez-Geppert, S., Eichele, T., Specht, K., Kugel, H., Pantev, C., & Huster, R. J. (2013). Functional parcellation of the inferior frontal and midcingulate cortices in a flanker-

463   stop-change paradigm. *Human Brain Mapping*, *34*(7), 1501–1514.

464 Eriksen, B. A., & Eriksen, C. W. (1974). Effects of noise letters upon the identification of a
465   target letter in a nonsearch task. *Perception & Psychophysics*, *16*(1), 143–149.
466   https://doi.org/10.3758/BF03203267

467 Falkenstein, M, Willemssen, R., Hohnsbein, J., & Hielscher, H. (2006). Effects of stimulus–
468   response compatibility in Parkinson's disease. A psychophysiological analysis. *Journal of*
469   *Neural Transmission*, *113*, 1449-1462.

470 Falkenstein, Michael, Christ, S., & Hohnsbein, J. (2000). ERP components on reaction errors
471   and their functional significance : a tutorial. *Biological Psychology*, *51*, 87–107.

472 Folstein, J. R., & Van Petten, C. (2008). Influence of cognitive control and mismatch on the
473   N2 component of the ERP: a review. *Psychophysiology*, *45*, 152–170.

474 Folstein, M. F., Folstein, S. E., & McHugh, P. R. (1975). "Mini-mental state": a practical
475   method for grading the cognitive state of patients for the clinician. *Journal of*
476   *Psychiatric Research*, *12*(3), 189–198.

477 Galán, L., Biscay, R., Rodríguez, J. L., Pérez-Abalo, M. C., & Rodriguez, R. (1997). Testing
478   topographic differences between event related brain potentials by using non-
479   parametric combinations of permutation tests. *Electroencephalography and Clinical*
480   *Neurophysiology*, *102*(3), 240–247.

481 Galan, L., Biscay, R., Valdes, P., Neira, L., & Virues, T. (1994). Multivariate statistical brain
482   electromagnetic mapping. *Brain Topography*, *7*(1), 17–28.
483   https://doi.org/10.1007/BF01184834

484 Hamilton, M. A. (1960). A Rating Scale for Depression. *J Neurol Neurosurg Psychiatry*, *23*,
485   56–62.

486 Herrmann, C. S., & Knight, R. T. (2001). Mechanisms of human attention: event-related
487   potentials and oscillations. *Neuroscience & Biobehavioral Reviews*, *25*(6), 465–476.

488 Hillyard, S. A., & Anllo-Vento, L. (1998). Event-related brain potentials in the study of visual
489   selective attention. *Proceedings of the National Academy of Sciences*, *95*(3), 781–787.

490 Hughes, A. J., Daniel, S. E., Kilford, L., & Lees, A. J. (1992). Accuracy of clinical diagnosis of
491   idiopathic Parkinson's disease: a clinico-pathological study of 100 cases. *Journal of*
492   *Neurology, Neurosurgery & Psychiatry*, *55*(3), 181–184.

493 Jasper, H. H. (1958). Report of the committee on methods of clinical examination in
494   electroencephalography. *Electroencephalography and Clinical Neurophysiology*, *10*(2),
495   370–375. https://doi.org/10.1016/0013-4694(58)90053-1

496 Kehagia, A. A., Barker, R. A., & Robbins, T. W. (2010). Neuropsychological and clinical
497   heterogeneity of cognitive impairment and dementia in patients with Parkinson's
498   disease. *The Lancet Neurology*, *9*(12), 1200–1213.

499 Luck, S. J., Woodman, G. F., & Vogel, E. K. (2000). Event-related potential studies of
500   attention. *Trends in Cognitive Sciences*, *4*(11), 432–440.

501 Mangun, G. R., & Hillyard, S. A. (1990). Allocation of visual attention to spatial locations:
502   tradeoff functions for event-related brain potentials and detection performance.
503   *Perception & Psychophysics*, *47*(6), 532–550.

McKinley, J. W., Shi, Z., Kawikova, I., Hur, M., Bamford, I. J., Sudarsana Devi, S. P., … Bamford, N. S. (2019). Dopamine Deficiency Reduces Striatal Cholinergic Interneuron Function in Models of Parkinson's Disease. *Neuron*, *103*(6), 1056-1072.e6. https://doi.org/10.1016/j.neuron.2019.06.013

Nichols, T., & Holmes, A. P. (2001). Nonparametric permutation tests for functional neuroimaging: A primer with examples. *Hum. Brain Mapp.*, *15*, 1–25.

Obeso, I., Wilkinson, L., Casabona, E., Bringas, M. L., Alvarez, M., Alvarez, L., … Jahanshahi, M. (2011). Deficits in inhibitory control and conflict resolution on cognitive and motor tasks in Parkinson's disease. *Exp Brain Research*, *212*, 371–384. https://doi.org/10.1007/s00221-011-2736-6

Pascual-Marqui, R. D., Lehmann, D., Koenig, T., Kochi, K., Merlo, M. C. G., Hell, D., & Koukkou, M. (1999). Low resolution brain electromagnetic tomography (LORETA) functional imaging in acute, neuroleptic-naive, first-episode, productive schizophrenia. *Psychiatry Research: Neuroimaging*, *90*(3), 169–179.

Pedroso, I., Bringas, M. L., Aguiar, A., Morales, L., Alvarez, M., Alvarez, L., & Valdes-Sosa, P. A. (2012). Use of Cuban Recombinant Human Erythropoietin in Parkinson's Disease Treatment. *MEDICC Review*, *14*(1), 11–17.

Picton, T. W., Bentin, S., Berg, P., Donchin, E., Hillyard, S. A., Johnson, R., … Taylor, M. J. (2000). Guidelines for using human event-related potentials to study cognition : Recording standards and publication criteria. *Psychophysiology*, *37*, 127–152.

Pires, L., Leitai, J., Guerrini, C., & Simoes, M. R. (2014). Event-related brain potentials in the study of inhibition: Cognitive control, source localization and age-related modulations. *Neuropsychological Review*, *24*, 461–490. https://doi.org/10.5956/jriet.33.327

Praamstra, P, Stegeman, D. F., Cools, A. R., & Horstink, M. W. (1998). Reliance on external cues for movement initiation in Parkinson's disease. Evidence from movement-related potentials. *Brain: A Journal of Neurology*, *121*(1), 167–177.

Praamstra, Peter, Plat, E. M., Meyer, A. S., & Horstink, M. W. I. M. (1999). Motor cortex activation in Parkinson's disease: Dissociation of electrocortical and peripheral measures of response generation. *Movement Disorders: Official Journal of the Movement Disorder Society*, *14*(5), 790–799.

Schrag, A., Jahanshahi, M., & Quinn, N. (2000). What contributes to quality of life in patients with Parkinson's disease? *Journal of Neurology, Neurosurgery & Psychiatry*, *69*(3), 308–312.

Seer, C., Lange, F., Georgiev, D., Jahanshahi, M., & Kopp, B. (2016). Event-related potentials and cognition in Parkinson's disease : An integrative review. *Neuroscience & Biobehavioral Reviews*, *69*, 691–714.

Seer, C., Lange, F., Loens, S., Wegner, F., Schrader, C., Dressler, D., … Kopp, B. (2017). Dopaminergic modulation of performance monitoring in Parkinson's disease: An event-related potential study. *Scientific Reports*, *7*(December 2016), 1–13. https://doi.org/10.1038/srep41222

Sirén, A., Faßhauer, T., Bartels, C., & Ehrenreich, H. (2009). Therapeutic Potential of Erythropoietin and its Structural or Functional Variants in the Nervous System.

Neurotherapeutics, *6*(January), 108–127.

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., …
Joliot, M. (2002). Automated anatomical labeling of activations in SPM using a
macroscopic anatomical parcellation of the MNI MRI single-subject brain. *Neuroimage*,
15(1), 273–289.

van Eimeren, T., Monchi, O., Ballanger, B., & Strafella, A. P. (2009). Dysfunction of the
default mode network in Parkinson disease: a functional magnetic resonance imaging
study. *Archives of Neurology*, *66*(7), 877–883.

Veen, V. van, & Carter, C. S. (2002). The timing of action-monitoring processes in the
anterior cingulate cortex. *Journal of Cognitive Neuroscience*, *14*(4), 593–602.

Verleger, R., Hagenah, J., Weiss, M., Ewers, T., Heberlein, I., Pramstaller, P. P., … Klein, C.
(2010). Responsiveness to distracting stimuli, though increased in Parkinson's disease,
is decreased in asymptomatic PINK1 and Parkin mutation carriers. *Neuropsychologia*,
48(2), 467–476. https://doi.org/10.1016/j.neuropsychologia.2009.10.004

Wylie, S A, Wildenberg, W. P. M. Van Den, Ridderinkhof, K. R., Bashore, T. R., Powell, V. D.,
Manning, C. A., & Wooten, G. F. (2009). The effect of Parkinson ' s disease on
interference control during action selection. *Neuropsychologia*, *47*, 145–157.
https://doi.org/10.1016/j.neuropsychologia.2008.08.001

Wylie, Scott A, Stout, J. C., & Bashore, T. R. (2005). Activation of conflicting responses in
Parkinson's disease: evidence for degrading and facilitating effects on response time.
Neuropsychologia, *43*(7), 1033–1043.
https://doi.org/10.1016/j.neuropsychologia.2004.10.008

Xue, Y., Zhao, L., & Guo, W. (2007). Intrastriatal administration of erythropoietin protects
dopaminergic neurons and improves neurobehavioral outcome in a rat model of
Parkinson's disease. *Neuroscience*, *146*, 1245–1258.
https://doi.org/10.1016/j.neuroscience.2007.02.004

Zhang, R., Brandt, M. D., Schrempf, W., Beste, C., & Stock, A.-K. (2017). Neurophysiological
mechanisms of circadian cognitive control in RLS patients-an EEG source localization
study. *NeuroImage: Clinical*, *15*, 644–652.