Grapevine rootstocks affect growth-related phenotypes

Running title: Grapevine rootstocks affect growth

Zoë Migicovsky[1*], Peter Cousins[2], Lindsay M. Jordan[3], Sean Myles[1], Richard Keith Striegler[3],

Paul Verdegaal[4], Daniel H. Chitwood[5,6*]

[1]Department of Plant, Food and Environmental Sciences, Faculty of Agriculture, Dalhousie

University, Truro, NS B2N 5E3, Canada

[2]E. & J. Gallo Winery, Modesto, CA, 95354, USA

[3]E. & J. Gallo Winery, Acampo, CA 95220, USA

[4]University of California Cooperative Extension, San Joaquin Valley, Stockton, CA, 95206,

USA

[5]Department of Horticulture, Michigan State University, East Lansing, Michigan, 48824, USA

[6]Department of Computational Mathematics, Science and Engineering, Michigan State

University, East Lansing, Michigan, 48824, USA

*Corresponding authors

Abstract

Grape growers use rootstocks to provide protection against pests and pathogens and to modulate

viticulture performance such as shoot growth. Our study examined two grapevine varieties

('Chardonnay' and 'Cabernet Sauvignon') grafted to 15 different rootstocks and determined the

effect of rootstocks on eight traits important to viticulture. We assessed the vines across five

years and identified both year and variety as contributing strongly to trait variation. The effect of

rootstock was relatively consistent across years and varieties, explaining between 8.99% and

9.78% of the variation in growth-related traits including yield, pruning weight, berry weight, and

Ravaz index (yield to pruning weight ratio). Increases in yield due to rootstock were generally

the result of increases in berry weight, likely due to increased water uptake by vines grafted to a

particular rootstock. We determined that it is possible to achieve an over 50% increase in yield,

pruning weight, or Ravaz index by choosing the optimal rootstock, indicating that rootstock

choice is crucial for grape growers looking to improve vine performance.

Introduction

Grafting joins two distinct plant parts: a scion (shoot system) from a donor plant and a rootstock

(root system) from a second plant to which the scion is attached. The practice of grafting chiefly

enables clonal propagation but can also have many other benefits, such as reducing the juvenility

period (increasing precocity) or size (dwarfing) in fruit trees[1–3].

In grapevines (*Vitis vinifera* L.), widespread use of grafting began in the late 1800s, following

the introduction of phylloxera (*Daktulosphaira vitifoliae* Fitch) to Europe from North America.

While *V. vinifera* is highly susceptible to phylloxera, which feeds on the roots of grapevines,

eastern North American *Vitis* species evolved in the presence of phylloxera and are tolerant

and/or resistant to it. By grafting *V. vinifera* scions to rootstocks of other *Vitis* species, *V. vinifera*

could be grown in European soils containing phylloxera, rescuing the wine industry[4].

Ten years after its detection in Europe, own-rooted (ungrafted) grapevines with phylloxera were first identified in California. The inter-continental spread of the pest was likely due to the importation of vines from European nurseries or from eastern North America[5]. However, due to the sandy soils of California's Central Valley (or San Joaquin Valley, specifically), phylloxera infections were not as severe and did not require the immediate use of rootstocks[6]. By the 1950s, less than 30% of California grapevines were grafted onto phylloxera-resistant rootstocks[7]. Still, over time, the California grapevine industry transitioned primarily to grafted vines. Currently, more than 80% of vineyards worldwide grow grafted vines[4].

In addition to allowing *V. vinifera* vines to grow in phylloxera-infested soils, grapevine rootstocks can provide tolerance to several other damaging pests and diseases including root-knot and dagger nematodes[8–10]. Rootstocks may also be used to improve resilience to abiotic stresses such as salinity[11] and drought[12]. Grafting grapevines to a particular rootstock can influence a wide range of traits in the scion including mineral composition[13], berry chemistry[14], and berry maturation[15].

Of particular interest to grape growers is the observation that rootstock choice can affect vine size and yield[16]. While an increase in yield is desirable, increasing vine size or vegetative growth also increases the cost of managing the vine, due to additional labour for vine training, pruning, and fruit thinning. An ideal rootstock will increase reproductive growth, or yield, without an accompanying increase in vegetative growth, which is assessed by measuring pruning weight or the amount of one-year-old dormant cuttings removed during the winter. The Ravaz index, or

yield divided by pruning weight from the following dormant season, can be calculated to

determine the relative ratio of reproductive to vegetative growth.

With all the potential benefits offered by a rootstock, deciding which one to use is an important

choice. While other changes to vineyard management can be made throughout the lifespan of the

vines–such as altering irrigation, fertilizer, pesticides, and pruning–rootstock choice is made only

once. In this study, we assessed eight traits of viticultural importance across two scion varieties

('Chardonnay' and 'Cabernet Sauvignon') grafted to 15 different rootstocks. The vines were

grafted near Lodi in San Joaquin County, California, in 1992 and evaluated from 1995 to 1999 in

order to determine the relative contributions of variety, year, and rootstock to phenotypic

variation.

Results

There was strong variation in phenotypes across the years of the study, with reduced yield and

fewer clusters in 1996 and 1998. Vines generally had higher pruning weights in 1996. However,

the relative rankings of the rootstocks were generally consistent across years (Figure 1).

Figure 1. Phenotypic variation across years (1995 to 1999) for each rootstock by scion

combination. Loess smoothing lines are plotted.

91 Using a linear model, we identified year as the largest source of variation captured by our data

92 (Figure 2). Year was a significant term for all phenotypes, ranging from 8.46% (pH) to 52.04%

93 (pruning weight). Year explained over 40% of the variation in pruning weight, Ravaz index, and

94 yield.

97 **Figure 2. Phenotypic variation explained by factors of interest estimated using a linear**

98 **model.** For each phenotype, the linear model was optimized by removing non-significant

99 interaction effects. For factors which explain a significant amount of variance ($p < 0.05$), the

100 percent variance explained is indicated using colour. Position in the vineyard (column) was

101 included in the model but is not plotted. Phenotypes are sorted in order of the most variance

102 explained by rootstock.

105 Grapevine variety explained a significant amount of variation in all traits except Ravaz index,

106 with the strongest effect for TA (32.12%), cluster number (31.98%), berry weight (26.91%), and

SSC (21.32%). The interaction between year and variety was significant for all traits, and over

20% of the variation in pH, SSC, and TA could be explained by this term.

Rootstock had a significant effect on all phenotypes and explained between 8.99% and 9.78% of

the variation in yield, berry weight, pruning weight, and Ravaz index. For half the phenotypes

assessed, the interaction between rootstock and year was removed from the model because it did

not explain a significant amount of variation in the trait. For the remaining traits, the interaction

between rootstock and year explained 4.31% to 7.68% of the variation (Figure 2).

While the interaction between variety and rootstock was retained as a significant term for all

phenotypes except TA and Ravaz index, it explained less than 4% of the variation in any given

phenotype.

Focusing on the phenotypes in which rootstock showed the strongest effect, we plotted the

distributions for yield (9.78%), berry weight (9.62%), pruning weight (9.14%), and Ravaz index

(8.99%) and compared each of the 15 rootstocks using a Tukey test (Figure 3). Across these

phenotypes, 'Ramsey' had among the highest yields, berry weights, and pruning weights, and

one of the lowest Ravaz indexes. The yield for 'Ramsey' was significantly higher than eight of

the other rootstocks evaluated. Similarly, 'Freedom' ranked within the top four for yield, berry

weight, and pruning weight measurements, but ranked 11th for Ravaz index. However,

'Freedom' and 'Ramsey' only had a significantly lower Ravaz index than '420 A'.

Figure 3. Variation in (A) yield, (B) berry weight, (C) pruning weight, and (D) Ravaz index

across vines grafted to 15 different rootstocks. Rootstocks are ordered from highest to lowest

median values. Tukey test results are reported from a linear model accounting for variation in

133 variety, year, position in the vineyard (column), and interaction effects. Rootstocks with the

134 same letter (indicated inside the plot) are not significantly different from each other.

136 The roostock '775 P' generated the lowest yields and smallest berries, with only 'SO4' and

137 '5BB' not differing significantly for these two phenotypes. In contrast, '775 P' was ranked 6th

138 for pruning weight, which resulted in a significantly lower Ravaz index than all other rootstocks

139 except '3309 C', '1103 P', and 'Freedom', although this trend is likely due primarily to the low

140 yield of 'Chardonnay' grafted to '775 P' (Figure S3). In comparison, '420 A' ranked 5th for

141 yield and had the lowest pruning weight, thus resulting in a Ravaz index which was significantly

142 higher than all other rootstocks.

144 For both 'Chardonnay' and 'Cabernet Sauvignon', Ravaz index was significantly correlated with

145 most other phenotypes with the exception of TA ('Chardonnay') and SSC ('Cabernet

146 Sauvignon') (Figure 4, Table S4). Ravaz index was positively correlated with cluster number for

147 'Chardonnay' ($r = 0.250$, $p = 3.398 \times 10^{-4}$) and 'Cabernet Sauvignon' ($r = 0.671$, $p < 1 \times 10^{-15}$).

148 Yield was not significantly correlated with pruning weight for either variety but it was positively

149 correlated with cluster number ('Chardonnay': $r = 0.634$, $p < 1 \times 10^{-15}$; 'Cabernet Sauvignon': r

150 $= 0.722$, $p < 1 \times 10^{-15}$) and berry weight ('Chardonnay': $r = 0.489$, $p < 1 \times 10^{-15}$; 'Cabernet

151 Sauvignon': $r = 0.579$, $p < 1 \times 10^{-15}$).

Figure 4. Spearman's correlations among phenotypes for (A) 'Chardonnay' and (B)

'Cabernet Sauvignon'. P-values were Bonferroni-corrected for multiple comparisons with a

particular variety.

Lastly, we evaluated the percent change between the best and worst performing rootstocks for

each phenotype (Figure 5, Table S5). The percent change ranged from 3.10% for SSC to 93.78%

for Ravaz index. In addition to Ravaz index (93.78%), cluster number (35.42%), pruning weight

(70.82%), and yield (77.35%) all increased by over 30%, while the remaining phenotypes

increased by less than 15%.

Figure 5. Percent change in each phenotype from rootstock with the lowest median to the

rootstock with the highest median. Phenotypes are ordered from largest percent change to

lowest percent change. Raw values are also listed.

Discussion

In California, most wine regions annually receive sufficient rainfall during the dormant season to

support desired canopy growth. However, there are years where low dormant season rainfall may

reduce canopy growth[25]. Previous work examining 'Merlot' vines across two years in California

found that soil moisture level during the dormant season impacted both vegetative and

reproductive growth, even when irrigation is applied after budbreak[25]. In our study, there was

less rainfall during the 1994 and 1997 dormant seasons when floral initiation would have

occurred. While the reduction in yield we observed in 1998 may be due to a dry 1997 dormant

season, the dormant season prior to 1996 had higher rainfall and it's unclear why the yield was

not higher. Thus, at least for the 1998 growing season, a reduction in rainfall the previous year

may have had a more severe impact on reproductive growth in contrast to vegetative growth.

In contrast to our study, previous work found that the reduction to pruning mass due to dormant

season rainfall was more severe than the reduction to yield, increasing the Ravaz index for vines

which did not receive rainfall[25]. While this is not consistent with our results, we do find that year

is the largest source of variation in growth-related traits, confirming that variable environmental

conditions between years, such as access to water during the dormant season, plays a crucial role

in plant growth and development.

During the growing season, grape growers can use management practices such as irrigation to

partly buffer against year-to-year variation[26]. The vines in this study were all irrigated using the

same management practices across five years, with a moderate RDI program of 80% from berry

set to veraison, therefore reducing the impact of weather fluctuations during the growing season,

When included in a linear model, variety explained over 20% of the variation we observed for

TA, berry weight, and SSC, indicating that there is a strong variety-specific effect on many berry

characteristics. Year, or vintage, had a significant interaction with variety for all phenotypes and

explained over 20% of the variation in berry chemistry measurements such as pH, SSC, and TA.

Even with consistent water management, berries from each variety responded differently to

environmental conditions. In comparison, for growth measurements such as yield and pruning

weight there was less variation explained by year by variety interaction, indicating the years with

low or high growth for 'Chardonnay' had a similar impact on 'Cabernet Sauvignon'. Thus, the

effect of year on growth was relatively similar across different grapevine varieties, while the

effect on berry chemistry differed between varieties.

In contrast to variety, the effect of rootstock rarely varied across years: the interaction between

rootstock and year was not a significant term for yield, pruning weight, pH or TA, and explained

less than 8% of the variation for the remaining phenotypes. In addition, we found that the effect

of a rootstock was generally consistent between 'Chardonnay' and 'Cabernet Sauvignon', with

the interaction between variety and rootstock explaining very little phenotypic variation (less

than 4%). This suggests that grape growers should place great emphasis on rootstock choice as a

critical decision during vineyard planning as performance of one rootstock, relative to others, is

generally consistent over time and between varieties.

The choice of rootstock is particularly important for growth-related traits such as yield, pruning

weight, berry size, and Ravaz index, where rootstock effects explained at least 9% of the

variation. In contrast to our study, previous work examining nine grape varieties grown

ungrafted and grafted to four different rootstocks found that yield and berry weight were not

affected by rootstock[27]. However, similar to our work, the study identified that vine and yield

components were more responsive to rootstock than fruit composition variables[27]. Our results are

also consistent with previous work identifying a significant difference in yield, pruning weight,

and berry weight of 'Shiraz' vines grafted to different rootstocks and measured across six

years[16].

In long-lived perennial plants where significant year-to-year variation can occur, the collection

of data across multiple years is a valuable tool for untangling the effect of the environment. By

evaluating the vines in this study across five years, we were able to account for the variation due

to year in our model and determine how much of the variation was due to rootstock. Similarly, a

recent seven year study examined 'Cabernet Sauvignon' grafted to three different rootstocks.

The study found no significant effect of rootstock on pruning weight, although yield and berry

weight did differ significantly[28]. When comparing the rootstocks which overlapped with our

study, the authors found similar results: '101-14 Mgt' and '420 A' did not differ significantly for

yield and berry weight, but '420 A' had a significantly higher Ravaz index[28]. Another study that

measured 'Cabernet Sauvignon' grafted to three different rootstocks across 25 years also found

that Ravaz index was significantly affected by rootstock choice, but only after 7 years of

planting. Similarly, yields across rootstocks only differed after 15 years[29]. Although we detect

variation in vines which had been planted for three to seven years, our dataset includes a much

broader representation of rootstocks. Given that grapevines may remain in the ground for at least

20 years, additional long term studies are needed in order to determine how the effect of

rootstocks differs over time.

Generally, rootstocks resulting in large values of one growth-related phenotype also resulted in

large measures of other growth-related phenotypes. For example, rootstocks that generated

higher yields generally also produced larger berries. While cluster number was more highly

correlated with yield than berry weight, much more variation in berry weight could be explained

by rootstock, indicating that increased yields due to rootstock were primarily a result of

increased berry weight and not additional clusters. This suggests that rootstock choice does not

influence floral initiation, but rather influences water uptake, which leads to variation in berry

weight. While high yields are generally desirable, the ratio of skin-to-pulp is an important

consideration for vinification, and this ratio is reduced when berries take on more water.

Previous work also demonstrated that in addition to decreasing fruit size, reducing water in

'Cabernet Sauvignon' increased desirable characteristics such as the concentrations of skin

tannin and anthocyanins[30]. Therefore, while the use of a rootstock to increase yields is beneficial,

this has to be balanced with ensuring that the berries maintain a desirable size, possibly through a

reduction in irrigation for more vigorous rootstocks.

While increased reproductive growth leading to increased yields is economically beneficial, if

the vegetative growth increases at the same rate, the Ravaz index, or crop load, of the vine will

remain consistent. Increased vegetative growth results in higher vine management costs, such as

pruning and leaf thinning. We demonstrated that Ravaz index was correlated with most of the

other phenotypes we measured. This suggests that the balance between reproductive and

vegetative growth in a vine is associated with many other characteristics of that vine. However,

pruning weight and yield were not correlated, likely because all vines were pruned to a similar

size and shoot number to prevent overcropping, but the number of clusters per shoot differed. As

a result, higher yields were positively correlated with both berry weight and cluster number, but

the strengths of these correlations indicate that the primary source of increased yield was more

clusters and not larger berries. In some instances, therefore, rootstock choice may increase

reproductive growth of a vine without an increase in vegetative growth and its associated costs.

While we were unable to evaluate it directly, we find it likely that much of the variation in

growth that can be attributed to rootstock in our study is due to increased water uptake by vines

grafted to certain rootstocks. Variation in water uptake is generally the result of some

combination of water uptake efficiency, the size and surface area of the root system, and

stomatal regulation to reduce water loss, among other factors[12]. For example, 'Ramsey' and

'Freedom' generally had high yields, large berries, and high pruning weights. Similarly, in an

Australia study, 'Shiraz' vines grown with irrigation and grafted to 'Ramsey' or 'Freedom'

rootstocks yielded more fruit than ungrafted vines and than vines grafted on the other five

rootstock varieties assessed, indicating that these rootstocks tend to increase yield and pruning

weight[31]. Other work found a rootstock-dependent effect of irrigation on some yield components

such as cluster number and berry weight, but not on yield itself [32]. In our study, all vines were

irrigated equally, which may have lead to a rootstock-specific effect on water uptake which

ultimately contributed to variation in yield and could be further controlled with rootstock-

specific irrigation regimes.

In addition to variation water uptake, it is possible some variation in growth is due to variation in

disease resistance. While phylloxera is a concern in the region, all vines were grafted to

rootstocks which should provide protection. Additionally, there is the potential for grapevine

fanleaf virus at this site. One of the key symptoms of fanleaf degeneration is a decrease in fruit

set which leads to a lower yield[33]. Given that only vines grafted to '039-16' would have fanleaf

protection in this study, and the yield of vines grafted to '039-16' is not significantly higher than

other rootstocks which do not offer protection, indicating that it is likely not a severe concern in

this vineyard. Thus, while there may be some variation in rootstock tolerance to other pests and

pathogens, this is unlikely to be a major factor in this study.

Lastly, we determined that the choice of one roostock over another has little effect on berry

chemistry, but can result in nearly a two-fold difference in growth-related traits like yield, Ravaz

index and pruning weight. Increasing yield, especially during the early years of production, can

have a dramatic influence on the profitability of a vineyard and the results of this study clearly

indicate that selection of the right rootstock is a valuable tool that grape growers can use to help

control vine size and yield. These results should be taken into account when considering which

rootstock to select, particularly in the San Joaquin Valley where this study was performed.

Future work can explore if the early advantage provided by rootstock is maintained throughout

the life of a vineyard.

Materials and Methods

Experimental design

In 1991, dormant field grown rootstocks were planted in a Tokay fine sandy loam soil[17]. On

April 10th, 1992, scionwood was whip-grafted to the planted rootstock. Rows were oriented east-

west with vine spacing of 2.13 m by 3.05 m (Figure S1). The trellis system was a bilateral

cordon with fixed foliage wires. The cordon wire was at 1.07 m height with single foliage wire

about 40.6 cm above. There were two wires 45.7 cm above the foliage catch wire at either ends

of a 63.5 cm cross arm. The vines were cordon trained and spur pruned.

Prior to vineyard establishment, wine grapes were grown at the site for over 75 years. Initial

plantings on this site were ungrafted *V. vinifera* vines. Because of this production history,

various pests were considered to be endemic. These included several species of nematodes,

phylloxera, many grape associated viruses, and oak root fungus (*Armillaria mellea*)[18]. All of

these soil pests and pathogens can cause considerable economic losses to growers. For this

reason, ungrafted vines were not included as a control in this study.

Vines were grafted to the following rootstocks: 'Freedom', 'Ramsey', '1103 Paulsen', '775

Paulsen', '110 Richter', '3309 Couderc', 'Kober 5BB', 'SO4', 'Teleki 5C', '101-14 Mgt', '039-

16', '140 Ruggeri', 'Schwarzman', '420 A', and 'K51-32'. The two scion varieties were

'Chardonnay' (selection FPS 04) and 'Cabernet Sauvignon' (selection FPS 07).

The experimental design was a randomized complete block design, split between 'Chardonnay'

and 'Cabernet Sauvignon'. There were four replications per treatment (rootstock). Each plot

consisted of eight or nine vines per plot, except for 'Kober 5BB' and 'SO4', which had four or

five vines each, to fit all treatments in the block. Data were collected for five years from 1995-

1999.

Vine management

Canopy management practices were consistent with regional guidelines and included shoot

thinning and leaf removal. Shoot thinning was performed pre-bloom and consisted of removal of

non-count shoots (shoots not originating from spur positions). Leaf removal was performed at

berry set on the north side of the vine only, to avoid excessive exposure and possible sunburn on

the south side. Four to six leaves were removed to open a window in the fruiting zone.

Irrigation and vine nutrition programs were standard Best Management Practice for the Lodi

District. The irrigation strategy followed a moderate Regulated Deficit Irrigation (RDI) program

of about 80% estimated crop evapotranspiration (ETc) losses, from berry set to veraison[19].

During the post-harvest period, vineyard irrigation was increased to 100% ETc. The vine

nutrition program consisted of the application of approximately 30 lbs of actual nitrogen (N) and

60 lbs of actual potassium (K) per acre at post bloom annually. Zinc (Zn) was applied in some

years, as local soils tend to be low in native levels of Zn[20]. All irrigation and nutrients were

applied through a drip system, composed of two 0.5 gph emitters per vine.

Data collection

Prior to harvest, a 100 berry sample was collected for each plot. The berries were counted and

weighed to determine average berry weight. Berries were crushed by hand in plastic collection

bags, then strained through cheesecloth to provide juice for analysis of soluble solids content

(SSC) (°Brix), pH, and titratable acidity (TA) (g/L). Juice samples were titrated to an endpoint of

pH 8.2 to determine TA[21]. SSC was determined by a temperature compensating Atago N1

refractometer (20 °C) and pH was measured using Beckman 200 pH meter with a dual KCl

electrode. Grapes were harvested once they reached an acceptable commercial level for SSC,

approximately 24.5 °Brix for 'Cabernet Sauvignon' and 23 °Brix for 'Chardonnay'. Within a

particular variety ('Cabernet Sauvignon' or 'Chardonnay') all vines were harvested on the same

day (Table S1).

The number of clusters per vine and total fruit yield were recorded. In late winter, vines were

pruned to retain two node fruiting spurs with a target of 50 nodes retained per vine (Table S1).

Dormant pruning weights were measured.

Weather data from 1994 to 1999 were downloaded from the National Environmental Satellite,

Data, and Information Service for Lodi, California, US (USC00045032) on September 30, 2019.

Minimum temperature, maximum temperature, and cumulative precipitation for 1994 to 1999

were plotted (Figure S2).

Statistical analysis

We calculated Ravaz index, a measurement of crop load, by dividing yield by pruning weight

from the following dormant season. As a result, our dataset consisted of eight traits, measured

across five years, for two scion varieties grafted to 15 different rootstocks (Table S2).

The following linear model was evaluated for each phenotype:

372   Phenotype ~ Year + Variety + Rootstock + Column + Year x Rootstock + Variety x

373   Rootstock + Year x Variety + Year x Variety x Rootstock

The model was optimized for each phenotype, which included the removal of the three-way

interaction in all cases as well as non-significant two-way interactions. All main effects were

retained. The percent variation was calculated for all terms by calculating the Sum of Squares for

a particular term, divided by the Total Sum of Squares, then multiplied by 100. The results for

significant terms ($p < 0.05$), except column (position in the vineyard), were plotted. We included

column in our model to account for variation due to position in the vineyard, but we do not

discuss those results here. They are included in our supplemental files and explain up to 10.72%

of the variation in a trait (Table S3).

We visualized phenotype data for 'Chardonnay' and 'Cabernet Sauvignon' separately using a

loess smoothing line to plot variation across years. For the four traits where rootstock explained

the largest amount of variation (i.e. yield, berry weight, pruning weight, and Ravaz index),

rootstocks were compared using a Tukey Test on the model results. For each phenotype, the raw

data and a corresponding boxplot were plotted for each rootstock. To visualize the variety-

specific rootstock effects, we plotted the median values (+/- standard deviation) for 'Cabernet

Sauvignon' and 'Chardonnay' separately for each phenotype (Figure S3).

Since there are large differences between the two grape varieties used in this study, we calculated

the correlation between phenotypes for 'Chardonnay' and 'Cabernet Sauvignon' separately,

using a Spearman's correlation in R v.3.60[22]. To correct for multiple testing, p-values within a

particular variety were Bonferroni-corrected. Heatmaps were generated using the heatmap.2

function in the gplots R package[23]. All remaining figures were plotted using ggplot2 in R[24].

Lastly, we determined the potential range of variation induced by rootstock choice by calculating

the percent change possible from the lowest rootstock median value to the highest rootstock

median value within a particular phenotype. These results were visualized with phenotypes

ordered from highest to lowest possible percent change.

Supplemental information

Figure S1. Vineyard map of rootstock evaluation trial

Figure S2. Variation in maximum temperature (°F), minimum temperature (°F) and

cumulative precipitation (inches) measured from January 1994 to December 1999.

Figure S3. Median values (+/- standard deviation) for each phenotype for 'Chardonnay'

and 'Cabernet Sauvignon' grafted to each rootstock.

Table S1. Harvest and pruning dates for 'Chardonnay' and 'Cabernet Sauvignon' vines

sampled from 1995-1999.

Table S2. Phenotype data collected from 1995 to 1999 for 'Chardonnay' and 'Cabernet

Sauvignon' vines grafted to 15 different rootstocks.

Table S3. Linear model results for each phenotype. Each model was optimized for each

phenotype: the main effects were retained in all cases but non-significant interactions were

removed.

Table S4. Results of Spearman's correlation between phenotypes for 'Chardonnay' and

'Cabernet Sauvignon'. P-values were Bonferroni-corrected for comparison within each variety.

Table S5. Variation across phenotypes based on median rootstock values. The maximum median, minimum median, average median are included as well as the maximum percent change (from minimum to maximum median) and average percent change across rootstocks.

Data Availability

All data supporting the results of this manuscript have been included as supplementary materials (Table S2).

Acknowledgments

ZM was supported by the National Science Foundation Plant Genome Research Program 1546869. SM was supported by the National Sciences and Engineering Council of Canada. We thank Matthew Rubin (Donald Danforth Plant Science Center) for statistical expertise. We acknowledge all of the individuals involved in maintaining the vineyard evaluated for this study, in particular Ernie and Jeff Dosio (Pacific AgriLands Inc., Modesto, CA).

Conflicts of interest

PC, LMJ, and RKS were employed by E. & J. Gallo Winery. The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Authors' Contributions

PV conceived this work and led the collection of the data. LMJ organized the data. ZM analyzed

the data and wrote the manuscript. PC, SM, and DHC provided feedback in data analysis and

interpreting the results. PC, RKS, and DHC provided project oversight and guidance. All authors

reviewed the manuscript.

References

1.　Webster, A. D. Rootstock and interstock effects on deciduous fruit tree vigour, precocity, and yield

productivity. *N. Z. J. Crop Hortic. Sci.* **23**, 373–382 (1995).

2.　Fazio, G. *et al.* Dw2, a new dwarfing locus in apple rootstocks and its relationship to induction of

early bearing in apple scions. *J. Am. Soc. Hortic. Sci.* **139**, 87–98 (2014).

3.　Warschefsky, E. J. *et al.* Rootstocks: Diversity, Domestication, and Impacts on Shoot Phenotypes.

Trends Plant Sci **21**, 418–37 (2016).

4.　Ollat, N., Bordenave, L., Tandonnet, J. P., Boursiquot, J. M. & Marguerit, E. Grapevine rootstocks:

origins and perspectives. *Acta Hortic.* 11–22 (2016) doi:10.17660/ActaHortic.2016.1136.2.

5.　Lider, L. A., Andrew Walker, M. & Wolpert, J. A. Grape rootstocks in California vineyards: the

changing picture. in *International Symposium on Viticulture and Enology 388* 13–18 (1995).

6.　Riaz, S. *et al.* Genetic diversity and parentage analysis of grape rootstocks. *Theor. Appl. Genet.*

(2019) doi:10.1007/s00122-019-03320-5.

7.　Lider, L. A. The performance of vigorous nematode resistant rootstocks in field trials in the San

Joaquin Valley. *Am. J. Enol. Vitic.* **10**, 147–151 (1959).

8.　Cousins, P. & Walker, M. Genetics of resistance to Meloidogyne incognita in crosses of grape

rootstocks. *Theor. Appl. Genet.* **105**, 802–807 (2002).

9. Hwang, C.-F. *et al.* Cloning and characterization of XiR1, a locus responsible for dagger nematode

resistance in grape. *Theor. Appl. Genet.* **121**, 789–799 (2010).

10. Ferris, H., Zheng, L. & Walker, M. Resistance of grape rootstocks to plant-parasitic nematodes. *J.*

Nematol. **44**, 377 (2012).

11. Zhang, X., Walker, R. R., Stevens, R. M. & Prior, L. D. Yield-salinity relationships of different

grapevine (Vitis vinifera L.) scion-rootstock combinations. *Aust. J. Grape Wine Res.* **8**, 150–156

(2002).

12. Serra, I., Strever, A., Myburgh, P. A. & Deloire, A. Review: the interaction between rootstocks and

cultivars (Vitis vinifera L.) to enhance drought tolerance in grapevine. *Aust. J. Grape Wine Res.* **20**,

1–14 (2014).

13. Migicovsky, Z. *et al.* Rootstock effects on scion phenotypes in a 'Chambourcin' experimental

vineyard. *Hortic. Res.* **6**, 64 (2019).

14. Cheng, J., Wei, L., Mei, J. & Wu, J. Effect of rootstock on phenolic compounds and antioxidant

properties in berries of grape (Vitis vinifera L.) cv. 'Red Alexandria'. *Sci. Hortic.* **217**, 137–144

(2017).

15. Walker, R. R., Read, P. E. & Blackmore, D. H. Rootstock and salinity effects on rates of berry

maturation, ion accumulation and colour development in Shiraz grapes. *Aust. J. Grape Wine Res.* **6**,

227–239 (2000).

16. Jones, T. H., Cullis, B. R., Clingeleffer, P. R. & Rühl, E. Effects of novel hybrid and traditional

rootstocks on vigour and yield components of Shiraz grapevines. *Aust. J. Grape Wine Res.* **15**, 284–

292 (2009).

17. UC-Davis. LAWR- California Soil Resource Lab. https://casoilresource.lawr.ucdavis.edu/ (2019).

18. Bettiga, L. J. *Grape pest management*. vol. 3343 (UCANR Publications, 2013).

19. Prichard, T., Hanson, B., Schwankl, L., Verdegaal, P. & Smith, R. Deficit irrigation of quality

winegrapes using micro-irrigation techniques. *Publ. Univ. Calif. Co-Oper. Ext. Dep. Land Air Water*

Resour. Univ. Calif. Davis 89 (2004).

20. Christensen, L. P., Kasimatis, A. N. & Jensen, F. L. *Grapevine nutrition and fertilization in the San*

Joaquin Valley. (Division of Agriculture and Natural Resources. University of California, 1978).

21. Amerine, M. A. & Ough, C. S. *Methods for analysis of musts and wines*. (1980).

22. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical

Computing. https://www.R-project.org/ (2019).

23. Warnes, G. R. *et al.* gplots: Various R Programming Tools for Plotting Data. https://CRAN.R-

project.org/package=gplots (2016).

24. Wickham, H. *ggplot2: elegant graphics for data analysis*. (Springer, 2016).

25. Mendez☐Costabel, M. P. *et al.* Effect of winter rainfall on yield components and fruit green aromas

of Vitis vinifera L. cv. Merlot in California. *Aust. J. Grape Wine Res.* **20**, 100–110 (2014).

26. Medrano, H., Escalona, J. M., Cifre, J., Bota, J. & Flexas, J. A ten-year study on the physiology of

two Spanish grapevine cultivars under field conditions: effects of water availability from leaf

photosynthesis to grape yield and quality. *Funct. Plant Biol.* **30**, 607–619 (2003).

27. Reynolds, A. G. & Wardle, D. A. Rootstocks Impact Vine Performance and Fruit Composition of

Grapes in British Columbia. *HortTechnology* **11**, 419–427 (2001).

28. Hickey, C. C., Hatch, T. A., Stallings, J. & Wolf, T. K. Under-Trellis Cover Crop and Rootstock

Affect Growth, Yield Components, and Fruit Composition of Cabernet Sauvignon. *Am. J. Enol. Vitic.*

67, 281–295 (2016).

29. Ollat, N., Tandonnet, J. P., Lafontaine, M. & Schultz, H. R. Short and long term effects of three

rootstocks on Cabernet Sauvignon vine behaviour and wine quality. in *Workshop on Rootstocks*

Performance in Phylloxera Infested Vineyards 617 95–99 (2003).

30. Roby, G., Harbertson, J. F., Adams, D. A. & Matthews, M. A. Berry size and vine water deficits as

factors in winegrape composition: anthocyanins and tannins. *Aust. J. Grape Wine Res.* **10**, 100–107

(2004).

31. Mccarthy, M. G., Cirami, R. M. & Furkaliev, D. G. Rootstock response of Shiraz (Vitis vinifera)

grapevines to dry and drip-irrigated conditions. *Aust. J. Grape Wine Res.* **3**, 95–98 (1997).

32. Stevens, R. M., Pech, J. M., Gibberd, M. R., Walker, R. R. & Nicholas, P. R. Reduced irrigation and

rootstock effects on vegetative growth, yield and its components, and leaf physiological responses of

Shiraz. *Aust. J. Grape Wine Res.* **16**, 413–425 (2010).

33. Walker, M. A., Wolpert, J. A. & Weber, E. Viticultural characteristics of VR hybrid rootstocks in a

vineyard site infected with grapevine fanleaf virus. *Vitis* **33**, 19–23 (1994).