Non-clonal emergence of colistin resistance associated with mutations in the BasRS

two-component system in *Escherichia coli* bloodstream isolates

Axel B. Janssen[a], Toby L. Bartholomew[b], Natalia P. Marciszewska[a], Marc J.M. Bonten[a], Rob

J.L. Willems[a], Jose A. Bengoechea[b], Willem van Schaik[a,c#]

[a] Department of Medical Microbiology, University Medical Center Utrecht, Utrecht University,

Utrecht, The Netherlands.

[b] Wellcome-Wolfson Institute for Experimental Medicine, Queen's University Belfast, Belfast,

United Kingdom.

[c] Institute of Microbiology and Infection, College of Medical and Dental Sciences, University of

Birmingham, Birmingham, United Kingdom.

Address correspondence to Willem van Schaik, w.vanschaik@bham.ac.uk.

Running title: Colistin resistance in *Escherichia coli* bloodstream isolates

Abstract

Infections by multidrug-resistant Gram-negative bacteria are increasingly common,

prompting the renewed interest in the use of colistin. Colistin specifically targets Gram-negative

bacteria by interacting with the anionic lipid A moieties of lipopolysaccharides, leading to

membrane destabilization and cell death. Here, we aimed to uncover the mechanisms of colistin

resistance in nine colistin-resistant *Escherichia coli* strains and one *E. albertii* strain. These were

the only colistin-resistant strains out of 1140 bloodstream *Escherichia* isolates collected in a

tertiary hospital over a ten-year period (2006 - 2015). Core genome phylogenetic analysis

showed that each patient was colonised by a unique strain, suggesting that colistin resistance was

acquired independently in each strain. All colistin-resistant strains had lipid A that was modified

with phosphoethanolamine. In addition, two *E. coli* strains had hepta-acylated lipid A species,

containing an additional palmitate compared to the canonical hexa-acylated *E. coli* lipid A. One

E. coli strain carried the mobile colistin resistance (mcr) gene *mcr-1.1* on an IncX4-type plasmid.

Through construction of chromosomal transgene integration mutants, we experimentally

determined that mutations in *basRS*, encoding a two-component signal transduction system,

contributed to colistin resistance in four strains. We confirmed these observations by reversing

the mutations in *basRS* to the sequences found in reference strains, resulting in loss of colistin

resistance. While the *mcr*-genes have become a widely studied mechanism of colistin resistance

in *E. coli*, sequence variation in *basRS* is another, potentially more prevalent but relatively

underexplored, cause of colistin resistance in this important nosocomial pathogen.

Importance

Multidrug resistance among Gram-negative bacteria has led to the use of colistin as a last-resort drug. The cationic colistin kills Gram-negative bacteria through electrostatic interaction with the anionic lipid A moiety of lipopolysaccharides. Due to increased use in clinical and agricultural settings, colistin resistance has recently started to emerge. In this study, we used a combination of whole genome sequence analysis and experimental validation to characterise the mechanisms through which *E. coli* strains from bloodstream infections can develop colistin resistance. We found no evidence of direct transfer of colistin-resistant isolates between patients. The lipid A of all isolates was modified by the addition of phosphoethanolamine. In four isolates, colistin resistance was experimentally verified to be caused by mutations in the *basRS* genes, encoding a two-component regulatory system. Our data show that chromosomal mutations are an important cause of colistin resistance among clinical *E. coli* isolates.

Introduction

Escherichia coli is a Gram-negative opportunistic pathogen that is a common cause of

bloodstream, urinary tract, and enteric infections (1). The rising prevalence of antibiotic

resistance in *E. coli*, in part due to the increasing global spread of the successful

multidrug-resistant clade C lineage of ST131, may limit options for future treatments of

infections (2, 3). Due to the emergence and spread of multidrug-resistant clones of *E. coli* and

other Enterobacteriaceae, and the lack of new antibiotics targeting Gram-negative bacteria,

colistin (polymyxin E) is increasingly used, despite its neuro- and nephrotoxic side effects, in the

treatment of clinical infections with multidrug-resistant and carbapenem-resistant *E. coli* and

other Enterobacteriaceae (4–6).

Colistin is a cationic, amphipathic molecule consisting of a non-ribosomal synthesized

decapeptide and a lipid tail (7, 8). Colistin specifically targets Gram-negative bacteria by binding

to the anionic phosphate groups of the lipid A moiety of lipopolysaccharides (LPS) through

electrostatic interactions (7–9). Colistin destabilizes the outer membrane, but the subsequent

disruption of the inner membrane ultimately leads to cell death (9, 10). Acquired colistin

resistance has been reported in various Gram-negative bacteria that were isolated from clinical,

veterinary, and environmental sources (11–13). The best-documented mechanism of colistin

resistance involves the modification of lipid A with cationic groups to counteract the electrostatic

interactions between colistin and lipid A (9). Lipid A modifications in Enterobacteriaceae may

be mediated by the acquisition of mutations in chromosomally located genes or the acquisition of

a mobile genetic element carrying one of the mobile colistin resistance (*mcr*)-genes, which

encode phosphoethanolamine transferases that catalyse the addition of a cationic

phosphoethanolamine group to lipid A (14–16).

Among Enterobacteriaceae, colistin resistance has been most intensively studied in

Salmonella and *Klebsiella pneumoniae* in which mutations in the regulatory genes *mgrB*, *phoPQ*

and *pmrAB* are important mechanisms leading to resistance (15, 17–19). In *E. coli* however,

mutations in *mgrB* and *phoPQ* have not been reported to lead to colistin resistance. This may be

caused by the increased rate of dephosphorylation of PmrA (BasR in *E. coli*) by PmrB (BasS in

E. coli) in *E. coli* compared to other Enterobacteriaceae, which effectively negates the possible

activating effects of mutations in *phoPQ* or *mgrB,* through PmrD, on the levels of

phosphorylated BasR. This may explain why not all of the previously described mutations

reported to lead to colistin resistance in *Salmonella* and *Klebsiella* confer resistance in *E. coli*

(14, 20–22). In addition, *phoPQ* expression in *E. coli* is not only controlled by MgrB but also by

the sRNA MicA, adding to the mechanisms controlling PhoPQ activation and making it less

likely that the deletion or inactivation of *mgrB* can contribute to colistin resistance in *E. coli* (14,

23). This may explain why colistin resistance in clinical *E. coli* strains has only been linked to

mutations in *basRS* (24–28), although experimental validation of the role of these mutations in

colistin resistance is currently mostly lacking.

The PmrAB (BasRS) two-component system plays a crucial role in mediating the

modification of LPS that lead to colistin resistance in Gram-negative bacteria (14, 17). Normally,

this two-component system is activated by environmental stimuli, such as the presence of

antimicrobial peptides or a low pH. Activation can increase virulence and survival through

evasion of the host immune system by upregulating genes associated with modification of LPS,

which is the predominant immunogenic molecule of Gram-negative bacteria (29, 30). In *E. coli*,

the activation of BasRS leads to increased expression of various operons, including its own. This

operon also includes *eptA*, which encodes a lipid A-specific phosphoethanolamine transferase

(11, 14, 31).

Relatively little is known about colistin resistance mechanisms in *E. coli*, other than the

acquisition of *mcr*-genes (32). Therefore, we studied a collection of colistin-resistant *E. coli*

strains from bloodstream infections by a combination of whole genome sequencing and

matrix-assisted laser desorption-ionisation time-of-flight (MALDI-TOF) analysis of their lipid A,

to identify colistin resistance mechanisms in *E. coli*. The role of mutations in *basRS* was

investigated through the construction of chromosomal integration mutants of different *basRS*

alleles.

Materials and methods

Ethical statement

Approval to obtain data from patient records was granted by the Medical Ethics Review Committee of the University Medical Center Utrecht, in Utrecht, The Netherlands (project numbers 16/641 and 18/472).

Colistin-resistant *E. coli* strains were isolated as part of routine diagnostic procedures. This aspect of the study did not require consent or ethical approval by an institutional review board.

Bacterial strains, growth conditions, and chemicals

Colistin-resistant *E. coli* strains from bloodstream infections were obtained retrospectively from the strain collection of the clinical microbiology laboratory of the University Medical Center Utrecht in Utrecht, The Netherlands. In initial routine diagnostic procedures, blood cultures were plated on TSA plates with 5% sheep blood. Strains collected up to 2011 were identified and their antibiogram was determined using the BD Phoenix automated identification and susceptibility testing system (Becton Dickinson, Vianen, The Netherlands). From 2011 onwards, species determination was performed by MALDI-TOF on a Bruker microflex system (Leiderdorp, The Netherlands). *E. coli* strain BW25113 and the BW25113-derived Δ*basRS* strain BW27848 from the Keio collection were obtained from the Coli Genetic Stock Center (33, 34). Strains were grown in Lysogeny Broth (LB; Oxoid, Landsmeer, The Netherlands) at 37°C with agitation at 300 rpm unless otherwise noted, with exception of strains containing pGRG36, which were grown at 30°C (35). When appropriate,

kanamycin (50 mg/L; Sigma-Aldrich, Zwijndrecht, The Netherlands), and ampicillin (100 mg/L;

Sigma-Aldrich) were used. Colistin sulphate was obtained from Duchefa Biochemie (Haarlem,

The Netherlands). L-(+)-arabinose was obtained from Sigma-Aldrich. Plasmids were purified

using the GeneJET Plasmid Miniprep kit (Thermo Fisher Scientific, Landsmeer, The

Netherlands). PCR products were purified from gel using GeneJET Gel Extraction and DNA

Cleanup Micro Kit (Thermo Fisher Scientific).

Determination of minimal inhibitory concentration

Minimal inhibitory concentrations (MICs) to colistin were determined as previously

described (36) in line with the recommendations of a joint working group of the Clinical &

Laboratory Standards Institute and the European Committee on Antimicrobial Susceptibility

Testing (EUCAST)

(http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/General_documents/Recomme

ndations_for_MIC_determination_of_colistin_March_2016.pdf), using BBL™ Mueller Hinton

II (cation-adjusted) broth (MHCAB; Becton Dickinson), untreated Nunc 96-wells round bottom

polystyrene plates (Thermo Scientific), and Breathe-Easy sealing membranes (Sigma-Aldrich).

The breakpoint value of an MIC > 2 µg/ml for colistin resistance in *E. coli* was obtained from

EUCAST (http://www.eucast.org/clinical_breakpoints/).

Genomic DNA isolation and whole-genome sequencing

Genomic DNA was isolated using the Wizard Genomic DNA purification kit (Promega,

Leiden, The Netherlands) according to the manufacturer's instructions. DNA concentrations of

the genomic DNA preparations were measured with the Qubit dsDNA Broad Range Assay kit

and the Qubit 2.0 fluorometer (Life Technologies, Bleiswijk, The Netherlands) and were all

higher than 20 ng/µl.

Sequence libraries for Illumina sequencing were prepared using the Nextera XT kit

(Illumina, San Diego, CA) according to the manufacturer's instructions with 1 ng genomic DNA

as input. Libraries were sequenced on an Illumina MiSeq system with a 500-cycle (2×250 bp)

MiSeq reagent kit v2.

For strain E3090, we performed long-read sequencing using the MinION platform

(Oxford Nanopore Technologies) to fully resolve the *mcr-1.1* plasmid. MinION library

preparation for barcoded 2D long-read sequencing was performed using the SQK-LSK208 kit

(Oxford Nanopore Technologies, Oxford, England, United Kingdom), according to the

manufacturer's instructions, with G-tube (Covaris, Woburn, Massachusetts, United States of

America) shearing of 1 µg chromosomal DNA for 2 x 120 seconds at 1500 *g*. Sequencing was

performed on the MinION sequencer (Oxford Nanopore Technologies) using 2D barcoded

sequencing through a SpotON Flow Cell Mk I (R9.4; Oxford Nanopore Technologies).

Genome assembly, MLST typing, and identification of antibiotic resistance genes

The quality of Illumina sequence data was assessed using FastQC v0.11.5

(https://github.com/s-andrews/FastQC). Raw Illumina sequencing reads were trimmed for quality

using nesoni v0.115 (https://github.com/Victorian-Bioinformatics-Consortium/nesoni) using

standard settings with the exception of a minimum read length of 100 nucleotides. *De novo*

genome assembly of the trimmed Illumina short-read data was performed using SPAdes v3.6.2

with the following settings: kmers used: 21, 33, 55, 77, 99, or 127, "careful" option turned on

and cut-offs for final assemblies: minimum contig/scaffold size = 500 bp, minimum

contig/scaffold average Nt coverage = 10-fold (37).

MinION sequence read data in FastQ format was extracted from Metrichor base-called

raw FAST5 read-files using Poretools (38). A hybrid assembly for strain E3090 was generated

with trimmed Illumina short-read data and Oxford Nanopore Technologies MinION long-read

data by using SPAdes v3.6.2 with the same settings as the Illumina short-read assemblies, and

specifying the long-read data with the --nanopore flag.

Gene prediction and annotation was performed using Prokka (39), using standard

settings. MLST typing was performed using the mlst package v2.10

(https://github.com/tseemann/mlst), using standard settings. Assembled contigs were assessed for

antibiotic resistance genes using ResFinder 3.2 (40), using standard settings.

Core genome phylogenetic analysis and determination of mutations in candidate colistin

resistance determinants

Genome assemblies generated in this study with Illumina data were aligned with 178

complete *E. coli* genomes and 32 *E. albertii* genomes that were available from NCBI databases

on 24 June 2016 (Supplemental Table S1) using ParSNP v1.2 (41). MEGA6 was used to

midpoint root and visualize the phylogenetic tree (42). We identified whether non-synonymous

mutations were present in *basRS* by pairwise comparison of the gene sequences of

colistin-resistant isolates to their closest matching publicly available genome from the

phylogenetic tree using BLAST (43). Mutations that were identified in the genome sequences

were confirmed through PCR (oligonucleotide primer sequences are provided in Supplemental

Table S2) and subsequent Sanger sequencing of the PCR product by Macrogen (Amsterdam, The

Netherlands).

Isolation and analysis of lipid A

Isolation of lipid A molecules and subsequent analysis by negative-ion MALDI-TOF

mass spectrometry was performed as previously described (19, 44, 45). Briefly, *Escherichia*

strains were grown in LB (Oxoid) and the lipid A was purified from stationary cultures using the

ammonium hydroxide/isobutyric acid method described earlier (46). Mass spectrometry analyses

were performed on a Bruker autoflex™ speed TOF/TOF mass spectrometer in negative

reflective mode with delayed extraction using as matrix an equal volume of dihydroxybenzoic

acid matrix (Sigma-Aldrich) dissolved in (1:2) acetonitrile-0.1% trifluoroacetic acid. The

ion-accelerating voltage was set at 20 kV. Each spectrum was an average of 300 shots. A peptide

calibration standard (Bruker) was used to calibrate the MALDI-TOF. Further calibration for

lipid A analysis was performed externally using lipid A extracted from *E. coli* strain MG1655

grown in LB medium at 37°C.

Construction of chromosomal *basRS* transgene insertions

Chromosomal transgene insertions of *basRS* were constructed in BW27848 by utilizing

the Tn7 transposon system on the pGRG36 plasmid (35). The strategy to generate the constructs

is illustrated in Supplemental Figure S1. The promoter of the *eptA-basRS* operon was fused to

the *basRS* coding sequence by separate PCRs for the promotor region and the *basRS* amplicon,

with high fidelity Phusion Green Hot Start II DNA Polymerase (Thermo Fisher Scientific) using

strain-specific primers (Supplemental Table S2; oligonucleotides were obtained from Integrated

DNA Technologies, Leuven, Belgium). The promoter and the *basRS* amplicon were

subsequently fused by overlap PCR. Fused PCR products were cloned into pCR-Blunt II-TOPO

using the Zero Blunt TOPO PCR Cloning kit (Thermo Fisher Scientific), and subsequently

subcloned into pGRG36 (35). Electrocompetent BW25113 and BW27848 *E. coli* cells were

prepared as described previously (47) and transformed using the following settings: voltage

1800V, capacitance 25 µF, resistance 200Ω, with a 0.2 cm cuvette using the Gene Pulser Xcell

electroporation system (Bio-Rad Laboratories, Veenendaal, The Netherlands). Transformants

were grown at 30°C. After confirming integration of the Tn7 transposon at the *att*Tn7 site by

PCR (primers listed in Supplemental Table S2) and Sanger sequencing (Macrogen), the pGRG36

plasmid was cleared by culturing at 37°C.

Inverse PCR site-directed mutagenesis was performed on amplicons cloned in pCR-Blunt

II-TOPO to reverse the mutations that were identified in colistin-resistant strains to the

sequences of *basR* or *basS* in the closest matching publicly available genome (48). After gel

purification of the amplified fragments, (hemi)methylated fragments were digested using DpnI

(New England Biolabs (NEB), Ipswich, Massachusetts, United States of America). Subsequently

the vector was recircularized using the Rapid DNA Ligation kit (Thermo Fisher Scientific) after

phosphorylation using T4 Polynucleotide kinase (NEB). The constructs were then transformed

into chemically competent DH5α *E. coli* cells (Invitrogen, Landsmeer, The Netherlands).

Mutated sequences were subsequently subcloned to pGRG36 as described above.

Data availability

Sequence data has been deposited in the European Nucleotide Archive (accession number

PRJEB27030).

Results

Low prevalence of colistin resistance in invasive *Escherichia* bloodstream isolates

A total of 1140 bloodstream isolates (collected from January 2006 to December 2015) for

which species identification had been performed and automated antibiotic susceptibility profiles

had been determined, were available for this study. Twelve isolates were deemed resistant to

colistin through routine diagnostic procedures. Two of those isolates were isolated from the same

patient, on the same day, and were thus considered duplicates, and only one of these was

included in this study. In ten of the eleven remaining isolates, colistin resistance, defined as an

MIC > 2 µg/ml colistin, was confirmed through broth microdilution (Table 1). Strain A783 was a

false positive for colistin resistance during automated susceptibility testing in routine diagnostic

procedures, and was excluded from subsequent analyses, leaving ten isolates for further

investigation.

The estimated prevalence of colistin resistance in *E. coli* strains causing bloodstream

infections isolated from January 2006 to December 2015 was thus determined to be 0.88%.

Three patients had received colistin in the three months before isolation of the colistin-resistant

strain (Table 1). Two of these patients received colistin to treat infections, but all three patients

were also administered colistin as part of selective digestive or oropharyngeal decontamination

(SDD/SOD), a prophylactic antibiotic treatment widely used in Dutch intensive care units (49).

The ten colistin-resistant strains were analyzed further in this study to determine their relatedness

and mechanism through which they had developed colistin resistance.

Colistin resistance was independently acquired by each individual bloodstream *E. coli*

isolate

To assess the phylogenetic relationships between the colistin-resistant strains, a

phylogenetic tree was generated based on the genome assemblies of the colistin-resistant strains

and 210 publicly available complete genome sequences (Supplemental Table S1). Based on a

core genome alignment of 874 kbp, we did not observe direct transmission of colistin-resistant

strains between patients (Figure 1A). Three colistin-resistant strains (strains I1121, H2129, and

G821) belonged to the globally disseminated ST131 clone, and all three were dispersed

throughout the multidrug-resistant clade C of ST131(Figure 1A, 1B) (3, 50). This indicates that

the ST131 strains in this study have independently acquired colistin resistance. Strain A2361

clustered among *E. albertii* (Figure 1A), although it had been typed as *E. coli* in routine

diagnostic procedures.

By screening for acquired antibiotic resistance genes through ResFinder 3.2, we found

that only strain E3090 carried the *mcr* gene *mcr-1.1* (0.086% of all bloodstream isolates; Figure

1C). After long-read sequencing and hybrid assembly, the *mcr-1.1*-gene in this strain appeared to

be located as the sole antibiotic resistance gene on a 32.7 kbp IncX4-type plasmid. This *mcr-1.1*

carrying IncX4-type plasmid from E3090 shares 99% identity to the previously reported *mcr-1.1*

carrying IncX4-type plasmid pMCR-1_Msc (GenBank accession MK172815.1) harboured by

E. coli isolated from patients in Russia (51), confirming the global dissemination of this plasmid

(52). In all strains studied here, a variety of acquired resistance genes was observed (Figure 1C),

reflecting the non-clonal nature of the colistin-resistant strains. The three colistin-resistant ST131

strains possessed different repertoires of acquired resistance genes, further excluding recent

transmission between patients of the ST131 strains studied here. Strain F2745 and E2372 carried

292   only one, and two resistance genes respectively, while the *E. albertii* strain A2361 did not

293   possess any acquired resistance genes.

295   ***Escherichia* isolates exclusively acquire colistin resistance by modification of phosphate**

296   **groups of lipid A**

297   To determine which modifications to lipid A are affecting colistin resistance in *E. coli* we

extracted lipid A from the clinical strains and the colistin-susceptible control *E. coli* strain

MG1655, and subjected them to MALDI-TOF mass spectrometry. The lipid A produced by all

E. coli strains showed lipid A species with a mass-to-charge ratio (*m/z*) of 1797 (Figure 2A),

corresponding to the canonical unmodified *E. coli* hexa-acylated lipid A (Figure 2B).

Colistin-resistant strains showed additional lipid A species at *m/z* 1921, consistent with the

addition of phosphoethanolamine (*m/z* 124) to the hexa-acylated species. Additional species were

detected in the lipid A produced by strains E650 and Z821. Species *m/z* 2036 indicated the

addition of palmitate (*m/z* 239) to the hexa-acylated species *m/z* 1797, whereas species *m/z* 2160

was consistent with the addition of palmitate to the hexa-acylated lipid A species containing

phosphoethanolamine (*m/z* 1910).

308   The *E. alberti* strain A2361 produced lipid A distinct from *E. coli.* Species *m/z* 1825 is

309   likely to represent a hexa-acylated species corresponding to two glucosamines, two phosphates,

310   four 3-OH-C_{14}, and two C_{14} (Figure 2C) Species *m/z* 1948 is consistent with the addition of

311   phosphoethanolamine to the hexa-acylated species, with a further addition of palmitate to

312   produce lipid A species *m/z* 2187. Species *m/z* 1868 and *m/z* 2107 could correspond to the loss of

313   the second phosphate group, compared to *m/z* 1948, and *m/z* 2187.

Identification of mutations in *basRS* as candidate mutations involved in colistin resistance.

Because chromosomal mutations in *basRS*, but not in other regulatory systems, have

previously been suggested to cause colistin resistance in *E. coli* (24–28), we next aimed to

establish the contribution of the *basRS* alleles in the colistin-resistant phenotype of these

bloodstream isolates. Due to the multidrug-resistant nature of the clinical isolates (Figure 1C),

we were unable to generate targeted mutations in these strains. Therefore, we made

chromosomal transgene insertion mutants of the different *basRS* alleles in the *att*Tn7 site in the

BW25113-derived Δ*basRS* strain BW27848 using the Tn7 transposon system. By making

chromosomal transgenes insertions, rather than using an *in trans* complementation method, we

excluded copy number effects by plasmids, and the need to use antibiotics to select for the

presence of a plasmid used for *in trans* complementation. Since BW27848 still possesses the

gene encoding for the phosphoethanolamine transferase EptA, we constructed sequences that

consisted of the fused sequences of the promotor region of the *eptA-basR-basS* operon and the

basRS coding sequences in order to prevent *eptA* gene dosage-dependent effects. We were

unable to generate the construct for strain E650, presumably due to the toxicity of the insert.

The colistin MIC determination of the generated *basRS* chromosomal transgene insertion

mutants from strains I1121, H2129, G821, and Z821 had higher colistin MIC values than the

BW27848::Tn7-empty strain, with observed MIC values ≥16-fold higher than that of the

BW27848::Tn7-empty strain (Table 2). As expected, the *basRS* allele of the *mcr-1.1* positive

strain E3090 did not lead to colistin resistance. We were unable to show the contribution of

basRS to colistin resistance in the additional four colistin-resistant strains (F2745, E2372,

D2373, A2361) that lacked *mcr-1.1*.

Mutations in the *basRS* genes contribute to colistin resistance in *E. coli*.

By construction of the chromosomal transgene insertion mutants, we identified the ability

of the *basRS* sequences of four strains (I1121, H212, G821, and Z821) to cause colistin

resistance in BW27848. To identify the mutations in the *basRS* alleles of these strains that

contribute to resistance, we compared the *basRS* encoding sequences of those strains causing

resistance to the phylogenetically most closely related publicly available *E. coli* genome

sequences used in the construction of Figure 1A. None of these reference strains were reported to

be colistin-resistant, or carried any of the *mcr*-genes. This comparison revealed four distinct

mutations: a L10R substitution in BasS in I1121, a G53S substitution in BasR in H2192, the

duplication of the HAMP-domain in BasS in G821, and a A159P substitution in BasS in Z821

(Supplemental Figure S2). As expected, in the *mcr-1.1* positive strain E3090 no mutations in

basRS were identified.

We hypothesised that the observed mutations were impacting the normal functioning of

the BasRS two-component system. To assess whether the mutations in *basRS* identified by

comparing the *basRS* sequences of the clinical strains I1121, H2129, G821, and Z821, and their

closest match in the set of 178 publicly available *E. coli* genome sequences (Supplemental

Figure S2) were causal to the development of colistin resistance, the identified mutations were

reversed through site-directed inverse PCR mutagenesis to match the *basRS* alleles of the

publicly available genome sequences. The MIC values of these mutants returned to levels similar

to that of the colistin-susceptible BW27848::Tn7-empty strain (Table 2). These experiments

support the involvement of *basRS* sequence variation in colistin resistance in *E. coli*.

Discussion

In the present study, we set out to characterise the the mechanisms through which *E. coli* bloodstream isolates can develop colistin resistance through a combination of whole genome sequence analysis and experimental validation. We did not find evidence for transfer of colistin-resistant strains between patients, suggesting that colistin resistance has been acquired independently in all cases. In seven patients colistin-resistant strains were isolated without the patients being previously exposed to the drug. All colistin-resistant strains had LPS that was modified by the addition of phosphoethanolamine to the lipid A moiety of LPS. Resistance in one of the bloodstream isolates could be explained by the acquisition of *mcr-1.1*. In four other strains, we identified mutations in *basRS* that contribute to colistin resistance. Although colistin-susceptible strains that were isogenic to the resistant strains were not available, we were able to pinpoint the mutations in *basRS* leading to resistance in these strains by matching the genomic sequences of our nosocomial isolates with publicly available genomes, none of which were reported to be colistin-resistant, and subsequent construction of chromosomally integrated *basRS* transgene alleles in the Δ*basRS* strain BW27848. The mechanisms of colistin resistance in the remaining five strains remain to be characterized.

Some of the mutations we experimentally link to colistin resistance in this study, have previously been associated with colistin resistance or the functioning of the BasRS two-component system. In this study, we demonstrated that the amino acid change L10R in BasS (strain I1121) also confers colistin resistance. An amino acid substitution in the same position of BasS (L10P) was previously experimentally proven to cause colistin resistance in *E. coli* (26). The glycine in position 53 of BasR has previously been reported to be altered in colistin-resistant

Enterobacteriaceae (53, 54) including in *E. coli* (55). The G53S change specifically, as in isolate

H2192, has been experimentally proven to contribute to colistin resistance in

Klebsiella (previously *Enterobacter*) *aerogenes* (56, 57) and *Salmonella enterica* subsp. *enterica*

serovar Typhimurium (58) and we extend those findings to *E. coli* here. The previously

unidentified duplication of 162 nucleotides in *basS* (strain G821) leads to the introduction of a

second HAMP domain in BasS and confers colistin resistance in the BW27848 background. The

HAMP domain is widespread in bacteria and is commonly involved in signal transduction as part

of two-component systems (59). We hypothesise that the addition of an extra HAMP domain in

BasS may change signal transduction in the protein, leading to the constitutive activation of the

histidine kinase domain of BasS, increased phosphorylation of BasR and upregulated expression

of *eptA*, ultimately resulting in the addition of phosphoethanolamine to lipid A. Finally, we

demonstrate that the A159P substitution in BasS (observed in strain Z821) contributes to colistin

resistance. A mutation leading to a A159V substitution was found in an *in vitro* evolution study

in which *E. coli* was evolved towards colistin resistance (60), and in clinical colistin-resistant

E. coli isolates (61), but experimental confirmation of the role of alterations in A159 in colistin

resistance in *E. coli* was so far lacking. Our data suggest that the *basRS* alleles of three *E coli*

strains (F2745, E2372, and D2373), and the *E. albertii* strain A2361, do not confer resistance in

the BW25113 *E. coli* background. Because *E. albertii* is phylogenetically distinct from *E. coli*,

its *basRS* allele may not function optimally in an *E. coli* background, explaining the inability of

the transgene insertion complementation in the *basRS* deletion of BW25113 *E. coli* strain to

cause colistin resistance (62). We are unable to explain the colistin resistance mechanisms of the

clinical isolates F2745, E2372, and D2373. It is likely that these strains have become resistant to

colistin through other mutations that finally lead to the modification of lipid A by

phosphoethanolamine.

The observed modification of lipid A with phosphoethanolamine in all isolates underlines

the crucial role of phosphoethanolamine transferases in the ability of *Escherichia* to become

resistant to polymyxins (14). The lipid A of three of the colistin-resistant strains was also

modified with palmitate, but the contribution of lipid A palmitoylation to colistin resistance in

clinical *E. coli* strains is currently unknown. We did not observe modifications of lipid A by

4-amino-4-deoxy-L-arabinose in the colistin-resistant isolates. While this modification was

shown to contribute to polymyxin B resistance under low Mg^{2+} conditions in a laboratory isolate

of *E. coli* (20), it may be rare in clinical *E. coli* isolates. Indeed, Sato *et al*. also exclusively found

phosphoethanolamine-modified lipid A in colistin-resistant clinical *E. coli* isolates (24). The

reliance of *Escherichia* on the modification of lipid A by phosphoethanolamine to acquire

colistin resistance, suggests that the inhibition of this class of enzymes by blocking the conserved

catalytic site (31) could be a target for future drug development and opens the possibility of

combination therapy with colistin and an inhibitor of phosphoethanolamine transferase (63).

With the increasing clinical issues posed by infections with multidrug-resistant Gram-negative

bacteria, there is an urgent need to better understand resistance mechanisms to last-resort

antibiotics like colistin. While the discovery of the *mcr*-genes have generated considerable

interest in transferable colistin resistance genes, our data suggest that chromosomal mutations

remain an important cause of colistin resistance among clinical isolates in the genus *Escherichia*.

Acknowledgements

We thank Eline A.M. Majoor for technical support and L. Marije Hofstra and Lidewij W.

Rümke for their review of patient records. We also thank the Utrecht Sequence Facility and Ivo

Renkens for their expertise in MinION Nanopore sequencing.

W.v.S. was funded through an NWO-Vidi grant (grant 917.13.357), and a Royal Society

Wolfson Research Merit Award. Work in J.A.B. laboratory was supported by Biotechnology and

Biological Sciences Research Council (BBSRC, BB/P020194/1) and Queen's University Belfast

start-up. T.L.B. is the recipient of a PhD fellowship funded by the Department for Employment

and Learning (Northern Ireland, UK). The funders had no role in study design, data collection

and interpretation, or the decision to submit the work for publication.

A.B.J. conceived and designed experiments, performed experiments, analysed data, and

wrote the manuscript. T.B.L. performed experiments, and analysed data. N.P.M. performed

experiments, and analysed data. M.J.M.B. wrote the manuscript. R.J.L.W. wrote the manuscript.

J.A.B. analysed data, and wrote the manuscript. W.v.S. conceived and designed experiments,

wrote the manuscript, and supervised the study. All authors reviewed and approved the final

version of the manuscript.

Competing interests

The authors declare no conflicts of interest.

References

1.   Kaper JB, Nataro JP, Mobley HLT. 2004. Pathogenic *Escherichia coli*. Nat Rev Microbiol
2:123–140.

2.   Kallonen T, Brodrick HJ, Harris SR, Corander J, Brown NM, Martin V, Peacock SJ,
Parkhill J. 2017. Systematic longitudinal survey of invasive *Escherichia coli* in England
demonstrates a stable population structure only transiently disturbed by the emergence of
ST131. Genome Res 27:1437–1449.

3.   McNally A, Kallonen T, Connor C, Abudahab K, Aanensen DM, Horner C, Peacock SJ,
Parkhill J, Croucher NJ, Corander J. 2019. Diversification of colonization factors in a
multidrug-resistant *Escherichia coli* lineage evolving under negative frequency-dependent
selection. MBio 10:e00644-19.

4.   Klein EY, Van Boeckel TP, Martinez EM, Pant S, Gandra S, Levin SA, Goossens H,
Laxminarayan R. 2018. Global increase and geographic convergence in antibiotic
consumption between 2000 and 2015. Proc Natl Acad Sci 115:E3463–E3470.

5.   Lim LM, Ly N, Anderson D, Yang JC, Macander L, Jarkowski A, Forrest A, Bulitta JB,
Tsuji BT. 2010. Resurgence of colistin: a review of resistance, toxicity,
pharmacodynamics, and dosing. Pharmacotherapy 30:1279–1291.

6.   Nation RL, Li J. 2010. Colistin in the 21st Century. Curr Opin Infect Dis 22:535–543.

7.   Domingues MM, Inácio RG, Raimundo JM, Martins M, Castanho MARB, Santos NC.
2012. Biophysical characterization of polymyxin B interaction with LPS aggregates and
membrane model systems. Biopolymers 98:338–344.

8.   Landman D, Georgescu C, Martin DA, Quale J. 2008. Polymyxins revisited. Clin
Microbiol Rev 21:449–465.

9.   Velkov T, Thompson PE, Nation RL, Li J. 2010. Structure-activity relationships of
polymyxin antibiotics. J Med Chem 53:1898–1916.

10.   Sabnis A, Klöckner A, Becce M, Evans LE, Furniss RCD, Mavridou DAI, Stevens MM,
Edwards AM. 2018. Colistin kills bacteria by targeting lipopolysaccharide in the
cytoplasmic membrane. bioRxiv 479618.

11.   Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, Doi Y, Tian G, Dong B, Huang
X, Yu LF, Gu D, Ren H, Chen X, Lv L, He D, Zhou H, Liang Z, Liu JH, Shen J. 2016.
Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and
human beings in China: a microbiological and molecular biological study. Lancet Infect
Dis 16:161–168.

12.   Halaby T, Kucukkose E, Janssen AB, Rogers MRC, Doorduijn DJ, van der Zanden AGM,
al Naiemi N, Vandenbroucke-Grauls CMJE, van Schaik W. 2016. Genomic
characterization of colistin heteroresistance in *Klebsiella pneumoniae* during a nosocomial
outbreak. Antimicrob Agents Chemother 60:6837–6843.

13.   Islam A, Rahman Z, Monira S, Rahman MA, Camilli A, George CM, Ahmed N, Alam M.
484   2017. Colistin resistant *Escherichia coli* carrying *mcr-1* in urban sludge samples: Dhaka,
485   Bangladesh. Gut Pathog 9:77.

486   14.   Baron S, Hadjadj L, Rolain J-M, Olaitan AO. 2016. Molecular mechanisms of polymyxin
487   resistance: knowns and unknowns. Int J Antimicrob Agents 48:583–591.

488   15.   Poirel L, Jayol A, Nordmann P. 2017. Polymyxins: Antibacterial activity, susceptibility
489   testing, and resistance mechanisms encoded by plasmids or chromosomes. Clin Microbiol
490   Rev 30:557–596.

491   16.   Zhang H, Srinivas S, Xu Y, Wei W, Feng Y. 2019. Genetic and biochemical mechanisms
492   for bacterial lipid A modifiers associated with polymyxin resistance. Trends Biochem Sci
493   44:973–988.

494   17.   Falagas ME, Rafailidis PI, Matthaiou DK. 2010. Resistance to polymyxins: mechanisms,
495   frequency and treatment options. Drug Resist Updat 13:132–138.

496   18.   Cannatelli A, D'Andrea MM, Giani T, Di Pilato V, Arena F, Ambretti S, Gaibani P,
497   Rossolini GM. 2013. In vivo emergence of colistin resistance in *Klebsiella pneumoniae*
498   producing KPC-type carbapenemases mediated by insertional inactivation of the
499   PhoQ/PhoP *mgrB* regulator. Antimicrob Agents Chemother 57:5521–5526.

500   19.   Kidd TJ, Mills G, Sá-Pessoa J, Dumigan A, Frank CG, Insua JL, Ingram R, Hobley L,
501   Bengoechea JA. 2017. A *Klebsiella pneumoniae* antibiotic resistance mechanism that
502   subdues host defences and promotes virulence. EMBO Mol Med 9:430–447.

503   20.   Rubin EJ, Herrera CM, Crofts AA, Trent MS. 2015. PmrD is required for modifications to
504   *Escherichia coli* endotoxin that promote antimicrobial resistance. Antimicrob Agents
505   Chemother 59:2051–2061.

506   21.   Winfield MD, Groisman EA. 2004. Phenotypic differences between *Salmonella* and
507   *Escherichia coli* resulting from the disparate regulation of homologous genes. Proc Natl
508   Acad Sci 101:17162–17167.

509   22.   Chen HD, Jewett MW, Groisman EA. 2011. Ancestral genes can control the ability of
510   horizontally acquired loci to confer new traits. PLoS Genet 7:e1002184.

511   23.   Coornaert A, Lu A, Mandin P, Springer M, Gottesman S, Guillier M. 2010. MicA sRNA
512   links the PhoP regulon to cell envelope stress. Mol Microbiol 76:467–479.

513   24.   Sato T, Shiraishi T, Hiyama Y, Honda H, Shinagawa M, Usui M, Kuronuma K, Masumori
514   N, Takahashi S, Tamura Y, Shin-ichi Y. 2018. Contribution of novel amino alterations in
515   PmrA or PmrB to colistin resistance in *mcr*-negative *Escherichia coli* clinical isolates,
516   including major multidrug-resistant lineages O25b:H4-ST131-H30Rx and non-x.
517   Antimicrob Agents Chemother 62:e00864-18.

518   25.   Froelich JM, Tran K, Wall D. 2006. A *pmrA* constitutive mutant sensitizes *Escherichia*
519   *coli* to deoxycholic acid. J Bacteriol 188:1180–1183.

520   26.   Cannatelli A, Giani T, Aiezza N, Pilato V Di, Principe L, Luzzaro F, Galeotti CL,
521   Rossolini GM. 2017. An allelic variant of the PmrB sensor kinase responsible for colistin
522   resistance in an *Escherichia coli* strain of clinical origin. Sci Rep 7:5071.

523   27.   Delannoy S, Le Devendec L, Jouy E, Fach P, Drider D, Kempf I. 2017. Characterization
524   of colistin-resistant *Escherichia coli* isolated from diseased pigs in France. Front
525   Microbiol 8:2278.

526   28.   Bourrel AS, Poirel L, Royer G, Vuillemin X, Kieffer N, Clermont O, Denamur E,
527   Nordmann P, Decousser J. 2019. Colistin resistance in Parisian inpatient faecal
528   *Escherichia coli* as the result of two distinct evolutionary pathways. J Antimicrob
529   Chemother 74:1521–1530.

530   29.   Park BS, Song DH, Kim HM, Choi B-S, Lee H, Lee J-O. 2009. The structural basis of
531   lipopolysaccharide recognition by the TLR4-MD-2 complex. Nature 458:1191–1196.

532   30.   Maeshima N, Fernandez RC. 2013. Recognition of lipid A variants by the TLR4-MD-2
533   receptor complex. Front Cell Infect Microbiol 3:3.

534   31.   Xu Y, Wei W, Lei S, Lin J, Srinivas S, Feng Y. 2018. An evolutionarily conserved
535   mechanism for intrinsic and transferable polymyxin resistance. MBio 9:e02317-17.

536   32.   Luo Q, Yu W, Zhou K, Guo L, Shen P, Lu H, Huang C, Xu H, Xu S, Xiao Y, Li L. 2017.
537   Molecular epidemiology and colistin resistant mechanism of *mcr*-positive and *mcr*-
538   negative clinical isolated *Escherichia coli*. Front Microbiol 8:2262.

539   33.   Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, Datsenko KA, Tomita M,
540   Wanner BL, Mori H. 2006. Construction of *Escherichia coli* K-12 in-frame, single-gene
541   knockout mutants: the Keio collection. Mol Syst Biol 2:2006.0008.

542   34.   Zhou L, Lei X, Bochner BR, Wanner BL. 2003. Phenotype microarray analysis of
543   *Escherichia coli* K-12 mutants with deletions of all two-component systems. J Bacteriol
544   185:4956–4972.

545   35.   McKenzie GJ, Craig NL. 2006. Fast, easy and efficient: site-specific insertion of
546   transgenes into Enterobacterial chromosomes using Tn7 without need for selection of the
547   insertion event. BMC Microbiol 6:39.

548   36.   Andrews JM. 2001. Determination of minimum inhibitory concentrations. J Antimicrob
549   Chemother 48:5–16.

550   37.   Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM,
551   Nikolenko SI, Pham S, Prjibelski AD, Pyshkin A V, Sirotkin A V, Vyahhi N, Tesler G,
552   Alekseyev MA, Pevzner PA. 2012. SPAdes: a new genome assembly algorithm and its
553   applications to single-cell sequencing. J Comput Biol 19:455–477.

554   38.   Loman NJ, Quinlan AR. 2014. Poretools: a toolkit for analyzing nanopore sequence data.
555   Bioinformatics 30:3399–3401.

556   39.   Seemann T. 2014. Prokka: rapid prokaryotic genome annotation. Bioinformatics 30:2068–

557   2069.

558   40.   Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM,
559   Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. J Antimicrob
560   Chemother 67:2640–2644.

561   41.   Treangen TJ, Ondov BD, Koren S, Phillippy AM. 2014. The Harvest suite for rapid core-
562   genome alignment and visualization of thousands of intraspecific microbial genomes.
563   Genome Biol 15:524.

564   42.   Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. 2013. MEGA6: Molecular
565   evolutionary genetics analysis version 6.0. Mol Biol Evol 30:2725–2729.

566   43.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic Local Alignment
567   Search Tool. J Mol Biol 215:403–410.

568   44.   Llobet E, Martínez-Moliner V, Moranta D, Dahlström KM, Regueiro V, Tomás A, Cano
569   V, Pérez-Gutiérrez C, Frank CG, Fernández-Carrasco H, Insua JL, Salminen TA,
570   Garmendia J, Bengoechea JA. 2015. Deciphering tissue-induced *Klebsiella pneumoniae*
571   lipid A structure. Proc Natl Acad Sci U S A 112:E6369–E6378.

572   45.   Llobet E, Campos MA, Giménez P, Moranta D, Bengoechea JA. 2011. Analysis of the
573   networks controlling the antimicrobial-peptide-dependent induction of *Klebsiella*
574   *pneumoniae* virulence factors. Infect Immun 79:3718–3732.

575   46.   El Hamidi A, Tirsoaga A, Novikov A, Hussein A, Caroff M. 2005. Microextraction of
576   bacterial lipid A: easy and rapid method for mass spectrometric characterization. J Lipid
577   Res 46:1773–1778.

578   47.   Sharma RC, Schimke RT. 1996. Preparation of electro-competent *E. coli* using salt-free
579   growth medium. Biotechniques 20:42–44.

580   48.   Silva D, Santos G, Barroca M, Collins T. 2017. Inverse PCR for point mutation
581   introduction, p. 87–100. *In* Methods in Molecular Biology. Humana Press Inc.

582   49.   Oostdijk EAN, Kesecioglu J, Schultz MJ, Visser CE, De Jonge E, Van Essen EHR,
583   Bernards AT, Purmer I, Brimicombe R, Bergmans D, Van Tiel F, Bosch FH, Mascini E,
584   Van Griethuysen A, Bindels A, Jansz A, Van Steveninck FAL, Van Der Zwet WC, Fijen
585   JW, Thijsen S, De Jong R, Oudbier J, Raben A, Van Der Vorm E, Koeman M, Rothbarth
586   P, Rijkeboer A, Gruteke P, Hart-Sweet H, Peerbooms P, Winsser LJ, Van Elsacker-Niele
587   AMW, Demmendaal K, Brandenburg A, De Smet AMGA, Bonten MJM. 2014. Effects of
588   decontamination of the oropharynx and intestinal tract on antibiotic resistance in ICUs a
589   randomized clinical trial. J Am Med Assoc 312:1429–1437.

590   50.   Ben Zakour NL, Alsheikh-Hussain AS, Ashcroft MM, Khanh Nhu NT, Roberts LW,
591   Stanton-Cook M, Schembri MA, Beatson SA. 2016. Sequential acquisition of virulence
592   and fluoroquinolone resistance has shaped the evolution of *Escherichia coli* ST131. MBio
593   7:e00347-16.

51.   Ageevets V, Lazareva I, Mrugova T, Gostev V, Lobzin Y, Sidorenko S. 2019. IncX4

plasmids harbouring *mcr-1* genes: further dissemination. J Glob Antimicrob Resist
18:166–167.

52. Wang R, van Dorp L, Shaw LP, Bradley P, Wang Q, Wang X, Jin L, Zhang Q, Liu Y,
Rieux A, Dorai-Schneiders T, Weinert LA, Iqbal Z, Didelot X, Wang H, Balloux F. 2018.
The global distribution and spread of the mobilized colistin resistance gene *mcr-1*. Nat
Commun 9:1179.

53. Nordmann P, Jayol A, Poirel L. 2016. Rapid detection of polymyxin resistance in
Enterobacteriaceae. Emerg Infect Dis 22:1038–1043.

54. Olaitan AO, Diene SM, Kempf M, Berrazeg M, Bakour S, Gupta SK, Thongmalayvong B,
Akkhavong K, Somphavong S, Paiboun P, Chaisiri K, Komalamisra C, Adelowo OO,
Fagade OE, Banjo OA, Oke AJ, Adler A, Assous MV, Morand S, Raoult D, Rolain JM.
2014. Worldwide emergence of colistin resistance in *Klebsiella pneumoniae* from healthy
humans and patients in Lao PDR, Thailand, Israel, Nigeria and France owing to
inactivation of the PhoP/PhoQ regulator *mgrB*: an epidemiological and molecular stu. Int J
Antimicrob Agents 44:500–507.

55. Trent MS, Ribeiro AA, Doerrler WT, Lin S, Cotter RJ, Raetz CRH. 2001. Accumulation
of a polyisoprene-linked amino sugar in polymyxin-resistant *Salmonella typhimurium* and
Escherichia coli: structural characterization and transfer to lipid A in the periplasm. J Biol
Chem 276:43132–43144.

56. Diene SM, Merhej V, Henry M, El Filali A, Roux V, Robert C, Azza S, Gavory F, Barbe
V, La Scola B, Raoult D, Rolain JM. 2013. The rhizome of the multidrug-resistant
Enterobacter aerogenes genome reveals how new "killer bugs" are created because of a
sympatric lifestyle. Mol Biol Evol 30:369–383.

57. Tindall BJ, Sutton G, Garrity GM. 2017. *Enterobacter aerogenes* Hormaeche and
Edwards 1960 (Approved Lists 1980) and *Klebsiella mobilis* Bascomb *et al*. 1971
(Approved Lists 1980) share the same nomenclatural type (ATCC 13048) on the
Approved Lists and are homotypic synonyms, with consequences for the name *Klebsiella*
mobilis Bascomb *et al*. 1971 (Approved Lists 1980). Int J Syst Evol Microbiol 67:502–
504.

58. Sun S, Negrea A, Rhen M, Andersson DI. 2009. Genetic analysis of colistin resistance in
Salmonella enterica serovar Typhimurium. Antimicrob Agents Chemother 53:2298–2305.

59. Schultz JE, Kanchan K, Ziegler M. 2015. Intraprotein signal transduction by HAMP
domains: a balancing act. Int J Med Microbiol 305:243–251.

60. Lee J-Y, Choi M-J, Choi HJ, Ko KS. 2015. Preservation of acquired colistin resistance in
Gram-negative bacteria. Antimicrob Agents Chemother 60:609–612.

61. Olaitan AO, Morand S, Rolain JM. 2016. Emergence of colistin-resistant bacteria in
humans without colistin usage: a new worry and cause for vigilance. Int J Antimicrob
Agents 47:1–3.

62.   van der Putten BC, Maramoros S, COMBAT consortium, Schultz C. 2019. Genomic
634   evidence for revising the *Escherichia* genus and description of *Escherichia ruysiae* sp.
635   nov. bioRxiv 781724.

636   63.   Son SJ, Huang R, Squire CJ, Leung IKH. 2018. MCR-1: a promising target for structure-
637   based design of inhibitors to tackle polymyxin resistance. Drug Discov Today 24:206–
638   216.

639   64.   McNally A, Cheng L, Harris SR, Corander J. 2013. The evolutionary path to
640   extraintestinal pathogenic, drug-resistant *Escherichia coli* is marked by drastic reduction
641   in detectable recombination within the core genome. Genome Biol Evol 5:699–710.

642   65.   Huys G, Cnockaert M, Janda JM, Swings J. 2003. *Escherichia albertii* sp. nov., a
643   diarrhoeagenic species isolated from stool specimens of Bangladeshi children. Int J Syst
644   Evol Microbiol 53:807–810.

647 **Tables**

Strain	Colistin MIC (μg/ml)	MLST	Date of isolation	History of colistin use
I1121	16	131	22 April 2015	Yes; inhalation and oral
H2129	8	131	22 July 2014	No
G821	16	131	19 March 2013	No
F2745	4	73	2 November 2012	No
E3090	8	10	12 November 2011	No
E2372	4	59	25 August 2011	No
E650	8	162	11 March 2011	No
D2373	8	6901	20 October 2010	Yes; oral
A2361	8	5268	3 November 2007	No
Z821	4	167	2 April 2006	Yes; oral

Table 1: Colistin-resistant *Escherichia* strains isolated from bloodstream infections.

Overview of colistin-resistant bloodstream isolates, including the MIC of colistin, MLST type

determined through whole genome sequencing, date of isolation, and information on the use of

colistin three months before the isolation of the colistin-resistant isolate, and if applicable, route

of administration. The MIC values represent the median of three independent replicate

experiments performed in triplicate.

Strain	Colistin MIC (µg/ml)
BW25113	0.25
BW27848	0.125
BW25113::Tn7 Empty	0.25
BW27848::Tn7 Empty	0.125
BW27848::Tn7 BW25113	0.125
BW27848::Tn7 I1121	2
BW27848::Tn7 I1121m	0.25
BW27848::Tn7 H2129	4
BW27848::Tn7 H2129m	0.25
BW27848::Tn7 G821	4
BW27848::Tn7 G821m	0.25
BW27848::Tn7 F2745	0.25
BW27848::Tn7 E3090	0.125
BW27848::Tn7 E2372	0.25
BW27848::Tn7 D2373	0.5
BW27848::Tn7 A2361	0.125
BW27848::Tn7 Z821	2
BW27848::Tn7 Z821m	0.125

Table 2. MICs of colistin of strains generated in this study.

E. coli strain BW27848 is the Δ*basRS* mutant of BW25113 (34). The *basRS* alleles of

colistin-resistant strains from this study were inserted into the *att*Tn7 site of BW27848. The

addition of "m" to a strain name indicates that the construct has been modified through inverse

PCR site-directed mutagenesis to reverse the mutation associated with colistin resistance. The

values represent the median of three independent replicate experiments performed in triplicate.

Figure legends

Figure 1: Colistin-resistant strains are not clonally related, and carry diverse acquired

antibiotic resistance genes

A) The phylogenetic tree represents the core genome alignment (874 kbp) of the

colistin-resistant strains and 210 publicly available *E. coli* and *E. albertii* genome sequences.

One representative reference strain per *E. coli* phylogroup is indicated (64). For *E. albertii*, the

LMG20976 type-strain is indicated (65). The different phylogroups of *E. coli* are indicated with

coloured backgrounds. Phylogroup A is coloured yellow; B1, grey; B2, orange; D, blue; E,

purple; F, green. The ST131 lineage of *E. coli* in phylogroup B2 is indicated by a dark orange

background. The *E. albertii* branch is indicated by a pink background. The colistin-resistant

strains characterized in this study are depicted in red, and highlighted by a red filled circle. **B)**

The phylogenetic tree represents the core genome alignment (3.55 Mbp) of the three

colistin-resistant ST131 strains and 19 publicly available ST131 *E. coli* strains genome

sequences. The colistin-resistant strains characterized in this study are depicted in red. Clades A,

B, and C of ST131 are indicated by purple, blue, and orange backgrounds respectively. **C)**

Antibiotic resistance genes in the genome sequences were detected by ResFinder 3.2 (40).

Classes of antibiotic resistance genes are abbreviated as follow: PMX, polymyxin resistance;

AGC, aminoglycoside resistance; TET, tetracycline resistance; BLA, beta-lactam resistance;

SUL, sulfonamide resistance; TMP, trimethoprim resistance; PHE, phenicol resistance; QLN,

quinolone resistance; MCL, macrolide.

Figure 2: MALDI-TOF spectra of lipid A from colistin-resistant nosocomial *Escherichia*

strains.

Negative ion MALDI-TOF mass spectrometry spectra of lipid A purified from **A)**

colistin-resistant strains and colistin-susceptible MG1655. Data represent the mass to charge

(*m/z*) ratios of each lipid A species detected and are representative of three extractions. **B)**

Proposed lipid A structures of the species produced by *E. coli* strains. **C)** Proposed lipid A

structures of *E. albertii* strain A2361. Modifications to unmodified lipid A are depicted in red.