Title: Loss of Aurora kinase signaling allows lung cancer cells to adopt an alternative

cell cycle and form multinucleated polyploid giant cells that resist anti-mitotic drugs

Author list: Vural Tagal[1*], Michael G. Roth[1,2]

Affiliations: Departments of Biochemistry[1], Harold Simmons Comprehensive Cancer

Center [2], UT Southwestern Medical Center, Dallas TX 75390 USA

Corresponding author*:

Vural Tagal vural.tagal@utsouthwestern.edu

Abstract

Polyploid multinucleated (PP/MN) giant cancer cells are common in tumors and have been associated with resistance to cancer therapy, tumor relapse, malignancy, immunosuppression, metastasis, cancer stem cell production and modulation of the tumor microenvironment. However, the molecular mechanisms that cause these cells to form are yet known. In this study, we discover that Aurora kinases are synergistic determinants of a switch from the proliferative cell cycle to polyploid growth and multinucleation in lung cancer cell lines. When Aurora kinases are inhibited together, lung cancer cells uniformly grow into PP/MN giant cells. These cells have adopted an endocycle in which the genome replicates, mitosis is omitted and cells grow in size. Consequently, such cells continue to safely grow in the presence of antimitotic agents. These polyploid multinucleated cancer cells can re-enter the proliferative cell cycle and grow in cell number when the treatment is terminated.

Main

The goal of cancer medicine is to eradicate tumors entirely. However, a major challenge in treating cancer is posed by intratumor heterogeneity[1,2]. The body of a tumor includes a diverse collection of cells harboring distinct molecular profiles with differential levels of sensitivity to treatment. The tumor population as a whole has the potential to access all the cellular mechanisms encoded in the genome and drug treatment selects for those cells that happen to have adopted an escape mechanism. To combat this, a goal of cancer medicine is to better understand the compositional and functional complexity of tumors in order to find more effective strategies to overcome the disease.

A well-known cancer cell type in tumor populations is called polyploid multinucleated (PP/MN) giant cancer cells [3-5]. These cells contain multiple nuclei with significantly elevated genomic content (sometimes more than 100 fold) when compared to other cancer cells within the same tumor. The frequency of giant cells can increase markedly in high-grade malignant cancers or following anticancer treatment. They can remain viable and metabolically active for extended periods of time. They can secrete an array of growth

factors, cytokines and chemokines [3,6]. Immunosuppressive proteins including programmed

death-ligand 1 (PD-L1) were also found to be overexpressed in these cells [7]. Therefore, they

are responsible of contributing to a microenvironment advantageous for tumor growth and

survival.

PP/MN cancer cells can repopulate, and form macroscopic colonies and spheroids *in vitro*

as they generate tumors when inoculated into mice [8-10]. These cells gain the properties of

cancer stem cells and produce daughter cells through asymmetric cell division patterns [11-13].

Compared with diploid cancer cells, these daughter cells express fewer epithelial markers

and acquire a mesenchymal phenotype, which is a key transformation for cancer

development, progression and metastasis [10,14,15].

Commonly, the ratio of PP/MN cancer cells is higher in high-grade malignant tumors than in

low-grade tumors, in relapsing tumors after chemotherapy than in tumors before

chemotherapy, and in the metastatic foci than in the primary tumor [16]. Such cells are known

to escape from cytotoxicity induced by major anti-mitotic agents including taxanes and vinca

alkaloids [17-21]. Emerging evidence has demonstrated that PP/MN cancer cells arise in lung

cancer [20], cervical carcinoma [22], ovarian cancer [10,23], prostate cancer [24,25], gliobastoma [26],

colorectal cancer [11,16,27], and breast cancer [15,28,29]. The role of PP/MN giant cancer cells and

their daughter cells in resistance to therapy, tumor relapse, metastasis, immunosuppression,

modulation of the tumor microenvironment and generation of cancer stem cells is well

documented. Therefore, revealing the molecular events that cause PP/MN giant cells to form

can lead to clinically relevant approaches for treating recurrent and metastatic disease,

which remains a major global health issue despite extensive research over the past century.

In the course of studying hypersensitivity to inhibitors of Aurora kinase A [30], we previously

reported an unusual dose response relationship to Aurora kinase inhibitors (AURKi). In

some non-small cell lung cancer (NSCLC) lines, more cells survived the inhibitor treatment

and kept growing at high than at lower drug concentrations. We show in the work presented

below that many NSCLC cell lines grow and uniformly form PP/MN giant cancer cells by

adopting an endocycle when Aurora kinases are inhibited. Endocycles omit mitosis, do not

require the mitotic machinery and therefore, provide a safe path for growth by avoiding the

catastrophic consequences of anti-mitotic perturbagens. Thus, at concentrations that induce

endocycling, we confirmed that Aurora kinase inhibitors protect NSCLC cells from the

cytotoxicity induced by all anti-mitotic cancer agents tested; but not agents targeting

mechanisms functional in interphase. These endocycling NSCLC cells can re-enter the

proliferative cell cycle if Aurora kinase inhibitors are withdrawn. We also demonstrate that

chemical programming of the cells to endocycle with Aurora kinase inhibitors can serve as a

methodology to identify antimitotic agents in a simple, high-throughput fashion.

Results

NSCLC cells exhibit three distinct dose response phenotypes to Aurora kinase A

inhibitors

Aurora kinase A is an important regulator of mitosis and functions in the assembly of the

mitotic spindle [31-33]. Several different Aurora A inhibitors have been developed and are

currently in clinical trials as anti-cancer drug candidates [34]. In a previous study, we

investigated these inhibitors and observed an unusual drug response relationship with an

Aurora kinase A inhibitor, MLN8237/Alisertib (MLN8237 henceforth) [35]. In contrast to the

standard sigmoidal or monophasic dose-response phenotypes (Fig. 1a, c), a subgroup of

cell lines treated with MLN8237 exhibited a multiphasic "camelback" drug-response pattern

(Fig. 1e). VX-689, another Aurora kinase A inhibitor, also caused a similar response pattern

in the same panel of non-small cell lung cancer (NSCLC) lines (Fig. 1b, d, f).

This unexpected drug response pattern of increased cell survival at higher drug

concentrations suggested that secondary targets of these inhibitors might control a

mechanism leading to drug resistance. This would be of concern, since peak plasma

concentrations of MLN8237 in humans reach 3 µM, well above the concentration at which it

selectively inhibits its primary target [36]. Thus, it was of interest to identify the secondary

targets of Aurora A inhibitors and the respective drug resistance mechanism.

Inactivation of Aurora kinases in combination restores cell survival and growth

NSCLC line NCI-H1693 was selected as representative of cell lines capable of this unusual

dose-response because it had the most dramatic phenotype with 35% of the cells growing at

87% of the rate of an untreated population in the presence of higher, potentially non-

selective concentrations of MLN8237 (Fig. 2a). NCI-H1693 cells had a first phase EC50 of

23 nM for MLN8237 and 64 nM for VX-689 (Fig. 2a, c). The EC50s of MLN8237 and VX-689

for the second, "restorative" phase in which the viability increased, was 180 nM and 26 μM,

respectively. At the highest and clinically irrelevant concentrations (Phase 3), MLN8237

gains a "general toxin" characteristic, which was not investigated in this study. In parallel, we

measured Caspase 3/7 activity to determine if apoptotic cell death occurred in the presence

of MLN8237 and VX-689 at any of concentrations measured. Apoptosis was first significantly

detected at 16 nM for MLN8237 and 47 nM for VX-689. However, caspase activation was

not detected during the second phase of the response to each drug (Fig. 2b, d). In an

orthogonal assay for cell growth, NCI-H1693 cells were stained with crystal violet after seven

days treatment with either MLN8237 or VX-689 and cell densities were compared with the

density of cells stained on day 0. We found that growth of NCI-H1693 cells was inhibited at

Aurora A-selective concentrations and was restored at 410 nM and 33.3 μM of MLN8237

and VX-689, respectively (Fig. 2e, g). Thus, the second phase of response to these two

inhibitors not only helped NCI-H1693 cells survive, but also restored their growth.

Kinase inhibitors frequently are selective for their intended target only over a narrow

concentration range [37-39]. Aurora kinases B and C are two orthologues of Aurora A in

humans and their high degree of homology is a well-known challenge for creating inhibitors

specific to Aurora kinase A. Therefore, Aurora B and Aurora C were obvious candidates for

the proteins responsible for reversing cytotoxic effects at higher concentrations of Aurora A

inhibitors. Aurora kinases auto-phosphorylate when bound to their activators and their

phosphorylation levels directly indicate their activation status in cells. In addition, Histone H3

is a well-characterized substrate of Aurora kinases B and C, and its phosphorylation at

Serine 10 is a direct reporter of Aurora B or C activity [40-43]. For these reasons, the

phosphorylation statuses of Aurora kinases A, B, C and Histone H3 were measured 48

hours after treating NCI-H1693 cells with MLN8237 or VX-689 at a series of concentrations

covering their two distinct phases of response. In these experiments, Paclitaxel at its EC100

(20 nM) was used as a positive control to inhibit the cell cycle at the mitotic stage where

Aurora kinases are active. We observed increased phosphorylation of Aurora B and C, and

Histone H3 starting at concentrations 50 nM of MLN8237 and 140 nM of VX-689. However,

starting at 410 nM of MLN8237 and 33.3 µM of VX-689, phosphorylation of Aurora B and C,

and Histone H3 was completely undetectable (Fig. 2f, h). Thus, there was a precise overlap

between cell death, apoptosis and selective inhibition of Aurora A and between lack of

apoptosis, continued cell growth and inhibition of Aurora B and C (Fig. 2i, j).

To test the hypothesis that Aurora B and C are the secondary targets causing the increase

in survival and growth of cancer cells, we used two selective Aurora B and C inhibitors,

AZD1152/Barasertib (AZD1152 henceforth) and GSK1070916 [44,45]. We first verified their

selectivity for Aurora B and C compared to Aurora A by measuring auto-phosphorylation

levels of Aurora kinases A, B and C, and phosphorylation of Histone H3 in NCI-H1693 cells.

Paclitaxel at its EC100 was used as a positive control. At 50 nM AZD1152 or 15 nM

GSK1070916 auto-phosphorylation of Aurora B, C and Histone H3, was not detected; but

Aurora A was phosphorylated (Fig. 3a, b). This indicated that at those concentrations, NCI-

H1693 cells were arrested in mitosis at a point after activation of Aurora A. However, starting

at 1.23 µM for AZD1152 or 140 nM for GSK1070916, phosphorylation of Aurora A was not

detected, indicating that both drugs also inhibit Aurora A at higher concentrations.

The dose-response pattern of viability of NCI-H1693 cells to AZD1152 and GSK1070916

resembled the multiphasic "camelback" response of those cells to Aurora A selective

inhibitors MLN8237 and VX-689 (Fig. 3c-f). AZD1152 and GSK1070916 each caused

significant reduction in viability of NCI-H1693 cells at concentrations selective for Aurora B

and C. However, growth was restored when NCI-H1693 cells were treated with

concentrations above 1.23 µM and 140 nM of AZD1152 and GSK1070916, respectively (Fig.

3g-h), where they began to antagonize Aurora A as a secondary target. These data

suggested that inhibiting Aurora A or Aurora B and C activities alone caused significant

mitotic catastrophe leading to cytotoxicity; but inhibition of all three Aurora kinases together

did not (Fig. 3i, j).

To ensure that Aurora kinases A, B and C, but not a previously unrecognized fourth target,

are solely responsible of restoring viability induced with individual treatments of each of

these four inhibitors, we treated NCI-H1693 cells with these four inhibitors as doublet

combinations at concentrations where they are selective for their primary targets (Fig. 4a, b).

Combination therapy of Aurora A and Aurora B, C inhibitors did not cause additive or

synergistic cytotoxicity; but instead allowed cell survival and growth (Fig. 4c, e). We

measured the activities of Caspase 3/7 (Fig. 4d, f) and used crystal violet staining (Fig. 4g, i)

to assess apoptosis and cell growth. Inhibiting the Aurora kinases together allowed NCI-

H1693 cells to survive and grow whereas inhibiting Aurora A or B selectively was inhibitory.

Consequently, concentrations of Aurora B, C inhibitors selective for their target could

prevent toxicity by Aurora A inhibitors when combined together in NCI-H1693 cells (Fig. 4h,

j).

Based on these observations, we hypothesized that pan-Aurora kinase inhibitors would be

non- or minimally effective on NCI-H1693 cells when compared with the first response

phases of MLN8237, VX-689, AZD1152 and GSK1070916. We performed biochemical

assays identical to the ones performed with the four Aurora A or B and C selective inhibitors.

We observed that pan-Aurora inhibitors, VX-680/Tozasertib (VX-680 henceforth) and PHA-

739358/Danusertib (PHA-739358 henceforth) have a resistance response pattern

resembling the second phase of multiphasic response of the first four selective inhibitors

(Extended Data Fig. 1a-h).

To determine how frequently NSCLC cell lines survived simultaneous inhibition of all Aurora

kinases, we measured the dose-responses to MLN8237 in a panel of 77 NSCLC and 4

immortalized human bronchial epithelial cell (HBEC) lines. More than fifty percent of these

cell lines had a multiphasic camel-back drug response pattern with more cell survival and

growth at the pan-Aurora kinase inhibitory phase in comparison to the responses at lower

Aurora A-selective doses (Fig. 5a). In addition, the rest of tested NSCLC lines were either

uniformly sensitive or uniformly resistant to the effective doses of MLN8237 with a sigmoidal

or monophasic response, respectively (Fig. 5a). An identical screen with VX-689 produced

results similar to those obtained with the screen of MLN8237 (Fig. 5b). In these studies,

Erlotinib, an EGFR-targeted therapy agent and FDA-approved drug for NSCLCs, was used

as control in parallel to MLN8237 and VX-689, on the same day at identical conditions and

no multiphasic "camelback" response pattern was observed (Fig. 5c). Overall, our data

indicated that the multiphasic response pattern against Aurora A inhibitors is common for

NSCLC cell lines, and more cells survived and continued to grow when all Aurora kinases

were inhibited in comparison to selective Aurora A inhibition.

Inactivation of Aurora kinase signaling allows cells to develop into polyploid

multinucleated (PP/MN) giant cancer cells

To determine what type of growth cells adopt after the loss of Aurora kinase signaling, we

examined cellular morphology and DNA content after inhibiting Aurora kinases individually or

in combination. After the inhibition of Aurora kinases with the non-selective high

concentration of MLN8237, we observed that NCI-H1693 cells grow substantially in size

instead of numbers and therefore, cellular ATP levels increase similar to untreated

proliferating cells (Fig. 6a-b). Their nuclear morphology together with enlarged size and

elevated DNA content, demonstrated that these cells develop into polyploid multinucleated

(PP/MN) giant cancer cells (Fig. 6b), a cancer cell type frequently found in drug-resistant

tumor populations and high-grade tumors. The average area of cells treated with pan-Aurora

kinase inhibitor for four days increased and cells had much more nuclear DNA in

comparison to the untreated ones (Fig. 6c). Polyploid genome was also compartmentalized

in multiple nuclei in these cells (Fig. 6c). Cell numbers were not increasing (Fig. 6d) in the

treated samples, but were balanced by increasing the cellular size. Therefore, total ATP

levels present in the treated wells remained similar to the wells with untreated cells. DNA

replication also continued (Fig. 6d). Increasing DNA content and cell size are characteristics

of an endocycle [46]. Overall, our findings suggested that after inhibiting Aurora kinase

signaling, cells might omit all or most of mitosis, continue with intact Gap (G) and Synthesis

(S) phases of cell cycle, and develop into PP/MN giant cancer cells (Fig. 6e).

Induced PP/MN giant cancer cells resist anti-mitotic agents

If an endocycle variant avoids mitosis when Aurora kinase signaling is turned off, we

hypothesized that mitotic machinery including spindle-related components and regulators

should partially or completely be dispensable in the induced PP/MN giant cells. Thus, cells

would be non-responsive to anti-mitotic agents under these conditions instead of causing

additive or synergistic effects whereas perturbing interphase cellular machineries should

remain cytotoxic (Fig. 7a). We tested NCI-H1693 cells with 17 anti-cancer or cytotoxic

agents including 9 acting during interphase (Gemcitabine, Carboplatin, Cisplatin, Etoposide,

Topotecan, Doxorubucin, Pemetrexed, Methotrexate, 5-Fluorouracil, Aphidicollin and

Bafilomycin A) and 8 inhibiting mitosis (Paclitaxel, Docetaxel, Vinorelbine, Vinblastine, PLK1

inhibitor BI-6727 (Volasertib), CENPE inhibitor GSK923295, EG5 inhibitors Ispinesib and

VS-83). For the cell viability assessments, the EC100 of each agent was used (Extended

Data Table 1). Aurora kinase inhibition prevented toxicity from all anti-mitotic drugs whereas

cells were sensitive to interphase perturbagens regardless of the presence of Aurora

inhibitors (Fig. 7b). In addition to individual treatments with these anti-cancer agents, we also

added a cocktail of the eight anti-mitotic agents to our survey and observed that PP/MN

giant cancer cells generated by pan-Aurora kinase inhibition were able to avoid death even

in that extreme condition (Fig. 7b). In parallel to the cell viability assays, we stained NCI-

H1693 cells with crystal violet to check the status of cell growth on day 7 in comparison to

the cell density on Day 0. Our results confirmed that NCI-H1693 cells survived and

continued to grow in the presence of all anti-mitotics and their cocktail, but not the anti-

interphase perturbagens (Fig. 7c). Lastly, we quantified the extent of tolerance of the PP/MN

giant cells to individual anti-mitotic agents by performing cell viability assays with serial

concentrations of eight anti-mitotic drugs in the presence and absence of pan-Aurora kinase

inhibitors. We observed at least 4 logarithmic (log) magnitude of drug resistance to

Paclitaxel, 5 logs to Docetaxel (Fig. 7d), and at least 2 logs of significant resistance to others

(Extended Data Fig. 2a-f).

Finally, we investigated whether the NCI-H1693 PP/MN cells that had switched to endocycle

could reenter the proliferative cell cycle and regenerate daughter cells. Four days after

inhibiting Aurora kinase signaling with combinations of Aurora A inhibitor, MLN8237 and

Aurora B, C inhibitors AZD1152 or GSK1070916 at concentrations selective for their primary

targets, the media with the combination treatments was washed out and NCI-H1693 cells

were allowed to grow in the absence of any drug treatments for 14 days. NCI-H1693 cells

recovered from the combination treatment and were able to form colonies, indicating the

ability of cells to reenter mitotic cell cycle (Fig. 7e). In comparison, NCI-H1693 cells treated

with Paclitaxel had significantly fewer colonies. These observations indicate that loss of

Aurora kinase signaling initiates an endocycle, which maintains growth in cell size instead of

increases in cell numbers. These cells grow into PP/MN giant cancer cells and can re-enter

a proliferative cell cycle when Aurora kinase signaling is restored (Fig. 7f).

Identification of the target and mechanism of action for biologically active molecules remains

a major task for drug discovery. To explore the possible use of a chemically-induced

endocycle as a high-throughput assay to identify antimitotic chemicals, NCI-H2122 cells, one

of the NSCLC lines with the most dramatic multiphasic camelback response to MLN8237

(Fig. 1e), was used to profile 12 cytotoxic chemicals in the presence or absence of 0.5 uM of

MLN8237, a concentration that induces endocyling (Fig. 8a, b). The chemicals used in this

test had been identified as toxic for subsets of our NSCLC cell line panel in a previous high-

throughput screen, but their mode of action was unknown [47]. Of the 12 chemicals,

MLN8237-induced NCI-H2122 PP/MN giant cells were protected from one, SW172170. To

determine if SW172170, but not the other chemical probes, had an anti-mitotic mechanism

of action as predicted by our assay, 48 hours after treating with the compounds, we checked

for increased phosphorylation of Histone H3 as a sign of mitotic arrest. SW172170 was the

only compound showed increase in Histone H3 phosphorylation levels (Fig. 8d). As an

orthogonal assay, NCI-H2122 cells treated with SW172170 as a single agent were stained

for DNA and tubulin. These cells had condensed DNA and multipolar spindles, which are the

typical characteristics of mitotic arrest and catastrophe (Fig. 8e). This demonstrates that

chemical inhibition of Aurora kinases in a cell line that responds by endocyling can be used

as a screen for anti-mitotic mode of action and therefore, can be employed as a tool to

assess the cell cycle phase (interphase vs mitosis), in which a toxic chemical acts.

Discussion

Tumors and tumor-derived cell lines contain polyploid giant cells with significantly increased

genomic content, often within multiple nuclei. Their frequency increases after exposure to

therapeutic agents. These PP/MN cells are not dormant, and form macroscopic colonies and

tumors in immunodeficient mice [8-10]. Their daughter cells harbor stemness, self-renewal and

mitotic capacities [11-13]. Consistent with these attributes, a significant number of recent

reports with different biological systems (tumor-derived cell lines, animal models and

specimens from cancer patients) have underlined the key roles played by PP/MN giant

cancer cells in tumorigenesis, metastasis, immunosuppression and disease relapse after

conventional cancer treatments [3,4,48]. PP/MN cells contribute to solid tumor heterogeneity

and are the main histological feature of malignant tumor in pathologic diagnosis. Our

findings demonstrate that inactivation of Aurora kinases A, B and C combinatorially is a

signal to switch to an endocycle that forms PP/MN giant cancer cells.

Our results indicate that in NSCLC cells Aurora kinases function together to keep the cancer

cell in a proliferative cell cycle. In many cell types, only Aurora A and B kinases are

expressed and the extent to which Aurora C plays a role in this activity is not determined by

our work. In a majority of NSCLC cell lines we tested, chemical inhibition of all family

members of Aurora kinases allowed the cells to adopt an alternative cell cycle where cells

continue to replicate DNA, become multinucleate and grow in size, but do not divide. In this

state, cells are resistant to all anti-mitotic drugs tested because the mitotic machinery is no

longer engaged. Our findings together suggest that Aurora kinases may operate as a cell

cycle switch and their off status generates PP/MN giant cancer cells tolerant to anti-mitotic

therapy in those cell lines where the alternative cell cycle is available.

In nature there are numerous examples of alternative cell cycles, called endocycles (also

known as endoreplication, endoreduplication or endomitosis), that generate healthy

polyploid cells for specialized functions. Endocycling is essential for the proper development

of specialized cell types in mammalian systems like megakaryocytes, placental trophoblast

cells, hepatocytes and cardiomyocytes [49]. The level of ploidy varies dramatically among

tissues, organs and species. Mammalian placental trophoblast cells can exhibit ploidy levels

of 4096 times the haploid number of chromosomes (4096C) [50,51]. Megakaryocytes generate

nuclei with DNA content up to 128C [52]. The highest recorded level for hepatocytes and

cardiomyocytes is 32C. By adulthood and with ageing, up to 70% of human and 85% of

rodent cardiac myocytes are polyploid [53,54]. The highest polyploidy level recorded in any

animal is ~500,000C, reported from the silk-producing glands of silk moth *Bombyx mori* [55].

The neurons of mollusks exhibit remarkable variations of ploidy, ranging from a low 32C in

the land snail *Triodopsis divesta* to an astonishing 200,000C in the giant neurons of the sea

hare *Aplysia californica* [56,57]. A range of 4C to 64C is typical in plants although the highest

level of polyploidy observed in plants is 24,576C in the endosperm of *Arum maculatum* [58,59].

Cancers can take advantage of endocycles by creating heterogeneity of cell types, some of

which are more drug resistant than the parental cells [60].

It is possible that some cancer cells do this by adopting an alternate cell cycle program

similar to those available to specialized cell types. In the brain, a blood-brain barrier is

formed by endothelial cells and these specialized cell types are known to be decorated with

efflux pumps and upregulate drug metabolizing enzymes [61,62]. The glial cells forming the

barrier in fruit fly *Drosophila melanogaster* was shown to require polyploidization for proper

differentiation [63,64]. In cancers, drug resistance mechanisms to lower the cytosolic drug

concentrations with these two systems are well-documented and could originate from cancer

cells that were capable of growing in endothelial cell-like endocycles.

How mitotic cycles are molecularly routed towards endocycles is largely unknown. However,

a recent study on cells from *Drosophila melanogaster* uncovered a role for Aurora kinase B

in endoreplication and its morphogenesis[65]. Reduced expression of Aurora kinase

orthologues in plants was also shown to control the switch from meristematic cell

proliferation to endoreduplication and differentiation[66]. Furthermore, Aurora kinase A was

found downregulated during the megakaryocyte endomitosis and differentiation in mice[67]. In

our experiments, chemically inhibiting Aurora kinases A or B alone did not induce

endocycling in human lung cancer cells, but led to growth arrest and reduced cell numbers.

Whether this is a difference between animal, plant and insect cells, or cancer and normal

cells remains to be determined. Unexpected adverse interactions in clinical trials with

combination therapies occur and mostly the reasons remain unknown. Currently, 60 clinical

trials with MLN8237, 10 with VX-689, 12 with AZD1152, 14 with GSK1070916, 11 with VX-

680 and 8 with PHA-739358 have been reported and among those many are performed in

combination with other anti-mitotic drugs such as Taxanes and Vinca alkaloids. Among the

concluded trials, some revealed that targeted mitotic inhibitors, including Aurora kinase

inhibitors, were less effective than expected [68,69]. Secondly, enlargement of cells in non-

responsive cancers or tumor models after treatments with various chemo- or -targeted

therapies has been reported [60]. Increased cell size was also observed in studies with Aurora

kinase-targeted therapies including MLN8237 [70-72]. However, the cause for this unexpected

cell enlargement remained unknown. Our study presents a potential mechanistic explanation

for these clinical results.

Lastly, the multiphasic dose-response pattern that we observed with inhibitors of Aurora

kinase signaling is not appropriately recognized by the standard approaches for drug dose-

response curve fitting. It was recently reported that the failure rate of the classical Hill

equation is more than 25% in large scale screens [73]. In experiments investigating the

response of NSCLC cell panels to synthetic chemicals [47], we have observed that dose-

responses poorly represented by sigmoidal curves are quite common. Some of these are

undoubtably due to chemical solubility or other experimental artifacts, but our current report

indicates that some indicate a biological response that is not considered by the assumptions

underlying most commonly used curve fitting programs.

In summary, our data suggest that Aurora kinase signaling functions at the intersection

between proliferative cell cycle and endocycle. Switching to endocycle variants creates

growing polyploid multinucleated cancer cells tolerant to anti-mitotic therapy. PP/MN cancer

cells also retain the capability to reenter the proliferative cell cycle after the treatments are

terminated.

Methods

Cell lines

NSCLC and immortalized HBEC lines were generated by the laboratories of John Minna and

Adi Gazdar and their identities have been confirmed by short tandem repeat analysis of

cellular DNA (Promega Corp, Madison, WI, USA). All cell lines were determined to be

mycoplasma free by testing with the e-Myco Plus Mycoplasma PCR Detection Kit (Bulldog

Bio, 2523448). NSCLC lines were cultured in RPMI-1640 medium (GIBCO, 11875)

supplemented with 10% fetal bovine serum (v/v) (Atlanta Biological, S11550). HBEC lines

were grown in Keratinocyte-SFM medium (GIBCO, 17005) supplemented with EGF and

keratinocyte extract (provided by the manufacturer). HBEC3-KT, HBEC30-KT and HBEC34-

KT are HBECs immortalized with stable expression of CDK4 and hTERT [74].

Reagents

MLN8237 (Alisertib) (ChemieTek, CT-M8237), VX-689 (Selleckchem, S2770), AZD1152

(Barasertib) (Selleckchem, S1147), GSK1070916 (Selleckchem, S2740), VX-680

(Tozasertib) (Chemietek, CT-VX680), PHA-739358 (Danusertib) (Selleckchem, S1107),

Paclitaxel (LC Labs, P-9600), Docetaxel (Sigma-Aldrich, 01885), Vinorelbine detartrate salt

hydrate (Sigma-Aldrich, V2264), Vinblastine sulfate (Sigma-Aldrich, V1377), BI 6727

(Volasertib) (Selleckchem, S2235), GSK923295 (Selleckchem, S7090), Ispinesib (Cayman

Chemical Company, 18014), VS-83 (Calbiochem, 178273), Gemcitabine (ChemieTek, CT-

GEM), Carboplatin (Sigma-Aldrich, C2538), Cisplatin (Sigma-Aldrich, 232120), Etoposide

(LC Labs, E-4488), Topotecan (Tocris, 4562), Doxorubucin hydrochloride (Sigma-Aldrich,

D1515), Pemetrexed (Tocris, 6185), Methotrexate (Tocris, 1230), 5-Fluorouracil (Sigma-

Aldrich, F6627), Aphidicollin (Calbiochem, 178273), Bafilomycin A (Sigma-Aldrich, B1793),

Erlotinib (LC Labs, E-4497) was purchased in powder form and stored at 10 mM or 20 mM

stock concentrations in dimethyl sulfoxide (DMSO) at −20 °C. Phospho-AURKA

(Thr288)/AURKB (Thr232)/AURKC (Thr198) (Cell Signaling, rabbit mAb, 2914, 1:1,000

dilution), Phospho-Histone H3 (Ser10) (Cell Signaling, rabbit mAb, 33770, 1:1,000 dilution)

and β-Actin (MP Biomedicals, mouse mAb, 69100, 1:10,000 dilution) antibodies were used

for immunoblotting assays. Texas Red™-X Phalloidin (ThermoFisher Scientific, T7471,

1:100 dilution), Tubulin antibody (Abcam, ab6160, 1:1000 dilution) and Hoechst 33342

reagent (ThermoFisher Scientific, H3570, 1:2000 dilution) were used for

immunofluorescence assays.

Microplate drug sensitivity assays

NCI-H1693 cells (3,000 per well) were seeded in 96-well plates and incubated at 37 °C for

24 h. Treatment with vehicle (DMSO) or serial dilutions of the indicated compounds in 9

concentrations from 1 to 33,300 nM were performed in triplicate (final DMSO

concentration=0.34% in all wells). After 96 h, media were removed and cell viability was

measured with a CellTiter-Glo assay (Promega, G7573).

A dose–response study of a panel of NSCLC and HBEC cell lines with MLN8237 (Alisertib),

VX-689 and Erlotinib was performed in 384-well plate format. Cells were dispensed at a

density of 1,000–1,500 cells per well with an automated liquid dispenser (BioTek MicroFlo;

Thermo-Fisher, Inc.). Each concentration was screened in triplicate. Cells were incubated for

24 h at 37 °C in the presence of 5% CO_2.with an automated liquid dispenser (BioTek

MicroFlo; Thermo-Fisher, Inc.) Drugs/chemicals were dosed in DMSO at 12 concentrations

ranging from 50 pM to 50 μM (half-log dilutions) in triplicate, using an Echo 555 acoustical

transfer instrument. After incubating the cells at 37 °C for 96 h, cell viability was measured

with a CellTiter-Glo assay. The comparative tests of three agents were performed on the

same days, in identical conditions. Dose responses of each cell line to each agent were

assessed at the indicated concentrations individually. Concentrations that caused increase

in the measured viability after the initial cytotoxic effect of respective agent were colored in

red.

Microplate apoptosis assay

421 Treatments of the indicated compounds were performed in the same manner with the

422 microplate drug sensitivity assays in 96-well plate format. 48 hours after treatments, culture

423 media were removed by centrifuging the assay plates upside down in liquid-collecting

424 containers at 30g for 1 min. Caspase 3 and/or 7 activity was assessed with Caspase-Glo 3/7

425 assay purchased from Promega (G8090).

Crystal violet staining assay

427 NCI-H1693 cells (1 × 10^5 per well) were seeded in 6-well plates and incubated at 37 °C for

428 24 h. Treatment with vehicle (DMSO) or serial dilutions of the indicated compounds in 10

429 indicated concentrations were performed (final DMSO concentration=0.34% in all wells).

430 After seven days, media were removed and cell growth was assessed with a crystal violet

431 staining assay solution (6% glutaraldehyde, 0.5% crystal violet (w/v) in water).

Western blotting

In all, 2 × 10^5 cells were treated with the indicated compounds at indicated concentrations in

six-well format. Forty-eight hours after treatments, the culture medium was aspirated and

cells were washed with PBS. Cells were lysed in RIPA lysis buffer (50 mM Tris buffer pH 8,

150 mM NaCl, 1% NP-40, 0.5% sodium, 0.1% SDS, 0.5% Sodium deoxycholate)

supplemented with protease and phosphatase inhibitors. The protein concentrations of cell

lysates were measured with the Protein DC assay (Bio-Rad, 500-0111). Phosphorylation of

histone H3 and Aurora kinases A, B, C, and β-Actin expression as internal control were

monitored by immunoblotting with monoclonal antibodies.

Light microscope and immunofluorescence assay

Cells in transparent bottom plates were fixed with 4% formaldehyde in PBS for 15 min at

room temperature. After washing with PBS, cells were permeabilized with 0.25% Triton-X in

PBS for 15 min at room temperature. After washing with PBS, cells were incubated with 10%

Bovine Serum Albumin (BSA) blocking solution for 1 h. Next, cells were immunostained with

Texas Red™-X Phalloidin or monoclonal Tubulin antibodies overnight. Hoechst 33342

reagent was used to stain DNA. Microscopic observations were performed on an EVOS FL

Cell Imaging System. Mean nuclear area of cells after the treatments with the indicated

compounds were calculated with the ImageJ software and graphed.

Data Availability

The data supporting this study are available from the corresponding author upon reasonable

request.

Acknowledgements

We thank Bruce Posner and the personnel of the Simmons Comprehensive Cancer Center

High-throughput Screening Core Facility for aid in dose-response experiments on the

NSCLC cell line panel. The human tumor-derived cell panel was created by Drs. John D.

Minna and Adi F. Gazdar. We thank our colleague Iryna Zubovych for critical comments.

This work was supported by grants CA176284 and CA221978 to MGR.

Author Information

Affiliations

Departments of Biochemistry, UT Southwestern Medical Center, Dallas, TX, USA

Vural Tagal, Michael G. Roth

Harold Simmons Comprehensive Cancer Center, UT Southwestern Medical Center, Dallas,

TX, USA

Michael G. Roth

Author Contributions

V.T. designed and performed the experiments. V.T. and M.G.R. conceived the study,

analyzed data and wrote the manuscript.

Ethics Declarations

Competing Interests

The authors declare no competing interests.

References

1   Dagogo-Jack, I. & Shaw, A. T. Tumour heterogeneity and resistance to cancer therapies. *Nat*
Rev Clin Oncol **15**, 81-94, doi:10.1038/nrclinonc.2017.166 (2018).
2   Marusyk, A., Almendro, V. & Polyak, K. Intra-tumour heterogeneity: a looking glass for
cancer? *Nat Rev Cancer* **12**, 323-334, doi:10.1038/nrc3261 (2012).
3   Mirzayans, R. & Murray, D. Intratumor Heterogeneity and Therapy Resistance: Contributions
of Dormancy, Apoptosis Reversal (Anastasis) and Cell Fusion to Disease Recurrence. *Int J Mol*
Sci **21**, doi:10.3390/ijms21041308 (2020).
4   Chen, J. *et al.* Polyploid Giant Cancer Cells (PGCCs): The Evil Roots of Cancer. *Curr Cancer*
Drug Targets **19**, 360-367, doi:10.2174/1568009618666180703154233 (2019).
5   Liu, J. The "life code": A theory that unifies the human life cycle and the origin of human
tumors. *Semin Cancer Biol* **60**, 380-397, doi:10.1016/j.semcancer.2019.09.005 (2020).
6   Bharadwaj, D. & Mandal, M. Senescence in polyploid giant cancer cells: A road that leads to
chemoresistance. *Cytokine Growth Factor Rev* **52**, 68-75, doi:10.1016/j.cytogfr.2019.11.002
(2020).
7   Diaz-Carballo, D. *et al.* A Distinct Oncogenerative Multinucleated Cancer Cell Serves as a
Source of Stemness and Tumor Heterogeneity. *Cancer Res* **78**, 2318-2331, doi:10.1158/0008-
5472.CAN-17-1861 (2018).
8   Fujiwara, T. *et al.* Cytokinesis failure generating tetraploids promotes tumorigenesis in p53-
null cells. *Nature* **437**, 1043-1047, doi:10.1038/nature04217 (2005).
9   Weihua, Z., Lin, Q., Ramoth, A. J., Fan, D. & Fidler, I. J. Formation of solid tumors by a single
multinucleated cancer cell. *Cancer* **117**, 4092-4099, doi:10.1002/cncr.26021 (2011).
10   Zhang, S. *et al.* Generation of cancer stem-like cells through the formation of polyploid giant
cancer cells. *Oncogene* **33**, 116-128, doi:10.1038/onc.2013.96 (2014).
11   Lopez-Sanchez, L. M. *et al.* CoCl2, a mimic of hypoxia, induces formation of polyploid giant
cells with stem characteristics in colon cancer. *PLoS One* **9**, e99143,
doi:10.1371/journal.pone.0099143 (2014).
12   Solari, F. *et al.* Multinucleated cells can continuously generate mononucleated cells in the
absence of mitosis: a study of cells of the avian osteoclast lineage. *J Cell Sci* **108 (Pt 10)**,
3233-3241 (1995).

506	13	Sundaram, M., Guernsey, D. L., Rajaraman, M. M. & Rajaraman, R. Neosis: a novel type of
507		cell division in cancer. *Cancer Biol Ther* **3**, 207-218, doi:10.4161/cbt.3.2.663 (2004).
508	14	Zhang, S., Mercado-Uribe, I., Hanash, S. & Liu, J. iTRAQ-based proteomic analysis of polyploid
509		giant cancer cells and budding progeny cells reveals several distinct pathways for ovarian
510		cancer development. *PLoS One* **8**, e80120, doi:10.1371/journal.pone.0080120 (2013).
511	15	Fei, F. *et al.* The number of polyploid giant cancer cells and epithelial-mesenchymal
512		transition-related proteins are associated with invasion and metastasis in human breast
513		cancer. *J Exp Clin Cancer Res* **34**, 158, doi:10.1186/s13046-015-0277-8 (2015).
514	16	Zhang, S., Zhang, D., Yang, Z. & Zhang, X. Tumor Budding, Micropapillary Pattern, and
515		Polyploidy Giant Cancer Cells in Colorectal Cancer: Current Status and Future Prospects.
516		*Stem Cells Int* **2016**, 4810734, doi:10.1155/2016/4810734 (2016).
517	17	Niu, N. *et al.* Linking genomic reorganization to tumor initiation via the giant cell cycle.
518		*Oncogenesis* **5**, e281, doi:10.1038/oncsis.2016.75 (2016).
519	18	Niu, N., Mercado-Uribe, I. & Liu, J. Dedifferentiation into blastomere-like cancer stem cells
520		via formation of polyploid giant cancer cells. *Oncogene* **36**, 4887-4900,
521		doi:10.1038/onc.2017.72 (2017).
522	19	Ogden, A., Rida, P. C., Knudsen, B. S., Kucuk, O. & Aneja, R. Docetaxel-induced
523		polyploidization may underlie chemoresistance and disease relapse. *Cancer Lett* **367**, 89-92,
524		doi:10.1016/j.canlet.2015.06.025 (2015).
525	20	Wertz, I. E. *et al.* Sensitivity to antitubulin chemotherapeutics is regulated by MCL1 and
526		FBW7. *Nature* **471**, 110-114, doi:10.1038/nature09779 (2011).
527	21	Finkin, S., Aylon, Y., Anzi, S., Oren, M. & Shaulian, E. Fbw7 regulates the activity of
528		endoreduplication mediators and the p53 pathway to prevent drug-induced polyploidy.
529		*Oncogene* **27**, 4411-4421, doi:10.1038/onc.2008.77 (2008).
530	22	Puck, T. T. & Marcus, P. I. Action of x-rays on mammalian cells. *J Exp Med* **103**, 653-666,
531		doi:10.1084/jem.103.5.653 (1956).
532	23	Lv, H. *et al.* Polyploid giant cancer cells with budding and the expression of cyclin E, S-phase
533		kinase-associated protein 2, stathmin associated with the grading and metastasis in serous
534		ovarian tumor. *BMC Cancer* **14**, 576, doi:10.1186/1471-2407-14-576 (2014).
535	24	Mittal, K. *et al.* Multinucleated polyploidy drives resistance to Docetaxel chemotherapy in
536		prostate cancer. *Br J Cancer* **116**, 1186-1194, doi:10.1038/bjc.2017.78 (2017).
537	25	Amend, S. R. *et al.* Polyploid giant cancer cells: Unrecognized actuators of tumorigenesis,
538		metastasis, and resistance. *Prostate* **79**, 1489-1497, doi:10.1002/pros.23877 (2019).
539	26	Bigner, S. H. *et al.* Specific chromosomal abnormalities in malignant human gliomas. *Cancer*
540		*Res* **48**, 405-411 (1988).
541	27	Fei, F. *et al.* Formation of Polyploid Giant Cancer Cells Involves in the Prognostic Value of
542		Neoadjuvant Chemoradiation in Locally Advanced Rectal Cancer. *J Oncol* **2019**, 2316436,
543		doi:10.1155/2019/2316436 (2019).
544	28	Sirois, I. *et al.* A Unique Morphological Phenotype in Chemoresistant Triple-Negative Breast
545		Cancer Reveals Metabolic Reprogramming and PLIN4 Expression as a Molecular
546		Vulnerability. *Mol Cancer Res* **17**, 2492-2507, doi:10.1158/1541-7786.MCR-19-0264 (2019).
547	29	Wang, X. *et al.* EMT-related protein expression in polyploid giant cancer cells and their
548		daughter cells with different passages after triptolide treatment. *Med Oncol* **36**, 82,
549		doi:10.1007/s12032-019-1303-z (2019).
550	30	Tagal, V. *et al.* SMARCA4-inactivating mutations increase sensitivity to Aurora kinase A
551		inhibitor VX-680 in non-small cell lung cancers. *Nat Commun* **8**, 14098,
552		doi:10.1038/ncomms14098 (2017).
553	31	Bury, L. *et al.* Plk4 and Aurora A cooperate in the initiation of acentriolar spindle assembly in
554		mammalian oocytes. *The Journal of cell biology* **216**, 3571-3590, doi:10.1083/jcb.201606077
555		(2017).

32 Hannak, E., Kirkham, M., Hyman, A. A. & Oegema, K. Aurora-A kinase is required for
centrosome maturation in Caenorhabditis elegans. *The Journal of cell biology* **155**, 1109-
1116, doi:10.1083/jcb.200108051 (2001).
33 Gopalan, G., Chan, C. S. & Donovan, P. J. A novel mammalian, mitotic spindle-associated
kinase is related to yeast and fly chromosome segregation regulators. *The Journal of cell
biology* **138**, 643-656 (1997).
34 Otto, T. & Sicinski, P. Cell cycle proteins as promising targets in cancer therapy. *Nat Rev
Cancer* **17**, 93-115, doi:10.1038/nrc.2016.138 (2017).
35 Tagal, V. *et al.* SMARCA4-inactivating mutations increase sensitivity to Aurora kinase A
inhibitor VX-680 in non-small cell lung cancers. *Nature communications* **8**, 14098,
doi:10.1038/ncomms14098
http://www.nature.com/articles/ncomms14098#supplementary-information (2017).
36 Cervantes, A. *et al.* Phase I pharmacokinetic/pharmacodynamic study of MLN8237, an
investigational, oral, selective aurora a kinase inhibitor, in patients with advanced solid
tumors. *Clin Cancer Res* **18**, 4764-4774, doi:10.1158/1078-0432.CCR-12-0571 (2012).
37 Bain, J., McLauchlan, H., Elliott, M. & Cohen, P. The specificities of protein kinase inhibitors:
an update. *Biochem J* **371**, 199-204, doi:10.1042/BJ20021535 (2003).
38 Bain, J. *et al.* The selectivity of protein kinase inhibitors: a further update. *Biochem J* **408**,
297-315, doi:10.1042/BJ20070797 (2007).
39 Davies, S. P., Reddy, H., Caivano, M. & Cohen, P. Specificity and mechanism of action of
some commonly used protein kinase inhibitors. *Biochem J* **351**, 95-105 (2000).
40 Walter, A. O., Seghezzi, W., Korver, W., Sheung, J. & Lees, E. The mitotic serine/threonine
kinase Aurora2/AIK is regulated by phosphorylation and degradation. *Oncogene* **19**, 4906-
4916, doi:10.1038/sj.onc.1203847 (2000).
41 Bolton, M. A. *et al.* Aurora B kinase exists in a complex with survivin and INCENP and its
kinase activity is stimulated by survivin binding and phosphorylation. *Molecular biology of
the cell* **13**, 3064-3077, doi:10.1091/mbc.e02-02-0092 (2002).
42 Giet, R. & Glover, D. M. Drosophila aurora B kinase is required for histone H3
phosphorylation and condensin recruitment during chromosome condensation and to
organize the central spindle during cytokinesis. *The Journal of cell biology* **152**, 669-682
(2001).
43 Crosio, C. *et al.* Mitotic phosphorylation of histone H3: spatio-temporal regulation by
mammalian Aurora kinases. *Molecular and cellular biology* **22**, 874-885 (2002).
44 Anderson, K. *et al.* Biochemical characterization of GSK1070916, a potent and selective
inhibitor of Aurora B and Aurora C kinases with an extremely long residence time1. *Biochem
J* **420**, 259-265, doi:10.1042/BJ20090121 (2009).
45 Mortlock, A. A. *et al.* Discovery, synthesis, and in vivo activity of a new class of
pyrazoloquinazolines as selective inhibitors of aurora B kinase. *Journal of medicinal
chemistry* **50**, 2213-2224, doi:10.1021/jm061335f (2007).
46 Lee, H. O., Davidson, J. M. & Duronio, R. J. Endoreplication: polyploidy with purpose. *Genes
Dev* **23**, 2461-2477, doi:10.1101/gad.1829209 (2009).
47 McMillan, E. A. *et al.* Chemistry-First Approach for Nomination of Personalized Treatment in
Lung Cancer. *Cell* **173**, 864-878 e829, doi:10.1016/j.cell.2018.03.028 (2018).
48 Liu, J. The dualistic origin of human tumors. *Semin Cancer Biol* **53**, 1-16,
doi:10.1016/j.semcancer.2018.07.004 (2018).
49 Edgar, B. A., Zielke, N. & Gutierrez, C. Endocycles: a recurrent evolutionary innovation for
post-mitotic cell growth. *Nat Rev Mol Cell Biol* **15**, 197-210, doi:10.1038/nrm3756 (2014).
50 Zybina, E. V. [Different modes of cell reproduction in the placental differentiation of
mammals]. *Tsitologiia* **25**, 1103-1119 (1983).

51 Zybina, E. V. & Chernogriadakaia, N. A. [A study of the polyploid nuclei of the giant
trophoblast cells of several species of rodents using phase contrast microscopy]. *Tsitologiia*
18, 161-165 (1976).
52 Ravid, K., Lu, J., Zimmet, J. M. & Jones, M. R. Roads to polyploidy: the megakaryocyte
example. *J Cell Physiol* **190**, 7-20, doi:10.1002/jcp.10035 (2002).
53 Mollova, M. *et al.* Cardiomyocyte proliferation contributes to heart growth in young
humans. *Proc Natl Acad Sci U S A* **110**, 1446-1451, doi:10.1073/pnas.1214608110 (2013).
54 Soonpaa, M. H., Kim, K. K., Pajak, L., Franklin, M. & Field, L. J. Cardiomyocyte DNA synthesis
and binucleation during murine development. *Am J Physiol* **271**, H2183-2189,
doi:10.1152/ajpheart.1996.271.5.H2183 (1996).
55 Perdrix-Gillot, S. DNA synthesis and endomitoses in the giant nuclei of the silkgland of
Bombyx mori. *Biochimie* **61**, 171-204, doi:10.1016/s0300-9084(79)80066-8 (1979).
56 Lasek, R. J. & Dower, W. J. Aplysia californica: analysis of nuclear DNA in individual nuclei of
giant neurons. *Science* **172**, 278-280, doi:10.1126/science.172.3980.278 (1971).
57 Mandrioli, M., Mola, L., Cuoghi, B. & Sonetti, D. Endoreplication: a molecular trick during
animal neuron evolution. *Q Rev Biol* **85**, 159-169 (2010).
58 Barow, M. Endopolyploidy in seed plants. *Bioessays* **28**, 271-281, doi:10.1002/bies.20371
(2006).
59 Scholes, D. R. & Paige, K. N. Plasticity in ploidy: a generalized response to stress. *Trends Plant*
Sci **20**, 165-175, doi:10.1016/j.tplants.2014.11.007 (2015).
60 Puig, P. E. *et al.* Tumor cells can escape DNA-damaging cisplatin through DNA
endoreduplication and reversible polyploidy. *Cell Biol Int* **32**, 1031-1043,
doi:10.1016/j.cellbi.2008.04.021 (2008).
61 Arvanitis, C. D., Ferraro, G. B. & Jain, R. K. The blood-brain barrier and blood-tumour barrier
in brain tumours and metastases. *Nat Rev Cancer*, doi:10.1038/s41568-019-0205-x (2019).
62 Schinkel, A. H. *et al.* Disruption of the mouse mdr1a P-glycoprotein gene leads to a
deficiency in the blood-brain barrier and to increased sensitivity to drugs. *Cell* **77**, 491-502,
doi:10.1016/0092-8674(94)90212-7 (1994).
63 Unhavaithaya, Y. & Orr-Weaver, T. L. Polyploidization of glia in neural development links
tissue growth to blood-brain barrier integrity. *Genes Dev* **26**, 31-36,
doi:10.1101/gad.177436.111 (2012).
64 Von Stetina, J. R., Frawley, L. E., Unhavaithaya, Y. & Orr-Weaver, T. L. Variant cell cycles
regulated by Notch signaling control cell size and ensure a functional blood-brain barrier.
Development **145**, doi:10.1242/dev.157115 (2018).
65 Rotelli, M. D. *et al.* A Cyclin A-Myb-MuvB-Aurora B network regulates the choice between
mitotic cycles and polyploid endoreplication cycles. *PLoS Genet* **15**, e1008253,
doi:10.1371/journal.pgen.1008253 (2019).
66 Petrovska, B. *et al.* Plant Aurora kinases play a role in maintenance of primary meristems
and control of endoreduplication. *New Phytol* **193**, 590-604, doi:10.1111/j.1469-
8137.2011.03989.x (2012).
67 Goldenson, B., Kirsammer, G., Stankiewicz, M. J., Wen, Q. J. & Crispino, J. D. Aurora kinase A
is required for hematopoiesis but is dispensable for murine megakaryocyte endomitosis and
differentiation. *Blood* **125**, 2141-2150, doi:10.1182/blood-2014-12-615401 (2015).
68 Benada, J. & Macurek, L. Targeting the Checkpoint to Kill Cancer Cells. *Biomolecules* **5**, 1912-
1937, doi:10.3390/biom5031912 (2015).
69 Manchado, E., Guillamot, M. & Malumbres, M. Killing cells by targeting mitosis. *Cell Death*
Differ **19**, 369-377, doi:10.1038/cdd.2011.197 (2012).
70 Kogiso, M. *et al.* Concurrent Inhibition of Neurosphere and Monolayer Cells of Pediatric
Glioblastoma by Aurora A Inhibitor MLN8237 Predicted Survival Extension in PDOX Models.
Clin Cancer Res **24**, 2159-2170, doi:10.1158/1078-0432.CCR-17-2256 (2018).

71 Haruki, T. *et al.* Relationship Between MYC Family Status and Sensitivity to Aurora Kinase
Inhibitors in Neuroendocrine and Other Lung Cancer Cell Lines. *J Thorac Oncol* **12**, S2044-
S2044 (2017).
72 Lee, J. W. *et al.* Combined Aurora Kinase A (AURKA) and WEE1 Inhibition Demonstrates
Synergistic Antitumor Effect in Squamous Cell Carcinoma of the Head and Neck. *Clin Cancer*
Res **25**, 3430-3442, doi:10.1158/1078-0432.CCR-18-0440 (2019).
73 Di Veroli, G. Y. *et al.* An automated fitting procedure and software for dose-response curves
with multiphasic features. *Sci Rep* **5**, 14701, doi:10.1038/srep14701 (2015).
74 Ramirez, R. D. *et al.* Immortalization of human bronchial epithelial cells in the absence of
viral oncoproteins. *Cancer Res* **64**, 9027-9034, doi:10.1158/0008-5472.CAN-04-3703 (2004).
665

Main Figure Titles and Legends

Figure 1. There are distinctly different response phenotypes to Aurora Kinase inhibition. a-f, Cells were treated with Aurora A selective inhibitors MLN8237 or VX-689 at the concentrations shown and cell viability was measured with a CellTiter-Glo assay, which measures cellular ATP as an indicator of living cells. Data are means of triplicate biological replicates and error bars are standard deviations (s.d). Some error bars are smaller than the data symbols. Cell lines representative of three distinct dose response curves are shown.

Figure 2. Aurora A inhibitors at higher concentrations antagonize their own cytotoxic activity and restore growth. **a,** NCI-H1693 cells were treated with serial concentrations of MLN8237 for four days and cell viability was measured with a CellTiter-Glo assay, which measures ATP. **b,** NCI-H1693 cell line was treated with serial concentrations of MLN8237 for two days and active apoptosis was measured with a CaspaseGlo 3/7 assay. **c, d,** CellTiter-Glo and CaspaseGlo 3/7 assays were repeated with VX-689 treatments. Data are means of triplicate biological replicates with s.d. Some error bars are smaller than the data symbols. **e,** NCI-H1693 cells were treated with serial concentrations of MLN8237 or **g,** VX-689 for seven days and cell growth quantified with a crystal violet staining assay. **f,** Cell lysates of NCI-H1693 cell line were collected 48-hours after treating with MLN8237 or **h,** VX-689 and immunoblotted to measure phosphorylated histone H3 and phosphorylated Aurora A, B, and C. Paclitaxel was used in both sets of immunoblots as positive control and B-Actin was used as internal immunoblotting control. **i, j,** Overlay of cell viability, caspase 3/7 activity and crystal violet staining images with immunoblotting results at Aurora A kinase selective and non-selective (pan-Aurora kinase) inhibitory concentrations of MLN8237 and VX-689.

Figure 3. Aurora B, C inhibitors exhibit a camelback drug response pattern similar to Aurora A inhibitors. a, Cell lysates of NCI-H1693 cells were collected 48-hours after treating with AZD1152 (Barasertib) or **b,** GSK1070916 and immunoblotted to monitor phosphorylated histone H3 and phosphorylated Aurora A, B, C levels. Paclitaxel was used in both sets of immunoblots as positive control and B-Actin was used as internal immunoblotting control. **c,** NCI-H1693 cells were treated with serial concentrations of

AZD1152 for four days and cell viability was measured with a CellTiter-Glo assay. **d,** NCI-

H1693 cells were treated with serial concentrations of AZD1152 for two days and cell death

was measured with a CaspaseGlo 3/7 assay. **e, f,** Assays were repeated with GSK1070916

treatments. Data are means of triplicate biological replicates with s.d. Some error bars are

smaller than the data symbols. **g,** NCI-H1693 cell line was treated with serial concentrations

of AZD1152 or **h,** GSK1070916 for seven days and cell growth was measured with a crystal

violet staining assay. **i, j,** Overlay of cell viability, caspase 3/7 activity and crystal violet

staining images with immunoblotting results at selective and non-selective concentrations of

AZD1152 and GSK1070916.

Figure 4. Combining Aurora A and B, C inhibitors at concentrations selective for their

primary targets prevents cytotoxicity and allows cell growth. a, Cell lysates of NCI-

H1693 cells were collected 48-hours after treating with combinations of MLN8237 and

AZD1152 or **b,** MLN8237 and GSK1070916 and immunoblotted to detect phosphorylated

histone H3 and phosphorylated Aurora A, B, C. Paclitaxel was used in both sets of

immunoblots as a positive control and β-Actin was used as an internal immunoblotting

control. **c,** NCI-H1693 cells were treated with combinations of MLN8237 and AZD1152 at

concentrations selective for their primary targets for four days and cell viability was

measured with a CellTiter-Glo assay. **d,** NCI-H1693 cells were treated with combinations of

MLN8237 and AZD1152 at their selective for two days and cell death was measured with a

CaspaseGlo 3/7 assay. **e, f,** Identical assays were repeated with GSK1070916 treatments.

g, NCI-H1693 cells were treated with combinations of Aurora A and B/C selective inhibitors

for seven days and cell growth was measured with a crystal violet staining assay. **h,** NCI-

H1693 cells were treated with serial concentrations of MLN8237 in combination with serial

dilutions of AZD1152 for four days and cell viability was measured with a CellTiter-Glo

assay. **i, j,** Identical assays were repeated with GSK1070916 treatments. Data from

CellTiter-Glo and CaspaseGlo 3/7 assays are presented as means of triplicate biological

replicates with s.d. Some error bars are smaller than the data symbols.

Figure 5. Survey of a NSCLC line panel reveals that more than 50% of cell lines have more cancer cell survival and growth at higher concentrations of Aurora kinase A inhibitors. a, A panel of NSCLC and HBEC lines was treated with MLN8237, **b,** VX-689 or **c,** Erlotinib. The results of dose–response experiments are shown in heatmap format. Decrease in viability in comparison to the untreated control is shown in shades of gray and white. Increase in viability in comparison to the cytotoxicity induced at lower concentrations is colored in shades of red.

Figure 6. Loss of Aurora kinase signaling allows cells to grow into polyploid multinucleated giant cancer cells. a, NCI-H1693 cells were treated with 0.41 μM of MLN8237 for four days and the ATP present was measured with a CellTiter-Glo assay. **b,** NCI-H1693 cells were treated with DMSO or 0.41 μM of MLN8237 for 20 days, stained with Phalloidin (red) and Hoechst (blue), and visualized at 10X magnification (Scale bar, 400 μM). **c,** NCI-H1693 cells were treated with 0.41 μM of MLN8237 for four days and visualized with bright-field or fluorescent microscopy at 10X magnification. Hoechst reagent was used to visualize the nuclei (Scale bar, 400 μM). **d,** The numbers of cells and their mean nuclear area from treated and untreated control samples are graphed. **e,** A proposed model of an endocycle induced with the loss of Aurora kinase signaling.

Figure 7. Inhibition of Aurora kinase signaling provides resistance to anti-mitotic toxins, but not interphase toxins. a, NCI-H1693 cells were co-treated with DMSO or 0.5 μM of MLN8237 and IC100 concentrations of interphase or mitotic perturbagens. **b,** Cell viability and growth were assessed with a CellTiter-Glo assay (after four days of treatment) or **c,** crystal violet staining assay (after seven days of treatment), respectively. **d,** NCI-H1693 cells were co-treated with DMSO or 0.5 μM of MLN8237 and serial concentrations of Paclitaxel or Docetaxel. Cell viability was measured with a CellTiter-Glo assay. **e,** NCI-H1693 cell line was treated with 0.1 μM of MLN8237 and 0.2 μM AZD1152 or 20 nM GSK1070916 for four days. Treatment media were replaced with drug-free media for 14 days. Colony formation was measured with a crystal violet staining assay. Data of CellTiter-Glo assay are means of triplicate biological replicates with s.d. Some error bars are smaller

than the data symbols. **f,** Proposed model of the cellular life cycle in the presence and

absence of Aurora kinase signaling.

Figure 8. Induction of polyploid multinucleated giant cancer cells through chemical

inhibition of Aurora kinase signaling can serve as a strategy to identify anti-mitotic

chemicals. a, Flow chart of the drug discovery project designed to identify probes

selectively cytotoxic to NCI-H2122 cells. **b,** Proposed model for the anticipated outcome of

screen identifying new anti-mitotic perturbagens. **c,** NCI-H2122 cells were treated with

DMSO or 0.5 μM of MLN8237 and each of 12 chemical hits identified in a compound library

screen. Cell viability was measured with a CellTiter-Glo assay. Paclitaxel was used as

positive control. Structure of SW172170, the anti-mitotic candidate probe is shown. **d,** Cell

lysates of NCI-H2122 cell line were collected 48-hours after treating with each chemical and

immunoblotted to measure phosphorylated histone H3. β-Actin was used as internal

immunoblotting control. Normalized band intensities of phosphorylated Histone H3 from

treated and untreated control samples are graphed. **e,** NCI-H2122 cells were treated with

SW172170 for four days and visualized with bright-field or fluorescent microscopy. Hoechst

reagent and monoclonal antibodies against Tubulin were used to visualize the nuclei and

mitotic spindles, respectively (Scale bar, 25 μM).

Extended Data Figure Titles and Legends

Extended Data Figure 1. Pan-Aurora kinase inhibitors do not inhibit growth of NCI-

H1693 cells. a, NCI-H1693 cells were treated with serial concentrations of VX-680 for four

days and cell viability was measured with a CellTiter-Glo assay. **b,** NCI-H1693 cells were

treated with serial concentrations of VX-680 for two days and apoptosis was measured with

a CaspaseGlo 3/7 assay. **c, d,** Assays were repeated with PHA-739358 treatments. Data

are means of triplicate biological replicates with s.d. Some error bars are smaller than the

data symbols. **e,** NCI-H1693 cell line was treated with serial concentrations of VX-680 or **f,**

PHA-739358 for seven days and cell growth was measured with a crystal violet staining

assay. A concentration of MLN8237 specific for Aurora kinase A and an EC100

concentration of Paclitaxel were used as controls. **g, h,** Overlay of cell viability, caspase 3/7

activity and crystal violet staining images with immunoblotting results.

Extended Data Figure 2. Endocycling cells resist anti-mitotic perturbagens. a,

Vinorelbine, **b,** Vinblastine, **c,** BI 6727 (Volasertib), **d,** GSK923295, **e,** SB-715992 (Ispinesib)

or **f,** VS-83 for four days and cell viability was measured with a CellTiter-Glo assay. Data are

means of triplicate biological replicates with s.d. Some error bars are smaller than the data

symbols.

Extended Data Table 1. EC100 values of all used FDA-approved chemotherapy agents

for NSCLCs or cytotoxic poisons in NCI-H1693.

Drug	EC100 (µM)
Paclitaxel	0.02
Docetaxel	0.02
Vinorelbine	0.02
Vinblastine	0.02
BI 6727 (Volasertib)	0.1
GSK923295	0.1
Ispinesib	0.02
VS-83	10
Gemcitabine	0.1
Topotecan-HCl	0.2
Methotrexate	0.2
Pemetrexed	1
Cisplatin	20
Carboplatin	100
Doxorubucin	2
Aphidicollin	5
Bafilomycin A	0.02
5-Fluorouracil	20

Sigmoidal drug-response relationship-Sensitive group

Monophasic drug-response relationship-Resistant group

Multiphasic "camelback" drug-response relationship

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Extended Data Figure 1

Extended Data Figure 2