May 27, 2020

Evolutionary Response to Resource Deprivation: Parallelism and Nonmonotonicity

Megan G. Behringer[§][*], Wei-Chin Ho[§][*], Samuel F. Miller, John C. Meraz, Gwyneth F. Boyer, Michael Lynch

Center for Mechanisms of Evolution, The Biodesign Institute; Arizona State University; Tempe, AZ 85287; United States of America.

[§]These authors contributed equally to this work.

*Please direct correspondence to:

Megan G. Behringer: megan.behringer@asu.edu
Wei-Chin Ho: weichinh@asu.edu

Abstract

Establishing reliable frameworks for predicting unknown outcomes from empirical

observations is of great interest to ecologists and evolutionary biologists. Strong predictability in

evolutionary responses has been previously demonstrated by the repeated observation of similar

phenotypes or genotypes across multiple natural or experimental populations in analogous

environments. However, the degree to which evolutionary outcomes can be predicted across

environmental gradients, or in fluctuating environments, remains largely unexplored. Presumably,

the phenotypic evolution in an intermediate environment could be interpolated from the evolved

phenotypes observed in two extreme environments, but this assumption remains to be fully tested.

Here, we report on the experimental evolution of *Escherichia coli* under three nutritional transfer

periods: every day, every 10 days, and every 100 days, representing increasing severity in

feast/famine cycles. After 900 days of experimental evolution, populations experiencing

intermediate durations of starvation had evolved longer times to reach maximum growth rate,

smaller colony sizes, higher biofilm formation, and higher mutation rates than populations

evolving in the other environmental extremes. Because the intermediately starved populations

exhibit significantly high molecular parallelism, these distinct phenotypes are likely due to non-

monotonic deterministic forces instead of increased stochastic forces commonly associated with

fluctuating environments. Our results demonstrate novel complexities associated with

evolutionary predictability across environmental gradients and highlight the risk of using

interpolation in evolutionary biology.

Keywords: *Escherichia coli*, Evolutionary Predictability, Experimental Evolution, Fluctuating
Environment, Mutation Rate, Parallel Evolution, Starvation, Subpopulation Structure

Evolution is driven by both deterministic (e.g., natural selection) and stochastic (e.g., genetic drift or environmental uncertainty) forces [1-8]. Across analogous environments, similar phenotypes or genotypes are commonly observed suggesting strong deterministic forces and a potential for evolutionary outcomes to be highly predictable [8-12]. Experimental evaluation suggests that higher environmental similarity leads to higher evolutionary predictability or parallelism, as two populations are more likely to have similar derived phenotypes and genotypes when evolved in more similar environments [13-15]. Despite several empirical tests of evolutionary predictability across multiple distinct constant environments, there is still a lack of sufficient examination across fluctuating environments that differ both in the intensity of the stress imposed and the periodicity of fluctuations. Generating such an understanding of evolutionary predictability across fluctuating environments is of particular importance, as they are more representative of what organisms often face in nature.

One set of experimental conditions with significant biological relevance is feast/famine cycles. Many microbial populations, including facultative pathogens with clinical or agricultural importance, frequently encounter environments with fluctuating resource availability [16,17]. *Escherichia coli*, for instance, has a broad habitat as an opportunistic pathogen, ranging from soil and wastewater to the lower gut, often oscillating between feast and famine [18-20]. Prior studies have demonstrated that *E. coli* populations are capable of surviving periods of prolonged starvation, during which they accumulate mutations in the genes *rpoS* [21] and *lrp* [22] that confer a phenotype known as growth advantage in stationary phase (GASP)[23-25]. Other phenotypes observed in *E. coli* populations during resource limitation have included temporarily increased mutation rates [26,27] and the evolution of trade-offs between maximum growth rate and sustainable growth [28]. Again, an understanding of how these genetic and phenotypic responses scale across

different durations of resource limitation is still lacking. Thus, conclusions that the phenotypic

response to repeated starvation, or any fluctuating selective pressure, scales progressively with

an increasing duration of stress is premature.

To address this issue, we surveyed the evolutionary response of *E. coli* populations along

a selective gradient of three resource replenishment cycles: every day, every 10 days, and every

100 days, and determined if constraints on the predictability of genotypes and phenotypes were

associated with repeated long-term starvation. If, for example, evolutionary predictability is

highly related to environmental similarity, phenotypic evolution in the intermediate environment

(10-days) should be predictable from the evolved phenotypes observed in the two environmental

extremes (1- and 100-days), i.e., the plot of phenotypic values against the environmental gradient

should be monotonic (a continuously increasing or decreasing function). As different molecular

and phenotypic traits have the potential to reveal distinct patterns in evolutionary responses, we

focus on evolved changes in growth dynamics, mutation rate, genomic evolution rate, and

evidence of subpopulation structure. We also examine the parallelism of derived mutations in

populations evolving under the same or different nutrient limitation conditions.

Results

To establish the evolution experiment, ancestral lines with two different initial genetic

backgrounds were used: a wild-type strain (WT) and a WT-derived strain with impaired methyl-

directed mismatch repair (MMR-) wherein an engineered deletion of *mutL* yields a ~150×

increase in the single nucleotide mutation rate [29]. Three distinct resource-limitation cycle

conditions (1-day, 10-days, and 100-days between transfers into fresh media) were deployed, and

each combination of genetic background and resource-limitation cycle condition was replicated

in eight parallel experiments, resulting in 48 experimental populations ($8\times3\times2$). Thus, this design

allows us to examine the parallelism within each treatment combination, as well as the

divergence between treatments. After 900 days of experimental evolution, changes in growth

behaviours were evaluated. Notably, populations subjected to 10-day cycles of resource

limitation take significantly longer to reach their maximum growth rate (T_{max}) than ancestral, 1-

and 100-day populations (**Fig. 1a**). In addition, 10-day populations present significantly smaller

colonies on LB agar than ancestral, 1-day, and 100-day populations (**Fig. 1b-e**).

Previously, we observed that early biofilm formation was one of the major growth

phenotypes that emerges in our culture conditions [30]. Examination of culture tubes after 24 h

revealed that 1-day and 10-day populations produce visible biofilms at the surface-air interface,

while ancestral and 100-day populations generally do not (**Fig. 1f**). Further, quantification of

biofilm formation reveals that 10-day populations produce significantly thicker biofilms than

ancestor, 1-day, and 100-day populations (**Fig. 1g**). This languid early growth combined with

increased capacity for biofilm formation in 10-day populations illustrates a pattern of non-

monotonic evolution for growth phenotypes in response to chronic resource limitation.

Next, as the mutation rate has been reported to increase during acute starvation, we

performed fluctuation tests for rifampicin resistance [31] on clones isolated from evolved

populations to check for evolved changes in the mutation rate. For populations originating from

WT backgrounds, all 12 clones isolated from 10-day resource-limitation cycles exhibited

significantly higher rates of rifampicin resistance than the WT ancestor (**Fig. 2a**). This evolved

widespread increase in mutation rate is unique to populations from 10-day resource-limitation

cycles, as significant increases were only observed for three clones isolated from 1-day

populations and none from 100-day populations. In MMR- backgrounds, mutation-rate patterns

were similar, as six out of twelve clones isolated from 10-day resource-limitation cycles evolved

higher rifampicin resistance rates than the MMR- ancestor, while only two and three tested

clones show such significance from 1-day and 100-day populations, respectively (**Fig. 2a**). Thus,

a trend of non-monotonic evolutionary responses is also observable in mutation rates --

cultivation in the intermediate, 10-day resource-limitation cycle often results in the evolution of

an increased mutation rate, while cultivation at the other extremes commonly results in relatively

little change in mutation rate.

As the effects of individual mutations can vary, it is possible for phenotypes and

genotypes to evolve at different rates and produce different evolutionary response patterns [32]. To

determine if the rate of genomic evolution also exhibits a similar non-monotonic pattern as

observed in growth dynamics and mutation rate, we assessed genomic-sequence divergence

throughout 900 days of evolution. Using high-throughput population-genomic data collected

every 100 days (mean coverage $> 100\times$, **Supplemental Table 1**), genomic divergence was

estimated by summing the derived-allele frequencies for all single nucleotide polymorphisms

(SNPs) at each time point. The rate of genomic evolution is then defined as the slope of the

linear regression of summed derived-allele frequencies against the estimated number of

generations for each resource-limitation treatment. Consistent with the magnitudes of change

observed in growth dynamics and mutation rate, 10-day populations also experienced the highest

genomic evolution rates (**Fig. 2b**). This pattern is true regardless of initial genetic background,

and alternative measures of genomic divergence do not qualitatively affect the results

(**Supplemental Fig. 1a-d**). In addition to the genomic evolution rate, comparison of the fixation

probabilities for detectable nonsynonymous, intergenic, and synonymous mutations, suggests

that positive selection is not significantly stronger in 10-day populations than the other

evolutionary conditions. Instead, it is more likely that the increased mutation rates that evolve in

response to 10-day resource-limitation cycles are the greatest contributing factor to the observed

increases in the genomic evolution rate (**Supplemental Fig. 1e-g**).

Although genomic and phenotypic evolutionary responses to resource availability diverge

from monotonically increasing or decreasing functions, the interaction of these responses may

result in different patterns for intraspecific diversity and community structure. For 1-day cycles

of resource limitation, our culture environment is known to support subpopulation structure [30].

However, how increases in the length of resource-limitation cycles differentially affects the

ability to create and maintain subpopulation structure remains unknown. We used a clade-aware

hidden Markov chain (caHMM), which assumes the coexistence of two clades (major and minor),

to infer whether each mutation belongs to the basal clade (i.e., mutations that sweep through the

entire population before the establishment of subpopulations), major clade, or minor clade. This

approach also infers if a mutation remains polymorphic or reaches fixation within its assigned

clade at each sample time point [33].

Application of caHMM revealed that resource-limitation cycle length affects the

persistence of subpopulation structure, as the durations of time that major and minor clade are

inferred to coexist (**Fig. 3a**) are significantly shorter in 100-day populations than in 1-day and

10-day populations (**Fig. 3b**). Consistent with this observation, the proportion of fixed mutations

assigned to the basal clade in a population is negatively correlated with the duration of

subpopulation structure persistence (**Supplemental Fig. 2a**), being greatest in 100-day

populations (**Supplemental Fig. 2b**). Plotting the temporal distribution of derived-allele

frequencies reveals that mutations with intermediate frequencies (0.3-0.7) are less abundant in

100-day populations (**Supplemental Fig. 2c**), again consistent with a lower tendency to form

and further maintain subpopulation structure.

To ensure that a lower tendency to support subpopulation structure is unlikely to be an

artefact of insufficient evolutionary time, we normalized our comparisons by the number of

derived mutations observed in each evolutionary treatment. For example, 1-day populations have

experienced the same amount of genomic evolution after 300-days of culture as 100-day

populations have experienced after 500-days of culture (calculated based on **Fig. 2b**). However,

by 300-days, fixation of mutations in the basal clade slowed and subpopulation structure

stabilized in 13 of 16 daily-transferred populations. Conversely in 100-day populations, although

the same amount of molecular evolution occurred by 500-days, subpopulation structure only

appeared to stabilize in 2 of 16 populations. Thus, evolutionary outcomes relating to the

maintenance of subpopulation structure in response to resource limitation can be described by a

decreasing monotonic function – such that, as the duration between resource replenishment

increases the ability to sustain subpopulation structure decreases. This also reveals that the

interaction of forces shaping the previously mentioned phenotypic and genotypic evolution in

response to resource-limitation may result in discordant patterns with the evolution of

subpopulation structure.

Lastly, to distinguish which mutations are likely positively selected and adaptive from

mutations that are likely promoted to fixation due to genetic draft [34], we focused on the

nonsynonymous mutations which were inferred to be fixed within either the basal, major, or

minor clades of the evolving populations. Here, two metrics were calculated which indicate the

degree of parallel evolution within a treatment combination and further identify genes which are

likely selective targets due to an overrepresentation of fixed nonsynonymous mutations: sum of

G-scores [35] and Bray–Curtis similarity [15]. Moreover, if the fixed non-synonymous mutations are

concentrated in a smaller subset of genes within a resource-limitation/genetic background

combination then the resulting values for these two metrics will further increase. For all

treatment combinations, both the sum of *G*-scores and the Bray-Curtis similarity values were

significantly greater than expected based on a simulated null distribution (**Fig. 4a-b**). Thus,

revealing a significant amount of parallel evolution within treatments and suggesting that

positive selection is likely shaping these populations. Interestingly, the statistical significance of

both metrics for 10-day and 100-day populations is also much greater than for 1-day populations,

suggesting a smaller pool of adaptive targets and that ability of the 10-day and 100-day

populations to explore the adaptive landscape is more limited, possibly due to their

environmental harshness.

To more deeply investigate the observed parallelism in likely adaptive mutations, we

conducted gene ontology analyses focusing on genes that were determined to significantly

contribute to the increased sum of *G*-scores in each treatment combination (**Fig. 4c;**

Supplemental Table 2). We additionally investigated genes which were overrepresented for

disruptive 'structural' mutations such as indels or IS-element insertions (**Fig. 4d; Supplemental**

Table 3). Furthermore, an additional set of 115 nonsynonymous mutations which were parallel

at the nucleotide level – that is, these mutations independently arose and are responsible for

identical amino-acid substitutions in at least two experimental populations within a treatment

combination (**Supplemental Fig. 3a-b; Supplemental Table 4**). Results of the gene ontology

analyses revealed that genes associated with DNA repair (GO:0006281) or replication

(GO:0006260) more commonly experienced mutations in 10-day populations than in 1- or 100-

day populations (**Supplemental Table 5**), consistent with the evolution of increased mutation-

rates in 10-day populations. However, some parallelism was observed which may indicate a

general response to resource limitation as genes related to global gene expression (regulation,

termination, and fidelity of transcription and translation), resource import, and metabolism

(**Supplemental Table 6-9**) were enriched for mutations in both 10- and 100-day populations.

Since populations evolving in response to 10-day and 100-day resource-limitation cycles

exhibited parallelism in biological processes that are enriched for likely adaptive mutations but

differed so greatly in their resulting phenotypes, we hypothesized that treatment-specific

differences must exist in the individual genes or amino acids experiencing mutations. Here,

notable differences in parallel amino-acid substitutions occurred in the stringent response

regulator *crp* (Y42C, 10-day populations; E56K, 100-day populations), the transcription

termination factor *rho* (R109H, 100-day populations), RNA polymerase subunit *rpoB* (D915G,

10-day populations; E859K, 100-day populations), and the translation elongation/ribosome

recycling factor, *fusA* (R437C, I654N, 10-day populations) (**Supplemental Table 4**). Although

these genes are all involved in processes that have broad effects on gene expression, it is

plausible that the mutation of these genes may result in widely-different phenotypic effects.

Particularly, in the case of *crp*, as the mutations that arise in 10-day and 100-day populations are

located in or proximal to the RNA polymerase-interacting domain. Here, mutation variants have

been previously characterized to have significantly different effects on *crp*'s interaction with

RNA polymerase and the activation of promoters [36,37]. Thus, further studies are needed to

understand how the alteration of genes that are responsible for these basic molecular processes is

beneficial to populations experiencing distinct levels of repeated starvation – both in terms of

differential expression and the resulting phenotypes.

Discussion

Several conclusions can be drawn from the reported results. First, we observed that the

pattern of evolutionary responses for many molecular and phenotypic traits do not follow simple

increasing or decreasing functions across an environmental gradient of resource limitation, i.e.,

they are non-monotonic. This is at least qualitatively consistent with previous theoretical and

experimental studies of fluctuating environments, suggesting that periodic stress results in higher

phenotypic variability than longer, more sustained stress – increasing their potential to evolve

novel adaptations [38,39]. In addition to changes in biofilm density, growth rate, and colony size, we

also notably see this phenomenon extend to the evolution of increased mutation rates, which are

often induced in stressful environments [40]. Consistent with our results, population-genetic

models have suggested that higher mutation rates can evolve in fluctuating environments and

increase in prevalence due to second-order selection [41-43]. In these cases, mutators are expected to

be the most prevalent in environments where conditions fluctuate at intermediate frequencies [44].

Thus, taking into account these observed non-monotonic patterns, our results highlight the risk of

assuming that expected evolutionary outcomes in response to intermediate levels of resource

limitation can be inferred from observations in more extreme environments.

Second, evolutionary patterns can be discordant between phenotypic, genomic, and

intraspecific community-structure outcomes. While we observed a non-monotonic pattern in the

evolution of growth dynamics, mutation rate, and genome-evolution rate, persistence in

subpopulation structure instead declined monotonically. Specifically, we observed that the

duration during which multiple clades coexisted within a population decreased with longer

resource-limitation cycles. This echoes the findings that resource-deficient environments

promote detachment of biofilm formers [45] or metastasis of cancer cells [46], both of which increase

population homogeneity. A pattern of decreasing subpopulation structure may arise due to a

reduced pool of beneficial mutations combined with increased population bottlenecks in more

resource-limited populations, limiting the ability to explore the adaptive landscape [47]. In effect,

this would result in a decreased potential for clonal interference and adaptive diversification, two

phenomena that would otherwise cause the appearance of coexisting clades.

Lastly, albeit surprisingly, repeated cycles of resource limitation do not appear to

promote the fixation of mutations affecting genes commonly associated with the growth

advantage in stationary phase (GASP) phenotype, such as *rpoS* [21,48] and *lrp* [22]. In fact, other

global regulators and genes with broad effects on transcript and protein expression (*crp*, *rho*,

rpoB, and *fusA*) are instead targets for likely adaptive mutations. Further investigation is needed

to determine if the absence of known GASP-associated mutations is due to strain-specific/culture

environment differences or because these mutations are transient, with benefits that are no longer

realized after repeated exposure to extreme starvation.

Materials and Methods

Construction of ancestral strains and experimental evolution

Experimental populations originated from one of four ancestral strains. Populations

denoted as WT are descendants of PMF2, a prototrophic derivative of *E. coli* K-12 str. MG1655,

while populations denoted as MMR- are descendants of PMF5 (PMF2, Δ*mutL*) provided by the

Foster Lab [29]. To introduce a neutral marker that would allow for the periodic screening of

contamination between populations, lambda red recombineering was used to introduce a 3513 bp

deletion removing the *araBAD* operon from one-half of both the WT and MMR-starting strains.

This deletion was confirmed by the appearance of red colonies after plating on TA agar (1%

Arabinose, 1%Tryptone, 0.5% NaCl, 0.1% Yeast Extract, 0.005% TTC (Sigma T8877)) and

next-generation sequencing.

Long-term cultures were established by inoculating 10 mL of LB-Miller broth (Dot

Scientific) with a single-isolated progenitor colony cultivated overnight at 37 °C on LB agar

plates. Resulting cultures were maintained in 16- $\times$ 100 mm glass culture tubes containing 10 mL

LB broth shaking at 175 rpm at 37°C. Selection conditions involved subjecting cultures to

varying degrees of repeated starvation in which 1 mL of thoroughly-vortexed culture was

transferred into 10 mL of fresh LB broth every day, every 10-days, or every 100-days. To

maintain an historical record of the populations and to revive populations in the case of

contamination or extinction, 1 mL of culture was collected pretransfer every 30 days from daily

and 10-day populations, frozen with liquid nitrogen, and stored at -80°C. In order to not disrupt

any potential ecological structure in 100-day populations, copies of these cultures were only

collected pretransfer every 100-days. In addition, 100-day populations were maintained in

triplicate (1 main culture and 2 backup cultures) with only the main culture being used to seed

new triplicate cultures every 100 days. In the case of a whole population extinction, one backup

culture would replace the main culture and the experiment would continue. This practice was

only necessary during the first 200-days as past this time point, no additional whole-population

extinction events occurred. Populations were screened for cross-contamination on days 90, 200,

300, 400, 500, 600, 700, 800, and 900 by streaking on MacConkey Agar (BD Difco)

supplemented with 0.4% Arabinose.

Evaluation of growth phenotypes

Growth curves were measured using an Epoch2 Microplate Spectrophotometer (Biolog).

To prepare cultures for growth rate measurement, 900-day populations and day-0 progenitor

strains were revived frozen -80°C stock by transferring a scrape of ice to 16- x 100 mm culture

tubes containing 10 mL of LB broth. Cultures were incubated overnight at 37°C and shaking at

175 rpm. Following overnight culture, 1.5 μL of each revived population or progenitor were

transferred into a well in a 96-well flat bottom culture plate (Corning/Falcon, 353072) containing

150 μL of LB broth. Evolved populations were measured as an average of eight replicates and

progenitor strains were measured as an average of 16 replicates, with an additional eight wells

dedicated to negative controls consisting solely of LB broth. After inoculation, lids were placed

on 96-well plates and were sealed with parafilm before placement in the spectrophotometer.

Throughout the duration of growth curve measurement, cultures were maintained at 37°C, with

double orbital shaking at 548 cpm. Growth measurements were taken every 15m over the course

of 16h at an OD of 600 nm. Maximum growth rate (r_{max}) was calculated for all measurable

replicates (WT: $n_{ans} = 32$, $n_{1-day} = 16$, $n_{10-day} = 16$, $n_{100-day} = 16$; MMR: $n_{ans} = 29$, $n_{1-day} = 16$, n_{10-day}

$= 12$, $n_{100-day} = 16$) using the Gen5 software (Biotek) as the maximum slope between two

timepoints and time to r_{max} (T_{max}) was recorded as the midpoint between r_{max} determining

timepoints. For daily lines experiencing diauxic growth, calculations of r_{max} and T_{max} were

restricted to the first logistic growth phase.

To measure colony size, revived 900-day populations and day-0 progenitor strains were

grown overnight at 37°C and shaking at 175 rpm in 16- x 100 mm culture tubes containing 10

mL of LB broth. Following overnight incubation, serial dilutions of 100 μL of culture in 900 μL

of phosphate buffered saline were made down to dilutions of 10^{-5}, before plating 100 μL of

diluted culture on to TA agar (1.6% agar, 1% arabinose, 1% tryptone, 0.5% NaCl, 0.1% yeast

extract, 0.0005% TTC) and incubating for 24h at 37°C. TA plates were then photographed with a

20 mm scale in frame using an iPhone XS. Photos were then analysed with ImageJ (NIH,

https://imagej.nih.gov/ij/), by first setting the pixel to distance ratio using the in-frame scale

before flatting the photo to 8-bit black and white, setting the threshold to highlight the colonies,

and using the 'measure particles' tool. Measurements for six arbitrary colonies per analysed

population were recorded.

Presence and absence of biofilm at the surface-air interface was determined by reviving

900-day populations and day-0 progenitor strains in 16- x 100 mm culture tubes containing 10

mL of LB broth overnight at 37°C and shaking at 175 rpm. Following incubation, culture tubes

were visually examined for the presence of biofilm attached to the culture tube at the surface-air

interface. Quantification of biofilm formation in evolved populations was conducted using a

microtiter plate biofilm assay [49]. In a non-treated 96-well plate (Corning/Falcon, 351172), 15 μL

of overnight culture was inoculated into 150 μL of LB broth. Each population was measured for

a total of eight replicates with each 96-well plate containing two columns (16-wells) of *araBAD*-

WT ancestor as a control and at least two columns (16-wells) of blanks. Inoculated 96-well

plates were incubated for 24 h at 37°C. Following incubation, plates were washed with 1x PBS,

stained with 0.1% crystal violet for 10 m (200 μL per well), washed again with 1x PBS, and left

to dry overnight. Crystal violet bound to biofilms was then solubilized with 30% acetate for 15 m

(200 μL per well), transferred to a fresh 96-well plate, and absorbance was quantified with a

using an Epoch2 Microplate Spectrophotometer at 550 nm.

DNA isolation and high-throughput sequencing

High resolution population tracking was conducted by collecting 1 mL of culture at day

90, 200, 300, 400, 500, 600, 700, 800, and 900 of experimental evolution. DNA was extracted

using the DNeasy UltraClean Microbial Kit (Qiagen 12224; formerly MO BIO UltraClean

Microbial DNA Kit) and submitted to either the Hubbard Center for Genomic Analysis at the

University of New Hampshire, the Center for Genomics and Bioinformatics at Indiana

University, or the CLAS Genomics Facility at Arizona State University for library preparation

and sequencing. Libraries were generated with the Nextera DNA Library Preparation Kit

(Illumina, FC-121-1030) following an augmented protocol for optimization of reagent use[50]

before being pooled and sequenced as paired-end reads to a target depth of 100x on an Illumina

HiSeq 2500 (UNH) or an Illumina NextSeq 500 (Indiana; ASU).

Sequencing analysis

Sequencing analysis was performed on the Mason and Carbonate high-performance

computing clusters at Indiana University. Sequencing reads were quality controlled using

Cutadapt v.1.9.1[51] to remove residual adapters and trim low quality sequences. QCed sequences

were then mapped to the *Escherichia coli* K-12 substr. MG1655 reference genome

(NC_000913.3) and all mutations and their frequencies were identified using Breseq v.0.30.2

with the predict-polymorphisms parameter setting[52]. All mutations that were identified to

previously exist in ancestral lines were discarded from the following analysis. Further, our

analysis only included the samples which passed the following additional four quality checks: 1)

we required mean sequencing depths > 10; 2) any WT sample identified to contain the 1,830 bp

deletion in *mutL* from the PMF5 progenitor strain was discarded; 3) regions lacking sequencing

coverage (i.e. depth = 0) must be smaller than 5% of the genome; and 4) the sequencing result

should reflect the correct genetic background in terms of *ara* markers, including a

nonsynonymous SNP at position 66528, an intergenic SNP at position 70289, and a multiple

base substitution mutation (SUB) at position 66533. For an *ara+* line, we required either of two

SNPs showing DAF < 0.2. For an *ara-* line, we required either of two SNPs showing DAF > 0.8

or the SUB is detected. In the end, 395 genomic profiles passed QC and were included in the

following analysis (**Supplemental Table 1**).

In order to remove the mutations that originated from the starting clone before

experimental evolution, we discarded any mutations with a DAF = 100% at one time point for at

least 11 experimental populations with the same genetic background from the analysis. In

addition, as the highly repetitive sequences in *rsx* genes are known to cause errors in SNP

calling[53], they were also removed from the analysis.

Guaranteed generations

The calculation of guaranteed number of generations is based on the following two

observations. First, pre-transfer population density in the culture tube was measured at day 200,

300, 400, 500, 600, 700, 800, 900, by serial dilution in phosphate-buffered saline (PBS) before

plating on LB agar for CFU counts and revealed that the population density does not

significantly change throughout experimental evolution (**Supplemental Fig. 4**). This suggests

the carrying capacity of experimental populations is recovered within a resource-limitation cycle

after 1:10 dilution and the populations must have experienced at least 3.3 ($\log_2 10$) generations

between resource-limitation cycles. Therefore, the first estimate of guaranteed generations (g_1)

per day equals 3.3, 0.33, and 0.033 when the resource-limitation cycle is one day, 10 days, and

100 days, respectively. Second, we found that between resource-limitation cycles novel

mutations arise and are able to be fixed within one sequencing interval. Therefore, given the

carrying capacity K, the population should at minimum experience $\log_2 K$ cell divisions in 100

days. Using estimated $K = 1.21$ x 10^{10}, 2.13 x 10^8, and 3.71 x 10^7 (**Supplemental Fig. 4**), the

second estimate of guaranteed generations (g_2) per day equals 0.30, 0.28, and 0.25 when the

resource-limitation cycle is one day, 10 days, and 100 days, respectively. The final number of

guaranteed generations (g) for each resource-limitation cycle is then determined as the larger

value between the corresponding g_1 and g_2.

376   Given that K was only estimated pre-transfer, when the feeding cycle is long it is possible

that populations have reached to a density K^* higher than K and decreased back to K within one

feeding cycle. If this is true, the first estimated guaranteed generations (g_1') per day should be

$[\log_2(K^*/K) + 3.3]$/RC, where RC is the length of the resource-limitation cycle. While g_1' is

likely to be larger than g, the observed fold-changes in the genomic evolutionary rates between

in longer resource-limitation cycles cannot be fully explained by this possibility. For example, if

the 9.4-fold change in 10-day lines compared to daily lines were entirely due to this possible

underestimation, g_1' needs to be 9.4-fold higher than g_1, which requires an implausible $K^* = 4.7$

x 10^{16}. Similarly, if the 8.8-fold changes in 100-day lines compared to daily lines were entirely

due to this possible underestimation, g_1' needs to be 8.8-fold higher than g, which also requires

an implausible $K^* = 7.2$ x 10^{64}. Therefore, even if we had tracked the population density

throughout the entire experimental evolution, it is likely that we still could not find substantial

effects of changing population densities on our observations.

Rate of genomic evolution

We quantified the level of genomic divergence for each experimental population at each

time point by summing all DAFs of detected mutations. Then we calculated the mean genomic

divergence across all eligible experimental populations in each resource-limitation cycle/genetic

background combination (**Supplemental Table 1**). When estimating the rate of genomic

evolution, the linear regression was performed by the function "lm" in R with formula "mean

genomic divergence ~ guaranteed generations + 0" (linear model), which enforces the y-intercept

as 0.

To explore the possibility of nonlinear relationship between the genomic divergence and

time, we also performed regression using the formula "mean genomic divergence ~ guaranteed

generations + square root of guaranteed generations + 0" (nonlinear model), which was

previously proposed to catch the trend of diminishing returns[35]. However, the results of nested

ANOVA test suggest no strong evidence supporting that the nonlinear model is substantially

better than the linear model (**Supplemental Fig. 1d**).

Fluctuation test and mutation rate estimation

To quantify the genetic mutation rate of our evolved populations and their ancestors, we

performed fluctuation tests as described[22] on clones isolated from each evolved population at 900

days. Briefly, fluctuation tests measure the rate of resistance to the antimicrobial rifampicin

which is conferred by mutations to *rpoB*. For each resource-limitation cycle/genetic background

combination, the four *ara-* populations were assayed. For the first two populations with smaller

population numbers, four independent clones were picked; for the last two populations with

larger population numbers; two independent clones were picked. For each of the WT or MMR-

ancestor, two independent clones were used. For each clone, 40 replicate experiments were

performed. Number of mutants as determined by CFU counts/mL were converted to an estimated

mutation rate and a corresponding 95% confidence interval by the function "newton.LD"

function in the R package "rSalvador"[54] for each clone.

When performing the permutation test for 36 WT clones (12 from 1-day populations, 12

from 10-day populations, and 12 from 100-day populations), we randomly shuffled their labels

of population in 10,000 times. After each shuffle, we counted the number of clones exhibiting

95% confidence interval with the label of 10-day populations. The final *P*-value was determined

by the proportion of these numbers larger or equal to the genuinely observed number exhibiting

95% confidence interval with the label of 10-day populations. The permutation test for 36 MMR-

clones was similarly implemented.

Clade-aware hidden Markov chain analysis (caHMM) and longest coexistence time

To test coexistence time in each experimental population, we performed clade-aware

hidden Markov model (caHMM) using a modified version of previously released code[33]. Our

modifications included: 1) modifying the code in "calculate_clade_hmm_wrapper.py," in order

to continue the iteration of experimental populations, even when the major clade frequency of all

time points for an experimental population cannot be estimated; 2) removing the original cut-off

for minimum coexistence time, as that parameter is too long for our experiments; and 3)

increasing the iteration number for hidden Markov chain from 5 to 50. When performing

caHMM, we constructed an annotated time course of mutations using the original format. If a

mutation is a structural mutation (DEL, INS, and MOB), the sequencing depth is determined by

the mean sequencing depth across the mapped adjacent nucleotide ("coverage_plus" or

"coverage_minus") of associated junction candidates (JCs) in breseq annotated file. The

sequencing depth of time 0 was set as 100.

Using the major clade frequency (f_M) and minor clade frequency (f_m) estimated by

caHMM, we calculated the longest coexistence time as follows: if there is no time point showing

$0.2 < f_M < 0.8$ and $0.2 < f_m < 0.8$, its longest length of coexistence is defined as zero; otherwise, we

found the longest time intervals within which every time point shows $0.01 < f_M < 0.99$ or $0.01 <$

$f_m < 0.99$. After calculating the longest length of coexistence for each population, to test whether

length of resource-limitation cycle and genetic background affects the longest length of

coexistence, we performed a Scheirer-Ray-Hare test, (nonparametric version of two-way

ANOVA) using the function "scheirerRayHare" in the R package "rcompanion"[55]. The sample

size for each resource-limitation cycle/genetic background combination is eight. The succeeding

post hoc analysis is done by Mann-Whitney *U* test with the function "wilcox.test" in R.

Identification of candidate mutations for positive selection

For the populations where caHMM does not finish, we instead performed well-mixed

hidden Markov chain (wmHMM) using the code "calculate_well_mixed_hmm_wrapper.py,"

which also infers the fixation status of a mutation under a one-clade model. The single clade in

wmHMM is defined as the basal clade. Candidate mutations for adaptation are then defined as

mutations that satisfy either of the following two criteria: (1) the mutations recognized as fixed

mutations in any of the basal, major, or minor clade in the output file from caHMM; (2) the

mutations that show DAF > 0.5 in at least two time points.

Calculation of G-scores

To quantify the parallelism of the set of candidate nonsynonymous mutations in a

resource-limitation cycle/genetic background combination, we calculated G-score for each

gene[35]. A larger G-score means more overrepresentation for a gene. To calculate G-scores, we

first counted the observed number of candidate nonsynonymous mutations in gene i per

resource-limitation cycle/genetic background combination (O_i), and the expected number for

gene i (E_i) was calculated by $O_{tot}(L_i/L_{tot})$, where $O_{tot} = \Sigma_i O_i$, L_i is the number of nonsynonymous

sites for gene i, and $L_{tot} = \Sigma_i L_i$. The G-score for gene i (G_i) was then calculated by $2O_i\ln(O_i/E_i)$

or defined as zero when $O_i = 0$. Note that $2O_i\ln(O_i/E_i)$ could be smaller than zero when $0 < O_i <$

E_i. This can lead to a paradox that G_i is smaller when a gene was hit by a few times compared to

the situation that the same gene was not hit at all. Therefore, we also defined $G_i = 0$ when

$2O_i\ln(O_i/E_i) < 0$ in the following analysis.

Because the null expectation of G-scores varies with total number of fixed

nonsynonymous mutations (**Supplemental Fig. 3b**), we performed 20,000 simulations in each of

which O_{tot} hits are randomly distributed among all L_{tot} sites across all genes in the genome. To

evaluate the significance of the sum of G-scores (**Fig. 4a**), we calculated the corresponding z

score by (the observed sum - mean of simulated sums) / (standard deviation of simulated sums).

Calculation of mean Bray-Curtis similarity

To quantify the parallelism of the set of candidate nonsynonymous mutations, we also

calculate the mean Bray-Curtis similarity across all pairs of experimental populations for a

resource-limitation cycle/genetic background combination. Specifically, for a pair of populations

j and k, their Bray-Curtis similarity is calculated by

$$1 - \frac{\Sigma_i|o_{ij}-o_{ik}|}{\Sigma_i(o_{ij}+o_{ik})},$$

where o_{ij} and o_{ik} is the observed number of candidate nonsynonymous mutations in gene i for

population j and k, respectively.

We also performed 1,000 simulations to acquire its null distribution for each resource-

limitation cycle/genetic background combination. In each simulation, we randomly sample the

nonsynonymous sites up to the number of observed candidate nonsynonymous mutations for

each population and calculated mean Bray-Curtis similarity as described above. After acquiring

the null distribution, to evaluate the significance of the mean Bray-Curtis similarity (**Fig. 4b**), we

calculated the corresponding z score by (the observed value - mean of simulated values) /

(standard deviation of simulated values).

Overrepresentation of the genes affected by nonsynonymous mutations

To evaluate the significance of G-score for gene i, we directly compared the G_i to the

distribution of 20,000 simulated G_i. The P-value was determined by the proportion of simulated

G_i larger or equal to the observed G_i. The multiple test correction was performed by multiply

each gene's P-value by the number of genes with at least one hit by the set of fixed

nonsynonymous mutations (Bonferroni correction). Only the genes with Bonferroni corrected P-

value < 0.05 are called significant.

Enrichment test of GO terms and KEGG pathways

Using this set of significant genes for each resource-limitation cycle/genetic background

combination, we performed enrichment tests of gene ontology terms and KEGG pathways using

the function "enrichGO" and "enrichKEGG", the organismal database org.EcK12.eg.db, and a q-

value cut-off $= 0.05$ in R package "DOSE"[56].

Overrepresentation of the genes affected by structural mutations

To test whether the set of genes that were affected by structural mutations (indels and IS-

element insertions) in a resource-limitation cycle/genetic background combination were

influenced by positive selection, we also calculated a *G*-score for each gene[35]. To calculate *G*-

scores, we first counted the observed number of populations with any structural mutations in

gene *i* per resource-limitation cycle/genetic background combination (O_i), and the expected

number for gene *i* (E_i) was calculated by $O_{tot}(L_i/L_{tot})$, where $O_{tot} = \Sigma_i O_i$, L_i is the gene length for

gene *i,* and $L_{tot} = \Sigma_i L_i$. The *G*-score for gene *i* (G_i) was then calculated by $2O_i \ln(O_i /E_i)$,

following the methods described in the above section.

We also performed 20,000 simulations and determined the Bonferroni corrected *P*-value

for each gene *i* following the methods described in the above section. As a result, we found all

the genes with $O_i \geq 2$ show Bonferroni corrected *P*-value < 0.05.

Data and Code Availability

Sequencing data generated during this study are available at NCBI's Sequencing Read Archive:

BioProject PRJNA532905 (Embargoed data accessible to reviewers: https://tinyurl.com/tsynzor).

Code generated to analyze sequencing data and generate figures is available at

https://github.com/LynchLab/ECEE_Starvation

Acknowledgements

We thank B. Choi, T. Doak, D.A. Drummond, P. Foster, W. Guo, J. T. Lennon, H. Long, J.

McKinlay, S. Snyder, E. Thorley, A. Urquidez, S. Vidrios, and S. Walls for their assistance and

helpful comments. High Performance Computing Resources were provided and maintained by

the National Center for Genome Analysis Support at Indiana University. This work was

supported by Army Research Office Grants ARO65308-LS-MU and W911NF-14-1-0411 and

National Institutes of Health Grant F32GM123703 and R35GM122566.

Author Contributions

Conceptualization, M.G.B., W-C.H., and M.L.; Methodology, M.G.B., W-C.H., and S.F.M.;

Investigation, M.G.B., W-C.H., S.F.M., G.F.B., and J.C.M.; Formal Analysis, M.G.B., and W-

C.H.; Writing – Original Draft, M.G.B., and W-C.H.; Writing – Review & Editing, M.G.B., W-

C.H., and M.L.; Visualization, M.G.B., and W-C.H.; Funding Acquisition, M.G.B. and M.L.

Declaration of Interests

The authors declare no competing interests.

 Figures and Legends

Figure 1 Phenotypic divergence of evolved lines after 900 days. Each resource-limitation

cycle/genetic background combination is represented by two populations; 1-day (orange) - WT:

P103, P108; MMR-: P106, P111; 10-day (blue) - WT: P403, P408; MMR-: P406, P411; 100-day

(green) - WT: P503, P508; MMR-: P506, P511. Significant differences in growth phenotypes

were determined by two-sided, pairwise Welch's t-tests adjusted by FDR and all error bars

represent ± s.e.m. **(a)** Growth curves of ancestor (black) and evolved populations cultured in LB

broth at 37 °C. Circles denote average absorbance at 600 nm measured every 15 minutes for 14

hours. Nested table displays the averages ± s.e.m. for maximum growth rate (r_{max}) and time to

maximum growth rate (T_{max}). T_{max} is presented in units of time (hours:minutes). Populations

evolving to 10-day resource-limitation cycles take significantly longer to reach T_{max} (WT: t_{anc} =

14.91, P_{anc} = 4.5 x 10^{-10}; $t_{1\text{-}day}$ = 11.65, $P_{1\text{-}day}$ = 4.6 x 10^{-10}; $t_{100\text{-}day}$ = 1.99, $P_{100\text{-}day}$ = 0.057; MMR-:

t_{anc}= 7.24, P_{anc} = 2.5 x 10^{-5}; $t_{1\text{-}day}$ = 7.81, $P_{1\text{-}day}$ = 8.4 x 10^{-6}; $t_{100\text{-}day}$ = 1.87, $P_{100\text{-}day}$ = 0.073;). **(b)**

Average colony sizes of evolved populations and ancestors after 24 h of growth on LB agar.

Populations evolving to 10-day resource-limitation cycles produce significantly smaller colonies

(WT: t_{anc} = 44.99, P_{anc}< 1 x10^{-16}; $t_{1\text{-}day}$ = 5.56, $P_{1\text{-}day}$ = 3.3 x 10^{-4}; $t_{100\text{-}day}$ = 4.85, $P_{100\text{-}day}$ = 5.6 x

10^{-4}; MMR-: t_{anc} = 16.10, P_{anc} = 5.4 x 10^{-10}; $t_{1\text{-}day}$ = 6.16, $P_{1\text{-}day}$ = 4.5 x 10^{-5}; $t_{100\text{-}day}$ = 2.22, $P_{100\text{-}day}$

= 0.039). Photos of **(c)** 1-day (P108), **(d)** 10-day (P408), and **(e)** 100-day (P508) populations co-

cultured with the WT ancestor on TA agar and incubated at 37°C for 24 h to show variation in

colony size. Colonies from evolved populations appear red while WT ancestor colonies appear

beige. Inset in **(d)** provides a magnified look at ancestor and 10-day colonies; the black arrow

points to a 10-day population colony, highlighting the evolved extreme reduction in colony size.

(f) Presence/absence of biofilm in evolved cultures after 24 h. Surface biofilm can be observed in

daily and 10-day populations, but is absent in 100-day populations and the ancestor. **(g)** Average

density of biofilm produced by evolved populations and ancestors in 96-well plate assay after 24

hours of growth, quantified via absorbance at 550 nm. Populations evolving to 10-day resource

limitation cycles produce significantly more biofilm (WT: t_{anc} = 10.63, P_{anc}= 7.0 x 10^{-10}; $t_{1\text{-}day}$ =

8.97, $P_{1\text{-}day}$ = 5.0 x 10^{-9}; $t_{100\text{-}day}$ = 10.25, $P_{100\text{-}day}$ = 7.0 x 10^{-10}; MMR-: t_{anc} = 15.37, P_{anc} = 1.6 x

10^{-15}; $t_{1\text{-}day}$ = 8.22, $P_{1\text{-}day}$ = 7.7 x 10^{-11}; $t_{100\text{-}day}$ = 14.21, $P_{100\text{-}day}$ = 7.5 x 10^{-15}).

Figure 2 Evolutionary dynamics of evolved populations. (a) Each panel shows mutation rates

of populations after 900 days of experimental evolution in different combinations of resource-

limitation cycle length (1, 10, or 100-days) and genetic background (MMR- for mismatch repair

defective or WT for wild-type). In each combination, four evolved replicate lines were tested,

and two or four clones per evolved line were isolated and measured. As mutation rates are

estimated via maximum-likelihood, the open circle and error bar represent the mean and the 95%

confidence interval for each clone. The grey dashed line represents the mutation-rate

measurement of the corresponding ancestor. The transparent coloured lines represent the mean

mutation-rate measurement of each combination. For both WT and MMR- backgrounds,

populations evolving to 10-day resource limitation cycles exhibit greater mutation rates than the

other treatments (WT: $P < 10^{-4}$; MMR-: 0.081; permutation test). **(b)** Each panel shows the mean

rate of genomic evolution among the parallel experimental populations for each resource-

limitation cycle length (1, 10, or 100 days) in MMR- (mismatch repair defective) or WT (wild-

type) genetic backgrounds. At each time point, the circle and the bar represent the mean and the

standard error. From left to right represent the sequencing profiles sampled at days 0, 90, 200,

300, 400, 500, 600, 700, 800, and 900. Note that the data for populations with 100-day resource-

limitation cycles are missing at day 500. The line represents the linear regression against the

number of generations. The estimated slope (b) and corresponding standard error are also shown

in each panel.

Fig.3

Figure 3 Ecological dynamics in experimental evolution. Muller plots of major (yellow) and

minor (pink) clade abundance for (**a**) Population 101 (1-day, WT) and Population 503 (100-day,

WT). Black arrows denote the maximum duration that two clades are observed to co-exist based

on the presence of associated SNPs identified with metagenomic sequencing. (**b**) The

distribution of durations of coexistence of major and minor clades, or subpopulation structure, as

determined by the caHMM among the parallel experimental populations for each resource-

limitation cycle/genetic background combination – 1-day (orange), 10-day (blue) and, 100-day

(green). The duration of coexistence for multiple clades in a population decreases as the duration

of resource limitation cycles increase (P = 6.1 x 10^{-6}, Scheirer-Ray-Hare test), with populations

evolving to 100-day resource limitation cycles exhibiting the least subpopulation structure (1-day

v. 100-day: P = 3.1 x 10^{-5}; 10-day v. 100-day: P = 4.2 x 10^{-5}, Mann-Whitney U test).

Fig.4

Figure 4 Evolutionary parallelism as evidence for positive selection. Within each resource-

limitation cycle/genetic background combination, the sum of G-scores was computed - a

goodness of fit test that can indicate the extent of parallel evolution and identify genes that are

likely selective targets. **(a)** The arrow and vertical dashed line show the observed sum of G-

scores for each treatment combination: 1-day (orange), 10-day (blue), and 100-day (green). The

histogram represents the null distribution of evolutionary parallelism as determined by 20,000

simulated sums of G-scores, where the position of mutations were randomized throughout the

genome. The significance of the observed sums can be evaluated by z-scores which quantify the

extent the observed value deviates from the null distribution ($z > 1.65$ for one-tailed $P < 0.05$). **(b)**

Similarly, the arrow and vertical dashed line show the observed mean of Bray–Curtis similarity

across the pairs of evolved populations evolved in a resource-limitation cycle/genetic

background combination. The histogram represents the null distribution of evolutionary

parallelism as determined by 1,000 simulated means of Bray–Curtis similarity, where the

position of mutations were randomized throughout the genome. The significance of the observed

sums can again be evaluated by z-scores ($z > 1.65$ for one-tailed $P < 0.05$). **(c)** List of genes with

at least one nonsynonymous mutation that significantly contribute to the increased sum of G-

scores and are likely under positive selection. Significance levels (simulated P-values with

Bonferroni correction) are shown by the different non-black colours of tiles. Genes with no such

hits in a particular resource-limitation cycle/genetic background combination are shown by black

tiles. **(d)** List of genes significantly overrepresented for structural mutations (IS-element

insertions and indels) that are likely under positive selection. Yellow tiles highlight genes

observed in at least two populations and yielding simulated P-values < 0.05 after Bonferroni

correction.

 Supplemental Information Titles and Legends

Supplemental Figure 1. Genomic diversity changing over time and probability of fixation.

Genomic diversity was measured by **(a)** number of SNPs per population, **(b)** nucleotide diversity,

(c) synonymous nucleotide diversity, and **(d)** number of SNPs per clone, respectively. In **(a)**-**(c)**,

the y-axis shows mean estimates among the parallel populations in a combination of one

resource-limitation cycle (1 day, 10 days, or 100 days) and one genetic background (MMR- for

mismatch repair defected, or WT for wild-type). Across the x-axis from left to right, circles

represent the sequencing profiles sampled at day 90, 200, 300, 400, 500, 600, 700, 800, and 900

(the data of populations with resource-limitation cycle = 100days at day 500 are missing). At

each time point, the circle and the bar represent the mean +/- standard error of mean. The dashed

lines represent the linear regression lines of number of SNPs against guaranteed generations, and

the estimated slopes (b) and corresponding standard errors are also shown. In (**d**), the results of

linear and nonlinear regression are compared. At each time point, a circle represents the number

of SNPs per clone in a parallel population. The solid line represents the result of linear regression

($y \sim 0 + x$); The dotted line represents the result of nonlinear regression ($y \sim 0 + x + x^{0.5}$).

Model comparison were performed by nested ANOVA: $P = 0.028$, 0.69, and 0.23 for resource-

limitation cycle = 1, 10, and 100, respectively in MMR- background; $P = 0.40$, 0.063, and 0.40

for resource-limitation cycle = 1, 10, and 100, respectively in WT background. In (**e**), similar

probabilities of synonymous, nonsynonymous, and intergenic mutations fixed in any clade were

found in populations with longer resource-limitation cycles. The dots and error bars represent

mean probability and its 95% confidence interval (estimated by 1.96 times of standard errors)

across parallel experimental populations. The fixed probably here is conditioning on the

detectability of mutations. In (**f**)-(**g**), populations with longer resource-limitation cycles do not

show significant higher fixation rates of nonsynonymous mutations and intergenic mutations

compared to the fixation rates of synonymous mutations. If positive selection is strong, one

would expect to see that mutations with a greater potential to change protein functionality or

levels (e.g. nonsynonymous mutations and intergenic mutations) are more likely to be fixed

within a clade than mutations with lesser potential (e.g. synonymous mutations). Therefore,

higher ratios of the number of nonsynonymous mutations fixed in any clade (F_N) to the number

of synonymous mutations fixed in any clade (F_S), or higher ratios of the number of intergenic

mutations fixed in any clade (F_I) to F_S, is expected. However, the above expectation was not

observed. In **(f)**, the value represented by a cross is calculated by the summed F_N and F_S of all

parallel experimental populations with a specific resource-limitation cycle length and genetic

background. The value represented by a dot is calculated by the F_N and F_S in each parallel

experimental population, and the dots with $F_S = 0$ are not shown. The dashed lines show the ratio

of the number of nonsynonymous mutations to the number of synonymous mutations detected in

mutation accumulation experiments. In **(g)**, the value represented by a cross is calculated by the

summed F_I and F_S of all parallel experimental populations with a specific resource-limitation

cycle length and genetic background. The value represented by a dot is calculated by the F_I and

F_S in each parallel experimental population, and the dots with $F_S = 0$ are not shown. The dashed

lines show the ratio of the number of intergenic mutations to the number of synonymous

mutations detected in mutation accumulation experiments. Therefore, according to **(e)**-**(g)**, no

evidence for significantly different levels of positive selection across resource-limitation cycles

was found.

Supplemental Figure 2. Additional evidence for less population substructure in 100-day

populations (a) Negative correlation between the proportion of fixation occurring in basal clades

and the longest length of coexistence across all experimental populations. Each dot is an

experimental population. ρ: Spearman's rank correlation coefficient. P: p-value of Spearman's

rank correlation. **(b)** The distribution of proportion of fixation occurred in basal clades for each

resource-limitation cycle/genetic background combination. We found resource-limitation cycle

length significantly affects the proportion of fixation occurred in basal clades ($P = 2.5 \times 10^{-4}$,

Scheirer-Ray-Hare test). The difference is significant between 1-day and 100-day populations (P

680     $= 5.5 \times 10^{-5}$, Mann-Whitney U test), 10-day and 100-day populations ($P = 0.016$, Mann-Whitney

681   U test), and daily and 10-day populations ($P = 1.6 \times 10^{-3}$, Mann-Whitney U test). **(c)** Temporal

682   distribution of mean derive allele frequencies (DAF) in evolved populations. Each dot shows a

683   mean fraction of genome showing a range of DAF with bin size 0.1 in different time point of

684   experimental evolution. The different time points of experimental evolution are shown by colors.

a

b

Supplemental Figure 3. Observed and expected parallelism in experimental evolution. (a)

shows the nucleotide-level parallelism observed in our experimental evolution. For each

combination of a resource-limitation cycle (1 day, 10 days, or 100days) and a genetic

background (MMR- for mismatch repair defected, or WT for wild-type), we plotted the

distribution of number of parallel experiments where the same fixed nonsynonymous SNPs are

identified. Due to a very low probability that any one nucleotide is mutated at least twice ($< 10^{-4}$

by simulation with a random sample of 10^4 mutations), our observed nucleotide-level parallelism

696 again suggests positive selection in action. **(b)** illustrates that the null expectation of G-scores

697 across genes vary with total number of mutations considered. The dashed line and dotted lines

698 summarize the mean and 95% range across 1000 simulations of hypothetical number of fixed

699 nonsynonymous mutations = 1, 50, 100, 500, 1000, 1500, 2000, 2500, 3000, 4000, 5000, 6000,

700 7000, 8000, 9000, 10000.

Supplemental Figure 4. Density measurement of experimental populations during

experimental evolution. Pre-transfer population density of 1- (orange), 10- (blue) and 100-day

(green) populations as determined by CFU counts. Bars represent mean population density of

WT and MMR- populations after 200, 300, 400, 500, 600, 700, 800, and 900 days of evolution

and error bars represent $\pm$s.e.m.

Supplemental Table 1. List of sequencing profiles used in the analysis

Supplemental Table 2. Molecular functions of genes overrepresented for fixed nonsynonymous

SNPs in each combination of resource-limitation cycle length and genetic background.

Supplemental Table 3. Molecular functions of genes overrepresented for fixed structural

mutations in each resource-limitation cycle length and genetic background.

Supplemental Table 4. List of candidate mutations found in genes which may alter mutation

rates

Supplemental Table 5. List of significantly enriched Gene Ontology terms and KEGG

pathways of genes with significant G-scores in MMR- populations with 10-day cycles but not in

populations with daily cycles.

Supplemental Table 6. List of significantly enriched Gene Ontology terms and KEGG

pathways of genes with significant G-scores in MMR- populations with 100-day cycles but not

in populations with daily cycles.

Supplemental Table 7. List of significantly enriched Gene Ontology terms and KEGG

pathways of genes with significant G-scores in WT populations with 10-day cycles but not in

populations with daily cycles.

Supplemental Table 8. List of significantly enriched Gene Ontology terms and KEGG

pathways of genes with significant G-scores in WT populations with 100-day cycles but not in

populations with daily cycles.

Supplemental Table 9. List of fixed nonsynonymous SNPs which hit identical nucleotides in

multiple experimental populations for each combination of resource-limitation cycle and genetic

background.

References

1   Lassig, M., Mustonen, V. & Walczak, A. M. Predicting evolution. *Nat Ecol Evol* **1**, 77

2   Jain, K. & Krug, J. Deterministic and stochastic regimes of asexual evolution on rugged
fitness landscapes. *Genetics* **175**, 1275-1288

3   Rouzine, I. M., Rodrigo, A. & Coffin, J. M. Transition between stochastic evolution and
deterministic evolution in the presence of selection: General theory and application to
virology. *Microbiol Mol Biol R* **65**, 151-185

4   Kimura, M. Stochastic processes and distribution of gene frequencies under natural
selection. *Cold Spring Harb Sym* **20**, 33-53

5   Fisher, R. A. On the dominance ratio. *B Math Biol* **52**, 297-318

6   Nosil, P. *et al.* Natural selection and the predictability of evolution in Timema stick
insects. *Science* **359**, 765-770

7   Chaturvedi, S. *et al.* The predictability of genomic changes underlying a recent host shift
in Melissa blue butterflies. *Mol Ecol* **27**, 2651-2666

8   Blount, Z. D., Lenski, R. E. & Losos, J. B. Contingency and determinism in evolution:
Replaying life's tape. *Science* **362**, eaam5979

9   Rosenblum, E. B., Parent, C. E. & Brandt, E. E. The molecular basis of phenotypic
convergence. *Annu. Rev. Ecol. Evol. Syst* **45**, 203-226

10   Parker, J. *et al.* Genome-wide signatures of convergent evolution in echolocating
mammals. *Nature* **502**, 228-231

11   Stern, D. L. The genetic causes of convergent evolution. *Nat Rev Genet* **14**, 751-764

12   Losos, J. B. Convergence, adaptation, and constraint. *Evolution* **65**, 1827-1840

13   Bailey, S. F., Rodrigue, N. & Kassen, R. The effect of selection environment on the
probability of parallel evolution. *Molecular biology and evolution* **32**, 1436-1448

14   Deatherage, D. E., Kepner, J. L., Bennett, A. F., Lenski, R. E. & Barrick, J. E. Specificity
of genome evolution in experimental populations of Escherichia coli evolved at different
temperatures. *Proceedings of the National Academy of Sciences* **114**, E1904-E1912

15   Turner, C. B., Marshall, C. W. & Cooper, V. S. Parallel genetic adaptation across
environments differing in mode of growth or resource availability. *Evolution letters* **2**,
355-367

16   Barton, I. S., Fuqua, C. & Platt, T. G. Ecological and evolutionary dynamics of a model
facultative pathogen: *Agrobacterium* and crown gall disease of plants. *Environmental*
Microbiology **20**, 16-29

17   Brown, S. P., Cornforth, D. M. & Mideo, N. Evolution of virulence in opportunistic
pathogens: generalism, plasticity, and control. *Trends in Microbiology* **20**, 336-342

18   Johnson, J. R. & Russo, T. A. Extraintestinal pathogenic Escherichia coli: "The other bad
E coli". *Journal of Laboratory and Clinical Medicine* **139**, 155-162

19   Smith, J. L., Fratamico, P. M. & Gunther, N. W. Extraintestinal pathogenic Escherichia
coli. *Foodborne Pathogens and Disease* **4**, 134-163

20   Jang, J. *et al.* Environmental *Escherichia coli*: ecology and public health implications-a
review. *J Appl Microbiol* **123**, 570-581

21   Farrell, M. J. & Finkel, S. E. The growth advantage in stationary-phase phenotype
conferred by *rpoS* mutations is dependent on the pH and nutrient environment. *J*
Bacteriol **185**, 7044-7052

783	22	Zinser, E. R. & Kolter, R. Prolonged stationary-phase incubation selects for *lrp* mutations
784		in *Escherichia coli* K-12. *J Bacteriol* **182**, 4361-4365
785	23	Zambrano, M. M. & Kolter, R. GASPing for life in stationary phase. *Cell* **86**, 181-184
786	24	Finkel, S. E. & Kolter, R. Evolution of microbial diversity during prolonged starvation. *P*
787		*Natl Acad Sci USA* **96**, 4023-4027
788	25	Finkel, S. E. Long-term survival during stationary phase: evolution and the GASP
789		phenotype. *Nature Reviews Microbiology* **4**, 113-120
790	26	Sniegowski, P. Evolution: bacterial mutation in stationary phase. *Curr Biol* **14**, R245-
791		R246
792	27	Loewe, L., Textor, V. & Scherer, S. High deleterious genomic mutation rate in stationary
793		phase of *Escherichia coli*. *Science* **302**, 1558-1560
794	28	Ying, B. W. *et al.* Evolutionary consequence of a trade-off between growth and
795		maintenance along with ribosomal damages. *Plos One* **10**, e0135639
796	29	Lee, H., Popodi, E., Tang, H. X. & Foster, P. L. Rate and molecular spectrum of
797		spontaneous mutations in the bacterium *Escherichia coli* as determined by whole-genome
798		sequencing. *Proc Natl Acad. Sci. USA* **109**, E2774-E2783
799	30	Behringer, M. G. *et al. Escherichia coli* cultures maintain stable subpopulation structure
800		during long-term evolution. *Proceedings of the National Academy of Sciences of the*
801		*United States of America* **115**, E4642-E4650
802	31	Foster, P. L. Methods for determining spontaneous mutation rates. *Methods Enzymol* **409**,
803		195-213
804	32	Zhang, J. Neutral theory and phenotypic evolution. *Molecular biology and evolution* **35**,
805		1327-1331
806	33	Good, B. H., McDonald, M. J., Barrick, J. E., Lenski, R. E. & Desai, M. M. The
807		dynamics of molecular evolution over 60,000 generations. *Nature* **551**, 45-+
808	34	Gillespie, J. H. Genetic drift in an infinite population. The pseudohitchhiking model.
809		*Genetics* **155**, 909-919
810	35	Tenaillon, O. *et al.* Tempo and mode of genome evolution in a 50,000-generation
811		experiment. *Nature* **536**, 165-170
812	36	Rhodius, V. A. & Busby, S. J. Transcription activation by the Escherichia coli cyclic
813		AMP receptor protein: determinants within activating region 3. *Journal of molecular*
814		*biology* **299**, 295-310
815	37	Niu, W., Kim, Y., Tau, G., Heyduk, T. & Ebright, R. H. Transcription activation at class
816		II CAP-dependent promoters: two interactions between CAP and RNA polymerase. *Cell*
817		**87**, 1123-1134
818	38	Canino-Koning, R., Wiser, M. J. & Ofria, C. Fluctuating environments select for short-
819		term phenotypic variation leading to long-term exploration. *PLoS computational biology*
820		**15**
821	39	Melnyk, A. H., McCloskey, N., Hinz, A. J., Dettman, J. & Kassen, R. Evolution of cost-
822		free resistance under fluctuating drug selection in Pseudomonas aeruginosa. *mSphere* **2**,
823		e00158-00117
824	40	Williams, A. B. & Foster, P. L. Stress-Induced Mutagenesis. *EcoSal Plus* **5**
825	41	Lukacisinova, M., Novak, S. & Paixao, T. Stress-induced mutagenesis: Stress diversity
826		facilitates the persistence of mutator genes. *PLoS Comput Biol* **13**, e1005609
827	42	Ishii, K., Matsuda, H., Iwasa, Y. & Sasaki, A. Evolutionarily stable mutation rate in a
828		periodically changing environment. *Genetics* **121**, 163-174

829	43	Tanaka, M. M., Bergstrom, C. T. & Levin, B. R. The evolution of mutator genes in bacterial populations: the roles of environmental change and timing. *Genetics* **164**, 843-854
832	44	Travis, J. & Travis, E. Mutator dynamics in fluctuating environments. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **269**, 591-597
834	45	Hunt, S. M., Werner, E. M., Huang, B. C., Hamilton, M. A. & Stewart, P. S. Hypothesis for the role of nutrient starvation in biofilm detachment. *Appl Environ Microb* **70**, 7418-7425
837	46	Garcia-Jimenez, C. & Goding, C. R. Starvation and pseudo-starvation as drivers of cancer metastasis through translation reprogramming. *Cell Metab* **29**, 254-267
839	47	de Visser, J. A. G. & Rozen, D. E. Limits to adaptation in asexual populations. *Journal of evolutionary biology* **18**, 779-788
841	48	Lange, R. & Hengge-Aronis, R. The nlpD gene is located in an operon with rpoS on the Escherichia coli chromosome and encodes a novel lipoprotein with a potential function in cell wall formation. *Mol Microbiol* **13**, 733-743
844	49	Merritt, J. H., Kadouri, D. E. & O'Toole, G. A. Growing and analyzing static biofilms. *Current protocols in microbiology* **22**, 1B. 1.1-1B. 1.18
846	50	Baym, M. *et al.* Inexpensive multiplexed library preparation for megabase-sized genomes. *PLoS One* **10**, e0128036
848	51	Martin, M. Cutadapt Removes Adapter Sequences From High-Throughput Sequencing Reads. *EMBnet.journal* **17**, 10-12
850	52	Deatherage, D. E. & Barrick, J. E. Identification of mutations in laboratory-evolved microbes from next-generation sequencing data using breseq. *Methods Mol Biol* **1151**, 165-188
853	53	McCloskey, D. *et al.* Evolution of gene knockout strains of *E. coli* reveal regulatory architectures governed by metabolism. *Nat Commun* **9**, 3796
855	54	Zheng, Q. rSalvador: An R package for the fluctuation experiment. *G3-Genes Genom Genet* **7**, 3849-3856
857	55	Mangiafico, S. S. *Summary and analysis of extension program evaluation in R.* (2016).
858	56	Yu, G. C., Wang, L. G., Yan, G. R. & He, Q. Y. DOSE: an R/Bioconductor package for disease ontology semantic and enrichment analysis. *Bioinformatics* **31**, 608-609

Fig.1

Fig.2

Fig.3

Fig.4