An updated antennal lobe atlas for the yellow fever mosquito *Aedes aegypti*

Shruti Shankar [1] and Conor J. McMeniman [1, 2, *]

[1] W. Harry Feinstone Department of Molecular Microbiology and Immunology, Johns Hopkins Malaria Research Institute, Johns Hopkins Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD 21205, USA. [2] The Solomon H. Snyder Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, MD 21205, USA.

*Corresponding author: cmcmeni1@jhu.edu

Abstract

The yellow fever mosquito *Aedes aegypti* is a prolific vector of arboviral and filarial diseases that largely relies on its sense of smell to find humans. To facilitate in-depth analysis of the neural circuitry underlying *Ae. aegypti* olfactory-driven behaviors, we generated an updated *in vitro* atlas for the antennal lobe olfactory brain region of this disease vector using two independent neuronal staining methods. We performed morphological reconstructions with replicate fixed, dissected and stained brain samples from adult male and female *Ae. aegypti* of the LVPib12 genome reference strain and determined that the antennal lobe in both sexes is comprised of approximately 80 discrete glomeruli. Guided by landmark features in the antennal lobe, we found 63 of these glomeruli are stereotypically located in spatially invariant positions within these *in vitro* preparations. A posteriorly positioned, mediodorsal glomerulus denoted MD1 was identified as the largest spatially invariant glomerulus in the antennal lobe. Spatial organization of glomeruli in a recently field-derived strain of *Ae. aegypti* from Puerto Rico was conserved, despite differences in antennal lobe shape relative to the inbred LVPib12 strain. This model *in vitro* atlas will serve as a useful community guide and resource to improve antennal lobe annotation and anatomically map projection patterns of neurons expressing target genes in this olfactory center. It will also facilitate the development of chemotopic maps of odor representation in the mosquito antennal lobe to decode the molecular and cellular basis of *Ae. aegypti* attraction to human scent and other chemosensory cues.

Author Summary

The olfactory system of the yellow fever mosquito *Aedes aegypti* is highly tuned for the detection of human odorants, as well as other chemical cues influencing host and food-search behavior, egg-laying and mating. To provide insights into the neuroanatomical organization of the olfactory system of this globally important disease vector, we have generated an updated *in vitro* atlas for the primary smell processing center of the *Ae. aegypti* brain, called the antennal lobe. These new guide maps facilitate systematic interrogation of antennal lobe morphology and naming of associated substructures in dissected brain samples of this species labeled with two common neural staining methods. We report that landmark features of the *Ae. aegypti* antennal lobe morphology and spatial organization appear conserved between mosquito sexes and across geographically divergent strains of this mosquito species. An improved understanding of *Ae. aegypti* antennal lobe neuroanatomy and how attractive or repellent odorant stimuli are encoded in this brain center has the potential to rapidly accelerate reverse engineering of synthetic chemical blends that effectively lure, confuse or repel this major disease vector.

Introduction

Aedes aegypti is a prolific vector of yellow fever, dengue, chikungunya, Zika and filariasis in tropical and subtropical regions around the world. This highly anthropophilic mosquito species uses its sense of smell to track volatile chemicals present in human skin odor and breath and orientate towards us. For example, carbon dioxide (1) and other volatile constituents of human body odor, including lactic acid (2), trigger behavioral activation and steer directional flight of this mosquito species towards humans to acquire a blood (3). Olfaction also influences other critical aspects of *Ae. aegypti* biology including oviposition (4), mating (5) and nectar-seeking (6). Molecular and cellular substrates underlying the *Ae. aegypti* sense of smell therefore represent key targets for the control of mosquito-borne diseases transmitted by this cosmopolitan and urbanized disease vector.

In the *Ae. aegypti* adult, olfactory sensory neurons (OSNs) expressing chemoreceptors that detect odorants are primarily localized to peripheral olfactory organs (1, 7-10) including the antennae, maxillary palps and proboscis. Dendritic sensory nerve endings of OSNs are found in various morphological classes of specialized porous sensilla distributed along the length of these organs. Each *Ae. aegypti* sensillum typically houses 2-3 OSNs (7-9). The axonal processes of these

neurons project centrally to the mosquito brain, to the first center of olfactory information

processing called the antennal lobes. The antennal lobes are made up of clusters or foci of

synaptic connectivity called glomeruli, where olfactory information is transmitted from OSNs to

second order neurons (11-13).

In the vinegar fly *Drosophila melanogaster*, an individual glomerulus is the point at which the

axonal projections of OSNs typically expressing a unique complement of ligand-binding

chemoreceptors converge (13, 14) and synapse with the dendritic arbors of projection neurons

(PNs) and local interneurons (LNs) (15). Each glomerulus is therefore tuned to a unique subset

of chemicals. Anterograde dye-filling of peripheral OSNs that respond functionally to the same

subset of odors and subsequent tracing of their glomerular projections indicates this pattern is

likely to be conserved in mosquitoes (12, 16, 17).

An initial morphological study of the *Ae. aegypti* antennal lobe by Bausenwein and Nick (18)

denoted the presence of approximately 35 antennal lobe glomeruli in a single *Ae. aegypti* female

examined. However, it was noted that due to the staining methods employed in this study that

subdivision of antennal lobe glomerular neuropil in some areas was unclear. Subsequently, Ignell

and coauthors (11) estimated that the antennal lobes of this species were comprised of between

49-50 glomeruli using improved immunohistochemical techniques.

A significant volume of the *Ae. aegypti* antennal lobes was previously ascribed a non-olfactory

mechanosensory function by Ignell (11). This large core of neuropil on the anterior surface of the

antennal lobe was annotated as the Johnston's Organ Centre (JOC) based on silver nitrate Golgi

staining. This analysis revealed the putative sensory innervation of this antennal lobe region by

afferent fibers from the Johnston's Organ (JO), an observation mirrored shortly thereafter by the

same authors in *Anopheles gambiae* (11, 19). However, a recent reassessment of antennal lobe

morphology in *Anopheles coluzzii* failed to observe the JOC in the antennal lobes of this species,

instead identifying discrete glomeruli in this spatial region (19).

To assist basic and applied studies of *Ae. aegypti* olfaction, we have used improved neuropil

labeling and confocal imaging techniques to re-examine the neuroanatomy of the *Ae. aegypti*

antennal lobe. We present three-dimensional (3D) models based on manual segmentation and

reconstructions of antennal lobe glomeruli from replicate male and female brains from the

LVPib12 *Ae. aegypti* genome reference strain. To facilitate spatial orientation and stereotypical

identification of glomeruli within the antennal lobe, we also present detailed two-dimensional (2D)
maps wherein individual glomeruli are named based on their spatial arrangement in this brain
region using nc82- and phalloidin-based neuronal staining methods. Finally, we examine antennal
lobe neuroanatomy and organization in a recently field-collected strain of *Ae. aegypti* from the
municipality of Patillas, Puerto Rico. Altogether, this work describes a standardized framework for
glomerular nomenclature in the *Ae. aegypti* antennal lobe. This atlas will serve as a useful
reference for neurogenetic and functional imaging studies of the antennal lobe in this
anthropophilic disease vector.
111
112   **Results**
113
Three-dimensional reconstruction of the antennal lobes of male and female *Aedes aegypti*
mosquitoes
To define boundaries of the antennal lobe of the LVPib12 genome reference strain of *Ae. aegypti*
(20), we initially stained whole dissected and fixed mosquito brains using the monoclonal antibody
nc82 as a synaptic marker and acquired 1 μm confocal sections of the central brain starting from
the surface of the antennal lobes through the central complex. Similar to other dipteran insects
such as *Drosophila melanogaster* (21), the deutocerebrum (11) in *Ae. aegypti* is composed of two
paired neuropils: the antennal lobe olfactory center and the periesophageal neuropils which
include the antennal mechanosensory and motor center (AMMC).
123
The left antennal lobe (LAL) and the right antennal lobe (RAL) flank the esophageal foramen and
are positioned dorsal to the AMMC (Figures 1a and 1b). The antennal lobes are innervated by
the antennal (flagellar) (8) nerves that collectively include the axons of OSNs originating from the
third antennal segment (11, 16). In addition, maxillary afferents from the distal segment of the
maxillary palp terminate in the antennal lobes (16), supporting its role as the first center in the
central mosquito brain to receive signals from the peripheral olfactory organs.
130
Confocal sections of the LVPib12 antennal lobe revealed that its constituent glomeruli varied both
in terms of morphology and volume, and are arranged at different depths relative to i) the anterior
to posterior axis of the mosquito body and ii) along the dorsal to ventral neuraxis (21). To re-
ascertain glomerular counts relative to previously published studies, we initially reconstructed the
left and right antennal lobes from 5 male and 5 female LVPib12 brains that had been stained with
nc82 antibodies.

We used the software program Amira to perform segmentation of the antennal lobe by manually

painting over all the voxels corresponding to neuropil representing individual glomeruli, which

were identified based on their size, shape, spatial position and depth. We found that each

antennal lobe in all LVPib12 brains examined was comprised entirely of discrete demarcated

glomeruli and we were unable to identify the wedge shaped, JOC non-glomerular core which was

described previously (11). The number of glomeruli were found to vary slightly across our

reconstructions from 80 ± 2 in the female antennal lobe to 77 ± 2 glomeruli in the male antennal

lobe (mean ± s.d.) (Figures 1g and 1h).

Spatial arrangement of glomeruli in the *Aedes aegypti* antennal lobe

We further sought to understand the pattern of glomerular arrangement within the LVPib12

antennal lobe. Based on spatial position along the anterior-posterior (A-P), ventral-dorsal (V-D)

and medial-lateral (M-L) axes, we initially aimed to categorize glomeruli in the male and female

antennal lobe into spatial groups, with reference to a previous antennal lobe model (11) that

annotated 13 spatial groups of glomeruli in the *Ae. aegypti* antennal lobe. However, as we

identified approximately 30 additional glomeruli in our reconstructions relative to this extant map,

we first re-annotated spatial descriptors for each group in our atlas to better reflect their spatial

positions along these coordinate axes of the antennal lobe (Table S1). These revised groupings

organize the *Ae. aegypti* antennal lobe into 13 new glomerular clusters that aid in identifying and

naming all glomeruli in each lobe. Each of these groups was assigned a specific color in our

reconstructed 3D antennal lobe models (Figure 1c to f).

Viewing the antennal lobe along the A-P axis, on the anterior slices of the antennal lobe we

identified the Antero-Dorsal (AD) group (previously denoted Ventral (11)) surrounded by the

Antero-Lateral (AL) group, laterally; the Antero-Medial (AM) and Antero-Central (AC) groups,

medially; and the Ventro-Central (VC) group, ventrally. The groups moving towards the posterior

sections of the antennal lobe include the Postero-Medial (PM) group, Postero-Central (PC)

(consolidated with the previously denoted Medio-Central group (11)) and the deeper Medio-

Dorsal (MD) groups, along the medial axis; Postero-Lateral (PL) group located on the lateral side

(consolidated with the previously denoted Latero-Central group (11)); Centro-Dorsal (CD) group

located centrally; and Postero-Ventral (PV) group located ventrally. In the most posterior slices,

the dorsal group is named Dorsal (D) (previously denoted Antero-Dorsal (11)); and the ventrally

located Ventral (V) (previously denoted Postero-Dorsal (11)) group.

A two-dimensional neuroanatomical map of the *Aedes aegypti* antennal lobe

Based on each reconstructed three-dimensional LVPib12 antennal lobe model, we generated two-dimensional maps, showing all manually segmented glomeruli in each confocal slice as observed along the anterior to posterior axis of *in vitro* antennal lobe imaging. As a visual guide, we first selected eight glomeruli with salient features such as a large volume, shape and consistent position across all reconstructions to serve as landmarks for identifying the positions of other glomerular groups. These eight landmark glomeruli are shown in the 3D models (Figure 1c to 1f) and include AD1 and VC6 in anterior antennal lobe slices and D5, PL9, CD4, V1, V8 and MD1 in the more posterior antennal lobe. We did not select landmark glomeruli from the PC, PM, AM and AC groups, as the boundaries of these peripheral glomeruli cannot always be reliably demarcated by the *in vitro* staining methods used here.

To facilitate stereotypical identification of glomeruli based on their relative position to the eight landmark glomeruli, we generated a reference key detailing the typical spatial arrangement of glomeruli in the *Ae. aegypti* antennal lobe (Table 1). In each reconstructed antennal lobe, we assigned a name to each glomerulus based on its spatial group and depth. Glomeruli within the spatial groups are typically arrayed in (i) rows and are numbered based on depth along the medial lateral axis, (ii) in a vertical line (column) or (iii) in a circular manner and numbered along the dorsal-ventral axis. Glomeruli within some spatial groups may also appear to be randomly arrayed. Useful reference arrays of glomeruli within and between spatial groups that assist with identification and naming of glomeruli based on their spatial position relative to landmark glomeruli are detailed in Table 1 and overlaid upon an example antennal lobe reconstruction in Figure S1.

Using our reference key, we classified antennal lobe glomeruli as spatially 'invariant' or 'variant' based on the frequency of identification of individual glomeruli across all reconstructions. Invariant glomeruli were defined as glomeruli with the same assigned name found in the same spatial position in the left and right lobes of both sexes, at a frequency of $\geq$80%. Variant glomeruli were classified as glomeruli with the same assigned name found with a frequency of <80% across reconstructions, according to these criteria. The number of spatially invariant and variant glomeruli found in each of the 13 spatial groups are described as follows;

The anterior surface of the antennal lobe is comprised of five groups of glomeruli:

1. **Antero-Dorsal group**. In anterior slices of the antennal lobe, AD1 is a landmark glomerulus, distinguishable by its large size and dorsal position. It is located towards the center of the antennal lobe. AD1 is the first antennal lobe glomerulus visualized when stepping down a z-series along the A-P axis, and the typical depth of this glomerulus from its anterior surface (denoted 0 µm) to posterior surface is 15 µm. AM1-3, AC1, VC1, AL1-3 glomeruli routinely cluster around the AD group. AD2 and AD3 are found ventral and towards the medial and lateral side of AD1, respectively. AD2 and AD3 glomeruli are typically found in the space between the AD1 and the VC group and are characteristic satellite glomeruli flanking the AD1 landmark. Three residual variant glomeruli (AD 4-6) are sometimes observed in more dorsal and posterior slices relative to the AD invariant glomeruli.

2. **Ventro-Central group**: The VC group of glomeruli are found ventral to the AD group in the anterior most slices of the antennal lobe. Previously, a large part of the central antennal lobe was considered to comprise of the Johnston's Organ Center (JOC). We note that in our reconstructions, this region consists entirely of discrete glomeruli that we have reclassified as the VC and PV spatial groups. The VC group is comprised of an anterior surface layer consisting of the VC1, VC2 and VC3 glomeruli arranged in a row ordered along the medial to lateral axis, or alternatively in a triangle with VC1 positioned dorsal to VC2 and VC3; and the VC4, VC5 and VC6 glomeruli in a more posterior bottom row ordered along the medial to lateral axis. The most posterior layer of this group comprises VC7, VC8 and VC9 that are arranged centrally in this region of the antennal lobe (Table 1). VC1 is a large glomerulus positioned ventral to the landmark glomerulus AD1. VC6 is a large landmark glomerulus located on the lateral side of the antennal lobe and ventral to VC3. Two variant glomeruli (VC10-VC11) in a ventral to dorsal column are sometimes observed more posterior to the VC7-9 cluster.

3. **Antero-Lateral group**: The AL group is observed in anterior slices within the dorsal and lateral region of the antennal lobe. The AL1 glomerulus is located laterally, relative to the landmark AD1. AL1, AL2, AL3 glomeruli are positioned in a D-V column on the lateral side of the antennal lobe in order of increasing depth along the A-P axis. For example, AL1 is found dorsal and anterior to AL2. In some reconstructions, three variant glomeruli were sometimes found ventral and posterior to AL3 and were named AL4-AL6.

4. **Antero-Medial group**: The AM group is located medial to the AD group. It is comprised
of the AM1, AM2 and AM3 glomeruli arranged in a column along the D-V axis. AM1 is a
large glomerulus located above AD1 towards the medial axis of the antennal lobe. A
variant glomerulus AM4 was observed ventral to AM3 in some reconstructions.

5. **Antero-Central group**: The AC group is located medial to the AM group bisecting the
equator of the antennal lobe in a D-V column flanking the medial side. In more posterior
slices, the boundaries of some the AC glomeruli were difficult to demarcate as this spatial
group is sandwiched between the AM and the more posterior PM spatial groups. AC1 and
3 were spatially invariant glomeruli, whereas variant glomeruli AC2, AC4 and AC5 were
observed in some reconstructions.

In mid-sections of the antennal lobe, we describe the following groups:

6. **Postero-Medial group**: This group is located posterior to the AM glomeruli. It is comprised
of glomeruli along a D-V column on the medial side of the antennal lobe. The boundaries
of PM1, PM2, and PM3 are noted for their small size, and position between the anterior
AC and posterior D groups. PM4 is positioned more ventrally and is larger in volume
relative to PM1-PM3. In more posterior slices, the ventral PM5 and PM6 are classified as
variant glomeruli.

7. **Postero-Central group**: Included are a small number of glomeruli typically located in
between the PM and CD groups. PC1, PC2, PC3 and PC4 glomeruli are arranged in a D-
V column. PC1 is found posterior to AC1. In the anterior to posterior direction, AC1-PC1-
PM1 glomeruli form an array. A variant glomerulus PC5 can be observed at times ventral
to PC4 bordering the ventral group.

8. **Postero-Ventral group**: The PV group is visible from the anterior to central slices of the
antennal lobe. PV glomeruli are found ventral to the VC group. The 5 glomeruli that
comprise this group are arranged in two rows ordered along the medial-lateral axis. The
ventral row consists of PV1, PV2 and PV3 glomeruli. PV1 flanks the medial antennal lobe
margin, with PV2 in the center and PV3 laterally. The dorsal row consists of PV4 and the
more posterior PV5. In more posterior slices, variant glomeruli PV6-PV9 may be present.

This group of glomeruli were previously considered to be part of the JOC (11), along with
the VC spatial group.

The posterior region (-15 μm to -35 μm) of the antennal lobe is comprised of 5 groups of glomeruli.
The CD group is located centrally within posterior slices, and the D and V groups are located
dorsal and ventral to the CD group respectively. The PL and the MD groups flank the CD group
along the lateral and the medial margins of the antennal lobe.

9. **Ventral group**: The V group is found posterior and dorsal to the PV group, in the deepest
layers of the antennal lobe. The landmark glomerulus V1 is found ventral and posterior to
PV1. V1 flanks the medial margin of the antennal lobe. In the medial to lateral direction
the other glomeruli of this group are arranged in three layers. The first layer consists of
V1, V2, V3, and V4 aligned in a medial-lateral row. The second more posterior layer
consists of V5, V6, V7 and V8 are arranged in an approximate lateral-medial row. V8,
which is posterior and medial relative to V7, is a landmark glomerulus that is easy to
identify because of its lateral adjacent position to MD1. V8 is found posterior to V1. In
some reconstructions we identified more posterior variant V glomeruli (V9-V11).

10. **Centro-Dorsal group**: The CD group is the central group in dorsal regions of the
posterior-most slices of the antennal lobe. CD1, CD2, CD3, CD4, CD5, and CD6 are
arranged almost in a circle. CD1 and CD2 lie centrally in a dorsal position relative to other
CD glomeruli, yet ventral to the D group. CD3 and CD4 are more medial in position and
ventral to CD1 and CD2. CD5 and the more lateral CD6 are aligned towards the lateral
side of the antennal lobe. CD4 is a landmark, conspicuous because it is the most ventral
CD glomerulus and due to its large, irregular shape. This glomerulus is located dorsal to
the V group. Overall, the CD glomeruli can be distinguished from the other posterior
antennal lobe groups by their smaller volume. In some reconstructions, more than 6 CD
glomeruli were observed. These variant glomeruli were found in more posterior slices and
named CD7-CD12.

11. **Postero-Lateral group**: The PL group of glomeruli are located along the lateral axis and
form a distinct cluster on the dorso-lateral corner as confocal slices approach the posterior
surface of the antennal lobe. This is especially apparent in the male antennal lobe. The
PL group lies more posterior than the AL group in both sexes. PL1 is the most ventral

glomerulus in this group. Thereafter PL glomeruli are named with increasing numbers

based on dorsal position relative to PL1 and depth. PL9 is a landmark that is characterized

by its large size, visible through the most posterior sections of the antennal lobe. PL9 is

positioned laterally diagonal and opposite to the MD1 glomerulus. PL1-9 are spatially

invariant glomeruli. Four residual variant glomeruli are sometimes observed in more

posterior sections (PL10-PL13).

12. **Dorsal group**: The D group consist of 4 invariant glomeruli (D1, D2, D4 and D5) and are

the most dorsal group of glomeruli in posterior slices. D1, D2 and the typically observed

variant glomerulus D3 are arranged in a triangular manner in the female antennal lobe,

with D1 found at the apex and D2 and D3 found ventrally. D3 is sometimes observed

laterally to D2. In the male, D1, D2 and D3 form a row along the M-L axis. The second

group of glomeruli are constituted by D4 and D5. D5 is considered a landmark glomerulus

because of its large size and most dorsal position. D4 is found ventral and more medial

relative to D5. Two additional variant glomeruli (D6-D7) were identified in some

reconstructions in more posterior slices.

13. **Medio-Dorsal group**: The MD group typically comprises 3 glomeruli that flank the medial

margin of the antennal lobe. These glomeruli are positioned in the most posterior slices of

the antennal lobe from -28 µm to -35 µm, and are named MD1, MD2 and MD3. MD1 is

the most posterior and largest glomerulus in the antennal lobe, situated at the same depth

as the dorsal neuropil including the ellipsoid and the fan shaped bodies, on either side of

the esophagus. The landmark MD1 forms the core of this group. MD2 and MD3 are more

anterior but are positioned around MD1 like satellite glomeruli. MD3 is the most anterior

and dorsal of these glomeruli. A variant glomerulus MD4 is typically observed anterior to

MD1. The MD glomeruli are known to receive input from the maxillary palp neurons (12).

An easily identifiable array of posterior antennal lobe glomeruli with increasing depth is:

PL9-MD2-V8-MD1, with PL9, MD1 and V8 as landmarks. PL9 and MD1 have a distinct

shape and large surface area.

Representative confocal stacks and color-coded 2D maps detailing the typical spatial

arrangement of invariant and variant glomeruli in the female left and right antennal lobes as shown

by nc82 staining, are illustrated in Figure 2 and Figure S2, respectively. Examples for the left and

right antennal lobes of adult males are illustrated in Figure 3 and Figure S3, respectively. The

gross 3D shape of reconstructed nc-82 stained antennal lobes varied noticeably between
individual mosquitoes (see Figure S4). However, the eight characteristic landmark glomeruli could
reliably be identified in all reconstructed models we generated, facilitating assignment of names
to all remaining glomeruli based on their relative spatial position.

Volumetric analysis of the *Aedes aegypti* antennal lobe

We performed volumetric analyses on the left and right antennal lobes from all nc82-stained
LVPib12 brains that we reconstructed (n = 5 brains per sex, 20 antennal lobes total in this dataset).
The mean total antennal lobe volume is 123,319 μm^3 in females, and 85,370 μm^3 in males (*Mann-*
Whitney, P<0.0001). The total antennal lobe volume of females is therefore approximately 1.4
times larger than that of male mosquitoes.

For both sexes, there were no significant differences in total antennal lobe volume between the
left and right sides of the brain (Table S2). Within antennal lobes derived from females (Figure 4)
and males (Figure 5), the mean volumes of glomeruli varied significantly (one-way ANOVA,
P<0.0001 for each lobe – Figures S5-S8). The largest glomerulus in both the female and male
antennal lobe is MD1, with a mean volume of 5,222 μm^3 and 4,090 μm^3, which accounts for 4.26%
and 4.81% of their total antennal lobe volumes, respectively.

To identify candidate sexually dimorphic glomeruli, we then calculated relative volumes for all 63
invariant glomeruli that we annotated in females and males (Table S2). To do this, we determined
the percentage volume that each invariant glomerulus sampled occupies within its concordant
antennal lobe, and then compared grouped relative glomerular volumes by sex using the non-
parametric *Mann-Whitney U*-Test. Using this approach, we identified two glomeruli with
significantly larger relative volumes in females: AD1 (P = 0.005) and VC1 (P = 0.036). Of these,
the landmark glomerulus AD1 has a mean volume of 4,708 μm^3 in females and 2,456 μm^3 in
males which accounts for 3.82% and 2.89% of their total antennal lobe volumes, respectively. In
addition, four glomeruli were significantly larger in males: AM3 (P = 0.004), CD3 (P = 0.019), CD4
(P = 0.015) and MD3 (P = 0.016). AM3, which is positioned medially on the anterior surface of
the antennal lobe, has a mean volume of 1,315 μm^3 in females and 1,431 μm^3 in males which
accounts for 1.06% and 1.67% of their total antennal lobe volumes, respectively.

Validating the revised neuroanatomical model of the *Aedes aegypti* antennal lobe

To verify glomerular counts and the spatial organization of the antennal lobe model reconstructed
from nc82-staining, we utilized a complementary and independent neuronal labeling method to
visualize glomeruli. Since the antennal lobes across diverse insect species are known to contain
a rich aggregation of F-actin filaments in OSNs (22), the fluorophore-conjugated toxin phalloidin
was used as a cytoskeletal marker of OSN axonal processes to visualize antennal lobe
morphology. Using this method, we could observe strong staining of all the major neuropil in the
central *Ae. aegypti* brain (Figure 6a and 6b). We then performed reconstructions on the left and
right antennal lobe from 3 male and 3 female LVPib12 mosquitoes. We counted 79 ± 1 glomeruli
in females and 77 ± 3 glomeruli in males (mean ± s.d.) (Figure 6g and 6h).

Relative to pre-synaptic nc82-staining, the diffuse labeling pattern of phalloidin from overlapping
OSN axonal processes traversing the antennal lobe made delineation of glomeruli using this
method challenging at times, particularly in the anterior and central antennal lobe. As a result, the
confocal images had to be magnified 4-5 times in order to perform segmentation. Furthermore, at
lower magnification we observed in many reconstructions that PV and VC glomeruli appeared to
be stained as a single large cluster of neuropil, rather than as individual glomerular units that
could be clearly demarcated with antibody labeling. This cluster has a similar morphology to the
non-glomerular JOC that was described in an earlier *Ae. aegypti* antennal lobe model generated
using phalloidin staining (11).

We selected representative reconstructions of the male and female antennal lobes in which we
were able to easily demarcate all landmark glomeruli, as described for the nc82-stained model
(Figure 6c to 6f). We then created 2D anatomical maps of female left and right antennal lobes
(Figure S9 and S10) to comprehensively name all glomeruli in this dataset. Consistent with the
nc82 model, we found that the antennal lobe was divisible into 13 distinct spatial groups following
our reference key. Applying our thresholding criteria to categorize spatially invariant and variant
glomeruli within this data set, we identified the exact complement of spatially invariant glomeruli
(n=63) seen with nc82-stained brains. Remaining glomeruli found at a frequency of <80% in the
antennal lobe of both sexes were categorized as variant glomeruli, consistent with morphological
distortions being inherent to both nc82 and phalloidin-based neuronal staining methods. Of note,
the PL group of glomeruli was positioned more dorsally in reconstructions of the male left antennal
lobe (Figure 6f).

To further assist with user identification of glomeruli in the LVPib12 strain using nc82 and
phalloidin-based neuronal staining methods, we have generated a series of digital models that
allow users to interactively view the spatial arrangement of glomeruli in left antennal lobe of a
female brain sample stained with each method from different perspectives and depths
(Supplementary Movie S1-S4).

Conservation of antennal lobe organization across geographically divergent strains of
Aedes aegypti

Finally, we compared the antennal lobe neuroanatomy of the inbred LVPib12 strain (20) originally
of African origin, with a recently established strain of *Ae. aegypti* derived from Patillas, Puerto
Rico (Figure 7a) that is routinely supplemented with field-caught mosquitoes (23). We applied
nc82-staining to dissected and fixed Patillas brain samples and reconstructed the left and right
antennal lobes of three male and three female *Ae. aegypti* mosquitoes in total. Overall, the gross
shape of antennal lobes from the LVPib12 and Patillas strains differed subtly. In particular in the
Patillas strain, the female left and right antennal lobes were rounded, while the PV group from the
male left and right antennal lobes appeared to taper out of the central region. Using our LVPib12
antennal lobe models as a guide, we generated 3D models of the Patillas male and female
antennal lobe, in which we classified and color-coded glomeruli into the 13 glomerular spatial
groups (Figure 7b to 7e).

We then segmented 3D models from a representative Patillas male and female brain sample to
yield 2D maps to name individual glomeruli. Using our reference key (Table 1), we named all
antennal lobe glomeruli in these brain samples, and then applied our thresholding criteria to
identify spatially invariant and variant glomeruli within the data set. We identified all 63 spatially
invariant glomeruli found in the LVPib12 strain and denoted the presence of residual variant
glomeruli consistent with our previous models. In total, we counted 81 ± 4 glomeruli in the female
antennal lobes (Figure 7f) and 77 ± 2 in the male antennal lobes (mean ± s.d.) (Figure 7g). We
conclude that despite gross morphological differences in antennal lobe shape, the spatial
organization of antennal lobe glomeruli in these geographically diverse populations of *Ae. aegypti*
appears conserved.

436
437
438
439

Discussion

Aedes aegypti is an emerging model system for mosquito olfaction (3, 24-30). An accurate anatomical map of the antennal lobe is crucial towards studies aimed at understanding how odors are detected and represented by the olfactory system of this disease vector. In this study, we generated an updated atlas for the *Ae. aegypti* antennal lobe providing systematic nomenclature and a standardized framework for identifying glomeruli within this primary olfactory processing center of the mosquito brain.

We determined the *Ae. aegypti* antennal lobe is composed of approximately 80 morphologically discrete glomeruli in both sexes. We surmise that the advanced confocal imaging techniques employed in our study may have facilitated clearer delineation of glomerular boundaries, thus revealing additional glomeruli relative to previous studies that found 35 or 49-50 glomeruli in the *Ae. aegypti* antennal lobe, respectively (11, 18). In comparison with the *Drosophila melanogaster* antennal lobe which has 56 glomeruli (31), the *Aedes aegypti* antennal lobe has markedly increased subdivision making assignment of names to individual glomeruli not a trivial matter.

To facilitate stereotypical identification of glomeruli, we first subdivided the antennal lobe into 13 spatial groups with revised nomenclature, merging several extant groups from a previous antennal lobe model (11), and annotating others with new descriptors to better reflect their true spatial position and accommodate the expanded complement of glomeruli we found (Table S1). Subsequently, we identified eight landmark glomeruli based on their characteristic shape, volume and position across these groups, with all other glomeruli annotated by their spatial position relative to these landmarks. Guided by replicate morphological reconstructions, we annotated 63 spatially invariant glomeruli that can be reliably identified in the same geometric arrangement within dissected and stained antennal lobe preparations.

Despite intra- and inter-strain variability in the overall shape of antennal lobes in this study, the spatial organization of invariant glomeruli from LVPib12 and Patillas strains was shared and discernable. These results suggest that antennal lobe organization is likely conserved across geographically divergent strains of *Ae. aegypti*. Our revised atlas may therefore serve as a useful reference for studying antennal lobe organization in other *Ae. aegypti* strains supplemental to those reported here.

We determined the largest invariant glomerulus in *Ae. aegypti* was MD1. Anterograde tracing

experiments in this species previously revealed that MD1 likely receives projections from maxillary

palp-derived OSNs (12) which are responsive to the volatile gas carbon dioxide (CO_2). The

increased volume of this glomerulus in both sexes may be reflective of the critical role that CO_2

plays in *Ae. aegypti* host-seeking behavior (3, 29, 32). We additionally identified six candidate

glomeruli including AD1 and AM1 with sex-specific differences in relative volume. In the future,

identifying what odorants these glomeruli detect may help to further clarify their role in sexually

dimorphic olfactory behaviors in this mosquito species.

We noted considerable variation in the spatial position of approximately 20% of glomeruli

annotated within our *in vitro* antennal lobe reconstructions. These variant glomeruli tended to

have smaller volumes and were clustered around boundaries between spatial groups, making

them difficult to accurately name. Typically, variant glomeruli were positioned in more posterior

layers of the antennal lobe and were frequently noted in PM, PL, CD and V groups. Such spatial

variability in glomerular position is likely due to technical artifacts associated with dissection,

fixation, protocol duration and coverslip placement for imaging that commonly distort antennal

lobe morphology (33). The *Ae. aegypti* antennal lobes are also positioned in close proximity to

the anterior esophagus, a thick structure lined with cuticle, epithelial cells and muscle (34). The

esophagus runs through the esophageal foramen and it was removed during our dissections to

prevent it from occluding the antennal lobes, possibly causing distortion or disassociating

glomeruli, and altering the shape of each antennal lobe.

As recently described for a revised antennal lobe model for *Anopheles coluzzii* (19), we did not

observe any large clusters of neuropil orthologous to the Johnston's Organ Center (JOC) (11) in

any of our reconstructions from the LVPib12 or Patillas *Ae. aegypti* strains. Instead, we identified

morphologically discrete glomeruli in the anterior region of the antennal lobe where this structure

was previously annotated. The JOC was assigned based on evidence from silver nitrate Golgi

staining experiments (15) that suggested putative neuronal connectivity between the antennal

lobe and the sound-receptive Johnston's Organ (JO). However, other anterograde tracing studies

in *Ae. aegypti* and *D. melanogaster* indicate afferent JO neurons project to the antennal motor

and mechanosensory center (AMMC) (35, 36) of the deutocerebrum, rather than the antennal

lobe (37, 38). Further in-depth neuroanatomical studies are clearly warranted to definitively

validate any afferent or efferent neuronal connectivity between the JO and antennal lobe in *Ae.*

aegypti and other mosquito species.

The *Ae. aegypti* genome encodes large chemoreceptor gene families including Odorant Receptors, Ionotropic Receptors and Gustatory Receptors (27, 39). To date, chemoreceptors have only been localized to OSNs on the *Ae. aegypti* antennae, maxillary palps and proboscis using bulk transcriptomics and immunohistochemical approaches (24, 27, 40). We propose this model atlas will be useful to map projection patterns of neurons expressing chemoreceptors and other genes involved in olfactory signaling to their associated glomeruli in the *Ae. aegypti* antennal lobe using genetic labeling approaches (19, 33, 41, 42). Such studies will be critical to validate our antennal lobe model that is currently based on morphology alone, toward understanding the molecular and cellular organization of the *Ae. aegypti* olfactory system.

Finally, this updated atlas lays the foundation for improved spatial registration during functional imaging within the *Ae. aegypti* antennal lobe with genetically encoded calcium indicators and dyes (29, 43, 44). In particular, studies with activity-dependent ratiometric indicators such as CaMPARI (45, 46), that provide a permanent readout of neural activity and are also compatible with post-hoc neuronal staining to aid in spatial registration of such signals, will benefit greatly from using the *in vitro* antennal lobe atlas reported here as a reference. Such analyses may help to reliably identify glomeruli activated or inhibited by diverse odorants that are behaviorally relevant to *Ae. aegypti* such as human odors, plant odors, pheromones and insect repellents. This information may further accelerate reverse engineering of synthetic odorant blends mimicking patterns of neurophysiological activity elicited by these natural olfactory stimuli, to yield novel applied strategies to modulate *Ae. aegypti* olfactory behaviors.

Materials and Methods

Mosquito maintenance and rearing

Aedes aegypti LVPib12 (20) and Patillas (23) strains were maintained at 27°C, 80% relative humidity with a 12 hr light:dark photoperiod. Immature stages were reared using a standardized protocol optimized for yielding synchronous pupation. Briefly, on day 0 (hatching), 500mL of a 1% w/v autoclaved broth solution of finely milled TetraMin (Tetra, 16110) in dH_2O was placed in a clean polycarbonate pan (Cambro, 24CW135) with covering lid (Cambro, 20CWCH135). Oviposition papers containing desiccated eggs were then completely submerged in the broth solution inside each allocated pan to facilitate larval hatching. On day 1, 500 ml of additional dH_2O and 1g of finely milled Tetramin was added to the pan to bring the total volume to 1L. On day 2,

oviposition papers were removed, and hatched larvae thinned to a set density of 200 larvae in 2L
of fresh dH$_2$O per pan and fed one coarsely dissociated 1g Tetramin tablet. On day 3, larvae in
each pan were fed with an additional 1g Tetramin tablet. Thereafter, larvae in each pan were fed
two 1g Tetramin tablets per day until pupation which fully occurred on day 6. After emergence,
adult mosquitoes were provided constant access to a 10% w/v sucrose solution. For routine
colony maintenance, females were blood fed using anesthetized mice (Johns Hopkins Animal
Care and Use Committee protocol # MO18H391).

Neuronal staining
Neuropil in the central brain of mated, 5-10 day old, male and nulliparous female *Ae. aegypti* were
initially stained with the primary monoclonal antibody nc82 (DSHB, nc82-s, AB_2314866) that
labels the pre-synaptic active zone protein Bruchpilot. To dissect brains, mosquito heads were
severed from the thorax and tissue fixed for 3 hr at 4°C in Milonig's buffer: 0.1M PBS pH 7.2 with
4% v/v paraformaldehyde (PFA) as a fixative. Next, adult brains were dissected out into 0.1 M
PBS using fine forceps (Dumont No 5, 100 nm tips) to carefully remove the head capsule as well
as the pigmented ommatidia over the optic lobes and any floating air sacs connected to the brain.
Dissected brains were washed 3 times for 20 min each at room temperature in 0.25% PBST
consisting of 0.25% v/v Triton-X-100 detergent as a permeabilizing agent prepared in 0.1 M PBS.
The brains were then treated in a blocking solution consisting of 2% w/v normal goat serum (NGS)
and 4% v/v PFA in 0.1 M PBS and incubated on a nutator overnight at 4°C. This step was followed
by incubation in a primary antibody solution for 3 days at 4°C consisting of mouse anti-nc82 (1:50
v/v in 0.1 M PBS with 0.25% Triton-X and 2% NGS). Next, the brains were washed three times at
room temperature in 0.25% PBST and incubated for 3 days at 4°C in a secondary antibody
solution consisting of goat anti-mouse Cy3 (Jackson ImmunoResearch 115-165-062, 1:200 v/v in
0.1 M PBS with 0.25% Triton-X and 2% NGS). Following three final washes at room temperature
for 20 min each in 0.25% PBST, the brains were mounted within the space created by two
coverslip (Number 2 - 170 µm) bridges on glass slides in 20µl of Slow-Fade Gold Antifade
Mountant (Invitrogen, S36936), to preserve 3D structure.

As an alternative method to immunohistochemistry, adult brains were stained with fluorophore
conjugated-toxin phalloidin to mark cytoskeletal F-actin filaments present ubiquitously across
neuropilar structures including OSN axonal processes. Adult heads were first fixed in Milonig's
buffer for 3 hr at 4° C and brain tissue dissected in 0.1 M PBS as described above. After washing
3 times in 0.25% PBST, the dissected brains were incubated in 0.25% PBST with 1/80 (v/v) Alexa

Fluor 488 Phalloidin (Invitrogen, A12379) and 2μg/ml DAPI nuclear counterstain (Invitrogen,
D1306), for 3 days at 4° C. Brains were then mounted on glass slides as described and imaged.
578

Image acquisition settings

580 Images of nc82- and phalloidin stained brains were taken on a single-point laser scanning, Carl-
581 Zeiss LSM 780 confocal microscope. To capture images of the entire adult brain for both methods,
582 a 10X objective lens (0.3 NA, Plan-Apochromat) was used. To yield images to perform 3D
583 reconstructions of the antennal lobes, images were taken with a 20X objective lens (0.8 NA, Plan-
584 Apochromat). Images were processed using Imaris software (Oxford Instruments).
585

For nc82-stained brains, a solid-state laser line at 561 nm (1% laser power) was used to excite
the Cy3 signal and the gain of the GaAsP detector was set to 600. 100 z-slices, with a z-step size
of 1 μm and a 1024 X 1024 pixel size were acquired. For 3D reconstructions, 60 z-slices with a z
step size of 1 μm were acquired. The power of the 561 nm laser was adjusted to 2% with detector
gain at 620. These settings ensured the acquisition of high intensity images, allowing for the
accurate de-lineation of glomerular boundaries while scanning through the entire volumes of both
antennal lobes, including a few slices above the surface of the antennal lobe, and a few slices
below the deep-seated MD1 glomerulus, so as to closely replicate three-dimensionality of the AL
structure in the reconstructed model. Images were imported in the *.lsm-Zeiss, Zen format into
the Amira software program. We generated inverted images of the z-slices, in Image j, first using
the gray lookup table (LUT) to get a black and white image, followed by the invert LUT. No
brightness and contrast adjustments were made.
598

599 For phalloidin stained brains, a 488 nm laser line (2% laser power) was used to excite the Alexa-
600 488 phalloidin fluorophore, with the detector gain at 500. The DAPI stain was excited with a 405
601 nm diode laser (2% laser power and detector gain at 400). 60 z-slices with a z step size of 1 μm
602 and a 1024 X 1024 pixel size were acquired. Phalloidin and DAPI stained confocal slices were
603 inverted using Adobe Photoshop, without altering brightness or contrast.
604

Antennal lobe reconstructions

3D reconstructions of the antennal lobe were performed using Amira software (FEI Houston Inc).
Confocal files were opened in the '.lsm' format and viewed using the Image Segmentation Editor
tool. For each z-slice typically ranging from 0 μm to -60 μm through the anterior-posterior axis of
the antennal lobe, olfactory glomeruli were identified by shape and position. We defined the start

of the antennal lobe (denoted 0µm) as the first z-slice where the surface of the first landmark and

most anterior glomerulus AD1 is located. Next, an image segmentation step was carried out in

which the voxels occupied by individual glomeruli were first highlighted using the Amira paintbrush

tool and then assigned a unique label and color. After segmentation of the voxels corresponding

to all antennal lobe glomeruli, the image volume was surface rendered to create a 3D model to

allow visualization of antennal lobe morphology along x-y-z axes.

Volumes of individual glomeruli were obtained (in μm^3) using the Material Statistics tool in Amira.

We used the ortho slice tool to generate 2D cartoons of all the confocal slices and saved the

images in 'tif' format. Each antennal lobe glomerulus was named based on its spatial position

relative to other glomeruli. For each strain and staining method used, we classified glomeruli as

spatially 'invariant' or 'variant' based on their frequency of identification across combined male

and female, left and right antennal lobe datasets. A threshold frequency of ≥80% was designated

for the classification of spatially invariant glomeruli.

Statistics

Based on our 3D atlas of the antennal lobe of female and male LVPib12 *Aedes aegypti*, we

determined the volume of each annotated glomerulus and the total volume of the antennal lobe

(constituting the cumulative volume of all stained glomeruli in a given antennal lobe) from each

brain sample. To test whether mean volumes of invariant glomeruli differed significantly within

each lobe, we performed a *One-Way ANOVA* and *Tukey's HSD* post hoc test to correct for

multiple comparisons. To evaluate size differences of homologous invariant glomeruli between

the left and right antennal lobe and between sexes, the non-parametric *Mann-Whitney U*-test was

performed. Statistical analysis was performed using GraphPad Prism Software version 8.1.1

(GraphPad Software, Inc.). *P*< 0.05 were considered significant.

Acknowledgements

We thank Christopher Potter for initial access to Amira software; Roberto Barrera (CDC Dengue

Branch, San Juan, PR) for the Patillas stain; Darya Task and Benjamin Burgunder for expert

assistance with brain immunohistochemistry and advice on image reconstructions; and Olena

Riabinina and members of the McMeniman and Potter laboratories for helpful discussions and

comments on the manuscript. This research was supported by funding from the National Institutes

of Health NIAID (R21 AI139358-01), Centers for Disease Control and Prevention (200-2017-

93143) and USAID (AID-OAA-F-16-00061) to C.J.M. Microscopy infrastructure at Johns Hopkins

School of Medicine Microscope Core Facility used in this research was supported by the National

Institutes of Health NCRR (S10OD016374). We further acknowledge generous support to the

McMeniman laboratory from the Johns Hopkins Malaria Research Institute and Bloomberg

Philanthropies. We thank the Developmental Studies Hybridoma Bank at University of Iowa for

supplying the monoclonal nc82 antibody generously donated by Erich Buchner.

Data Availability

Raw imaging files and datasets analyzed in this study are available from the corresponding author

upon request.

References

1.   Kellogg FE. Water vapour and carbon dioxide receptors in Aedes aegypti. J Insect

Physiol. 1970;16(1):99-108.

2.   Sokolove EEDG. Lactic acid-sensitive receptors on the antennae of the mosquito,Aedes

aegypti. Journal of comparative physiology. 1976;Volume 105,(Issue 1,): pp 43–54.

3.   McMeniman CJ, Corfas RA, Matthews BJ, Ritchie SA, Vosshall LB. Multimodal

integration of carbon dioxide and other sensory cues drives mosquito attraction to humans. Cell.

2014;156(5):1060-71.

4.   Santana AL, Roque RA, Eiras AE. Characteristics of grass infusions as oviposition

attractants to Aedes (Stegomyia) (Diptera: Culicidae). J Med Entomol. 2006;43(2):214-20.

5.   Fawaz EY, Allan SA, Bernier UR, Obenauer PJ, Diclaro JW, 2nd. Swarming

mechanisms in the yellow fever mosquito: aggregation pheromones are involved in the mating

behavior of Aedes aegypti. J Vector Ecol. 2014;39(2):347-54.

6.   Peach DAH, Gries R, Zhai H, Young N, Gries G. Multimodal floral cues guide

mosquitoes to tansy inflorescences. Sci Rep. 2019;9(1):3908.

7.   McIver SB. Fine structure of antennal sensilla coeloconica of culicine mosquitoes.

Tissue Cell. 1973;5(1):105-12.

8.   McIver S. Structure of sensilla trichodea of female Aedes Aegypti with comments on

innervation of sensilla. J Insect Physiol. 1978;24:8.

9.   McIver SB. Fine structure of pegs on the palps of female culicine mosquitoes. Can J

Zool. 1972;50(5):571-6.

10.   McMeniman CJ. Disruption of Mosquito Olfaction. 2016 ed: Academic Press, Elsevier;

2014. p. 227-52.

11.   Ignell R, Dekker T, Ghaninia M, Hansson BS. Neuronal architecture of the mosquito

deutocerebrum. J Comp Neurol. 2005;493(2):207-40.

12.   Anton S, van Loon JJ, Meijerink J, Smid HM, Takken W, Rospars JP. Central projections

of olfactory receptor neurons from single antennal and palpal sensilla in mosquitoes. Arthropod

Struct Dev. 2003;32(4):319-27.

13.   Wilson RI. Early olfactory processing in Drosophila: mechanisms and principles. Annu

Rev Neurosci. 2013;36:217-41.

14.   Vosshall LB, Wong AM, Axel R. An olfactory sensory map in the fly brain. Cell.

2000;102(2):147-59.

15.   Chou Y-H, Spletter ML, Yaksi E, Leong JCS, Wilson RI, Luo L. Diversity and wiring

variability of olfactory local interneurons in the Drosophila antennal lobe. Nat Neurosci.

2010;13(4):439-49.

16.   Distler P, Boeckh J. Central projections of the maxillary and antennal nerves in the

mosquito Aedes aegypti. J Exp Biol. 1997;200(Pt 13):1873-9.

17.   Ghaninia M, Ignell R, Hansson BS. Functional classification and central nervous

projections of olfactory receptor neurons housed in antennal trichoid sensilla of female yellow

fever mosquitoes, Aedes aegypti. European Journal of Neuroscience. 2007;26(6):1611-23.

18.   Bausenwein B, Nick P. Three dimensional reconstruction of the antennal lobe in the

mosquito Aedes aegypti. In: Wehner R, Elsner N, editors. New neuroethology on the move.

Stuttgart: Thieme; 1998. p. 386.

19.   Riabinina O, Task D, Marr E, Lin CC, Alford R, O'Brochta DA, et al. Organization of

olfactory centres in the malaria mosquito Anopheles gambiae. Nat Commun. 2016;7:13010.

20.   Nene V, Wortman JR, Lawson D, Haas B, Kodira C, Tu ZJ, et al. Genome sequence of

Aedes aegypti, a major arbovirus vector. Science. 2007;316(5832):1718-23.

21.   Ito K, Shinomiya K, Ito M, Armstrong JD, Boyan G, Hartenstein V, et al. A systematic

nomenclature for the insect brain. Neuron. 2014;81(4):755-65.

22.   Rossler W, Kuduz J, Schurmann FW, Schild D. Aggregation of f-actin in olfactory

glomeruli: a common feature of glomeruli across phyla. Chem Senses. 2002;27(9):803-10.

23.   Poole-Smith BK, Hemme RR, Delorey M, Felix G, Gonzalez AL, Amador M, et al.

Comparison of vector competence of Aedes mediovittatus and Aedes aegypti for dengue virus:

implications for dengue control in the Caribbean. PLoS Negl Trop Dis. 2015;9(2):e0003462.

24.   DeGennaro M, McBride CS, Seeholzer L, Nakagawa T, Dennis EJ, Goldman C, et al.

orco mutant mosquitoes lose strong preference for humans and are not repelled by volatile

DEET. Nature. 2013;498(7455):487-91.

25.   McBride CS, Baier F, Omondi AB, Spitzer SA, Lutomiah J, Sang R, et al. Evolution of

mosquito preference for humans linked to an odorant receptor. Nature. 2014;515(7526):222-7.

26.   Vinauger C, Lahondere C, Wolff GH, Locke LT, Liaw JE, Parrish JZ, et al. Modulation of

Host Learning in Aedes aegypti Mosquitoes. Curr Biol. 2018;28(3):333-44 e8.

27.   Matthews BJ, McBride CS, DeGennaro M, Despo O, Vosshall LB. The

neurotranscriptome of the Aedes aegypti mosquito. BMC Genomics. 2016;17:32.

28.   Raji JI, Melo N, Castillo JS, Gonzalez S, Saldana V, Stensmyr MC, et al. Aedes aegypti

Mosquitoes Detect Acidic Volatiles Found in Human Odor Using the IR8a Pathway. Curr Biol.

2019;29(8):1253-62 e7.

29.   Vinauger C, Van Breugel F, Locke LT, Tobin KKS, Dickinson MH, Fairhall AL, et al.

Visual-Olfactory Integration in the Human Disease Vector Mosquito Aedes aegypti. Curr Biol.

2019;29(15):2509-16 e5.

30.   Liu MZ, Vosshall LB. General Visual and Contingent Thermal Cues Interact to Elicit

Attraction in Female Aedes aegypti Mosquitoes. Curr Biol. 2019;29(13):2250-7 e4.

31.   Tanaka NK, Endo K, Ito K. Organization of antennal lobe-associated neurons in adult

Drosophila melanogaster brain. J Comp Neurol. 2012;520(18):4067-130.

32.   van Breugel F, Riffell J, Fairhall A, Dickinson MH. Mosquitoes Use Vision to Associate

Odor Plumes with Thermal Targets. Curr Biol. 2015;25(16):2123-9.

33.   Grabe V, Strutz A, Baschwitz A, Hansson BS, Sachse S. Digital in vivo 3D atlas of the

antennal lobe of Drosophila melanogaster. J Comp Neurol. 2015;523(3):530-44.

34.   Charles C.Dapples AOL. Inner surface morphology of the alimentary canal in Aedes

aegypti (L.) (diptera: culicidae). International Journal of Insect Morphology and Embryology.

1974;Volume 3(Issues 3–4):433-4.

35.   Childress SA, McIver B. Morphology of the deutocerebrum of female Aedes aegypti

(Diptera: Culicidae). Can J Zool. 1984;62: 1320–8.

36.   Nyhof JM, McIver SB. Structural organization of the brain and subesophageal ganglion

of male Aedes aegypti (L.) (Diptera: Culicidae). Int J Insect Morphol Embryol. 1989;18:13–37.

37.   Steven A. Childress SBM. Morphology of the deutocerebrum of female *Aedes aegypti*

(Diptera: Culicidae). Canadian Journal of Zoology1984. p. 1320-8.

38.   Nyhof JM, Mciver SB. Structural Organization of the Brain and Subesophageal Ganglion

of Male Aedes-Aegypti (L) (Diptera, Culicidae). Int J Insect Morphol. 1989;18(1):13-37.

39.   Matthews BJ, Dudchenko O, Kingan SB, Koren S, Antoshechkin I, Crawford JE, et al.

Improved reference genome of Aedes aegypti informs arbovirus vector control. Nature.

2018;563(7732):501-7.

40.   Bohbot J, Pitts RJ, Kwon HW, Rutzler M, Robertson HM, Zwiebel LJ. Molecular

characterization of the Aedes aegypti odorant receptor gene family. Insect Mol Biol.

2007;16(5):525-37.

41.   Couto A, Alenius M, Dickson BJ. Molecular, anatomical, and functional organization of

the Drosophila olfactory system. Curr Biol. 2005;15(17):1535-47.

42.   Fishilevich E, Vosshall LB. Genetic and functional subdivision of the Drosophila antennal

lobe. Curr Biol. 2005;15(17):1548-53.

43.   Dana H, Sun Y, Mohar B, Hulse BK, Kerlin AM, Hasseman JP, et al. High-performance

calcium sensors for imaging activity in neuronal populations and microcompartments. Nat

Methods. 2019;16(7):649-57.

44.   Abdelfattah AS, Kawashima T, Singh A, Novak O, Liu H, Shuai Y, et al. Bright and

photostable chemigenetic indicators for extended in vivo voltage imaging. Science.

2019;365(6454):699-704.

45.   Moeyaert B, Holt G, Madangopal R, Perez-Alvarez A, Fearey BC, Trojanowski NF, et al.

Improved methods for marking active neuron populations. Nat Commun. 2018;9(1):4440.

46.   Fosque BF, Sun Y, Dana H, Yang CT, Ohyama T, Tadross MR, et al. Neural circuits.

Labeling of active neural circuits in vivo with designed calcium integrators. Science.

2015;347(6223):755-60.

Figure 1. Antennal lobe morphology of the LVPib12 *Aedes aegypti* strain as shown using nc82 staining. Confocal images of a representative adult (**a**) female and (**b**) male brain by nc82 antibody labeling (magenta); Scale bars, 100 μm. Right (RAL) and left antennal lobes (LAL) are indicated. Anterior and posterior views of three-dimensional reconstructive models of the right and left antennal lobe from a female (**c** & **e**) and male (**d** & **f**) mosquito are shown, respectively. Visible landmark glomeruli on these surfaces are denoted. Antennal lobe glomeruli are color-coded according to their classification into 13 groups based on their spatial position along anterior-posterior (a-p), dorsal-ventral (d-v), and medial-lateral (m-l) co-ordinate axes. The total number of glomeruli found in reconstructed (**g**) female and (**h**) male antennal lobes are shown (*n* = 5 brains per sex).

775   **Table 1**: Reference key describing the spatial arrangement of glomeruli in the *Aedes aegypti* antennal lobe.

Spatial group[1]	Landmark glomeruli	Glomerular Array[2]	Typical Array Depth[3] μm	Inter-Group Glomerular Array[4]
Antero-Dorsal (AD)[+]	AD1	AD1-AD2^V-AD3^L	AD1: 0-15 μm AD2, AD3: 3-14 μm	1. AD1-AM1^M-AC1^M 2. AD1-AL1^L-PL4^L 3. AD1-VC1^V
Ventro-Central (VC)[+]	VC6	VC1-VC2^L-VC3^{L‡} VC4^{V,M}-VC5^L-VC6^{L‡} VC7^M-VC8^L-VC9^{D§}	4-15 μm 5-15 μm 2-15 μm	VC6-PV4^D-PV3^V
Antero-Lateral (AL)	-	AL1-AL2^V-AL3^{V†}	6-15 μm	AD1-AL1^L-PL4^L
Antero-Medial (AM)	-	AM1-AM2^V-AM3^{V†}	2-14 μm	AD1-AM1^M-AC1^M
Antero-Central (AC)	-	AC1-AC2^{*V}-AC3^{V†}	6-15 μm	AD1-AM1^M-AC1^M
Postero-Central (PC)[+]	-	PC1-PC2^V-PC3^V-PC4^{V†}	12-15 μm	AM1-AC1^M-PM1^L-PC1^L
Postero-Medial (PM)	-	PM1-PM2^V-PM3^V-PM4^{V†}	20-25 μm	AM1-AC1^M-PM1^L-PC1^L
Postero-Ventral (PV)[+]	-	PV1-PV2^L-PV3^{L‡} PV4^D-PV5^{M‡}	8-20 μm 15-20 μm	PV1-PM6^M-V1^V
Ventral (V)[+]	V1, V8	V1-V2^L-V3^L-V4^{L‡} V5^D-V6^V-V7^M-V8^{M‡}	25-30 μm 25-35 μm	PV1-PM6^M-V1^V
Centro-Dorsal (CD)	CD4	CD1-CD2^V-CD3^V-CD4^V-CD5^D-CD6$^{D°}$	20-30 μm	PC4-CD4^L-V4^L
Postero-Lateral (PL)[+]	PL9	PL1-PL2^D-PL3^D-PL4^D-PL5^D-PL6^{D†} PL7^V-PL8^L-PL9^{L§}	14-18 μm 18-23 μm	1. AL1-PL4^D 2. PL9-MD1^M-MD2^V-V8^V
Dorsal (D)[+]	D5	D1-D2^V-D3^{L*} D4^M-D5^D	20-25 μm 25-30 μm	D5-D4^M-MD3^V
Medio-Dorsal (MD)	MD1	MD4-MD3^D-MD1^V-MD2^{V†}	28-35 μm	PL9- V8^{M,V}-MD2^D-MD1^D

[1] Spatial groups of antennal lobe glomeruli with new nomenclature and/or organization relative to Ignell et al. (2005) are indicated by

an addition symbol[+] as justified in Table S1.

[2] Description of the geometric arrangement of glomeruli within each spatial group. Glomeruli are typically arrayed in columns†, rows‡,

circles°, or randomly arrayed and numbered based on depth§. Each glomerulus may occupy a medial^M, lateral^L, dorsal^D or ventral^V

position relative to the preceding glomerulus in its array. Commonly observed variant glomeruli are indicated by an asterisk*.

[3] The depth of glomerular arrays in dissected brains is subject to variation based on the extent to which the antennal lobe is distorted

by removal of the esophagus, fixation and/or coverslip placement.

[4] Arrangement of guide arrays of glomeruli across spatial groups. Glomeruli within arrays are typically arranged as layers oriented

from the anterior to posterior sections of the antennal lobe. See also Figure S1.

Figure 2. Representative confocal stack from the left antennal lobe of an adult female LVPib12 *Aedes aegypti* as shown
by nc82 staining. Ten frontal planes from a total of 35 images taken at 1μm intervals were selected for illustration of
the typical geometric arrangement of glomeruli; Scale bar, 20 μm. The depth of each confocal slice is indicated.
Glomeruli within each reconstructed slice are color-coded according to their predicted spatial group. Glomeruli are
numbered, with 61 out of 63 spatially invariant glomeruli evident in the ten antennal lobe slices depicted here. Variant
glomeruli are indicated by asterisks.

794
795
Figure 3. Representative confocal stack from the left antennal lobe of an adult male LVPib12 *Aedes aegypti* as shown
by nc82 staining. Ten frontal planes from a total of 40 images taken at 1μm intervals were selected for illustration of
the typical geometric arrangement of glomeruli; Scale bar, 20 μm. The depth of each confocal slice is indicated.
Glomeruli within each reconstructed slice are color-coded according to their predicted spatial group. Glomeruli are
numbered, with 59 out of 63 spatially invariant glomeruli evident in the ten antennal lobe slices depicted here. Variant
glomeruli are indicated by asterisks.
802

Figure 4. Frequency and volume of glomeruli observed in three-dimensional reconstructions of the (**a**) left and (**b**) right
antennal lobe of adult female LVPib12 *Aedes aegypti* with nc82 staining. A heatmap of glomerular frequency (GF%)
and the associated classification of glomeruli assigned to each of the 13 glomerular spatial groups during replicate
reconstructions is shown. Invariant glomeruli are classified as those having a GF of $\geq$80% across all antennal lobe
reconstructions, while variant glomeruli below this threshold are indicated by asterisks (*n* = 5 brains).

Figure 5. Frequency and volume of glomeruli observed in three-dimensional reconstructions of the (**a**) left and (**b**) right antennal lobe of adult male LVPib12 *Aedes aegypti* with nc82 staining. A heatmap of glomerular frequency (GF%) and the associated classification of glomeruli assigned to each of the 13 glomerular spatial groups during replicate reconstructions is shown. Invariant glomeruli are classified as those having a GF of ≥80% across all antennal lobe reconstructions, while variant glomeruli below this threshold are indicated by asterisks (*n* = 5 brains).

815
816
Figure 6. Antennal lobe morphology of the LVPib12 *Aedes aegypti* strain as shown using phalloidin staining. Confocal
images of a representative adult (**a**) female and (**b**) male brain with F-actin filaments labeled with a phalloidin-
conjugated fluorophore (green) and nuclei labeled with DAPI dye counterstain (blue); Scale bars, 100 μm. Right (RAL)
and left antennal lobes (LAL) are indicated. Anterior and posterior views of three-dimensional reconstructive models of
the right and left antennal lobe from a female (**c** & **e**) and male (**d** & **f**) mosquito are shown, respectively. Visible
landmark glomeruli on these surfaces are denoted. Antennal lobe glomeruli are color-coded according to their
classification into 13 groups based on their spatial position along anterior-posterior (a-p), dorsal-ventral (d-v), and
medial-lateral (m-l) co-ordinate axes. The total number of glomeruli found in reconstructed (**g**) female and (**h**) male
antennal lobes are shown (*n* = 3 brains per sex).

Figure 7. Antennal lobe morphology of the Patillas *Aedes aegypti* strain as shown using nc82 staining. This mosquito strain was founded with field-collected *Ae. aegypti* from the municipality of Patillas, Puerto Rico (**a**). Anterior and posterior views of three-dimensional reconstructive models of the right and left antennal lobe from a female (**b** & **d**) and male (**c** & **e**) mosquito are shown, respectively. Visible landmark glomeruli on these surfaces are denoted. Antennal lobe glomeruli are color-coded according to their classification into 13 groups based on their spatial position along anterior-posterior (a-p), dorsal-ventral (d-v), and medial-lateral (m-l) co-ordinate axes. The total number of glomeruli found in reconstructed (**f**) female and (**g**) male antennal lobes are shown (*n* = 3 brains per sex).

835 **Table S1**: Changes to Nomenclature and/or Organization of Glomerular Spatial Group Classifiers in this updated
836 version of the *Aedes aegypti* antennal lobe atlas

Spatial Group – Updated Atlas[1]	Spatial Group – Ignell *et al*. (2005)	Justification for Changes
Antero-Dorsal (AD)[+]	= Ventral (V)	Renamed to more accurately reflect the anterior and dorsal position of these glomeruli
Ventro-Central (VC)[+]	= replaces the JOC	Ventro-Central (VC) and Postero-Ventral (PV) groups replace the previously annotated Johnston's Organ Center (JOC)
Antero-Lateral (AL)	= Same	-
Antero-Medial (AM)	= Same	-
Antero-Central (AC)	= Same	-
Postero-Central (PC)[+]	= Postero-Central (PC) + Medio-Central (MC)	Postero-Central (PC) and Medio-Central (MC) groups were consolidated due to their close spatial proximity in posterior and central antennal lobe sections
Postero-Medial (PM)	= Same	-
Postero-Ventral (PV)[+]	= replaces the JOC	Postero-Ventral (PV) and Ventro-Central (VC) groups replace the previously annotated Johnston's Organ Center (JOC)
Ventral (V)[+]	= Postero-Dorsal (PD)	Renamed to more accurately reflect the ventral position of these glomeruli in posterior antennal lobe sections
Centro-Dorsal (CD)	= Same	-
Postero-Lateral (PL)[+]	= Postero-Lateral (PL) + Latero-Central (LC)	Postero-Lateral (PL) and Latero-Central (LC) groups were consolidated due to their close spatial proximity in posterior and lateral antennal lobe sections
Dorsal (D)[+]	= Antero-Dorsal (AD)	Renamed to more accurately reflect the dorsal position of these glomeruli
Medio-Dorsal (MD)	= Same	-

839 [1] Spatial groups of glomeruli with new nomenclature and/or organization in our updated atlas relative to Ignell *et al*. (2005) are
840 indicated by an addition symbol[+].

841 **Table S2:** Quantitative comparison of glomerular volume in the female and male antennal lobe.

Glomerulus	Female		n	Male		n	U-test for right and left AL of female	U-test for right and left AL of male	U-test for female and male
	Volume (10³ μm³)	Relative volume (%)		Volume (10³ μm³)	Relative volume (%)		P-value	P-value	P-value
AC1	1.34 ± 0.46	1.07 ± 0.31	10	1.24 ± 0.52	1.48 ± 0.66	9	0.310	1.000	0.133
AC3	1.24 ± 0.47	0.99 ± 0.35	10	1.05 ± 1.04	1.20 ± 1.17	8	0.841	0.486	0.460
AL1	3.14 ± 1.03	2.56 ± 0.83	10	2.04 ± 1.15	2.38 ± 1.31	10	0.421	0.421	0.529
AL2	2.04 ± 0.73	1.66 ± 0.58	10	1.19 ± 0.72	1.38 ± 0.86	10	0.421	1.000	0.280
AL3	1.73 ± 0.74	1.37 ± 0.54	10	0.78 ± 0.54	0.88 ± 0.55	9	0.841	0.730	0.065
AM1	3.89 ± 1.31	3.10 ± 0.78	10	1.86 ± 1.00	2.19 ± 1.24	10	0.151	0.691	0.089
AM2	2.15 ± 1.07	1.73 ± 0.86	10	1.95 ± 0.75	2.29 ± 0.85	10	0.691	0.222	0.105
AM3	1.31 ± 0.61	1.06 ± 0.47	10	1.43 ± 0.36	1.67 ± 0.36	10	1.000	0.421	0.004*
AD1	4.71 ± 0.83	3.82 ± 0.56	10	2.46 ± 0.67	2.89 ± 0.76	10	0.421	0.056	0.005*
AD2	1.33 ± 0.68	1.09 ± 0.58	10	0.81 ± 0.64	0.93 ± 0.67	10	1.000	0.548	0.280
AD3	1.03 ± 0.58	0.84 ± 0.46	10	0.76 ± 0.55	0.90 ± 0.68	9	0.222	0.111	1.000
VC1	4.18 ± 1.72	3.33 ± 1.30	10	1.82 ± 0.53	2.14 ± 0.64	10	1.000	0.310	0.036*
VC2	2.44 ± 0.98	1.95 ± 0.72	10	1.51 ± 0.68	1.75 ± 0.79	10	0.691	0.151	0.029
VC3	3.39 ± 2.26	2.64 ± 1.57	10	1.31 ± 0.67	1.50 ± 0.64	10	0.548	0.691	0.123
VC4	2.53 ± 2.58	1.99 ± 2.05	10	1.27 ± 0.62	1.50 ± 0.76	10	0.841	0.841	0.912
VC5	2.38 ± 1.55	1.88 ± 1.03	10	1.60 ± 0.62	1.87 ± 0.72	10	0.008*	0.841	0.436
VC6	3.13 ± 2.15	2.52 ± 1.60	10	2.77 ± 1.15	3.23 ± 1.29	10	0.841	1.000	0.315
VC7	1.03 ± 0.85	0.84 ± 0.67	10	0.87 ± 0.54	1.02 ± 0.61	9	0.032*	0.016*	0.315
VC8	1.07 ± 0.69	0.87 ± 0.53	9	0.51 ± 0.26	0.60 ± 0.30	10	0.730	0.151	0.356
VC9	0.73 ± 0.48	0.61 ± 0.43	9	0.47 ± 0.27	0.55 ± 0.31	9	0.730	0.191	0.730
PC1	0.55 ± 0.41	0.46 ± 0.34	10	0.40 ± 0.32	0.47 ± 0.38	10	0.032	0.421	0.912
PC2	0.66 ± 0.37	0.55 ± 0.30	9	0.68 ± 0.60	0.78 ± 0.68	10	0.064	0.691	0.720
PC3	0.65 ± 0.36	0.53 ± 0.31	8	0.40 ± 0.20	0.46 ± 0.23	9	0.886	0.905	0.673
PC4	0.64 ± 0.41	0.53 ± 0.37	10	0.48 ± 0.26	0.55 ± 0.28	9	0.056	0.111	0.661
PV1	2.30 ± 0.71	1.86 ± 0.51	10	1.98 ± 0.59	2.35 ± 0.76	10	0.548	0.056	0.166
PV2	2.01 ± 1.21	1.61 ± 0.90	10	1.91 ± 1.44	2.20 ± 1.60	10	0.056	1.000	0.481
PV3	2.24 ± 1.43	1.80 ± 1.12	10	1.81 ± 1.25	2.07 ± 1.41	10	0.841	0.151	0.912
PV4	0.98 ± 0.51	0.79 ± 0.40	10	0.68 ± 0.38	0.79 ± 0.43	10	0.151	0.310	0.796
PV5	0.75 ± 0.62	0.59 ± 0.44	10	0.61 ± 0.25	0.71 ± 0.27	10	0.841	0.691	0.248
PM1	0.64 ± 0.44	0.53 ± 0.36	10	0.60 ± 0.27	0.69 ± 0.30	9	0.222	0.191	0.243
PM2	0.74 ± 0.45	0.61 ± 0.37	9	0.59 ± 0.27	0.68 ± 0.35	9	0.413	0.730	0.667
PM3	0.88 ± 0.53	0.73 ± 0.45	9	0.53 ± 0.44	0.63 ± 0.57	10	0.413	0.222	0.400
PM4	0.92 ± 0.46	0.77 ± 0.42	9	0.49 ± 0.30	0.58 ± 0.35	9	0.556	1.000	0.297
PL1	2.14 ± 1.32	1.77 ± 1.09	10	1.61 ± 1.04	1.85 ± 1.10	10	1.000	0.056	0.971
PL2	1.70 ± 1.07	1.40 ± 0.87	10	1.75 ± 1.29	2.05 ± 1.54	10	0.008*	0.095	0.481
PL3	2.95 ± 1.26	2.44 ± 1.07	10	1.70 ± 0.86	2.04 ± 1.12	10	1.000	0.548	0.393
PL4	2.10 ± 1.12	1.74 ± 0.92	10	1.22 ± 1.16	1.46 ± 1.41	10	0.421	0.421	0.248
PL5	1.21 ± 0.87	0.97 ± 0.62	10	1.06 ± 0.63	1.25 ± 0.76	10	0.421	0.222	0.481
PL6	1.05 ± 0.62	0.86 ± 0.55	10	1.29 ± 1.28	1.49 ± 1.39	10	0.151	1.000	0.166
PL7	0.96 ± 0.68	0.80 ± 0.61	10	0.81 ± 0.66	0.99 ± 0.80	8	0.310	0.686	0.829
PL8	1.86 ± 1.23	1.54 ± 1.08	10	0.85 ± 0.49	1.00 ± 0.57	9	0.548	0.905	0.243
PL9	2.40 ± 0.73	1.97 ± 0.68	9	2.23 ± 1.40	2.64 ± 1.59	9	1.000	0.413	0.340
D1	1.79 ± 1.74	1.42 ± 1.36	10	1.04 ± 0.39	1.24 ± 0.51	10	0.032*	0.222	0.529
D2	1.06 ± 0.78	0.86 ± 0.62	10	0.63 ± 0.42	0.76 ± 0.55	10	0.841	0.691	0.481
D4	1.21 ± 0.54	1.01 ± 0.47	9	1.39 ± 0.67	1.66 ± 0.86	10	0.905	0.421	0.113
D5	1.73 ± 1.33	1.47 ± 1.17	10	1.14 ± 0.64	1.31 ± 0.67	9	0.841	0.730	0.968
CD1	0.80 ± 0.41	0.64 ± 0.32	10	0.76 ± 0.53	0.91 ± 0.65	10	0.691	0.841	0.393
CD2	0.87 ± 0.51	0.72 ± 0.45	10	0.60 ± 0.43	0.73 ± 0.55	10	0.548	0.691	0.853
CD3	0.59 ± 0.32	0.50 ± 0.29	10	0.78 ± 0.30	0.93 ± 0.39	10	0.548	1.000	0.019*
CD4	1.19 ± 0.49	1.00 ± 0.49	10	1.52 ± 0.58	1.77 ± 0.64	10	0.151	1.000	0.015*
CD5	0.97 ± 0.73	0.80 ± 0.62	10	0.80 ± 0.50	0.92 ± 0.55	10	0.841	0.222	0.684
CD6	0.80 ± 0.62	0.66 ± 0.50	9	0.53 ± 0.19	0.63 ± 0.22	10	0.064	0.841	0.661
V1	2.29 ± 1.46	1.90 ± 1.28	10	1.68 ± 0.79	1.98 ± 0.99	10	0.421	0.841	0.579
V2	1.51 ± 0.90	1.20 ± 0.68	10	0.82 ± 0.38	0.96 ± 0.41	10	0.008*	0.095	0.579
V3	1.69 ± 0.95	1.33 ± 0.69	10	1.12 ± 1.02	1.25 ± 1.04	10	0.691	0.691	0.796
V4	1.38 ± 0.93	1.14 ± 0.80	10	0.62 ± 0.36	0.74 ± 0.42	10	0.421	0.691	0.315
V5	1.15 ± 0.52	0.94 ± 0.42	10	0.88 ± 0.57	1.02 ± 0.62	10	0.691	0.691	0.971
V6	1.04 ± 0.93	0.86 ± 0.80	10	1.23 ± 1.13	1.43 ± 1.30	10	0.691	0.222	0.143
V7	0.74 ± 0.41	0.60 ± 0.34	10	1.10 ± 0.96	1.26 ± 1.08	9	0.548	0.556	0.400
V8	0.94 ± 0.40	0.76 ± 0.31	9	0.96 ± 0.56	1.11 ± 0.60	10	1.000	0.691	0.182
MD1	5.22 ± 1.09	4.26 ± 0.94	10	4.09 ± 0.63	4.81 ± 0.81	10	0.691	0.032*	0.218
MD2	1.84 ± 1.03	1.52 ± 0.80	9	1.27 ± 0.64	1.48 ± 0.71	10	0.191	1.000	0.720
MD3	1.33 ± 0.44	1.10 ± 0.35	8	1.44 ± 0.61	1.70 ± 0.69	10	0.114	1.000	0.016*
Total AL Volume	123.32 ± 12.03		10	85.37 ± 7.65		10	0.548	0.548	<0.001*

842
843
844 Volume and relative volume are presented as Mean ± SD. Relative volume was used for statistical comparison to control for variation in the absolute volume in individuals, except for comparing total antennal lobe volumes. Asterisks indicate statistical significance at the level of P = 0.05.

Figure S2. Representative confocal stack from the right antennal lobe of an adult female LVPib12 *Aedes aegypti* as
shown by nc82 staining. Twelve frontal planes from a total of 35 images taken at 1µm intervals were selected for
illustration of the typical geometric arrangement of glomeruli; Scale bar, 20 µm. The depth of each confocal slice is
indicated. Glomeruli within each reconstructed slice are color-coded according to their predicted spatial group.
Glomeruli are numbered, with 61 out of 63 spatially invariant glomeruli evident in the twelve antennal lobe slices
depicted here. Variant glomeruli are indicated by asterisks.

Figure S3. Representative confocal stack from the right antennal lobe of an adult male LVPib12 *Aedes aegypti* as
shown by nc82 staining. Ten frontal planes from a total of 40 images taken at 1µm intervals were selected for illustration
of the typical geometric arrangement of glomeruli; Scale bar, 20 µm. The depth of each confocal slice is indicated.
Glomeruli within each reconstructed slice are color-coded according to their predicted spatial group. Glomeruli are
numbered, with 63 out of 63 spatially invariant glomeruli evident in the ten antennal lobe slices depicted here. Variant
glomeruli are indicated by asterisks.

Figure S4. Representative variation in antennal lobe shape in reconstructions from *in vitro* fixed, dissected and stained

brains. Three dimensional models of the right and left antennal lobes from 5 brain samples of LVPib12 females stained

with nc82 antibody are shown. Anterior and posterior perspectives of each antennal lobe reconstruction are illustrated.

Glomeruli are color-coded in the 3D models according to their spatial group. Landmark glomeruli are labelled on all

reconstructed antennal lobes and include AD1, VC6 located on the anterior surface and D5, V1, V8, MD1, CD4 and

PL9 on the posterior surface of the antennal lobe. Scale bars = 10 µm.

Figure S5. Probability heatmap visually representing the statistical significance of pairwise post-hoc Tukey's HSD tests

comparing the mean volumes of all 63 spatially invariant glomeruli in left antennal lobe of nc82-stained brains from

adult female LVPib12 *Aedes aegypti* (n = 5 brains). Multiplicity adjusted *P* values are plotted for each comparison. AD1,

VC1 and MD1 typically had larger volumes when compared with all other glomeruli. Glomerular means in this lobe

differed significantly as determined by one-way ANOVA: F (62, 247) = 5.412, *P*<0.0001. Abbreviations: NS (not

significant). N/A (comparisons between volumetric means of the same glomerulus are not applicable).

Figure S6. Probability heatmap visually representing the statistical significance of pairwise post-hoc Tukey's HSD tests

comparing the mean volumes of all 63 spatially invariant glomeruli in right antennal lobe of nc82-stained brains from

adult female LVPib12 *Aedes aegypti* (n = 5 brains). Multiplicity adjusted *P* values are plotted for each comparison. AM1,

AD1, VC1, VC3 and MD1 typically had larger volumes when compared with all other glomeruli. Glomerular means in

this lobe differed significantly as determined by one-way ANOVA: F (62, 242) = 6.521, *P*<0.0001. Abbreviations: NS

(not significant). N/A (comparisons between volumetric means of the same glomerulus are not applicable).

Figure S7. Probability heatmap visually representing the statistical significance of pairwise post-hoc Tukey's HSD tests

comparing the mean volumes of all 63 spatially invariant glomeruli in left antennal lobe of nc82-stained brains from

adult male LVPib12 *Aedes aegypti* (n = 5 brains). Multiplicity adjusted *P* values are plotted for each comparison. AD1,

VC6 and MD1 typically had larger volumes when compared with all other glomeruli. Glomerular means in this lobe

differed significantly as determined by one-way ANOVA: F (62, 241) = 4.472, *P*<0.0001. Abbreviations: NS (not

significant). N/A (comparisons between volumetric means of the same glomerulus are not applicable).

Figure S8. Probability heatmap visually representing the statistical significance of pairwise post-hoc Tukey's HSD tests

comparing the mean volumes of all 63 spatially invariant glomeruli in right antennal lobe of nc82-stained brains from

adult male LVPib12 *Aedes aegypti* (n = 5 brains). Multiplicity adjusted *P* values are plotted for each comparison. VC6

and MD1 typically had larger volumes when compared with all other glomeruli. Glomerular means in this lobe differed

significantly as determined by one-way ANOVA: F (62, 245) = 5.404, *P*<0.0001. Abbreviations: NS (not significant).

N/A (comparisons between volumetric means of the same glomerulus are not applicable).

911
912
Figure S9. Representative confocal stack from the left antennal lobe of an adult female LVPib12 *Aedes aegypti* as
shown by phalloidin staining. Fourteen frontal planes from a total of 45 images taken at 1µm intervals were selected
for illustration of the typical geometric arrangement of glomeruli; Scale bar, 20 µm. The depth of each confocal slice is
indicated. Glomeruli within each reconstructed slice are color-coded according to their predicted spatial group.
Glomeruli are numbered, with 63 out of the 63 spatially invariant glomeruli evident in the fourteen antennal lobe slices
depicted here. Variant glomeruli are indicated by asterisks.

919
920
921   **Figure S9** continued

Figure S10. Representative confocal stack from the right antennal lobe of an adult female LVPib12 *Aedes aegypti* as shown by phalloidin staining. Thirteen frontal planes from a total of 45 images taken at 1μm intervals were selected for illustration of the typical geometric arrangement of glomeruli; Scale bar, 20 μm. The depth of each confocal slice is indicated. Glomeruli within each reconstructed slice are color-coded according to their predicted spatial group. Glomeruli are numbered, with 63 out of the 63 spatially invariant glomeruli evident in the thirteen antennal lobe slices depicted here. Variant glomeruli are indicated by asterisks.

931 **Figure S10** continued